

# Age-Related Impairment in Insulin Release

## The Essential Role of $\beta_2$ -Adrenergic Receptor

Gaetano Santulli,<sup>1,2</sup> Angela Lombardi,<sup>3,4</sup> Daniela Sorriento,<sup>1</sup> Antonio Anastasio,<sup>1</sup> Carmine Del Giudice,<sup>1</sup> Pietro Formisano,<sup>4</sup> Francesco Béguinot,<sup>4</sup> Bruno Trimarco,<sup>1</sup> Claudia Miele,<sup>4</sup> and Guido Iaccarino<sup>5</sup>

In this study, we investigated the significance of  $\beta_2$ -adrenergic receptor ( $\beta_2$ AR) in age-related impaired insulin secretion and glucose homeostasis. We characterized the metabolic phenotype of  $\beta_2$ AR-null C57Bl/6N mice ( $\beta_2$ AR<sup>-/-</sup>) by performing in vivo and ex vivo experiments. In vitro assays in cultured INS-1E  $\beta$ -cells were carried out in order to clarify the mechanism by which  $\beta_2$ AR deficiency affects glucose metabolism. Adult  $\beta_2$ AR<sup>-/-</sup> mice featured glucose intolerance, and pancreatic islets isolated from these animals displayed impaired glucose-induced insulin release, accompanied by reduced expression of peroxisome proliferator-activated receptor (PPAR) $\gamma$ , pancreatic duodenal homeobox-1 (PDX-1), and GLUT2. Adenovirus-mediated gene transfer of human  $\beta_2$ AR rescued these defects. Consistent effects were evoked in vitro both upon  $\beta_2$ AR knockdown and pharmacologic treatment. Interestingly, with aging, wild-type ( $\beta_2$ AR<sup>+/+</sup>) littermates developed impaired insulin secretion and glucose tolerance. Moreover, islets from 20-month-old  $\beta_2$ AR<sup>+/+</sup> mice exhibited reduced density of  $\beta_2$ AR compared with those from younger animals, paralleled by decreased levels of PPAR $\gamma$ , PDX-1, and GLUT2. Overexpression of  $\beta_2$ AR in aged mice rescued glucose intolerance and insulin release both in vivo and ex vivo, restoring PPAR $\gamma$ /PDX-1/GLUT2 levels. Our data indicate that reduced  $\beta_2$ AR expression contributes to the age-related decline of glucose tolerance in mice. *Diabetes* 61:692–701, 2012

**I**mpairment of glucose metabolism with age represents a major determinant of type 2 diabetes epidemics within the elderly population. The molecular mechanisms underlying these changes have not been fully elucidated and are likely attributable to multiple causes (1,2). Aging per se is associated with a continuous decrease in basal insulin release (3). The size of this effect is sufficient to increase the likelihood of developing abnormalities in glucose tolerance and even overt diabetes (2,4). The consequence of aging on glucose tolerance occurs in different species, having been identified in rats

(5,6) as well as in humans (4,7,8). However, why insulin secretion deteriorates with aging remains a moot point.

The noradrenergic system provides fine-tuning to the endocrine pancreas activity through the function of  $\alpha$ - and  $\beta$ -adrenergic receptors (ARs) (9,10). The reciprocal regulation exerted by insulin and the adrenergic system has been well documented through a large number of studies (11–13). More recent evidence shows that mice with simultaneous deletion of the three known genes encoding the  $\beta$ ARs ( $\beta_1$ ,  $\beta_2$ , and  $\beta_3$ ) present a phenotype characterized by impaired glucose tolerance (14). Studies with  $\beta_2$ AR agonists further suggest that the  $\beta_2$ AR may play an important role in regulating insulin secretion (15). In addition, different human polymorphisms in the  $\beta_2$ AR gene have been associated with higher fasting insulin levels (16). Nevertheless, the impact of the  $\beta_2$ AR subtype on glucose tolerance and insulin secretion is still unclear.

Similar to glucose tolerance,  $\beta$ AR function and responsiveness deteriorate with aging (17–20), but the precise mechanisms involved are unknown. However, current evidence indicates that aging may downregulate  $\beta$ AR signaling,  $\beta_2$ AR in particular, by decreasing the expression of molecular components of the adrenergic signaling machinery (21–24). We have therefore hypothesized that age-dependent alterations in  $\beta$ AR function impair glucose-regulated insulin release by the pancreatic  $\beta$ -cells and may contribute to deterioration of glucose tolerance. To test this hypothesis, we explored the consequences of  $\beta_2$ AR knockout on insulin secretion in mice and investigated the significance of the age-related changes in  $\beta_2$ AR function with regard to glucose tolerance.

### RESEARCH DESIGN AND METHODS

**In vivo studies.** We studied male mice with a homozygous deletion of the  $\beta_2$ AR gene ( $\beta_2$ AR<sup>-/-</sup>) and backcrossed >12 generations onto C57Bl/6N background. Founders were provided by Brian Kobilka (Stanford University, Stanford, CA) (25). Wild-type littermates ( $\beta_2$ AR<sup>+/+</sup>) were used as controls. The animals were housed in a temperature-controlled (22°C) room with a 12-h light/dark cycle in accordance with the *Guide for the Care and Use of Laboratory Animals* published by the National Institutes of Health (NIH publication no. 85-23, revised 1996), and experiments were approved by the ethics committee of the Federico II University. Mice were killed by cervical dislocation. Pancreata were excised and collected rapidly after mice were killed. Samples were weighted, fixed by immersion in 4% paraformaldehyde for histology, homogenized for determination of total insulin content, or snap-frozen in liquid nitrogen and stored at -80°C for subsequent analyses. For determination of insulin or glucagon content, pancreatic tissue was homogenized in acid ethanol and extracted at 4°C overnight. The acidic extracts were dried by vacuum, reconstituted, and subjected to insulin and glucagon measurements.

**Glucose tolerance test and assessment of insulin secretion.** Glucose tolerance test (GTT) was performed as previously described (9,26). Briefly, mice were fasted overnight and then injected with glucose (2 g/kg i.p.). Blood glucose was measured by tail bleeding (Glucose Analyzer II; Beckman Coulter, Brea, CA) at indicated time points. The assessment of insulin secretion before

From the <sup>1</sup>Department of Clinical Medicine, Cardiovascular & Immunologic Sciences, “Federico II” University of Naples, Naples, Italy; <sup>2</sup>Columbia-Presbyterian Medical Center, College of Physicians & Surgeons, Columbia University, New York, New York; <sup>3</sup>Columbia University Medical Center, Columbia University, New York, New York; the <sup>4</sup>Department of Cellular and Molecular Biology and Pathology and Institute of Experimental Endocrinology and Oncology “Gaetano Salvatore,” “Federico II” University of Naples, Naples, Italy; and the <sup>5</sup>School of Medicine, University of Salerno, Salerno, Italy.

Corresponding authors: Guido Iaccarino, giaccarino@unisa.it, and Claudia Miele, c.miele@ieos.cnr.it.

Received 21 July 2011 and accepted 3 December 2011.

DOI: 10.2337/db11-1027

This article contains Supplementary Data online at <http://diabetes.diabetesjournals.org/lookup/suppl/doi:10.2337/db11-1027/-/DC1>.

G.S. and A.L. contributed equally to this work.

© 2012 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

and during glucose challenge was performed as previously described (9,27). Blood from the mandibular vein of overnight-fasted mice was collected at the indicated time for serum insulin assessment. The evaluation of glucagon secretion was performed by collecting blood from the mandibular vein of randomized mice before and after injection of insulin (0.75 IU/kg i.p.). Serum insulin and plasma glucagon were assayed by radioimmunoassay (Millipore, Billerica, MA). **Histological analysis.** Immunohistochemistry was carried out on paraffin sections using the H38 rabbit insulin antibody or the N-17 goat glucagon antibody (1:200 dilution; both from Santa Cruz Biotechnology, Santa Cruz, CA). Goat anti-rabbit serum coupled with peroxidase (Immunotech, Marseille, France) was used as secondary antibody. After primary antibody exposure, slides were incubated with biotinylated anti-rabbit or anti-goat IgG and peroxidase-labeled streptavidin (Dako Corporation). All reactions were revealed with diaminobenzidine. Sections were counterstained with hematoxylin and mounted (28,29).

#### Ex vivo studies

**Isolation of mouse pancreatic islets.** Islets of Langerhans were isolated by collagenase digestion (26,30). In brief, mice were killed as described above, the fur was soaked with ethanol, and the abdomen was opened to locate and excise the pancreas. Digestion was completed with collagenase P (Roche Applied Sciences, Penzberg, Germany) in a shaking water bath (37°C) for 5–8 min. The digested pancreas was treated with DNase I (New England Biolabs, Ipswich, MA). The islets were hand-picked under a stereomicroscope using a syringe with a 25-gauge needle and cultured at 37°C with 95% air and 5% CO<sub>2</sub> in complete RPMI-1640 supplemented with 5% heat-inactivated FBS, 1 mmol/L sodium pyruvate, 50 μmol/L 2-mercaptoethanol, 2 mmol/L glutamine, 10 mmol/L HEPES, 100 units/mL penicillin, and 100 μg/mL streptomycin (all from Sigma-Aldrich, Saint Louis, MO) as previously described (31,32).

**Evaluation of insulin secretion.** Isolated islets were preincubated at 37°C for 30 min in Krebs-Ringer bicarbonate buffer (120 mmol/L NaCl, 4.7 mmol/L KCl, 1.2 mmol/L MgSO<sub>4</sub>, 1.2 mmol/L KH<sub>2</sub>PO<sub>4</sub>, 2.4 mmol/L CaCl<sub>2</sub>, and 20 mmol/L NaHCO<sub>3</sub>) supplemented with 10 mmol/L HEPES and 0.2% BSA and gassed with a mixture of 95% O<sub>2</sub> and 5% CO<sub>2</sub> containing 2 mmol/L glucose. Twenty size-matched islets collected in each tube were incubated for 1 h in a 37°C water bath with 500 μL Krebs-Ringer bicarbonate buffer medium containing 2.8 mmol/L glucose, 16.7 mmol/L glucose, or 2.8 mmol/L glucose plus 33 mmol/L KCl. Islets were then pelleted by centrifugation (9,000g, 2 min, 4°C), and supernatants were collected for insulin secretion. Insulin concentrations were determined using radioimmunoassay (31).

#### Adenovirus-mediated reinstallation of β<sub>2</sub>AR in vivo and ex vivo.

Twenty-month-old β<sub>2</sub>AR<sup>+/+</sup> mice were anesthetized by isoflurane (4%) inhalation and maintained by mask ventilation (isoflurane 1.8%) (28). After laparotomy, we identified and mobilized the distal pancreas and we performed two injections (50 μL each) of adenovirus (7 × 10<sup>10</sup> plaque-forming units [pfu]/mouse) using a 30-gauge needle. Given the high homology between the human and mouse β<sub>2</sub>AR (33) and the validated use of the adenoviruses expressing the human β<sub>2</sub>AR (Adβ<sub>2</sub>AR) for in vivo gene transfer (34–36), we used the human Adβ<sub>2</sub>AR or, as control, an empty expression cassette derived from pcDNA3.2/V5/GW/D-TOPO (AdEmpty), kindly provided by Walter J. Koch, Jefferson University (Philadelphia, PA). Finally, abdominal incision was quickly closed in layers using 3–0 silk suture, and animals were observed and monitored until recovery. Isolated islets were infected (12 × 10<sup>4</sup> pfu/islet) with Adβ<sub>2</sub>AR or AdEmpty. After 3 h incubation with the adenoviruses, pancreatic islets were incubated with fresh medium and under normoxic conditions (95% air, 5% CO<sub>2</sub>) at 37°C for 24 h.

**In vitro studies.** The INS-1E β-cells (provided by P. Maechler, University of Geneva, Geneva, Switzerland), derived and selected from the parental rat insulinoma INS-1 β-cell line, were maintained in monolayer culture in RPMI-1640 medium as described above for the ex vivo experiments. For the insulin secretion assays, cells were seeded at a density of 5 × 10<sup>5</sup> cells/cm<sup>2</sup> for at least 96 h before use (31). Transient transfection of the PPARγ cDNA and control empty plasmid was performed using Lipofectamine 2000 (Invitrogen, Carlsbad, CA) (37). We also tested the effects of the specific pharmacological β<sub>2</sub>AR antagonist ICI 118,551 (0.1 μmol/L) or agonist (1 μmol/L fenoterol) (both from Sigma-Aldrich) (36).

#### Short hairpin RNA design, generation, and transfection and radioligand-binding assay.

Stealth short hairpin (sh)RNA oligoribonucleotides against β<sub>2</sub>AR and scramble were designed (sequences and efficiency shown in Supplementary Table 1 and Supplementary Fig. 2, respectively) and then synthesized by Invitrogen. The shRNA hairpin configuration is cleaved by the cellular machinery into small interfering RNA, which is then bound to the RNA-induced silencing complex. This complex binds to and cleaves mRNAs that match the small interfering RNA that is bound to it. We transfected INS-1E β-cells with 100 nmol/L sh-β<sub>2</sub>AR RNA or sh-scramble RNA using Lipofectamine 2000 (Invitrogen) according to the manufacturer's instructions. Assays were performed at least 3 days after shRNA transfection. Membrane fractions from pancreatic tissue, isolated islets, and INS-1E β-cells were used for

radioligand-binding studies to assess the density of ARs as previously described and validated (35,36,38,39).

**Real-time RT-PCR analysis.** Total cellular RNA was isolated from INS-1E β-cells, isolated islets, and pancreatic tissue samples with the RNeasy kit (Qiagen, Germantown, MD), according to the manufacturer's instructions, as previously described (31,40). PCRs were analyzed using SYBR Green mix (Invitrogen). Reactions were performed in triplicate using Platinum SYBR Green qPCR Super-UDG by means of an iCycler IQ multicolor Real Time PCR Detection system (Bio-Rad, Hercules, CA). Cyclophilin was used as an internal standard. Primer sequences are reported in Supplementary Table 2.

**Immunoblotting.** Immunoblot analysis was performed as previously described (31,37). Blots were probed with mouse monoclonal antibodies against adenylate cyclase type VI (AC-VI) (Abcam, Cambridge, MA), pancreatic and duodenal homeobox (PDX)-1, GLUT2, peroxisome proliferator-activated receptor (PPAR)γ, G-protein-coupled receptor (GRK)2, G protein α<sub>s</sub> (Gα<sub>s</sub>), clathrin heavy chain, and actin (Santa Cruz Biotechnology). Experiments were performed in triplicate to ensure reproducibility. Membrane extracts were obtained as previously described (29,38). Data are presented as arbitrary units using actin as internal control (clathrin heavy chain for membrane extracts) as indicated.

**Measurement of cAMP production in vitro and ex vivo.** Intracellular content of cAMP was determined using a cAMP<sup>125I</sup>-scintillation proximity assay (GE Healthcare, Piscataway, NJ) according to the manufacturer's instructions. Briefly, we used 20 size-matched islets (for the ex vivo assays) and 4,000 cells/well INS-1E (for the in vitro assays). Islets and β-cells were washed once and preincubated at 37°C in HEPES-buffered Krebs-Ringer solution containing 1 mmol/L glucose and 0.5 mmol/L isobutylmethylxanthine (a phosphodiesterase inhibitor) for 1 h and incubated for another 15 min in the same buffer with or without 100 μmol/L forskolin (MP Biomedicals, Solon, OH), 3 mmol/L NaF (Thermo Fisher Scientific, Pittsburgh, PA), or 1 μmol/L isoproterenol (Tocris Bioscience, Ellisville, MO). The reaction was stopped by addition of 50 mmol/L HCl and neutralized with NaOH. The cAMP levels were normalized to the protein concentration.

**Statistical analysis.** All data are presented as means ± SE. Statistical differences were determined by one-way or two-way ANOVA as appropriate, and Bonferroni post hoc testing was performed when applicable. A *P* value <0.05 was considered significant. Statistical analysis was performed using GraphPad Prism (version 5.01; GraphPad Software Inc., San Diego, CA).


## RESULTS

**Metabolic phenotype of β<sub>2</sub>AR<sup>-/-</sup> mouse.** To investigate in vivo the relevance of the β<sub>2</sub>AR gene in the regulation of insulin secretion, we compared the metabolic phenotype of adult (6 months old) β<sub>2</sub>AR<sup>-/-</sup> and β<sub>2</sub>AR<sup>+/+</sup> mice. Blood glucose was significantly higher in the null mice compared with that in their wild-type littermates both upon fasting and under random feeding conditions (Table 1). In addition, their fasting serum insulin levels were significantly reduced (Table 1). Upon glucose loading (GTT), the β<sub>2</sub>AR<sup>-/-</sup> mice displayed a marked reduction in glucose tolerance (Fig. 1A and B). In β<sub>2</sub>AR<sup>+/+</sup> mice, we observed a threefold increase in insulin secretion 3 min after intraperitoneal glucose injection, presumably corresponding with the peak of first-phase insulin release. This was followed by a decrease at 10

TABLE 1  
Metabolic characteristics of adult (6 months old) wild-type and knockout mice

	β <sub>2</sub> AR <sup>+/+</sup>	β <sub>2</sub> AR <sup>-/-</sup>
<i>n</i>	15	14
Body weight (g)	29.6 ± 1.1	28.2 ± 0.7
Food intake (g/day)	3.1 ± 0.6	3.1 ± 0.8
Water intake (mL/day)	5.8 ± 0.6	6.3 ± 0.9*
Random-fed blood glucose (mg/dL)	170.1 ± 12.3	198.4 ± 11.1*
Fasting blood glucose (mg/dL)	75.7 ± 8.2	135.3 ± 11.5*
Fasting serum insulin (ng/mL)	0.41 ± 0.03	0.30 ± 0.07*

Data are means ± SE unless otherwise indicated. \**P* < 0.05 vs. β<sub>2</sub>AR<sup>+/+</sup>.



**FIG. 1.** Metabolic profile of  $\beta_2AR^{-/-}$  mice. Six-month-old  $\beta_2AR^{-/-}$  mice and their wild-type littermates ( $\beta_2AR^{+/+}$ ) were fasted for 16 h and subjected to intraperitoneal glucose loading (2 g/kg body weight). Blood glucose (A and B) and serum insulin (C and D) were monitored for 120 min after glucose administration ( $n = 14-18$  animals per group).  $\beta_2AR^{-/-}$  mice displayed glucose intolerance (A) and impaired insulin secretion (C). We calculated the AUC from glucose (B) and insulin excursion (D) curves. Peak insulin-to-peak glucose ratio (E) represents  $\beta$ -cell function, as better described in RESEARCH DESIGN AND METHODS. Bars represent means  $\pm$  SE. \* $P < 0.05$  vs.  $\beta_2AR^{+/+}$ , Bonferroni post hoc test. AUC, area under the curve.


min and then a gradual increase over 30 min that may indicate a second-phase response (27,41).

In  $\beta_2AR^{-/-}$  mice, the early phase of insulin secretory response to glucose was reduced by more than twofold. The late response was also significantly impaired in the  $\beta_2AR^{-/-}$  compared with  $\beta_2AR^{+/+}$  mice (Fig. 1C and D). The peak insulin-to-peak glucose ratio was also decreased (Fig. 1E), further indicating impaired insulin response to hyperglycemia in the null mice.


To investigate whether the alterations in glucose tolerance identified in the  $\beta_2AR^{-/-}$  mice were contributed by deranged glucagon release, we further measured blood glucose and plasma glucagon levels 30 min after insulin administration. Indeed, insulin administration determines a fall in blood glucose and a counterregulatory rise in plasma glucagon (2,30). However,  $\beta_2AR^{-/-}$  and  $\beta_2AR^{+/+}$  mice exhibited comparable glucose and glucagon responses to insulin administration (Supplementary Fig. 1A and B). Pancreatic islet histology also did not show any significant difference in these mice (Fig. 2A), similar to total insulin and glucagon pancreatic content (Fig. 2B and C).

We then posed the further question of whether the reduced glucose insulin secretion observed in the  $\beta_2AR^{-/-}$  mice in vivo may represent the direct consequence of the  $\beta_2AR^{-/-}$  lack in the  $\beta$ -cells or whether it is indirectly mediated by other regulatory factors. To answer this question, we analyzed glucose effect on islets isolated from the null mice. As shown in Fig. 2D, these islets responded poorly to increased glucose concentration in the culture medium compared with the islets from their wild-type littermates but were fully responsive to KCl depolarization.

**Islets and  $\beta$ -cell profiling after  $\beta_2AR$  deletion.** To gain further insight into the mechanism leading to impaired insulin secretion in mice lacking  $\beta_2AR$ , we profiled the expression of different genes relevant to  $\beta$ -cell regulation by real-time RT-PCR of islet mRNA. As shown in Fig. 3A and B, mRNA levels of both *PDX-1* and *GLUT2*, two major genes involved in  $\beta$ -cell function, were decreased in islets from  $\beta_2AR^{-/-}$  mice by 75 and 60%, respectively. Also, mRNA levels of the *PDX-1/GLUT2* upstream regulator *PPAR $\gamma$*  were decreased by 54% compared with islets from wild-type mice (Fig. 3C). Reliable results were obtained in



**FIG. 2.** Comparison of  $\beta_2AR^{+/+}$  and  $\beta_2AR^{-/-}$  pancreatic islets. Immunohistochemical analysis (A) of the islets was carried out on paraffin sections using insulin (left panel) or glucagon (right panel) antibodies. Microphotographs are representative of images obtained from pancreas sections of five 6-month-old  $\beta_2AR^{+/+}$  (upper panel) or  $\beta_2AR^{-/-}$  (lower panel) mice. Insulin (B) and glucagon (C) content in isolated islets from  $\beta_2AR^{+/+}$  ( $n = 10$ ) or  $\beta_2AR^{-/-}$  ( $n = 13$ ) mice. Insulin secretion in response to basal (2.8 mmol/L) or high (16.7 mmol/L) glucose concentration and to KCl (33 mmol/L) was measured in isolated islets from  $\beta_2AR^{+/+}$  (□) and  $\beta_2AR^{-/-}$  (■) mice (D). Bars represent means  $\pm$  SE of data from 10 mice per group. \* $P < 0.05$  vs.  $\beta_2AR^{+/+}$ , Bonferroni post hoc test. (See also Supplementary Fig. 1.) (A high-quality digital representation of this figure is available in the online issue.)



**FIG. 3.** Gene expression profile in isolated Langerhans islets from  $\beta_2AR^{+/+}$  and  $\beta_2AR^{-/-}$  mice. The abundance of mRNAs for PDX-1 (A), GLUT2 (B), and PPAR $\gamma$  (C) was determined by real-time RT-PCR analysis of total RNA, using cyclophilin as internal standard. The mRNA levels in  $\beta_2AR^{-/-}$  mice are relative to those in control animals. Each bar represents means  $\pm$  SE of four independent experiments in each of which reactions were performed in triplicate using the pooled total RNAs from five mice/genotype. Proteins from a Western blot representative of three independent experiments were quantified by densitometry (D and E). \* $P < 0.05$  vs.  $\beta_2AR^{+/+}$ , Bonferroni post hoc test. AU, arbitrary units.

immunoblotting experiments (Fig. 3D and E). PDX-1 and GLUT2 mRNAs were also reduced to a similar extent in total pancreatic tissue from the  $\beta_2AR^{-/-}$  mice (data not shown).


We then sought to demonstrate whether these abnormalities in gene expression were directly caused by  $\beta_2AR$  silencing. To pursue this objective, we silenced with a specific shRNA (Supplementary Fig. 2) the  $\beta_2AR$  gene in the glucose-responsive INS-1E  $\beta$ -cell line (INS-1E $_{sh\beta_2AR}$ ) (Fig. 4A and B). As shown in Fig. 4C, this specific knock-down impaired glucose-induced insulin secretion by 58% in these cells. A similarly sized effect was achieved by treatment with the specific  $\beta_2AR$  antagonist ICI, while the  $\beta_2AR$  agonist fenoterol showed an opposite action (Supplementary Fig. 3A). Consistent with our ex vivo results, the INS-1E $_{sh\beta_2AR}$  displayed a reduction in PDX-1, GLUT2, and PPAR $\gamma$  mRNA (Fig. 4D–F) and protein levels (Fig. 4G and H). Interestingly, transient transfection of a PPAR $\gamma$  cDNA in INS-1E $_{sh\beta_2AR}$   $\beta$ -cells increased glucose-induced insulin secretion compared with the wild-type INS1-E control  $\beta$ -cells (Fig. 4C). In addition, overexpression of PPAR $\gamma$  prevented the downregulation of both PDX-1 and GLUT2 occurring in the INS-1E $_{sh\beta_2AR}$   $\beta$ -cells (Fig. 4D–H). Consistently, treatment of INS-1E  $\beta$ -cells with ICI decreased PDX-1 and GLUT2 mRNA and protein levels, while PPAR $\gamma$  overexpression completely prevented the effect of ICI (Supplementary Fig. 3B–D), suggesting that  $\beta_2AR$  controls insulin secretion through a PPAR $\gamma$ /PDX-1-mediated mechanism.

To better define the  $\beta_2AR$  downstream mechanism leading to PPAR $\gamma$  activation, we assessed the cAMP levels in these cells, observing an impaired production of cAMP in INS-1E $_{sh\beta_2AR}$   $\beta$ -cells both in basal condition and after stimulation with the  $\beta AR$  agonist isoproterenol (Supplementary

Fig. 4A). Accordingly, to rule out possible involvement of other components of  $\beta_2AR$  signaling machinery, we assessed the protein level of AC-VI, GRK2, and  $G\alpha_s$ , and we found no significant difference (Supplementary Fig. 4B and C). Parallel results were obtained in ex vivo experiments, performed to investigate the possible age-related alterations in the  $\beta_2AR$  transduction pathway, comparing pancreatic islets isolated from adult (6 months old) and old (20 months old)  $\beta_2AR^{+/+}$  mice (Supplementary Fig. 4D and E).

**$\beta_2AR$  overexpression rescued the age-related impairment in insulin release.** Based on radioligand binding and real-time RT-PCR analysis, the expression of both  $\beta_2AR$  protein and mRNA was significantly decreased in islets from aged (20 months old)  $\beta_2AR^{+/+}$  mice compared with those isolated from adult (6 months old) mice (Fig. 5A and B). PDX-1, GLUT2, and PPAR $\gamma$  expression (both in terms of mRNA and protein level) was also reduced, and insulin release in response to glucose, though not that evoked by KCl depolarization, was impaired in islets from the aged mice (Fig. 5C–H), suggesting that the reduced  $\beta_2AR$  density constrains islet glucose response in these animals. To prove this hypothesis, we used an adenoviral construct driving overexpression of human  $\beta_2AR$  in mouse islets. Interestingly, infection of islets isolated from wild-type old mice with this construct induced a twofold increase in  $\beta_2AR$  expression (Fig. 5A and B) and returned glucose-induced insulin secretion to levels comparable with those of islets from 6-month-old mice (Fig. 5C) accompanied by restored expression of PDX-1, GLUT2, and PPAR $\gamma$  (Fig. 5D–H).

In the in vivo setup, 20-month-old  $\beta_2AR^{+/+}$  mice exhibited a significant reduction in fasting serum insulin levels (Table 2) accompanied by impaired glucose tolerance and



**FIG. 4.**  $\beta_2$ AR levels, glucose-stimulated insulin secretion, and gene expression profile in silenced INS-1E  $\beta$ -cells. Treatment with a specific  $\beta_2$ AR-shRNA significantly decreased the density (by 73.7% [A]) and mRNA levels (by 59.1% [B]) of  $\beta_2$ AR in INS-1E  $\beta$ -cells.  $\beta_2$ AR-shRNA inhibited the insulin secretory response to 16.7 mmol/L glucose, which was rescued by the overexpression of PPAR $\gamma$  (C). KCl-induced insulin release (C) was not significantly different among the studied groups.  $\beta_2$ AR-shRNA also determined a significant reduction in mRNA level of PDX-1 (D), GLUT2 (E), and PPAR $\gamma$  (F) that was prevented by the overexpression of PPAR $\gamma$ . Bars represent means  $\pm$  SE from four to five independent experiments in each of which reactions were performed in triplicate (□, control, i.e. untreated INS-1E  $\beta$ -cells; ▨, sh-scramble; ▩, sh-scramble+empty vector; ▤, sh- $\beta_2$ AR; ▥, sh- $\beta_2$ AR+PPAR $\gamma$ ; \* $P$  < 0.05 vs. control, Bonferroni post hoc test; basal is glucose 2.8 mmol/L. Equal amount of proteins from three independent experiments was analyzed by Western blotting and quantified by densitometry (G and H). \* $P$  < 0.05 vs. sh-scramble. AU, arbitrary units. (See also Supplementary Figs. 2–4.)


insulin response upon GTT (Fig. 6A–E). We have therefore designed a gene therapy protocol aimed to prove that these abnormalities can be corrected by restoring  $\beta_2$ AR density. Accordingly, we infected the pancreas of aged mice by Ad $\beta_2$ AR injection. This injection effectively rescued  $\beta_2$ AR expression in the pancreatic tissue, returning it to levels comparable with those of 6-month-old mice (Supplementary Fig. 5A and B), and restored the expression of PDX-1, GLUT2, and PPAR $\gamma$  (Supplementary Fig. 5C–E). Injections in the distal pancreas did not induce  $\beta_2$ AR expression in other tissues, such as the liver (Supplementary Fig. 6A and B) or the skeletal muscle (Supplementary Fig. 6C and D).

These effects were paralleled by significant improvement in glucose tolerance and insulin secretion during GTT (Fig. 6A–E). Fasting insulin levels also increased, reaching values similar to those measured in 6-month-old mice (Table 2), further underlining the relevance of  $\beta_2$ AR function in enabling adequate pancreatic  $\beta$ -cell response to hyperglycemia.

## DISCUSSION

In the present work, we provide evidence that  $\beta_2$ AR gene deletion in mice causes reduction of glucose-stimulated insulin release by pancreatic  $\beta$ -cells. This phenotype is reminiscent of that observed in mice with targeted  $\beta$ -cell disruption of the  $G\alpha_s$  gene (30). In these mice, however, the impairment of  $G\alpha_s$  prevented response to multiple  $G\alpha_s$ -related receptors, causing a severe phenotype, with gross abnormalities in pancreatic islets. Interestingly, in islets from  $\beta_2$ AR $^{-/-}$  mice, PPAR $\gamma$  expression was reduced by 50%, leading to repression of the PPAR $\gamma$  downstream molecules PDX-1 and GLUT2, two key effectors of  $\beta$ -cell function (26,42,43). This downregulation resulted in a clear impairment in insulin release, though islet architecture and insulin content were not affected by the  $\beta_2$ AR gene deletion.

Rosen et al. (44) showed that islets from mice with targeted elimination of PPAR $\gamma$  in  $\beta$ -cells were approximately twice as large as those from control mice. Thus, we can speculate that in our model the 50% reduction in



**FIG. 5.**  $\beta_2AR$  ex vivo infection rescued age-dependent impairment of  $\beta$ -cell function. Density (A) and mRNA levels (B) of  $\beta_2AR$  were evaluated on cell membranes of islets isolated from  $\beta_2AR^{+/+}$  mice. Insulin release (C) was determined upon exposure to the indicated concentration of glucose or KCl as described in RESEARCH DESIGN AND METHODS. mRNA levels of PDX-1 (D), GLUT2 (E), and PPAR $\gamma$  (F) were determined by real-time RT-PCR using the pooled total RNAs from five mice/group with cyclophilin as internal standard. Each bar represents means  $\pm$  SE of five independent experiments in each of which reactions were performed in triplicate. Islets isolated from  $\beta_2AR^{+/+}$  mice were solubilized and aliquots of the lysates were blotted with PDX-1, GLUT2, and PPAR $\gamma$  antibodies. Actin was used as loading control. The autoradiographs shown (G) are representative of three independent experiments, which are quantified in H.  $\square$ , age 6 months (mos);  $\blacksquare$ , 20 months;  $\text{▨}$ , 20 months AdEmpty;  $\text{▩}$ , 20 months Ad $\beta_2AR$ . \* $P < 0.05$  vs.  $\beta_2AR^{+/+}$  6 months, Bonferroni post hoc test.

PPAR $\gamma$  levels is sufficient to restrain  $\beta$ -cell function without altering islet mass.


The mechanistic significance of  $\beta_2AR$  gene knockout was further sustained by in vitro studies in the INS-1E pancreatic  $\beta$ -cells, showing that the silencing of the  $\beta_2AR$  as well as the pharmacological treatment with a specific  $\beta_2AR$  antagonist impaired glucose response and down-regulated PPAR $\gamma$  expression, reducing both PDX-1 and GLUT2 levels. No alteration of  $\alpha ARs$  was observed instead

(data not shown). In addition, treatment with the  $\beta_2AR$  agonist fenoterol activated PPAR $\gamma$ /PDX-1/GLUT2 signaling, indicating that, at least in part,  $\beta_2AR$  controls insulin secretion through this pathway. Indeed, in this study we show that exogenous PPAR $\gamma$  expression in INS-1E  $\beta$ -cells silenced for  $\beta_2AR$  led to recovery of PDX-1/GLUT2 levels and glucose-stimulated insulin secretion. This finding is supported by recent evidence that directly relates  $\beta_2AR$  to PPAR $\gamma$  (42,45–47), a key element in the process of insulin

**TABLE 2**  
Metabolic effects of  $\beta_2AR$  overexpression in aged (20 months old)  $\beta_2AR^{+/+}$  mice

	20 months old			6 months old
	Untreated	AdEmpty	Ad $\beta_2AR$	Untreated
<i>n</i>	10	6	8	15
Body weight (g)	38.4 $\pm$ 1.7*	38.1 $\pm$ 2.4*	38.6 $\pm$ 2.1*	29.6 $\pm$ 1.1
Food intake (g/day)	4.0 $\pm$ 1.1*	4.2 $\pm$ 1.8*	3.9 $\pm$ 1.5*	3.1 $\pm$ 0.6
Water intake (mL/day)	6.8 $\pm$ 1.2*	6.9 $\pm$ 1.7*	6.6 $\pm$ 1.8*	5.8 $\pm$ 0.6
Random-fed blood glucose (mg/dL)	176.5 $\pm$ 8.6	178.4 $\pm$ 12.7	173.2 $\pm$ 10.4	170.1 $\pm$ 12.3
Fasting blood glucose (mg/dL)	84.2 $\pm$ 10.4	83.6 $\pm$ 11.9	77.7 $\pm$ 11.7	75.7 $\pm$ 8.2
Fasting serum insulin (ng/mL)	0.32 $\pm$ 0.04*	0.33 $\pm$ 0.05*	0.42 $\pm$ 0.1	0.41 $\pm$ 0.03

Data are means  $\pm$  SE unless otherwise indicated. \* $P < 0.05$  vs. adult (6 months old)  $\beta_2AR^{+/+}$  mice.



**FIG. 6.** Adenoviral vector-mediated  $\beta_2$ AR gene transfer in the mouse pancreas rescued age-related reduction in glucose tolerance. Blood glucose levels (A) and serum insulin (C) after 120 min of glucose administration ( $n = 14$ – $18$  animals per group). We calculated the AUC from glucose (B) and insulin excursion (D) curves. Twenty-month-old  $\beta_2$ AR $^{+/+}$  mice showed glucose intolerance (A and B), impaired insulin secretion (C and D), and also an impairment in  $\beta$ -cell function, evaluated measuring the peak insulin-to-peak glucose ratio (E). All of these parameters were restored after Ad $\beta_2$ AR in vivo infection. \* $P < 0.05$  vs.  $\beta_2$ AR $^{+/+}$  at 6 months (mos) of age, Bonferroni post hoc test. (See also Supplementary Figs. 5 and 6.)



secretion that has also recently been investigated in aging (43,48). Our results are consistent with these observations, sustaining also the hypothesis that cAMP levels could act as a connecting link through which  $\beta_2$ AR signaling leads to activation of PPAR $\gamma$  (49,50). Moreover, the cAMP assays, performed both in INS-1E<sub>sh</sub> $\beta_2$ AR pancreatic  $\beta$ -cells and in islets isolated from aged mice, showed an impairment in basal conditions and after stimulation with isoproterenol, while the responses to NaF and forskolin were not affected. Also, G $\alpha_s$  and AC-VI protein levels were not significantly different among the explored settings. This combination of events is usually observed in models of  $\beta_2$ AR gene deletion or impaired  $\beta_2$ AR signaling (18,34).

Whether and to what extent  $\beta_2$ AR gene knockout in liver and peripheral tissues affects glucose homeostasis in the  $\beta_2$ AR<sup>-/-</sup> mice remain to be conclusively addressed. Indeed, variations at the  $\beta_2$ AR locus have also been reported to associate with insulin resistance in type 2 diabetic patients (16). However, as shown in this work, the impaired glucose tolerance of  $\beta_2$ AR<sup>-/-</sup> mice is likely contributed by the defective  $\beta$ -cell function, as indicated by the major effect of  $\beta_2$ AR lack on glucose-evoked insulin secretion.

In humans, glucose tolerance declines with age, resulting in a high prevalence of type 2 diabetes and impaired glucose tolerance in the elderly population (2,8). How, at the individual level, glucose tolerance declines remains unclear, but it is likely determined by multiple factors including diminished insulin secretion (3,7). In rat models and in humans, a progressive decline in  $\beta$ -cell activity with age has been documented (4,6). In the present work, we show that the same occurs in the C57Bl/6N mouse and is paralleled by the development of abnormal glucose tolerance. Similar to previous findings in several human tissues (18–21,24), our results show that these changes are accompanied by reduced  $\beta_2$ AR levels in mouse pancreatic islets. The decreased  $\beta_2$ AR density in islets from aged mice recapitulates the mechanisms leading to the insulin secretory defect occurring in  $\beta_2$ AR-null mice, indicating that it may contribute to the age-related impairment in glucose tolerance. Indeed, both in vivo and ex vivo experiments of  $\beta_2$ AR gene transfer revealed that recovery of normal  $\beta_2$ AR levels rescued insulin release and glucose tolerance in aged mice. Thus, in the mouse model progressive decline of islet  $\beta_2$ AR density appears to contribute to the reduction in glucose tolerance that accompanies aging. Whether the same also occurs in humans needs to be clarified and is currently under investigation in our laboratory.

In conclusion, we have shown that  $\beta_2$ AR physiologically regulates pancreatic  $\beta$ -cell insulin secretion by modulating PPAR $\gamma$ /PDX-1/GLUT2 function. Reduced  $\beta_2$ AR expression contributes to the age-dependent deterioration of glucose tolerance.

#### ACKNOWLEDGMENTS

The financial support of Telethon and Fondazione Veronesi is gratefully acknowledged.

No potential conflicts of interest relevant to this article were reported.

G.S. conceived the project, performed experiments, analyzed data, and wrote the manuscript. A.L. performed experiments, analyzed data, and wrote the manuscript. D.S. performed experiments and contributed to discussion. A.A. performed experiments. C.D.G. performed experiments. P.F. analyzed data and contributed to discussion. F.B. analyzed data and wrote the manuscript. B.T. designed

research and supervised the project. C.M. analyzed data and wrote the manuscript. G.I. designed research, analyzed data, and wrote the manuscript. G.S. and G.I. are the guarantors of this work and, as such, had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

The authors thank Brian Kobilka (Stanford University, Stanford, CA) for providing the founders of  $\beta_2$ AR<sup>-/-</sup> mice, Pierre Maechler (University of Geneva, Geneva, Switzerland) for supplying INS-1E pancreatic  $\beta$ -cells, and Walter J. Koch (Center for Translational Medicine, and Thomas Jefferson University, Philadelphia, PA) for providing the Ad $\beta_2$ AR. The valuable technical assistance of Alfonso Anastasio ("San Giovanni di Dio" Hospital, Frattaminore, Italy) is also acknowledged.

#### REFERENCES

- Defronzo RA. Banting Lecture. From the triumvirate to the ominous octet: a new paradigm for the treatment of type 2 diabetes mellitus. *Diabetes* 2009;58:773–795
- Basu R, Breda E, Oberg AL, et al. Mechanisms of the age-associated deterioration in glucose tolerance: contribution of alterations in insulin secretion, action, and clearance. *Diabetes* 2003;52:1738–1748
- Gumbiner B, Polonsky KS, Beltz WF, Wallace P, Brechtel G, Fink RI. Effects of aging on insulin secretion. *Diabetes* 1989;38:1549–1556
- Ma X, Becker D, Arena VC, Vicini P, Greenbaum C. The effect of age on insulin sensitivity and insulin secretion in first-degree relatives of type 1 diabetic patients: a population analysis. *J Clin Endocrinol Metab* 2009;94:2446–2451
- Reaven E, Wright D, Mondon CE, Solomon R, Ho H, Reaven GM. Effect of age and diet on insulin secretion and insulin action in the rat. *Diabetes* 1983;32:175–180
- Perfetti R, Rafizadeh CM, Liotta AS, Egan JM. Age-dependent reduction in insulin secretion and insulin mRNA in isolated islets from rats. *Am J Physiol* 1995;269:E983–E990
- Ihm SH, Matsumoto I, Sawada T, et al. Effect of donor age on function of isolated human islets. *Diabetes* 2006;55:1361–1368
- Iozzo P, Beck-Nielsen H, Laakso M, Smith U, Yki-Järvinen H, Ferrannini E; European Group for the Study of Insulin Resistance. Independent influence of age on basal insulin secretion in nondiabetic humans. *J Clin Endocrinol Metab* 1999;84:863–868
- Rosengren AH, Jokubka R, Tojjar D, et al. Overexpression of alpha2A-adrenergic receptors contributes to type 2 diabetes. *Science* 2010;327:217–220
- Doyle ME, Egan JM. Pharmacological agents that directly modulate insulin secretion. *Pharmacol Rev* 2003;55:105–131
- Lembo G, Napoli R, Capaldo B, et al. Abnormal sympathetic overactivity evoked by insulin in the skeletal muscle of patients with essential hypertension. *J Clin Invest* 1992;90:24–29
- Seals DR, Esler MD. Human ageing and the sympathoadrenal system. *J Physiol* 2000;528:407–417
- Lembo G, Capaldo B, Rendina V, et al. Acute noradrenergic activation induces insulin resistance in human skeletal muscle. *Am J Physiol* 1994;266:E242–E247
- Asensio C, Jimenez M, Kühne F, Rohner-Jeanraud F, Muzzin P. The lack of beta-adrenoceptors results in enhanced insulin sensitivity in mice exhibiting increased adiposity and glucose intolerance. *Diabetes* 2005;54:3490–3495
- Haffner CA, Kendall MJ. Metabolic effects of beta 2-agonists. *J Clin Pharm Ther* 1992;17:155–164
- Ikarashi T, Hanyu O, Maruyama S, et al. Genotype Gly/Gly of the Arg16Gly polymorphism of the beta2-adrenergic receptor is associated with elevated fasting serum insulin concentrations, but not with acute insulin response to glucose, in type 2 diabetic patients. *Diabetes Res Clin Pract* 2004;63:11–18
- Scarpace PJ, Mooradian AD, Morley JE. Age-associated decrease in beta-adrenergic receptors and adenylate cyclase activity in rat brown adipose tissue. *J Gerontol* 1988;43:B65–B70
- Xiao RP, Tomhave ED, Wang DJ, et al. Age-associated reductions in cardiac beta1- and beta2-adrenergic responses without changes in inhibitory G proteins or receptor kinases. *J Clin Invest* 1998;101:1273–1282
- Schocken DD, Roth GS. Reduced beta-adrenergic receptor concentrations in aging man. *Nature* 1977;267:856–858

20. Feldman RD, Limbird LE, Nadeau J, Robertson D, Wood AJ. Alterations in leukocyte beta-receptor affinity with aging. A potential explanation for altered beta-adrenergic sensitivity in the elderly. *N Engl J Med* 1984;310:815–819
21. Bao X, Mills PJ, Rana BK, et al. Interactive effects of common beta2-adrenoceptor haplotypes and age on susceptibility to hypertension and receptor function. *Hypertension* 2005;46:301–307
22. Kang KB, Rajanayagam MA, van der Zyppe A, Majewski H. A role for cyclooxygenase in aging-related changes of beta-adrenoceptor-mediated relaxation in rat aortas. *Naunyn Schmiedeberg Arch Pharmacol* 2007;375:273–281
23. Ryall JG, Plant DR, Gregorevic P, Sillence MN, Lynch GS. Beta 2-agonist administration reverses muscle wasting and improves muscle function in aged rats. *J Physiol* 2004;555:175–188
24. White M, Roden R, Minobe W, et al. Age-related changes in beta-adrenergic neuroeffector systems in the human heart. *Circulation* 1994;90:1225–1238
25. Chruscinski AJ, Rohrer DK, Schauble E, Desai KH, Bernstein D, Kobilka BK. Targeted disruption of the beta2 adrenergic receptor gene. *J Biol Chem* 1999;274:16694–16700
26. Evans-Molina C, Robbins RD, Kono T, et al. Peroxisome proliferator-activated receptor gamma activation restores islet function in diabetic mice through reduction of endoplasmic reticulum stress and maintenance of euchromatin structure. *Mol Cell Biol* 2009;29:2053–2067
27. Vigliotta G, Miele C, Santopietro S, et al. Overexpression of the ped/pea-15 gene causes diabetes by impairing glucose-stimulated insulin secretion in addition to insulin action. *Mol Cell Biol* 2004;24:5005–5015
28. Santulli G, Basilicata MF, De Simone M, et al. Evaluation of the anti-angiogenic properties of the new selective  $\alpha v\beta 3$  integrin antagonist RGDechiHCit. *J Transl Med* 2011;9:7
29. Sorriento D, Santulli G, Fusco A, Anastasio A, Trimarco B, Iaccarino G. Intracardiac injection of AdGRK5-NT reduces left ventricular hypertrophy by inhibiting NF-kappaB-dependent hypertrophic gene expression. *Hypertension* 2010;56:696–704
30. Xie T, Chen M, Zhang QH, Ma Z, Weinstein LS. Beta cell-specific deficiency of the stimulatory G protein alpha-subunit Gsalpha leads to reduced beta cell mass and insulin-deficient diabetes. *Proc Natl Acad Sci USA* 2007;104:19601–19606
31. Lombardi A, Ulianich L, Treglia AS, et al. Increased hexosamine biosynthetic pathway flux dedifferentiates INS-1E cells and murine islets by an extracellular signal-regulated kinase (ERK)1/2-mediated signal transmission pathway. *Diabetologia* 2012;55:141–153
32. Fiory F, Lombardi A, Miele C, Giudicelli J, Béguinot F, Van Obberghen E. Methylglyoxal impairs insulin signalling and insulin action on glucose-induced insulin secretion in the pancreatic beta cell line INS-1E. *Diabetologia* 2011;54:2941–2952
33. McGraw DW, Forbes SL, Mak JC, et al. Transgenic overexpression of beta (2)-adrenergic receptors in airway epithelial cells decreases bronchoconstriction. *Am J Physiol Lung Cell Mol Physiol* 2000;279:L379–L389
34. Mutlu GM, Dumasius V, Burhop J, et al. Upregulation of alveolar epithelial active Na<sup>+</sup> transport is dependent on beta2-adrenergic receptor signaling. *Circ Res* 2004;94:1091–1100
35. Ciccarelli M, Sorriento D, Cipolletta E, et al. Impaired neoangiogenesis in  $\beta 2$ -adrenoceptor gene-deficient mice: restoration by intravascular human  $\beta 2$ -adrenoceptor gene transfer and role of NF $\kappa$ B and CREB transcription factors. *Br J Pharmacol* 2011;162:712–721
36. Iaccarino G, Ciccarelli M, Sorriento D, et al. Ischemic neoangiogenesis enhanced by beta2-adrenergic receptor overexpression: a novel role for the endothelial adrenergic system. *Circ Res* 2005;97:1182–1189
37. Sorriento D, Ciccarelli M, Santulli G, et al. The G-protein-coupled receptor kinase 5 inhibits NFkappaB transcriptional activity by inducing nuclear accumulation of IkappaB alpha. *Proc Natl Acad Sci USA* 2008;105:17818–17823
38. Perino A, Ghigo A, Ferrero E, et al. Integrating cardiac PIP3 and cAMP signaling through a PKA anchoring function of p110 $\gamma$ . *Mol Cell* 2011;42:84–95
39. Ciccarelli M, Santulli G, Campanile A, et al. Endothelial alpha1-adrenoceptors regulate neo-angiogenesis. *Br J Pharmacol* 2008;153:936–946
40. Oriente F, Iovino S, Cassese A, et al. Overproduction of phosphoprotein enriched in diabetes (PED) induces mesangial expansion and upregulates protein kinase C-beta activity and TGF-beta1 expression. *Diabetologia* 2009;52:2642–2652
41. Mauvais-Jarvis F, Virkamaki A, Michael MD, et al. A model to explore the interaction between muscle insulin resistance and beta-cell dysfunction in the development of type 2 diabetes. *Diabetes* 2000;49:2126–2134
42. Faisy C, Pinto FM, Blouquit-Laye S, et al. beta2-Agonist modulates epithelial gene expression involved in the T- and B-cell chemotaxis and induces airway sensitization in human isolated bronchi. *Pharmacol Res* 2010;61:121–128
43. Blalock EM, Phelps JT, Pancani T, et al. Effects of long-term pioglitazone treatment on peripheral and central markers of aging. *PLoS ONE* 2010;5:e10405
44. Rosen ED, Kulkarni RN, Sarraf P, et al. Targeted elimination of peroxisome proliferator-activated receptor gamma in beta cells leads to abnormalities in islet mass without compromising glucose homeostasis. *Mol Cell Biol* 2003;23:7222–7229
45. Collins S, Yehuda-Shnaidman E, Wang H. Positive and negative control of Ucp1 gene transcription and the role of  $\beta$ -adrenergic signaling networks. *Int J Obes (Lond)* 2010;34(Suppl. 1):S28–S33
46. Fogli S, Pellegrini S, Adinolfi B, et al. Rosiglitazone reverses salbutamol-induced  $\beta(2)$ -adrenoceptor tolerance in airway smooth muscle. *Br J Pharmacol* 2011;162:378–391
47. Tadaishi M, Miura S, Kai Y, et al. Effect of exercise intensity and AICAR on isoform-specific expressions of murine skeletal muscle PGC-1 $\alpha$  mRNA: a role of  $\beta$ -adrenergic receptor activation. *Am J Physiol Endocrinol Metab* 2011;300:E341–E349
48. Sung B, Park S, Yu BP, Chung HY. Modulation of PPAR in aging, inflammation, and calorie restriction. *J Gerontol A Biol Sci Med Sci* 2004;59:997–1006
49. Guri AJ, Hontecillas R, Bassaganya-Riera J. Abscisic acid synergizes with rosiglitazone to improve glucose tolerance and down-modulate macrophage accumulation in adipose tissue: possible action of the cAMP/PKA/PPAR  $\gamma$  axis. *Clin Nutr* 2010;29:646–653
50. Kim SP, Ha JM, Yun SJ, et al. Transcriptional activation of peroxisome proliferator-activated receptor-gamma requires activation of both protein kinase A and Akt during adipocyte differentiation. *Biochem Biophys Res Commun* 2010;399:55–59