

RESEARCH

Open Access

# Prolonged oral cannabinoid administration prevents neuroinflammation, lowers $\beta$ -amyloid levels and improves cognitive performance in Tg APP 2576 mice

Ana María Martín-Moreno<sup>1,2,7</sup>, Begoña Brera<sup>1,2</sup>, Carlos Spuch<sup>2,3,8</sup>, Eva Carro<sup>2,3</sup>, Luis García-García<sup>4</sup>, Mercedes Delgado<sup>4</sup>, Miguel A Pozo<sup>4,5</sup>, Nadia G Innamorato<sup>2,6</sup>, Antonio Cuadrado<sup>2,6</sup> and María L de Ceballos<sup>1,2\*</sup>

## Abstract

**Background:** Alzheimer's disease (AD) brain shows an ongoing inflammatory condition and non-steroidal anti-inflammatories diminish the risk of suffering the neurologic disease. Cannabinoids are neuroprotective and anti-inflammatory agents with therapeutic potential.

**Methods:** We have studied the effects of prolonged oral administration of transgenic amyloid precursor protein (APP) mice with two pharmacologically different cannabinoids (WIN 55,212-2 and JWH-133, 0.2 mg/kg/day in the drinking water during 4 months) on inflammatory and cognitive parameters, and on <sup>18</sup>F-fluoro-deoxyglucose (<sup>18</sup>FDG) uptake by positron emission tomography (PET).

**Results:** Novel object recognition was significantly reduced in 11 month old Tg APP mice and 4 month administration of JWH was able to normalize this cognitive deficit, although WIN was ineffective. Wild type mice cognitive performance was unaltered by cannabinoid administration. Tg APP mice showed decreased <sup>18</sup>FDG uptake in hippocampus and cortical regions, which was counteracted by oral JWH treatment. Hippocampal GFAP immunoreactivity and cortical protein expression was unaffected by genotype or treatment. In contrast, the density of Iba1 positive microglia was increased in Tg APP mice, and normalized following JWH chronic treatment. Both cannabinoids were effective at reducing the enhancement of COX-2 protein levels and TNF- $\alpha$  mRNA expression found in the AD model. Increased cortical  $\beta$ -amyloid (A $\beta$ ) levels were significantly reduced in the mouse model by both cannabinoids. Noteworthy both cannabinoids enhanced A $\beta$  transport across choroid plexus cells *in vitro*.

**Conclusions:** In summary we have shown that chronically administered cannabinoid showed marked beneficial effects concomitant with inflammation reduction and increased A $\beta$  clearance.

**Keywords:** Alzheimer's disease,  $\beta$ -amyloid peptide, cannabinoids, glial activation, interleukin 6, anti-inflammatories, tumor necrosis factor- $\alpha$

## Background

Alzheimer's disease (AD) is the major cause of dementia. The cognitive impairment is associated with the degeneration of particular subsets of neurons in regions involved in learning and memory processes. In addition another invariant feature of AD is neuroinflammation,

considered a consequence of glial activation and reflected as astrogliosis and microglial activation, in particular around senile plaques, one of the pathological hallmarks of the disease, along neurofibrillary tangles. Indeed, lots of inflammatory parameters are found in AD brains [1,2]. Once initiated the inflammatory process it may contribute independently to neural dysfunction and cell death, establishing a self-perpetuating vicious cycle by which inflammation induces further neurodegeneration. The recognition of inflammation as

\* Correspondence: mceballos@cajal.csic.es

<sup>1</sup>Neurodegeneration Group, Dept. of Cellular, Molecular and Developmental Neurobiology, Instituto Cajal, CSIC, Doctor Arce, 37, Madrid, 28002, Spain  
Full list of author information is available at the end of the article

an important component in the disease led to the discovery that prolonged treatment with non-steroidal anti-inflammatories (NSAIDs) had beneficial effects for AD. Indeed, several prospective works have shown that this kind of treatment markedly reduced the risk of suffering the neurologic condition, delayed its onset, ameliorated the symptomatic severity and slowed cognitive decline [3-5]. However their administration to already demented patients may be ineffective, suggesting the importance of early administration or, alternatively, the existence of additional targets of NSAIDs, besides cyclooxygenase inhibition. Nevertheless, other compounds with anti-inflammatory activity may be disease modifying drugs, which may delay onset or slow its progression, in contrast with the present AD palliative treatment.

Cannabinoids, whether plant derived, synthetic or endocannabinoids, interact with two well characterized cannabinoid receptors, CB<sub>1</sub> and CB<sub>2</sub> [6,7]. In addition, some cannabinoids may interact with other receptors, such as the TRPV<sub>1</sub> receptor or the orphan receptor GPR55 [8,9]. The CB<sub>1</sub> receptor is widely distributed, with a particularly high expression in brain, which contrasts with the limited expression of the CB<sub>2</sub> receptor, which is characteristic of immune organs and cells [10]. In fact, while CB<sub>1</sub> receptors are expressed by all types of cells in the brain (neurons and glial cells), CB<sub>2</sub> are mainly localized in microglial cells [6,9-11], the resident immune cell of the brain.

We and others have proposed cannabinoids as preventive treatment for AD [12-14], based on their neuroprotective [15,16] and anti-inflammatory effects [11,17,18]. Indeed, cannabinoids are able to decrease the release of cytokines and nitric oxide in cultured microglial cells induced by lipopolysaccharide [19,20] and A $\beta$  addition [12,21]. In several *in vitro* studies cannabidiol (CBD), the major non-psychotropic constituent of cannabis, has shown to be neuroprotective against  $\beta$ -amyloid (A $\beta$ ) addition to cultured cells. This action was a consequence of reduction of oxidative stress and blockade of apoptosis [22], tau-phosphorylation inhibition through the Wnt/ $\beta$ -catenin pathway [23] and decreased iNOS expression and nitrite generation [24]. *In vivo* experiments have shown that several cannabinoids were effective at preventing Alzheimer's disease related changes. In a previous work we have reported that synthetic cannabinoids, such as WIN55,212-2 and JWH-133, prevented the cognitive impairment, glial activation and neuronal marker loss in  $\beta$ -amyloid injected rats [12]. Enhancement of endocannabinoid levels by subchronic uptake inhibition reversed the increase in inflammatory parameters, such as COX-2, TNF- $\alpha$  and IL-1, in A $\beta$ -injected mice, although cognitive impairment was only prevented in early treated mice [25]. Further, we have recently reported that CBD and WIN55,212-2 (WIN) inhibited both glial activation and cognitive deficits, as judged in the water maze test, and

by a mechanism involving decreasing microglial activation, as shown in cultured microglial cells [20].


So far the possible effects of cannabinoids have not been studied in a genetic AD model, which mimics the amyloidosis and neuroinflammation [26,27] that occurs in the neurologic condition, in the absence of overt neurodegeneration [28]. Therefore in this work we have addressed the question whether cannabinoid agonists would ameliorate the cognitive deficits, neuroinflammation and altered A $\beta$  levels in this AD model. To that end we have used WIN 55,212-2 (WIN), a mixed CB<sub>1</sub>/CB<sub>2</sub> agonist [6] and JWH-133 (JWH) as selective CB<sub>2</sub> agonist [29]. The CB<sub>2</sub> selective agonist is devoid of psychoactivity, an advantage for its clinical endorsement, however we reasoned that a low dose of WIN may be free of psychoactive effects as well. The drugs were administered in drinking water [30] for a prolonged period, 4 months, to mimic a possible clinical setting. Given that cannabinoid treatment may be preventive, but not curative, administration of drugs was started at 7 months of age, when no cognitive dysfunction or plaque deposition exists, and it was prolonged for 4 months (eg until 11 months of age). We report here that such a treatment ameliorates cognitive performance, decreases neuroinflammation and A $\beta$  levels, likely by increasing its transport to the periphery.

## Results

### Effect of cannabinoid oral treatment on cognitive impairment and glucose uptake by PET

In a previous work we assessed the effects of cannabinoids following intraventricular administration concomitant to A $\beta$  injection for 7 days to rats [12]. This type of administration is not feasible in the Clinic, and is restricted to very few disease conditions. Other authors have used oral administration of 0.1 mg/kg/day tetrahydrocannabinol (THC) effectively [30]. Therefore, in this work we have chosen to administer cannabinoids in the drinking water to investigate their effects in the genetic model of AD. Given that we aimed a preventive treatment, it was initiated at 7 months of age when they do not exhibit A $\beta$  deposition in plaques [28].

First we examined whether continuous oral treatment with a low dose (0.2 mg/kg/day for 4 months) of cannabinoids was able to rescue the cognitive impairment of Tg APP eleven months old mice in the novel object recognition test. Exploration activity was monitored during the training trial and all the animals displayed similar attention to the two identical objects. Wild type mice treated with vehicle were able to discriminate between the familiar object (Figure 1) and a novel object, exploring the latter for a longer time. WIN and JWH did not affect this behavior in wild type mice, even after prolonged treatment. As expected Tg APP mice did not distinguish both objects, showing cognitive impairment. In contrast Tg



APP mice treated with the CB<sub>2</sub> selective cannabinoid JWH spent more time exploring the novel object, although WIN was ineffective at that respect.


<sup>18</sup>FDG uptake is significantly decreased in areas involved in learning and memory both in AD patients and in subjects showing mild cognitive impairment [31]. Moreover it is considered to be an early marker of the neurologic disease. Prolonged WIN treatment reduced the uptake in wild type mice (Figure 2 B, C, D), while JWH was without effect. As expected 12 month Tg APP vehicle treated mice had reduced <sup>18</sup>FDG uptake (25-30% depending on the region of interest; Figure 2 A, B, C, D). JWH continuous administration fully reversed the reduction in glucose uptake in Tg APP mice (Figure 2 A, B, C, D), which was slightly higher than that of wild type mice (approximately 15%), while WIN was ineffective on that respect. Therefore JWH, the CB<sub>2</sub> selective agonist, was capable of improving cognitive impairment and of reversing the reduction in <sup>18</sup>FDG uptake in cortical areas and hippocampus of Tg APP.

**Cannabinoids reduce microglial activation and inflammatory parameters in Tg APP mice cerebral cortex**  
Neuroinflammation is reflected in AD and its transgenic models brain as astrogliosis and microglial activation,


particularly surrounding senile plaques [26,28]. GFAP immunohistochemistry showed similar staining patterns in hippocampus in the different groups studied (Figure 3 A and 3B), irrespective of the genotype or the treatment received. Similarly GFAP protein levels, as assessed by Western blotting, were maintained across the different groups (Figure 3 C and 3D). Microglial cell staining and density was assessed by using ionized Ca<sup>2+</sup>-binding adaptor molecule-1 (Iba-1) antibody. The antibody labeled resting microglial cells in wild type mice, characterized by the fine tortuous branching, which was unaffected by either of the cannabinoids used in the oral chronic treatment (Figure 4, A). Tg APP vehicle treated mice showed an increase (53%) in Iba-1 positive microglial cells compared with wild type mice (Figure 4, A and 4B). While continuous WIN treatment did not alter this increased density, the CB<sub>2</sub> selective agonist JWH decreased cell density so that cell number accounted 83% of that of wild type mice (Figure 4 B). These results indicate that at this age Tg APP mice do not show astrogliosis, but they had microglial activation, which was effectively counteracted by JWH oral treatment.

Recent works have shown increased CB<sub>2</sub> microglial expression following a lesion or in a neurodegeneration context [32], including AD brain [13,33]. We also addressed this issue by means of double immuno-fluorescence. We revealed CB<sub>2</sub> expression in Iba-1 microglial cells (additional file 1 A1-3) in quinolinic acid injected striatum (additional file 2) and not in the contralateral striatum (additional file 1, B1-3), as expected. However there was no evident CB<sub>2</sub> staining in any region examined of Tg APP mice (additional file 1, C1-3). Regarding CB<sub>2</sub> protein expression Tg APP mice had similar levels compared to wild type mice which also received vehicle (Figure 5 A and 5B). JWH administration, but not WIN, to wild type mice reduced (45%) CB<sub>2</sub> protein levels (Figure 5, B). Both WIN and JWH significantly decreased CB<sub>2</sub> protein (Figure 5, B) in the AD model, which attained very low levels (around a 75%-80% reduction). We used Western analysis or qRT-PCR to assess possible changes in inflammatory parameters in cerebral cortex (Figure 5). JWH, but not WIN, decreased COX-2 expression in wild type mice (Figure 5 C and 5D). This parameter was markedly increased a 49% in Tg APP mice (Figure 5 D). Noteworthy, both cannabinoids effectively reversed COX-2 protein enhancement, reaching even lower levels than in wild type mice (40% and 60% respectively vs. wild type vehicle treated mice).

Different cytokines have been shown to be increased in AD brain and in its genetic model. IL-6 mRNA expression was similar across the different groups (Figure 5, E). As expected cortical TNF- $\alpha$  mRNA expression was dramatically increased (around 6 fold vs wild type vehicle treated mice; Figure 5 F). The cannabinoid agonists used



**Figure 2** JWH oral administration rescued the decreased  $^{18}\text{F}$ -DG uptake in TgAPP assessed by PET. A: representative images [MR,  $^{18}\text{F}$ -DG uptake and merged] in hippocampus of wild type (Wt) and Tg APP vehicle treated mice. Continuous oral treatment of wild type mice with WIN decreased uptake in all regions studied (B: hippocampus; C: Frontal cortex; D: Temporo-parietal cortex). Tg APP showed decreased uptake which was normalized by JWH oral treatment (A-D). Animals were treated with vehicle or cannabinoids (0.2 mg/kg) in the drinking water for 4 months, starting at 7 months of age. Results are mean  $\pm$  SEM (n = 4 mice/group) of the ratio of average radioactivity in a given region of interest (ROI) by the radioactivity in cerebellum (Cb) expressed as percentage. \*p < 0.05, \*\* p < 0.01 vs vehicle treated mice; #p < 0.05 vs wild type-vehicle mice (ANOVA followed by Student's t test).



**Figure 3 Cannabinoid oral administration did not alter GFAP immunostaining or protein expression.** GFAP hippocampal immunostaining (IR) or cerebral cortical protein expression was similar in all groups, irrespective of genotype or treatment. A: representative IR in hippocampus from wild type and Tg APP mice treated with vehicle or cannabinoids. The area assessed is shown (5 hippocampal areas) B: Optical density (OD) was measured by densitometry. C: representative GFAP Western blot (WB) and D: optical density (OD) in cortical samples. Animals were treated with vehicle or cannabinoids (0.2 mg/kg) in the drinking water for 4 months, starting at 7 months of age. Results are mean  $\pm$  SEM (n = 6-7 mice/group).

in this study prevented this increase, although this cytokine expression continued to be higher than that of wild type mice (Figure 5, F).


Taken together these results demonstrate that continuous cannabinoid administration has anti-inflammatory activity given that different mediators were effectively counteracted in Tg APP.

#### Cannabinoids oral administration decreases $\beta$ -amyloid levels and increases its transport through choroid plexus monolayers

The Tg APP model parallels the amyloidosis of the neurologic condition. Eleven month old Tg APP mice have no detectable insoluble (formic acid extracted) A $\beta$  levels

by ELISA. As expected no soluble human transgenic A $\beta$  levels were detected in wild type mice, while Tg APP mice had high levels of A $\beta$ <sub>1-40</sub> (Figure 6, A) and detectable levels of the more amyloidogenic A $\beta$ <sub>1-42</sub> fragment (Figure 6, B). JWH decreased by 27% the levels of A $\beta$ <sub>1-40</sub>, and a 30% similar reduction in A $\beta$ <sub>1-42</sub> was found following the administration of either WIN or JWH (Figure 6, A and 6B).

To determine the mechanism by which both cannabinoids reduce A $\beta$  levels first we examined the effect of both compounds on A $\beta$  release from cultured APP/PS1 glioma cells. There was no difference in the presence or the absence of drugs (data not shown) suggesting that neither the release nor the synthesis of A $\beta$  was altered. Next we



wondered whether the cannabinoids would reduce A $\beta$  levels by enhancing its clearance, a therapeutic strategy intensively studied for AD treatment. Therefore we assessed A $\beta$  transport across choroid plexus monolayers *in vitro*. Since there are contradictory reports on the presence of CB<sub>1</sub> receptors in choroid plexus [34,35] first we performed stainings with CB<sub>1</sub> and CB<sub>2</sub> antibodies. We revealed the presence of both receptors in rat choroid plexus (Figure 6, C). Then we examined the possible time course of the transport of A $\beta$ <sub>1-40</sub> (5  $\mu$ g/ml) added to the upper compartment, by measuring the amount of peptide in the lower compartment by Western blotting at different time points (Figure 6 D). In the absence of any drug the peptide was recovered at 12 h, and at longer times (24 and 48 h), after its addition to the upper compartment. However, the transport was much quicker in the presence of WIN or JWH (200 nM) and A $\beta$  was mainly transported at shorter times (1-3 h; Figure 6 D). To quantify this transport we took advantage of ELISA assays and assessed peptide levels at 1 and 3 h after the addition of A $\beta$  to the cell cultures (Figure 6 E). WIN and JWH promoted the same peptide transport both at 1 and 3 h, while almost no detectable A $\beta$  crossed the choroid plexus cell monolayer in their absence (Figure 6 E). In summary, prolonged oral administration of cannabinoids effectively reduced A $\beta$  levels in Tg APP brain, likely due to increased peptide clearance through the blood- or the CSF-barrier.

#### Effect of prolonged oral treatment with cannabinoids on GSK3- $\beta$ levels

AD is characterized by the existence of increased tau phosphorylation, mainly by the action of GSK3- $\beta$ , which is deregulated. We assessed by Western blotting total and


pSer9-GSK3- $\beta$  (inactive form) levels in cerebral cortex, from Wt and Tg APP mice following the different treatments. Cannabinoid agonist treatment did not change p-Ser9 GSK3- $\beta$  protein levels in wild type mice. In contrast, there was a reduction (around 40%) in p-GSK3- $\beta$  in the AD model which was normalized by WIN administration, but not JWH (28% decrease vs wild type vehicle treated mice). Total GSK3- $\beta$  protein levels remained largely unchanged. Similarly no changes in the levels of either p-Ser21 GSK3- $\alpha$  or total GSK3- $\alpha$  protein levels were observed (Figure 7). Therefore, the pathological pSer9-GSK3- $\beta$  activity was normalized by the CB<sub>1</sub>/CB<sub>2</sub> mixed agonist WIN.

#### Discussion

In this work we describe several beneficial effects of the prolonged oral treatment with two cannabinoid agonists with different pharmacological profiles. Both cannabinoids were effective at decreasing the inflammatory parameters and A $\beta$  levels. However, it was the CB<sub>2</sub> selective agonist JWH that was able to prevent cognitive deficits and glucose metabolism reduction.

Our results differ from a recent work [36] reporting variable effects on water maze performance and fear conditioning, no changes in A $\beta$  levels or plaque burden following subchronic treatment with the CB<sub>1</sub>/CB<sub>2</sub> mixed agonist HU-210. Methodological differences may account for the results, since they used male and female double APP23/PS1 transgenic mice, at 4 (young) or 6-9 weeks of age, and a very low dose of the cannabinoid agonist (10 or 50  $\mu$ g/kg) injected twice daily [14].

CB<sub>1</sub> activation induces psychoactive effects. In fact mixed CB<sub>1</sub>/CB<sub>2</sub> and CB<sub>1</sub> selective agonists after acute




**Figure 5** Cannabinoid oral treatment decreased inflammatory parameters of Tg APP mice. A, B: Cannabinoids decreased CB<sub>2</sub> protein levels by *Western blotting* in Tg APP. C, D: Cannabinoids counteracted the increase in COX-2 protein levels by *Western blotting*. E: IL6 mRNA expression (qRT-PCR) was similar in all groups studied. F: Cannabinoids blocked the increased TNF-α mRNA expression (qRT-PCR) in Tg APP. Animals were treated with vehicle or cannabinoids (0.2 mg/kg) in the drinking water for 4 months, starting at 7 months of age. Results are mean ± SEM (n = 6-7 mice/group) \*p < 0.05, \*\* p < 0.01 vs vehicle treated mice; #p < 0.05 vs wild type mice vehicle treated mice (ANOVA followed by Student's t test).

administration decrease motor activity and impair learning and memory [37-39], although at higher doses than the one we used in this study (0.2 mg/kg/day). However, chronic administration of those drugs induces tolerance to their acute effects in different behavioural tests [40,41]. Therefore, the fact that WIN did not affect learning in the novel object recognition test in wild type mice might suggest tolerance after prolonged administration. Similarly WIN was without effect on recognition memory of Tg APP mice. In contrast, CB<sub>2</sub> selective agonists, such as JWH, are devoid of psychoactivity after acute administration and does not alter motor activity [[42], Martín-

Moreno et al., in preparation] after systemic administration or object recognition memory following its intrahippocampal injection [43]. After prolonged administration to wild type mice JWH neither altered learning nor memory in the present work. Interestingly this compound effectively counteracted the cognitive impairment of Tg APP mice.

The brain is the organ with the highest glucose consumption, which is believed to be coupled to neural activity. The reduction in brain glucose uptake has been repeatedly demonstrated in AD patients [31,44], in particular in regions involved in memory, and it is highly




**Figure 6** Cannabinoids decreased Aβ levels of Tg APP mice and increased transport through choroid plexus cell monolayer. **A:** Both cannabinoids significantly decreased Aβ<sub>1-42</sub> cortical levels. **B:** JWH decreased Aβ<sub>1-40</sub> cortical levels in Tg APP mice. Animals were treated with vehicle or cannabinoids (0.2 mg/kg) in the drinking water for 4 months, starting at 7 months of age. Results are mean ± SEM (n = 6-7 mice/group). **C:** rat choroid plexus express both CB<sub>1</sub> and CB<sub>2</sub> receptors. **D:** time-course of Aβ<sub>1-40</sub> transport through choroid plexus cells. Both cannabinoids (200 nM) reduced the transport time compared to untreated cultures (representative *Western blot*). **E:** Cannabinoids enhanced Aβ<sub>1-40</sub> transport through choroid plexus cells at 1 and 3 h after addition vs untreated control cultures (mean ± SEM, n = 4). **E:** \*p < 0.05, \*\*\* p < 0.01 vs vehicle treated mice or cultures (Student's t test).

correlated with cognitive deficits. Prolonged oral WIN administration reduced glucose uptake, as measured by PET <sup>18</sup>F-DG, in wild type mice in cortical regions and hippocampus. This result may be deleterious, in spite of the observation of no memory impairment in the cognitive test selected. In fact, previous autoradiographic work has reported either normal <sup>3</sup>H-DG uptake in hippocampus or a decrease, depending on the dose of WIN [45] acutely administered. Decreased <sup>18</sup>F-DG uptake in Tg APP in the present work is in essential agreement with

previous reports, either by autoradiographic techniques [46] or PET [47]. Notably JWH administration for 4 months did not alter glucose uptake in wild type mice, while it completely abrogated the reduction observed in Tg APP. Brain activity in general, and cognition in particular rely on glucose metabolism [47], therefore the effects of the CB<sub>2</sub> agonist on both glucose utilization and recognition memory are of therapeutic interest.

Inflammation has several drawbacks including learning and memory impairment [48], in particular during ageing





**Figure 7** WIN oral treatment counteracted decreased phospho-Ser9 GSK3β in cerebral cortex of Tg APP mice. **A:** representative Western blot of p-Ser9 GSK3β. **B:** optical density (OD). **C:** total-GSK3β. **D:** optical density (OD). **E:** representative Western blot of p-Ser21 GSK3α. **F:** optical density (OD). **G:** total-GSK3α in cerebral cortex. **H:** optical density (OD). Wild type (Wt) and Tg APP mice were treated with vehicle or cannabinoids (0.2 mg/kg) in the drinking water for 4 months, starting at 7 months of age. Results are mean ± SEM (n = 6-7 mice/group) \*p < 0.05 vs wt-veh mice, # p < 0.05 vs Tg APP-vehicle treated mice; #p < 0.05 vs wild type vehicle treated mice (ANOVA followed by Student's t test).

[49,50]. The compounds under study behaved as anti-inflammatory agents, in agreement with previous reports [11,12,20,25,49-51]. Microglial activation, but not astrogliosis, was observed in 11 month old Tg APP mice. Previous works have shown prominent reactive astrogliosis and increased GFAP expression in transgenic mice, that appears to be age-dependent and related with disease progression [26,28,52], albeit being restricted to plaques. The lack of astrogliosis and changes in GFAP expression may be explained by the absence of plaques in the mice model at this age. This result is in agreement with the detection of GFAP mRNA, as assessed by non-radioactive *in situ* hybridization, in reactive astrocytes in close proximity with A $\beta$  plaques at 14 months of age, but not before, when plaques were absent [53]. Indeed we found no plaques with glial associated cells, although there was a significant enhancement in microglial cell density in Tg APP mice. Continuous JWH treatment for 4 months normalized this parameter, but WIN was ineffective. In different contexts (eg lesions) both astrocytes and microglia could be engaged in inflammation. However in light of those results we can ascribe the increase in inflammatory mediators to microglial activation, given that there was no overt astrogliosis. COX-2 protein levels and TNF- $\alpha$  mRNA expression is increased in AD and its transgenic model [54,55]. Both cannabinoids significantly decreased COX-2 and TNF- $\alpha$ , as expected, since both compounds share CB<sub>2</sub> receptor activation, as shown by the down-regulation of its expression in Tg APP mice. It should be noted that although CB<sub>2</sub> receptors could be expressed by some neurons [56] they are mainly expressed by microglial cells [6,9-11], and are involved in the modulation of several inflammatory mediators [12,19-21,25]. We did expect an increase in CB<sub>2</sub> receptor expression in the transgenic model given that A $\beta$  addition to microglial cultures enhance it [21]. However, in Tg APP mice we did not observe CB<sub>2</sub> co-localization with Iba-1, which contrasts with the microglial co-expression in AD brain [13,33]. We neither found an increase by Western blotting, in agreement with our previous results in AD patients or A $\beta$ -injected rats [12]. This supports the notion that at this age there is an ongoing glial activation of low magnitude in Tg APP and that cannabinoids down-modulate this response.

A $\beta$  removal is considered a therapeutic strategy in AD, promoted either by vaccination [57,58] or by enhancing its clearance towards the periphery [59,60]. One of the most interesting findings of the present work is the A $\beta$  lowering ability of both cannabinoids, which we report for the first time. Prolonged oral JWH treatment decreased A $\beta$ <sub>1-40</sub> levels in brain and both cannabinoids decreased the more amyloidogenic fragment, A $\beta$ <sub>1-42</sub>. Given that the drugs did not alter A $\beta$  release we speculated that APP cleavage was not altered, and

therefore studied whether cannabinoids changed the peptide transport *in vitro*. Rat choroid plexus expressed CB<sub>1</sub>, in agreement with [34], and also CB<sub>2</sub> receptor protein, making feasible their activation by the drugs under study. Cannabinoids favored A $\beta$  transport, that was mainly observed at shorter times (1-3 h) compared to control experiments. This interesting effect merits further study.

At variance with those effects oral treatment with WIN, but not JWH, normalized the levels of GSK3- $\beta$  in Tg APP mice. Neurofibrillary tangles (NFTs), resulting from an abnormal phosphorylation of microtubule-associated tau proteins, represent a key pathological hallmark of Alzheimer's brain. GSK3- $\beta$  is the kinase mainly responsible for tau hyperphosphorylation, therefore inhibiting its activity is considered of therapeutic interest. The effect of WIN after prolonged oral administration is in accordance with reports showing a CB<sub>1</sub> dependent increase in GSK3- $\beta$  phosphorylation in cultured cells [61], in brain after acute cannabinoid agonist administration [62], and with a reduction in tau phosphorylation [23]. Neurofibrillary tangles are intraneuronal elements and neurons are in general devoid of CB<sub>2</sub> receptors. Therefore the effects of WIN on GSK3- $\beta$ , which were not mimicked by the CB<sub>2</sub> selective agonist JWH, might be explained by its interaction with CB<sub>1</sub> receptors in neurons.

Given that our previous work had shown that cannabinoids were preventive against the A $\beta$  effects, both *in vitro* and *in vivo* [12,20] we decided to start the continuous oral treatment at 7 months of age. At this time Tg APP mice do not have plaques and show normal learning and memory compared to wild type mice. Nevertheless, the treatment ended at 11 months of age when Tg APP begin to show memory disruption. According to our results the prolonged drug treatment decreased microglial activation of Tg APP mice along several inflammatory mediators, which were increased. However, ageing alters microglial responsiveness (eg to A $\beta$  production and deposition), which is highly dynamic and context dependent [63]. Therefore a potential caveat of our results is that they may not be applicable to aged pathological microglia as occurs in severe AD. However, a preventive treatment at very early stages of the disease may be feasible and beneficial as has been shown with the anti-inflammatory trifusal, both in amnesic mild cognitively impaired patients [64] and in Tg APP mice [65].

Over the last decade important findings on the involvement of the endocannabinoid system in AD has been gathered. Indeed, in AD brain there is increased expression of CB<sub>2</sub> receptors in microglia and of fatty acid amide hydrolase, the enzyme responsible for anandamide degradation, in astrocytes around plaques [13]. However CB<sub>1</sub> localization is markedly altered and its protein expression and functionality diminished [12]. Furthermore, molecular

rearrangements in different endocannabinoid system elements suggest that 2-AG signaling is increased, possibly contributing to synaptic failure in AD [66], while anandamide levels are decreased and are inversely correlated with A $\beta$  levels [67]. Interestingly, the CB<sub>2</sub> receptor expression has been reported to be increased both in the brain of AD patients [68] and in peripheral blood, where a significant correlation was found with the dementia score [69]. Finally, a CB<sub>2</sub> PET radiotracer is accumulated in brains showing neuroinflammation (eg LPS injected and Tg APP/PS1 mice; [70]). These latter results suggest the importance of CB<sub>2</sub> receptor as a biomarker of the neurologic disease, but also as a therapeutic target. CB<sub>2</sub> receptor increased expression in AD appears to be a consequence of microglia activation, but more importantly they render microglia susceptible to cannabinoid modulation, decreasing the generation cytotoxic molecules and inhibiting microglial activation, while promoting its migratory activity [10,11,20].

## Conclusions

In summary, cannabinoid agonists, in particular CB<sub>2</sub> selective agonists, interfere with several interconnected events of importance in the pathophysiology of AD. These compounds by directly interacting with cannabinoid receptors, in particular CB<sub>2</sub>, decrease microglial activation thereby reducing inflammation and its consequences (eg cognitive deficits). At the same time they may indirectly have beneficial effects on microglial activation (eg decrease cytokine release) by lowering brain A $\beta$  levels.

## Methods

### Animals and treatments

Tg APP transgenic mice were obtained via heterozygous breeding of mice expressing the 695 aa long isoform of the human APP containing a double mutation Lys 670-Asn, Met 671-Leu [52] (swedish mutation) under transcriptional control of the hamster prion promoter on a C57/BL6 breeding background. Male Tg APP and wild type littermates, used as controls, were 7 months age at the beginning of the experiments. Mice were group-housed (4-5 animals per cage) with a 12:12 h light/dark cycle and with *ad libitum* access to food and water. All of the experiments were performed according to ethical regulations on the use and welfare of experimental animals of the European Union and the Spanish Ministry of Agriculture, and the procedures were approved by the bioethical committee of the CSIC.

WIN 55,212-2 (WIN) and JWH-133 (JWH) were administered in the drinking water at a dose of 0.2 mg/kg/day using ethanol (0.1%) as vehicle. The amount of water drunk by the animals was assessed every other day and the treatment was adjusted to their weight. There

was no difference in the body weight or the ingested water between groups, all along the experiment, discarding a possible reinforcing effect of cannabinoids.

Animals were sacrificed by cervical dislocation followed by decapitation at 11 months of age. The brain was sagittally divided. One hemisphere was rapidly dissected on a cold plate, frozen on dry ice and stored at -80°C until assayed. The other hemisphere was immersion fixed in PF 4% in PB 0.1 M for 24 h, cryoprotected in sucrose 15% (24 h) and 30% (24 h) in PB, snap frozen in hexane (-60°C) and stored at -20°C until cut with a sliding microtome.

### Novel object recognition test

The arena measured 40 × 40 cm, surrounded by 30-cm-high perimeter black walls, that was located in an isolated room that was novel to the animals. The floor of the arena was covered with used sawdust. The arena was monitored by a video-camera located above the arena. The procedure consisted in three visits to the arena in subsequent days. The first day mice were placed into the empty arena for 15 min (habituation). The second day they were allowed to explore two identical objects during two 10 min trials 5 min apart (training). On the third day (test), one of the objects was changed by a novel one, with different shape and color, and the mice explored the arena for 10 min. Data collection was carried out by the Ethovision software (Noldus, The Netherlands) and exploring was defined as “directing the nose at a distance equal to or less than 3 cm from the object or touching it with the nose”. The time spent exploring the familiar object and the new object was expressed as a percentage. Objects were cleaned before every exposure with acetic acid 0.1% to prevent any olfactory clues. Experiments were performed at the same time of the day (9.00-14.00 h).

### <sup>18</sup>Fdeoxyglucose (18FDG) Positron Emission Tomography (PET)

Fasted mice were anesthetized with isoflurane (2%) and injected (ip) with <sup>18</sup>FDG (11,1 MBq or 300  $\mu$ Ci/200  $\mu$ l saline, PET Technologic Institute, Madrid). Thirty min later <sup>18</sup>FDG uptake images of each mouse were acquired for 30 min by PET imaging (Albira PET, 8 detectors, Gem-Imaging, Spain; [71]). The regions of interest were previously delineated in magnetic resonance (MR) T2-weighted images (Bruker Biospin, Germany) of each animal. Quantification of the metabolic activity was performed by co-registering the PET images of the brains to their own MR image as described by [72]. In our case, the field of view (FOV) of the PET scanner is 80 × 80 × 40 mm and the number of pixels of the reconstructed tomographic image is 160 × 160 × 80 pixels, being the voxel size 0.5 mm<sup>3</sup>. Co-registration of the PET image to the MRI, leads to a reduction of the pixel size to 0.2 mm<sup>3</sup>, given the trilinear

interpolation done by the PMOD software (PMOD Technologies, version 2.9, Switzerland). The  $^{18}\text{F}$ FDG uptake of the different brain areas were normalized to the  $^{18}\text{F}$ FDG uptake in the cerebellum (considered as reference region).

#### Immunohistochemistry and image analysis

Immunostaining was performed on floating sections (35  $\mu\text{m}$ ) as described previously [12]. In brief, following several washes with PBS, the endogenous peroxidase was blocked (3% hydrogen peroxide in methanol), washed again, and incubated for 90 min in PB containing 0.2% Triton  $\times$  100 and 10% normal goat serum. Sections were incubated with the different antibodies in PB containing 0.2% Triton  $\times$  100 and 1% normal goat serum overnight at 4°C. Dilutions of antibodies were as follows: anti-GFAP (1:1500, Sigma); anti-Iba-1 (1:1000, WAKO) (additional file 2). Development was conducted by the ABC method (Pierce, Rockford, IL), and immunoreactivity was visualized by 3,3'-diaminobenzidine oxidation as chromogen, with or without nickel enhancement. Images were acquired with Zeiss Axiocam high resolution digital color camera, using the same settings and the segmentation parameters (MCID software, InterFocus Imaging, UK) were constant for a given marker and experiment. The mean value for each animal per region results from the analysis of 5-6 sections.

#### A $\beta$ levels

A $\beta$  measurements were performed by two ELISA kits, one for each fragment (A $\beta$ 1-40 and A $\beta$ 1-42), from Biosource following the manufacturer instructions. The samples were sonicated (5 sec) in 10 vol of protein lysis buffer containing protease inhibitors (see Western blotting for details). The lysate was centrifuged (18,000  $\times$  g, for 10 min at 4°C). The supernatant was considered soluble and the pellets were further extracted by sonicating with formic acid and centrifuged. Prior to the Elisa the insoluble samples were diluted 3 times with Tris 1 M, pH 10, while the soluble samples were diluted 5 times in Elisa buffer. The results were expressed as pg/mg of protein measured by the Bradford method, using BSA as standard.

#### A $\beta$ transport

A double-chamber choroid plexus epithelial cell culture system mimicking the blood-cerebrospinal fluid (CSF) interface was used for *in vitro* studies, as previously described [59]. After seeding, the cells were incubated for 24 h and thereafter A $\beta$ <sub>1-40</sub> (5  $\mu\text{g}/\text{ml}$ ) was added to the lower chamber in the absence or presence of the cannabinoid agonists (500 nM). At different time points the medium was collected from the upper chamber, and the A $\beta$ <sub>1-40</sub> content was determined by immunoblotting. In other experiments A $\beta$ <sub>1-40</sub> levels were assessed by selective Elisa kits (Biosource) as described above.

#### Western blotting

Western blot was performed as described previously [61]. In brief, tissues were sonicated in lysis buffer, samples were centrifuged at high speed for 10 min, and supernatants were collected. Total protein was assessed by the Bio-Rad (Hercules, CA) protein assay. An aliquot of each sample (40  $\mu\text{g}$  of protein) was separated by SDS-PAGE (10%), and proteins were transferred from the gels onto nitrocellulose membranes. The blots were blocked with 1% defatted dry milk for 1 h at room temperature and incubated overnight at 4°C with the following antibodies: anti-GFAP (1:10.000, DAKO), anti-CB<sub>2</sub> (1:10.000, Affinity Bioreagents); anti-COX2 (1:100, Abcam); anti-p-GSK3 $\beta$  (1:5000, BD); anti-GSK3 $\beta$  (1:1.500, Cell Signaling) (additional file 2). Finally, samples were subjected to enhanced chemiluminescence and densitometric analysis. Band densitometric analysis was performed by Quantity One quantitation software (version 5.0; Bio-Rad) from film exposures; the background was subtracted, and the optical density percentage was obtained considering 100% the control samples within the same film. Tubulin was used as loading control. Every membrane contained samples from each treatment group, for comparison purposes.

The specificity of the CB<sub>2</sub> antibodies used (additional file 2) was assessed by preincubating with the human antigenic peptide (2  $\mu\text{g}/\text{ml}$  overnight at 4°C under agitation), as stated in additional file 2, before its use in Western blotting, which resulted in blockade of the immunoreactive signal (see additional file 3).

#### Analysis of mRNA levels by quantitative real-time PCR

Total RNA from cortex was extracted using TRIzol reagent according to the manufacturer's instructions (Invitrogen). To avoid interference with potential genomic DNA amplification, we treated 1  $\mu\text{g}$  of total RNA with 1  $\mu\text{l}$  DNase I (Invitrogen) plus 1  $\mu\text{l}$  of 10 $\times$  Buffer (Invitrogen). The samples were incubated at RT for 15 min. EDTA (25 mM) was added to the mixture and the samples were incubated at 65°C for 15 min to heat inactivate the DNase I. For cDNA synthesis a total of 1  $\mu\text{g}$  of RNA from the different samples were reverse-transcribed for 75 min at 42°C using 5 U of avian myeloblastosis virus reverse transcriptase (Promega) in the presence of 20 U of RNasin (Promega). The real-time PCR reaction was performed in 25  $\mu\text{l}$  using the fluorescent dye SYBR Green Master mix (Applied Biosystems) and a mixture of 5 pmol of reverse and forward primers. The primers used were for TNF- $\alpha$  forward primer 5' CATCTTCTCAAATTCGAGTGACAA 3' and reverse primer 5' TGGGAGTAGACAAGGTACAACCC 3' and for IL-6 forward primer 5' GAGGATACCCTCCCAACAGACC 3' and reverse primer 5' AAGTGCATCATCGTTGTTCATACA 3'. Quantification was performed on an ABI PRISM 7900 sequence detection

system (Applied Biosystems). PCR cycles proceeded as follows: initial denaturation for 10 min at 95°C, then 40 cycles of denaturation (15 sec, 95°C), annealing (30 sec, 60°C), and extension (30 sec, 60°C). The melting-curve analysis showed the specificity of the amplifications. Threshold cycle, which inversely correlates with the target mRNA level, was measured as the cycle number at which the reporter fluorescent emission appears above the background threshold (data not shown). Data analysis is based on the  $\Delta$ CT method with normalization of raw data to a house-keeping gene ( $\beta$ -actin). All of the PCRs were performed in triplicate.

### Statistical analysis

Statistical significance analysis was assessed by using one-way or two-way analysis of variance (ANOVA) followed by unpaired Student's t test (version 5.0, Prism software, GraphPad, USA). A value of  $p < 0.05$  was considered significant.

### Additional material

**Additional file 1: Co-localization of CB<sub>2</sub> and Iba-1 immunostaining in quinolinic acid injected striatum.** Iba-1 positive microglia in striatum also express CB<sub>2</sub> receptor protein ipsilateral to the toxin injection, but was absent in the contralateral (unlesioned) striatum. Tg APP microglial cells are devoid of CB<sub>2</sub> immunostaining.

**Additional file 2: Additional methods.** (quinolinic acid striatal lesion and blockade of CB<sub>2</sub> antibodies with CB<sub>2</sub> blocking peptide) and additional file 1, which includes the antibodies, commercial source and dilution, used in immunohistochemistry and Western blotting.

**Additional file 3: Blockade of CB<sub>2</sub> antibodies 744 and 746 with CB<sub>2</sub> antigenic peptide.** The CB<sub>2</sub> peptide completely blocked immunostaining.

### Abbreviations

A $\beta$ :  $\beta$ -amyloid peptide; AD: Alzheimer's disease; COX-2: cyclooxygenase; GSK3- $\beta$ : glycogen synthase kinase 3- $\beta$ ; IL6: interleukin 6; TNF- $\alpha$ : tumor necrosis factor- $\alpha$

### Acknowledgements

We thank Dr. K. Mackie for anti-CB<sub>1</sub> antibody and M. E. Fernández de Molina and S. Plazuelo for excellent technical assistance. We acknowledge Dr. I. Lastres-Becker for thoughtful comments and constructive discussion on the manuscript. This work was supported by the Spanish Ministry of Science and Technology (SAF 2005-02845 to M.L.C.). A.M.M.-M. was recipient a fellowship from the Ministry of Education and Science.

### Author details

<sup>1</sup>Neurodegeneration Group, Dept. of Cellular, Molecular and Developmental Neurobiology, Instituto Cajal, CSIC, Doctor Arce, 37, Madrid, 28002, Spain. <sup>2</sup>Centre for Biomedical Research on Neurodegenerative Diseases (CIBERNED), Madrid, Spain. <sup>3</sup>Neuroscience Group, Research Institute Hospital 12 de Octubre, Av. de Córdoba s/n Madrid, 28041, Spain. <sup>4</sup>CAI de Cartografía Cerebral, Instituto Pluridisciplinar, UCM, Paseo Juan XXIII 1, Madrid, 28040, Spain. <sup>5</sup>PET Technology Institute, Manuel Bartolomé Cossío 10, Madrid, 28040, Spain. <sup>6</sup>Departamento de Bioquímica, Instituto de Investigaciones Biomédicas "Alberto Sols"UAM-CSIC, Facultad de Medicina, Universidad Autónoma de Madrid, Arturo Duperier 4, Madrid, 28029, Spain. <sup>7</sup>Current Address: MD Anderson Cancer Center, Arturo Soria 270, 28033 Madrid, Spain.

<sup>8</sup>Current Address: Department of Pathology and Neuropathology, University Hospital of Vigo (CHUVI), Vigo, Spain.

### Authors' contributions

AMM-M carried out the mice treatment, the behavioural experiments, Western blotting, ELISA studies ex vivo, the molecular genetic studies, and helped to draft the manuscript. BB and MLC performed the immunohistochemical studies. NI and AC assessed qPCR experiments and Western blotting. LGG, MD and MAP designed, carried out and analyzed the PET studies. CS and EC designed and conducted the transport experiments in choroid plexus cell monolayers, including the ELISAs assays. MLC conceived of the study, participated in its design and coordination and completed the manuscript. All authors read and approved the final manuscript.

### Competing interests

The authors declare that they have no competing interests.

Received: 19 September 2011 Accepted: 16 January 2012

Published: 16 January 2012

### References

1. Akiyama H, Barger S, Barnum S, Bradt B, Bauer J, Cole GM, Cooper NR, Eikelenboom P, Emmerling M, Fiebich BL, Finch CE, Frautschy S, Giffin WS, Hampel H, Hull M, Landreth G, Lue L, Mrak R, Mackenzie IR, McGeer PL, O'Banion MK, Pachter J, Pasinetti G, Plata-Salaman C, Rogers J, Rydel R, Shen Y, Strohmeyer R, Tooyoma I, Van Muiswinkel FL, Veerhuis R, Walker D, Webster S, Wegrzyniak B, Wenk G, Wyss-Coray T: **Inflammation and Alzheimer's disease.** *Neurobiol Aging* 2000, **21**:383-421.
2. Wyss-Coray T, Mucke L: **Inflammation in neurodegenerative disease - a double-edged sword.** *Neuron* 2002, **35**:419-432.
3. Breitner JC, Welsh KA, Helms MJ, Gaskell PC, Gau BA, Roses AD, Pericak-Vance MA, Saunders AM: **Delayed onset of Alzheimer's disease with nonsteroidal anti-inflammatory and histamine H2 blocking drugs.** *Neurobiol Aging* 1995, **16**:523-530.
4. In 't Veld BA, Launer LJ, Hoes AW, Ott A, Hofman A, Breteler MM, Stricker BH: **NSAIDs and incident Alzheimer's disease. The Rotterdam Study.** *Neurobiol Aging* 1998, **19**:607-611.
5. McGeer PL, Schulzer M, McGeer EG: **Arthritis and anti-inflammatory agents as possible protective factors for Alzheimer's disease: A review of 17 epidemiologic studies.** *Neurology* 1997, **47**:425-432.
6. Howlett AC, Barth F, Bonner TI, Cabral G, Casellas P, Devane WA, Felder CC, Herkenham M, Mackie K, Martin BR, Mechoulam R, Pertwee RG: **International Union of Pharmacology. XXVII. Classification of cannabinoid receptors.** *Pharmacol Rev* 2002, **54**:161-202.
7. Piomelli D: **The molecular logic of endocannabinoid signalling.** *Nat Rev Neurosci* 2003, **4**:873-884.
8. Begg M, Pacher P, Bátkai S, Osei-Hyiaman D, Offertáler L, Mo FM, Liu J, Kunos G: **Evidence for novel cannabinoid receptors.** *Pharmacol Ther* 2005, **106**:133-145.
9. Stella N: **Cannabinoid and cannabinoid-like receptors in microglia, astrocytes, and astrocytomas.** *Glia* 2010, **58**:1017-1030.
10. Stella N: **Cannabinoid signaling in glial cells.** *Glia* 2004, **48**:267-277.
11. Walter L, Stella N: **Cannabinoids and neuroinflammation.** *Brit J Pharmacol* 2004, **141**:775-785.
12. Ramírez BG, Blázquez C, Gómez del Pulgar T, Guzmán M, de Ceballos ML: **Prevention of Alzheimer's disease pathology by cannabinoids: neuroprotection mediated by blockade of microglial activation.** *J Neurosci* 2005, **25**:1904-1913.
13. Benito C, Núñez E, Pazos MR, Tolón RM, Romero J: **The endocannabinoid system and Alzheimer's disease.** *Mol Neurobiol* 2007, **36**:75-81.
14. Campbell VA, Gowran A: **Alzheimer's disease; taking the edge off with cannabinoids?** *Br J Pharmacol* 2007, **152**:655-662.
15. Mechoulam R, Spatz M, Shohami E: **Endocannabinoids and Neuroprotection.** *Sci STKE* 2002, re5.
16. van der Stelt M, Veldhuis WB, Maccarrone M, Bär PR, Nicolay K, Veldink GA, Di Marzo V, Vliegenghart JFG: **Acute neuronal injury, excitotoxicity, and the endocannabinoid system.** *Mol Neurobiol* 2002, **26**:317-346.
17. Correa FG, Mestre L, Docagne F, Borrell J, Guaza C: **The endocannabinoid anandamide from immunomodulation to neuroprotection. Implications for multiple sclerosis.** *Vitam Horm* 2009, **81**:207-230.

18. Klein WK: **Cannabinoid-based drugs as anti-inflammatory therapeutics.** *Nat Rev Immunol* 2005, **5**:400-411.
19. Cabral GA, Marciano-Cabral F: **Cannabinoid receptors in microglia of the central nervous system: immune functional relevance.** *J Leukoc Biol* 2005, **78**:1192-1197.
20. Martín-Moreno AM, Reigada D, Ramírez BG, Mechoulam R, Innamorado N, Cuadrado A, de Ceballos ML: **Cannabidiol and other cannabinoids reduce microglial activation in vitro and in vivo: relevance to Alzheimer's disease.** *Mol Pharmacol* 2011, **79**:964-973.
21. Ehrhart J, Obregon D, Mori T, Hou H, Sun N, Bai Y, Klein T, Fernandez F, Tan J, Shytle RD: **Stimulation of cannabinoid receptor 2 (CB2) suppresses microglial activation.** *J Neuroinflammation* 2005, **2**:29.
22. Iuvone T, Esposito G, Esposito R, Santamaria R, Di Rosa M, Izzo AA: **Neuroprotective effect of cannabidiol, a non-psychoactive component from Cannabis sativa, on beta-amyloid-induced toxicity in PC12 cells.** *J Neurochem* 2004, **89**:134-141.
23. Esposito G, De Filippis D, Carnuccio R, Izzo AA, Iuvone T: **The marijuana component cannabidiol inhibits  $\beta$ -amyloid-induced tau protein hyperphosphorylation through Wnt/ $\beta$ -catenin pathway rescue PC12 cells.** *J Mol Med* 2006, **84**:253-258.
24. Esposito G, De Filippis D, Maiuri MC, De Stefano D, Carnuccio R, Iuvone T: **Cannabidiol inhibits nitric oxide synthase protein expression and nitric oxide production in  $\beta$ -amyloid stimulated PC12 neurons through p38 MAP kinase and NF $\kappa$ B involvement.** *Neurosci Lett* 2006, **399**:91-95.
25. van der Stelt M, Mazzola C, Esposito G, Matias I, Petrosino S, De Filippis D, Micale V, Steardo L, Drago F, Iuvone T, Di Marzo V: **Endocannabinoids and beta-amyloid-induced neurotoxicity in vivo: effect of pharmacological elevation of endocannabinoid levels.** *Cell Mol Life Sci* 2006, **63**:1410-1424.
26. Frautschy SA, Yang F, Irizarry M, Hyman B, Saido TC, Hsiao K, Cole GM: **Microglial response to amyloid plaques in APPsw transgenic mice.** *Am J Pathol* 1998, **152**:307-317.
27. Masliah E, Sisk A, Mallory M, Mucke L, Schenk D, Games D: **Comparison of neurodegenerative pathology in transgenic mice overexpressing V717F beta-amyloid precursor protein and Alzheimer's disease.** *J Neurosci* 1996, **16**:5795-5811.
28. Irizarry MC, McNamara M, Fedorchak K, Hsiao K, Hyman BT: **APPsw transgenic mice develop age-related A beta deposits and neuropil abnormalities, but no neuronal loss in CA1.** *J Neuropathol Exp Neurol* 1997, **56**:965-973.
29. Huffman JW, Liddle J, Yu S, Aung MM, Abood ME, Wiley JL, Martin BR: **3-(1',1'-Dimethylbutyl)-1-deoxy-delta-8-THC and related compounds: synthesis and selective ligands for the CB2 receptor.** *Bioorg Med Chem Lett* 1999, **7**:2905-2914.
30. Steffens S, Veillard NR, Arnaud C, Pelli G, Burger F, Staub C, Karsak M, Zimmer A, Frossard JL, Mach F: **Low dose oral cannabinoid therapy reduces progression of atherosclerosis in mice.** *Nature* 2005, **434**:782-786.
31. Mosconi L: **Brain glucose metabolism in the early and specific diagnosis of Alzheimer's disease. FDG-PET studies in MCI and AD.** *Eur J Nucl Med Mol Imaging* 2005, **32**:486-510.
32. Palazuelos J, Aguado T, Pazos MR, Julien B, Carrasco C, Resel E, Sagredo O, Benito C, Romero J, Azcoitia I, Fernández-Ruiz J, Guzmán M, Galve-Roperh I: **Microglial CB2 cannabinoid receptors are neuroprotective in Huntington's disease excitotoxicity.** *Brain* 2009, **132**:3152-3164.
33. Halleskog C, Mulder J, Dahlström J, Mackie K, Hortobágyi T, Tanila H, Kumar Puli L, Färber K, Harkany T, Schulte G: **WNT signaling in activated microglia is proinflammatory.** *Glia* 2011, **59**:119-131.
34. Ashton JC, Appleton I, Darlington CL, Smith PF: **Cannabinoid CB1 receptor protein expression in the rat choroid plexus: a possible involvement of cannabinoids in the regulation of cerebrospinal fluid.** *Neurosci Lett* 2004, **364**:40-42.
35. Suárez J, Romero-Zerbo SY, Rivera P, Bermúdez-Silva FJ, Pérez J, De Fonseca FR, Fernández-Llebrez P: **Endocannabinoid system in the adult rat circumventricular areas: an immunohistochemical study.** *J Comp Neurol* 2010, **518**:3065-3085.
36. Chen B, Bromley-Brits K, He G, Cai F, Zhang X, Song W: **Effect of synthetic cannabinoid HU210 on memory deficits and neuropathology in Alzheimer's disease mouse model.** *Curr Alzheimer Res* 2010, **7**:255-261.
37. Baek J, Zheng Y, Darlington CL, Smith PF: **The CB1 receptor agonist, WIN 55,212-2, dose-dependently disrupts object recognition memory in adult rats.** *Neurosci Lett* 2009, **464**:71-73.
38. Lichtman AH, Varvel SA, Martin BR: **Endocannabinoids in cognition and dependence.** *Prostaglandins Leukot Essent Fatty Acids* 2002, **66**:269-285.
39. Yim TT, Hong NS, Ejaredar M, McKenna JE, McDonald RJ: **Post-training CB1 cannabinoid receptor agonist activation disrupts long-term consolidation of spatial memories in the hippocampus.** *Neuroscience* 2008, **151**:929-936.
40. Abood ME, Martin BR: **Molecular neurobiology of the cannabinoid receptor.** *Int Rev Neurobiol* 1996, **39**:197-221.
41. Hampson RE, Simeral JD, Kelly EJ, Deadwyler SA: **Tolerance to the memory disruptive effects of cannabinoids involves adaptation by hippocampal neurons.** *Hippocampus* 2003, **13**:543-556.
42. Sánchez C, de Ceballos ML, Gómez del Pulgar T, Rueda D, Corbacho C, Velasco G, Galve-Roperh I, Huffman JW, Ramón y Cajal S, Guzmán M: **Inhibition of glioma growth in vivo by selective activation of the CB2 cannabinoid receptor.** *Cancer Res* 2001, **61**:5784-5789.
43. Clarke JR, Rossato JI, Monteiro S, Bevilacqua LR, Izquierdo I, Cammarota M: **Posttraining activation of CB1 cannabinoid receptors in the CA1 region of the dorsal hippocampus impairs object recognition long-term memory.** *Neurobiol Learn Mem* 2008, **90**:374-381.
44. Berti V, Osorio RS, Mosconi L, Li Y, De Santi S, de Leon MJ: **Early detection of Alzheimer's disease with PET imaging.** *Neurodegener Dis* 2010, **7**:131-135.
45. Pontieri FE, Conti G, Zocchi A, Fieschi C, Orzi F: **Metabolic mapping of the effects of WIN 55212-2 intravenous administration in the rat.** *Neuropsychopharmacology* 1999, **21**:773-776.
46. Niwa K, Kazama K, Younkin SG, Carlson GA, Iadecola C: **Alterations in cerebral blood flow and glucose utilization in mice overexpressing the amyloid precursor protein.** *Neurobiol Dis* 2002, **9**:61-68.
47. Sadowski M, Pankiewicz J, Scholtzova H, Ji Y, Quartermain D, Jensen CH, Duff K, Nixon RA, Gruen RJ, Wisniewski T: **Amyloid-beta deposition is associated with decreased hippocampal glucose metabolism and spatial memory impairment in APP/PS1 mice.** *J Neuropathol Exp Neurol* 2004, **63**:418-428.
48. Chapman TR, Barrientos RM, Ahrensden JT, Maier SF, Patterson SL: **Synaptic correlates of increased cognitive vulnerability with aging: peripheral immune challenge and aging interact to disrupt theta-burst late-phase long-term potentiation in hippocampal area CA1.** *J Neurosci* 2010, **30**:7598-7603.
49. Marchalant Y, Brothers HM, Norman GJ, Karelina K, DeVries AC, Wenk GL: **Cannabinoids attenuate the effects of aging upon neuroinflammation and neurogenesis.** *Neurobiol Dis* 2009, **34**:300-307.
50. Marchalant Y, Cerbai F, Brothers HM, Wenk GL: **Cannabinoid receptor stimulation is anti-inflammatory and improves memory in old rats.** *Neurobiol Aging* 2008, **29**:1894-1901.
51. Esposito G, Scuderi C, Savani C, Steardo L Jr, De Filippis D, Cottone P, Iuvone T, Cuomo V, Steardo L: **Cannabidiol in vivo blunts beta-amyloid induced neuroinflammation by suppressing IL-1beta and iNOS expression.** *Br J Pharmacol* 2007, **151**:1272-1279.
52. Hsiao K, Chapman P, Nilsen S, Eckman C, Harigaya Y, Younkin S, Yang F, Cole G: **Correlative memory deficits, A beta elevation, and amyloid plaques in transgenic mice.** *Science* 1996, **274**:99-102.
53. Mehlhorn G, Hollborn M, Schliebs R: **Induction of cytokines in glial cells surrounding cortical beta-amyloid plaques in transgenic Tg2576 mice with Alzheimer pathology.** *Int J Dev Neurosci* 2000, **18**:423-431.
54. Colton CA, Mott RT, Sharpe H, Xu Q, Van Nostrand WE, Vitek MP: **Expression profiles for macrophage alternative activation genes in AD and in mouse models of AD.** *J Neuroinflammation* 2006, **3**:27.
55. Heneka MT, Sastre M, Dumitrescu-Ozimek L, Hanke A, Dewachter I, Kuiperi C, O'Banion K, Klockgether T, Van Leuven F, Landreth GE: **Acute treatment with the PPARgamma agonist pioglitazone and ibuprofen reduces glial inflammation and A beta1-42 levels in APPV7171 transgenic mice.** *Brain* 2005, **128**:1442-1453.
56. Van Sickle MD, Duncan M, Kingsley PJ, Mouhate A, Urbani P, Mackie K, Stella N, Makriyannis A, Piomelli D, Davison JS, Marnett LJ, Di Marzo V, Pittman QJ, Patel KD, Sharkey KA: **Identification and functional characterization of brainstem cannabinoid CB2 receptors.** *Science* 2005, **310**:329-332.
57. Bard F, Cannon C, Barbour R, Burke RL, Games D, Grajeda H, Guido T, Hu K, Huang J, Johnson-Wood K, Khan K, Kholodenko D, Lee M, Lieberburg J,

- Motter R, Nguyen M, Soriano F, Vasquez N, Weiss K, Welch B, Seubert P, Schenk D, Yednock T: **Peripherally administered antibodies against amyloid beta-peptide enter the central nervous system and reduce pathology in a mouse model of Alzheimer disease.** *Nat Med* 2000, **6**:916-919.
58. Schenk D, Barbour R, Dunn W, Gordon G, Grajeda H, Guido T, Hu K, Huang J, Johnson-Wood K, Khan K, Kholodenko D, Lee M, Liao Z, Lieberburg I, Motter R, Mutter L, Soriano F, Shopp G, Vasquez N, Vandeventer C, Walker S, Wogulis M, Yednock T, Games D, Seubert P: **Immunization with amyloid- $\beta$  attenuates Alzheimer disease-like pathology in the PDAPP mouse.** *Nature* 1999, **400**:173-177.
59. Carro E, Trejo JL, Gomez-Isla T, LeRoith D, Torres-Aleman I: **Serum insulin-like growth factor I regulates brain amyloid- $\beta$  levels.** *Nat Med* 2002, **8**:1390-1397.
60. Matsuoka Y, Saito M, LaFrancois J, Saito M, Gaynor K, Olm V, Wang L, Casey E, Lu Y, Shiratori C, Lemere C, Duff K: **Novel therapeutic approach for the treatment of Alzheimer's disease by peripheral administration of agents with an affinity to beta-amyloid.** *J Neurosci* 2003, **23**:29-33.
61. Molina-Holgado E, Vela JM, Arévalo-Martín A, Almazán G, Molina-Holgado F, Borrell J, Guaza C: **Cannabinoids promote oligodendrocyte progenitor survival: involvement of cannabinoid receptors and phosphatidylinositol-3 kinase/Akt signaling.** *J Neurosci* 2002, **22**:9742-9753.
62. Ozaita A, Puighermanal E, Maldonado R: **Regulation of PI3K/Akt/GSK-3 pathway by cannabinoids in the brain.** *J Neurochem* 2007, **102**:1105-1114.
63. Lucin KM, Wyss-Coray T: **Immune activation in brain aging and neurodegeneration: too much or too little?** *Neuron* 2009, **64**:110-122.
64. Gómez-Isla T, Blesa R, Boada M, Clarimón J, Del Ser T, Domenech G, Ferró JM, Gómez-Ansón B, Manubens JM, Martínez-Lage JM, Muñoz D, Peña-Casanova J, Torres F, TRIMCI Study Group: **A randomized, double-blind, placebo controlled-trial of triflusal in mild cognitive impairment: the TRIMCI study.** *Alzheimer Dis Assoc Disord* 2008, **22**:21-29.
65. Coma M, Serenó L, Da Rocha-Souto B, Scotton TC, España J, Sánchez MB, Rodríguez M, Agulló J, Guardia-Laguarta C, Garcia-Alloza M, Borrelli LA, Clarimón J, Lleó A, Bacskai BJ, Saura CA, Hyman BT, Gómez-Isla T: **Triflusal reduces dense-core plaque load, associated axonal alterations and inflammatory changes, and rescues cognition in a transgenic mouse model of Alzheimer's disease.** *Neurobiology of Disease* 2010, **38**:482-491.
66. Mulder J, Zilberter M, Pasquaré SJ, Alpár A, Schulte G, Ferreira SG, Köfalvi A, Martín-Moreno AM, Keimpema E, Tanila H, Watanabe M, Mackie K, Hortobágyi T, de Ceballos ML, Harkany T: **Molecular reorganization of endocannabinoid signalling in Alzheimer's disease.** *Brain* 2011, **134**:1041-1060.
67. Jung KM, Astarita G, Yasar S, Vasilevko V, Cribbs DH, Head E, Cotman CW, Piomelli D: **An amyloid  $\beta$ (42)-dependent deficit in anandamide mobilization is associated with cognitive dysfunction in Alzheimer's disease.** *Neurobiol Aging* 2011.
68. Grünblatt E, Zander N, Bartl J, Jie L, Monoranu CM, Arzberger T, Ravid R, Roggendorf W, Gerlach M, Riederer P: **Comparison analysis of gene expression patterns between sporadic Alzheimer's and Parkinson's disease.** *J Alzheimers Dis* 2007, **12**:291-311.
69. Grünblatt E, Bartl J, Zehetmayer S, Ringel TM, Bauer P, Riederer P, Jacob CP: **Gene expression as peripheral biomarkers for sporadic Alzheimer's disease.** *J Alzheimers Dis* 2009, **16**:627-634.
70. Horti AG, Gao Y, Ravert HT, Finley P, Valentine H, Wong DF, Endres CJ, Savonenko AV, Dannals RF: **Synthesis and biodistribution of [11C]A-836339, a new potential radioligand for PET imaging of cannabinoid type 2 receptors (CB2).** *Bioorg Med Chem* 2010, **18**:5202-5207.
71. Balcerzyk M, Kontaxakis G, Delgado M, Garcia-Garcia L, Correcher C, Gonzalez AJ, Gonzalez A, Rubio JL, Benlloch JM, Pozo MA: **Initial performance evaluation of a high resolution Albira small animal positron emission tomography scanner with monolithic crystals and depth-of-interaction encoding from a user's perspective.** *Meas Sci Technol* 2009, **20**:104011.
72. Jupp B, O'Brien TJ: **Application of coregistration for imaging of animal models of epilepsy.** *Epilepsia* 2007, **48**(Suppl 4):82-89.

doi:10.1186/1742-2094-9-8

**Cite this article as:** Martín-Moreno et al.: Prolonged oral cannabinoid administration prevents neuroinflammation, lowers  $\beta$ -amyloid levels and improves cognitive performance in Tg APP 2576 mice. *Journal of Neuroinflammation* 2012 **9**:8.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

