

RESEARCH ARTICLE

Open Access

Stigma toward schizophrenia among parents of junior and senior high school students in Japan

Hatsumi Yoshii^{1*}, Yuichiro Watanabe^{2,3†}, Hideaki Kitamura^{2†}, Zhang Nan^{1†} and Kouhei Akazawa^{1†}

Abstract

Background: Stigma toward schizophrenia is a substantial barrier to accessing care and adhering to treatment. Provisions to combat stigma are important, but in Japan and other developed countries there are few such provisions in place that target parents of adolescents. The attitudes of parents are important to address as first schizophrenic episodes typically occur in adolescence. In overall efforts to develop an education program and provisions against stigma, here we examined the relationship between stigma toward schizophrenia and demographic characteristics of parents of junior and senior high school students in Japan. The specific hypothesis tested was that contact and communication with a person with schizophrenia would be important to reducing stigma. A questionnaire inquiring about respondent characteristics and which included a survey on stigma toward schizophrenia was completed by 2690 parents.

Results: The demographic characteristics significantly associated with the Devaluation-Discrimination Measure were family income, occupation, presence of a neighbor with schizophrenia, and participation in welfare activities for people with mental illness ($p < 0.05$). The mean \pm SD score was 32.74 ± 5.66 out of a maximum of 48 points on the Link Devaluation-Discrimination Measure.

Conclusions: Stigma toward schizophrenia among parents of junior and senior high school students was in fact significantly stronger among members of the general public who had had contact with individuals with schizophrenia. In addition, stigma was associated with family income.

Background

The stigma attached to the psychosis of schizophrenia is the greatest obstacle to improving the lives of affected individuals and their families. Stigma can lead to a loss of opportunities, decreased self-esteem, discouraging experiences in the workplace or criminal justice system, and disparities in access to health care [1]. It can also drive individuals away from society and incline them towards nonadherence to medication [2]. Moreover, among various factors such as ignorance, denial, lack of motivation, absence of information on early psychosis, and lack of access to appropriate interventions, stigma is associated with psychosis remaining untreated [3].

People see the risks of intervention for schizophrenia as concerning two main issues: drug side-effects and

stigma or anxiety surrounding the use of the word 'psychosis' [4]. Many people shun individuals with symptoms of schizophrenia and do not accept them in their communities [5,6]. Therefore, someone with schizophrenia, or who has a family member with the illness, might be less willing to acknowledge the symptoms of the disorder and seek medical care. In this way, stigma becomes a substantial barrier to accessing care and adhering to treatment [1,2].

The purpose of our ongoing research in schizophrenia is ultimately to develop an educational program for early detection and intervention of the disease, and the program naturally must include provisions to counteract stigma. Several sophisticated educational programs to counteract stigma have already been developed to provide basic information on schizophrenia and its associated behaviors to various populations under appropriate conditions [7-9], but there have been no studies of the effectiveness of such programs among the parents of adolescents in Japan. The attitudes of parents

* Correspondence: yoshii@clg.niigata-u.ac.jp

† Contributed equally

¹Department of Medical Informatics and Statistics, Niigata University Graduate School of Medicine, 1-754 Asahimachi-dori, Chuo-ku, Niigata, Japan 951-8520

Full list of author information is available at the end of the article

are important to address as first schizophrenic episodes typically occur in adolescence. Our recent study suggested that such parents can learn basic information about schizophrenia and better discriminate schizophrenia symptoms by completing a quick (13-minute) web-based educational program [10]. Therefore, in this study, targeting the issue of stigma specifically, we evaluated stigma toward schizophrenia among a large sample of parents of adolescents and examined the association between stigma toward schizophrenia and several demographic characteristics of the respondents. We tested the hypothesis that contact and communication with a person with schizophrenia would be important to reducing stigma [11]. Through investigation of this hypothesis, we aimed to determine which factors, including that of contact and communication with a person with schizophrenia, influence stigma toward schizophrenia among parents of junior and senior high school students in Japan.

Methods

Setting and participants

The participants were 2,690 Japanese parents of junior and senior high school students (1,410 and 1,280, respectively) who were extracted from 1,370,000 candidates included in a database administered by a Japanese private company specializing in questionnaire research. Consent from the participants was obtained by the research company and the survey was conducted using an Internet website developed by the company. Stratified random sampling was adopted to prevent bias in the gathered sample, and sex and region were used as the stratifying variables, as previously described [10]. The study sample of 2,690 parents mostly comprised those aged 40-50 years (1,904; 70.8%). The study was approved by the Ethics Committee of the Niigata University School of Medicine.

Questionnaire

The participants completed a questionnaire that requested information on sociodemographic data, including age, occupation, and education. In addition, to survey stigma, the study questionnaire contained the Link's Devaluation-Discrimination Measure [12] (modified for schizophrenia), which uses a 4-point Likert scale (4, strongly agree; 3, tend to agree; 2, tend to disagree; and 1, strongly disagree). Items 1, 2, 3, 4, 8, and 10 were reverse scored. A higher score on the Devaluation-Discrimination Measure indicates greater stigma.

Statistical analysis

All analyses were performed using the Statistical Package for Social Science (SPSS) version 16.0. A *p* value less than 0.05 was considered to indicate statistical

significance, and all statistical tests were two-tailed. One-way analysis of variance and Student's *t*-test were used to examine associations between stigma toward schizophrenia and the demographic characteristics of the respondents. Multiple comparisons were performed with post hoc Bonferroni tests for continuous variables [13]. Cross-tabulation was used to examine associations between occupation and both family income and education.

Results

Participant characteristics

Regarding the highest level of educational attainment, 1,063 (39.5%) participants had completed college, 766 (28.5%) had completed high school, and 385 (14.3%) had completed junior high school. Eighty-eight (3.3%) participants reported contact with a close friend or family member who has schizophrenia and 222 (8.3%) reported participation in welfare activities for people with mental illness. The association between occupation and family income is shown in Table 1, and that between occupation and education is shown in Table 2. These findings are discussed in our previous article [10].

Link Devaluation-Discrimination Measure

The mean \pm SD Devaluation-Discrimination Measure score was 32.74 ± 5.66 , from a possible total score ranging from 12 to 48.

The demographic characteristics significantly associated with the Devaluation-Discrimination Measure were occupation, presence of a neighbor with schizophrenia, family income, and participation in welfare activities for people with mental illness (Student's or Welch's *t*-test and one-way analysis of variance) (Table 3). Specifically, regarding occupation, participants working in the agriculture, forestry, and fisheries sector had the highest score (mean \pm SD: 34.64 ± 3.72), and participants earning less than the equivalent of US\$11,000 had the highest score (mean \pm SD: 33.71 ± 4.92). The results of Bonferroni's correction revealed a significant difference between those participants categorized under professional and those categorized under production labor services and other (both *P* < 0.05). The mean score of respondents who reported having a neighbor with schizophrenia was higher than that of individuals who did not. In addition, the mean score of respondents who had taken part in welfare activities for people with mental illness was higher than that of individuals who had not (Table 3).

Discussion

Among parents of junior and senior high school students in Japan, the mean score for the Devaluation-

Table 3 Distribution of scores on the Devaluation-Discrimination Measure by demographic characteristics of the parents of adolescents

	Devaluation-Discrimination Measure	
	Mean ± SD	p*
Age(years)		P = 0.635
30-39	32.47 ± 4.29	
40-49	32.80 ± 4.43	
50-59	32.62 ± 4.55	
60-69	32.47 ± 4.42	
Occupation		p = 0.001
Agriculture, forestry, and fisheries	34.64 ± 3.72	
Production labor services	32.22 ± 4.27 ^a	
Transportation and communications	32.48 ± 4.43	
Sales and marketing	32.92 ± 4.42	
Service industry	32.74 ± 4.13	
Professional	33.63 ± 4.86 ^b	
Other	31.78 ± 3.93	
Family income, (US dollars)		p = 0.042
< 11,000	33.71 ± 4.92	
11,000-32,000	32.68 ± 4.71	
32,000-53,000	32.44 ± 4.26	
53,000-110,000	32.66 ± 4.36	
> 110,000	33.20 ± 4.70	
Proximity to person with schizophrenia		p = 0.004
Yes	34.39 ± 5.40	
No	32.68 ± 4.40	
Participation in welfare activities for people with mental illness		p = 0.010
Yes	33.47 ± 4.79	
No	32.67 ± 4.41	

* Student's or Welch's t-test, one-way analysis of variance.

a P < 0.05, production labor services versus professional

b P < 0.05, professional versus other

menacing, or simply exaggerated or incomprehensible, may produce aversion and fear. The findings of these studies are likely due to the nature of participant contact with schizophrenics. A study in a medical school in Turkey compared attitude changes in students participating in a 3-week psychiatric training program with those participating in an ophthalmology shift and reported no significant difference. The authors suggested that contact during job training might not affect attitudes because of the characteristics of the interaction [21]. Thus, the nature of contact with people with schizophrenia is likely to determine the extent of any stigma toward them. In addition, the disease view engenders a less favorable estimation of the mentally disordered [22]. Therefore, labeling behaviors 'schizophrenic' has a more pessimistic outlook [23]. In light of this, do any provisions created to counteract stigma require a sufficient base knowledge of schizophrenia? A Hong Kong study

found that as 'knowledge' based on the mental illness perspective increased, attitudes became more negative [24]. Therefore, unless a sufficient base knowledge is provided to participants in welfare activities for people with mental illness in the form of an educational program, they may become more negative in their attitude as higher knowledge.

Conclusions

In summary, stigma toward schizophrenia was stronger among members of the general public who had had contact with people with the disorder, as well as those with lower family income. These results contribute to our ultimate research aim of developing an educational program that can combat stigma toward the disease in Japan.

Acknowledgements

The authors thank the study participants. This work was supported by a 2010 Grant-in-Aid for Scientific Research (C) from the Japan Society for the Promotion of Science (22592581), entitled "Investigations of the Consciousness of Parents of Junior and Senior High School Students Concerning Schizophrenia and the Development of Educational Enlightenment Media".

Author details

¹Department of Medical Informatics and Statistics, Niigata University Graduate School of Medicine, 1-754 Asahimachi-dori, Chuo-ku, Niigata, Japan 951-8520. ²Department of Psychiatry, Niigata University Graduate School of Medical and Dental Sciences, 1-754 Asahimachi-dori, Chuo-ku, Niigata, Japan 951-8520. ³Health Administration Center, Headquarters for Health Administration, Niigata University, 8050 Ikarashi-nincho, Nishi-ku, Niigata, Japan 950-2181.

Authors' contributions

HY and KA designed the study and drafted the manuscript. YW, HK, and ZN made significant contributions to the content of the paper and were responsible for the final editorial revision. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 13 July 2011 Accepted: 22 December 2011

Published: 22 December 2011

References

1. Esterberg ML, Compton MT, McGee R, Shim R, Hochman K: Knowledge about schizophrenia and social distance toward individuals with schizophrenia: A survey among predominantly low-income, urban, African American community members. *J Psychiatr Pract* 2008, **14**:86-93.
2. Lysaker PH, Davis LW, Warman DM, Strasburger Amy, Beattie N: Stigma, social function and symptoms in schizophrenia and schizoaffective disorder: Associations across 6 months. *Psychiatry Res* 2007, **149**:89-95.
3. Chong SA, Lee C, Bird L, Verma S: A risk reduction approach for schizophrenia: The early psychosis intervention programme. *Ann Acad Med Singapore* 2004, **33**:630-5.
4. De Koning MB, N Bloemen OJ, van Amelsvoort TAMJ, Becker HE, Nieman DH, van der Gaag Mvan, Linszen DH: Early intervention in patients at ultra high risk of psychosis: benefits and risks. *Acta psychiatrica scandinavica* 2009, **119**:426-42.
5. Berge M, Ranney M: Self-esteem and stigma among persons with schizophrenia: Implications for mental health. *Care Manag J* 2005, **6**:139-144.

6. Corrigan PW, Green A, Lundin R, Kubiak MA, Penn DL: **Familiarity with and social distance from people who have serious mental illness.** *Psychiatr Serv* 2001, **52**:953-8.
7. Stuart H: **Reaching Out to High School Youth: The effectiveness of a video-based antistigma program.** *Can J Psychiatry* 2006, **51**:647-53.
8. Kadri N, Sartorius N: **The global fight against the stigma of schizophrenia.** *PLoS Med* 2005, **2**:0597-9.
9. Gaebel W, Zaske H, Baumann AE, Klosterkötter J, Maier W, Decker P, Möller HJ: **Evaluation of the German WPA "Program against stigma and discrimination because of schizophrenia-Open the doors": Results from representative telephone surveys before and after three years of antistigma interventions.** *Schizophr Res* 2008, **98**:184-93.
10. Yoshii H, Watanabe Y, Kitamura H, Chen J, Akazawa K: **Effect of an education program on improving knowledge of schizophrenia among parents of junior and senior high school students in Japan.** *BMC Public Health* 2011, **11**(1):323.
11. Tanaka G, Inadomi H, Kikuchi Y, Ohta Y: **Evaluating stigma against mental disorder and related factors.** *Psychiatry Clin Neurosci* 2004, **58**:558-66.
12. Link BG: **Understanding labelling effects in the area of mental disorders: an assessment of the effects of expectations of rejection.** *Am Sociol Rev* 1987, **52**:96-112.
13. Glantz SA: **Primer of biostatistics sixth edition.** New York: The McGraw-Hill Companies; 2005.
14. Mann CE: **Factors associated with stigmatization of persons with mental illness.** *Psychiatric Services* 2004, **55**:185-187.
15. Papadopoulos C, Leavey G, Vincent C: **Factors influencing stigma: A comparison of Greek-cypriot and English attitudes towards mental illness in north London.** *Soc Psychiatry Psychiatr Epidemiol* 2002, **37**:430-434.
16. Ministry of Education, Culture, Sports, Science and Technology. [<http://www.nta.go.jp/kohyo/tokei/kokuzeicho/minkan2009/pdf/001.pdf>].
17. Ministry of Education, Culture, Sports, Science and Technology. [http://www.mext.go.jp/b_menu/hakusho/html/hpad197501/hpad197501_3_106.html].
18. Dickerson FB, Sommerville J, Origoni AE, Ringel NB, Parente F: **Experiences of stigma among outpatients with schizophrenia.** *Schizophrenia Bulletin* 2002, **28**:143-155.
19. Shimotsu S, Sakamoto S, Horikawa N, Sakano Y: **Reliability and validity of the Japanese version of Link's Devaluation-Discrimination Scale.** *Seishinka Chiryogaku* 2006, **21**:521-528.
20. Penn DL, Kohlmaier JR, Corrigan PW: **Interpersonal factors contributing to the stigma of schizophrenia: social skills, perceived attractiveness, and symptoms.** *Schizophr Res* 2000, **45**:37-45.
21. Arkar H, Eker D: **Influence of a 3-week psychiatric training programme on attitudes toward mental illness in medical students.** *Soc Psychiatry Psychiatr Epidemiol* 1997, **32**:171-6.
22. Mehta S, Farina A: **Is being 'sick' really better? Effect of the disease view of mental disorder on stigma.** *J Soc Clin Psychol* 1997, **16**:405-419.
23. Cormack S, Furnham A: **Psychiatric labelling, sex role stereotypes and beliefs about the mentally ill.** *Int J Soc Psychiatry* 1998, **44**:235-247.
24. Chou K, Mak K: **Attitudes to mental patients among Hong Kong Chinese: a trend study over two years.** *Int J Soc Psychiatry* 1998, **44**:215-224.

doi:10.1186/1756-0500-4-558

Cite this article as: Yoshii et al.: Stigma toward schizophrenia among parents of junior and senior high school students in Japan. *BMC Research Notes* 2011 **4**:558.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

