

ERRATUM

African 2, a Clonal Complex of *Mycobacterium bovis* Epidemiologically Important in East Africa

Stefan Berg, M. Carmen Garcia-Pelayo, Borna Müller, Elena Hailu, Benon Asiimwe, Kristin Kremer, James Dale, M. Beatrice Boniotti, Sabrina Rodriguez, Markus Hilty, Leen Rigouts, Rebuma Firdessa, Adelina Machado, Custodia Mucavele, Bongo Nare Richard Ngandolo, Judith Bruchfeld, Laura Boschirolí, Annélle Müller, Naima Sahraoui, Maria Pacciarini, Simeon Cadmus, Moses Joloba, Dick van Soelingen, Anita L. Michel, Berit Djønne, Alicia Aranaz, Jakob Zinsstag, Paul van Helden, Françoise Portaels, Rudovick Kazwala, Gunilla Källenius, R. Glyn Hewinson, Abraham Aseffa, Stephen V. Gordon, and Noel H. Smith

VLA Weybridge, New Haw, Surrey KT15 3NB, United Kingdom; Division of Molecular Biology and Human Genetics, Faculty of Health Sciences, Stellenbosch University, P.O. Box 19063, Tygerberg 7505, South Africa; Armauer Hansen Research Institute, P.O. Box 1005, Addis Ababa, Ethiopia; Department of Medical Microbiology, Makerere University Medical School, P.O. Box 7072, Kampala, Uganda; Tuberculosis Reference Laboratory, National Institute for Public Health and the Environment (RIVM), Centre for Infectious Disease Control (CIB/LIS), P.O. Box 1, 3720 BA Bilthoven, Netherlands; Reparto Genomica, Istituto Zooprofilattico Sperimentale della Lombardia e dell'Emilia, Via Bianchi n. 9, 25124 Brescia, Italy; Dept. de Sanidad Animal, Facultad de Veterinaria, and Centro Vigilancia Sanitaria Veterinaria (MISAVET), Universidad Complutense, Avenida, Puerta de Hierro s/n, 28040 Madrid, Spain; Institute for Infectious Diseases, University of Bern, Friedbühlstrasse 51, CH-3010 Bern, Switzerland; Agence Française de Sécurité Sanitaire des Aliments, 23 Avenue du Général-de-Gaulle, 94706 Maisons-Alfort Cedex, France; Unit of Infectious Diseases, Department of Medicine, Solna, Karolinska Institutet, Karolinska University Hospital, 17177 Stockholm, Sweden; Department of Veterinary Public Health & Preventive Medicine, University of Ibadan, Ibadan, Nigeria; Laboratoire de Recherches Vétérinaires et Zootechniques de Farcha, BP 433, N'Djaména, Chad; Sokoine University of Agriculture, Morogoro, Tanzania; Université Saad Dahlab, Route de Soumaa, BP 270, Blida, Algeria; Department of Animal Health, National Veterinary Institute, BP 750 Sentrum, N-0106 Oslo, Norway; Faculdade de Veterinaria, Universidade Eduardo Mondlane, CP 257 Maputo, Mozambique; Department of Microbiology, Institute of Tropical Medicine, Nationalestraat 155, B-2000 Antwerp, Belgium; Faculty of Veterinary Science, University of Pretoria, Private Bag X04, and ARC-Onderstepoort Veterinary Institute, Private Bag X05, Onderstepoort 0110, South Africa; Department of Clinical Science and Education, Karolinska Institutet, Södersjukhuset, 11883 Stockholm, Sweden; Swiss Tropical and Public Health Institute, Socinstrasse 57, 4002 Basel, Switzerland; College of Life Sciences and UCD Conway Institute, University College Dublin, Dublin 4, Ireland; and VLA Weybridge, New Haw, Surrey KT15 3NB, United Kingdom, and Centre for the Study of Evolution, University of Sussex, Brighton BN1 9QL, United Kingdom

Volume 193, no. 3, p. 670–678, 2011. Page 671, column 2: Lines 7 through 14 should read as follows: “The status of the RDAf2 region (deleted or intact) was assessed by multiplex PCR with a set of three primers (primer set Af2): two primers targeting the flanking regions of RDAf2 (RDAf2_Fw, 5'-ACCGCCCTGTCCTATGTGAG, and RDAf2_Rev, 5'-TGACGGTTGCCTTCTTGAC) and one primer hybridizing with the internal region of RDAf2 (RDAf2_IntRev, 5'-CACTGTCTCCGCTCATCATG). A PCR product of 451 bp (RDAf2 intact) or 711 bp (RDAf2 deleted) was identified by agarose gel electrophoresis.”