



Published in final edited form as:

Mol Cancer Ther. 2012 March ; 11(3): 517–525. doi:10.1158/1535-7163.MCT-11-0454.

Mesothelin targeted agents in clinical trials and in preclinical development

Ronan J. Kelly¹, Elad Sharon¹, Ira Pastan², and Raffit Hassan²

¹Medical Oncology Branch, Center for Cancer Research, National Cancer Institute, National Institutes of Health, Bethesda, MD

²Laboratory of Molecular Biology, Center for Cancer Research, National Cancer Institute, National Institutes of Health, Bethesda, MD

Abstract

Mesothelin is a tumor differentiation antigen that is highly expressed in several human malignancies including malignant mesothelioma, pancreatic, ovarian and lung adenocarcinomas. The limited expression of mesothelin on normal human tissues and its high expression in many common cancers make it an attractive candidate for cancer therapy. Several agents including an immunotoxin, mAb, antibody drug conjugate, and tumor vaccine are in various stages of development to treat patients with mesothelin expressing tumors. This review article highlights ongoing clinical trials as well as other approaches to exploit mesothelin for cancer therapy that are in preclinical development.

Keywords

Mesothelin; Immunotoxin; SS1P; MORAb-009; CA-125

Introduction

Mesothelin is a 40-kDa cell surface glycoprotein that is present on normal mesothelial cells lining the pleura, peritoneum and pericardium (1). Mesothelin was originally identified as the antigen recognized by the mAb K1 that was produced by immunization of mice with the human ovarian cancer cell line OVCAR3 (2). The mesothelin gene encodes a precursor protein of 71-kDa that is processed to yield a 31-kDa shed protein named megakaryocyte-potentiating factor (MPF) and the 40-kDa cell bound fragment mesothelin (3). MPF was isolated from the culture supernatant of a pancreatic cancer cell line and in mouse bone marrow cultures demonstrated megakaryocyte colony-forming activity in the presence of interleukin-3 (4). By immunohistochemistry using anti-mesothelin mAbs, mesothelin expression in normal human tissues is noted only in the single layer of mesothelial cells that line the pleura, peritoneum and pericardium, surface epithelial cells of the ovary, tunica vaginalis, rete testis and the tonsillar and fallopian tube epithelial cells (2, 5). However, mesothelin is highly expressed in several human tumors, including epithelial mesotheliomas (~100% of cases) and lung (~50% of cases), ovarian (~70% of cases) and pancreatic/biliary adenocarcinomas (~100% of cases) (5–8). In the majority of these cancers there is diffuse, homogenous cell surface mesothelin expression. However, the pattern of mesothelin

Corresponding Author: Dr. Raffit Hassan, Laboratory of Molecular Biology, 37 Convent Dr., Rm. 5116, National Cancer Institute, Bethesda, MD 20892-4264 USA, Tel: (301) 451-8742; Fax: (301) 402-1344; hassanr@mail.nih.gov.

The authors declare no potential conflicts of interest

reactivity in lung cancer is different from the other tumors in that these tumors show mostly cytoplasmic staining rather than membranous staining (9). Mesothelin is also expressed in many other cancers including gastric cancer, biphasic synovial sarcoma and uterine adenocarcinoma (5). The frequency and characterization of mesothelin expression in different tumor types has been described in prior reviews (3, 10).

The normal biological function of mesothelin is unknown. Mutant mice that lack both copies of the mesothelin gene had no detectable phenotype and both male and female mice produced healthy offspring suggesting that mesothelin is not involved in normal growth and development (11). Mesothelin may play a role in cell adhesion and preclinical studies have demonstrated that mesothelin is the receptor for the tumor antigen CA-125 (also known as MUC16) and that they bind to each other with a high specificity (12, 13). Tumors expressing CA-125 may bind to mesothelin expressed on the surface of mesothelial cells that line the pleural or peritoneal cavity leading to an increase in heterotypic cell adhesion and promoting metastatic spread (12, 13). Studies in pancreatic cancer suggest a role for mesothelin in tumorigenesis by increasing cellular proliferation, migration and S-phase cell populations (14). The exact biological role of MPF is also unclear. Wang et al. have recently demonstrated that in cells over-expressing MPF, phosphorylation of MAP kinase, ERK1/2 was enhanced and the rate of cell death was decreased, leading to an increase in cell numbers (15). Although, the function of mesothelin in cancer is not fully understood at this time its limited expression on normal human tissues and high expression in many cancers makes it an attractive tumor associated antigen for cancer therapy.

Clinical Trials of Mesothelin Targeted Agents

Several different therapeutic agents are being evaluated for treatment of patients with mesothelin expressing cancers (Table 1) (16–25). These include antibody based therapies with SS1(dsFv)-PE38 (SS1P) a recombinant immunotoxin that targets mesothelin, a high affinity chimeric (mouse/human) mAb [IgG/κ] (MORAb-009) and an anti-mesothelin antibody drug conjugate (BAY-94 9343). In addition, a mesothelin tumor vaccine (CRS-207) and adoptive T cell immunotherapy using mesothelin specific chimeric antigen receptors are also being evaluated in clinical trials.

Anti-mesothelin Immunotoxin, SS1P

The immunotoxin SS1P [SS1(dsFv)PE38] consists of an anti-mesothelin Fv, which was obtained from a phage display library of mice immunized with recombinant mesothelin, genetically fused to a truncated form of the *Pseudomonas* exotoxin, PE38. After binding to cell surface mesothelin via the Fv, PE38 is internalized into the cell, undergoes processing and kills the cell due to inhibition of protein synthesis by ADP ribosylation and inactivation of elongation factor 2 (26). SS1P, which has high affinity for mesothelin (Kd, 0.72M), is highly active against several mesothelin expressing cell lines and causes regression of mesothelin expressing xenografts in nude mice (27, 28). SS1P also demonstrated cytotoxicity against tumor cells directly obtained from patients with ovarian cancer and mesothelioma (29, 30).

Preclinical studies have shown marked anti-tumor synergy when SS1P is combined with several commonly used chemotherapeutic agents such as paclitaxel, gemcitabine or cisplatin (31, 32). Although, the combination of SS1P with chemotherapeutic agents did not result in synergy in cell culture there was a marked increase in anti-tumor activity with the combination in tumor xenograft models. Using a mesothelin expressing cell line, A431-K5, grown as a tumor xenograft in athymic nude mice, treatment with SS1P plus paclitaxel resulted in increased anti-tumor activity with durable complete tumor regressions compared to treatment with paclitaxel or SS1P alone (31). This effect of paclitaxel to enhance the

cytotoxicity of SS1P in *in vivo* studies was not due to an indirect effect of paclitaxel by increasing tumor permeability due to damage of tumor endothelial cells. Rather, it is due to the direct effect of paclitaxel on tumor cells. The mechanism for this synergy was evaluated using a pair of mesothelin expressing cervical cancer cell lines, KB, that are sensitive or resistant to paclitaxel as tumor xenografts in mice. The synergistic effect of paclitaxel and SS1P was only seen in paclitaxel sensitive KB tumor xenografts (33). Killing of tumor cells by paclitaxel altered the tumor architecture and significantly decreased the concentration of shed mesothelin in the tumor extracellular fluid. These results suggest that the synergy between SS1P and chemotherapy is due to the ability of cytotoxics to decrease the shed mesothelin in the tumor extracellular space, which allows more of the administered SS1P to bind to tumor cells and results in increased cell killing. In addition, paclitaxel alters tumor architecture by killing tumor cells allowing increased tumor penetration. Based on these preclinical studies the combination of SS1P with chemotherapy is being evaluated in clinical trials involving mesothelin expressing cancers.

A limitation of immunotoxin-based therapies, such as SS1P, is the development of neutralizing antibodies to the toxin portion of the molecule, which limits repeated administration of the drug to patients. Previous efforts to decrease the immune response to the immunotoxins using different approaches, such as the use of the anti-B-cell mAb rituximab, have been unsuccessful (34). However, we have recently shown that immune depletion using the regimen of pentostatin plus cyclophosphamide completely abrogates the anti-immunotoxin immune response in immunocompetent BALB/c mice when repeat injections of SS1P were administered. This regimen resulted in host B-cell and T-cell depletion with minimal myeloid cell depletion (35). A pilot clinical trial to evaluate this approach to decrease the immunogenicity of SS1P in patients has just opened to patient accrual (36).

Phase I clinical trials of single agent SS1P

There have been two Phase I trials of SS1P reported utilizing different schedules of administration, either as a bolus *i.v.* infusion or as a continuous infusion. In a Phase I dose escalation study, 34 patients with advanced mesothelin expressing cancers, including 20 with mesothelioma, 12 with ovarian cancer and 2 with pancreatic cancer who had failed standard therapy, were treated with SS1P given as a 30 minute *i.v.* infusion every other day for either 3 or 6 doses (16). The first cohort of 17 patients received 6 doses of SS1P every other day for 6 doses with a maximum tolerated dose (MTD) of 18 $\mu\text{g}/\text{kg}/\text{dose}$. The dose limiting toxicities (DLT) were grade 3 urticaria (1 patient) and grade 3 vascular leak syndrome (2 patients). A second cohort of 17 patients received only 3 doses with a MTD of 45 $\mu\text{g}/\text{kg}/\text{dose}$ with grade 3 pleuritis being the DLT. At the MTD of 45 $\mu\text{g}/\text{kg}/\text{dose}$, the mean C_{max} of SS1P was 483 ng/ml and the half-life was 466 minutes. Thirty-three patients were considered evaluable for response and 4 patients had minor responses, 19 had disease stabilization and 10 had progressive disease.

In the second Phase I trial SS1P was administered as a continuous *i.v.* infusion over 10 days (17). Twenty-four patients that included 16 with mesothelioma, 7 with ovarian cancer and 1 with pancreatic cancer were enrolled at five different dose levels of SS1P. The MTD was 25 $\mu\text{g}/\text{kg}/\text{d}$ on a 10 day continuous infusion and the DLT was reversible vascular leak syndrome. As a single agent SS1P has demonstrated modest clinical activity and continuous infusion shows no significant benefit over bolus dosing. Due to the ease of administration, as well as pharmacokinetic data that show high blood levels and prolonged half-life of SS1P when given as a bolus dose, this schedule is being evaluated in combination studies with chemotherapy.

Phase I clinical trials of SS1P in combination with chemotherapy

Based on preclinical studies that showed that the activity of SS1P could be increased when it is given in combination with chemotherapy, an ongoing clinical trial is evaluating SS1P in combination with pemetrexed and cisplatin as frontline therapy for patients with advanced, unresectable malignant pleural mesothelioma. The primary objective of this study is to evaluate the MTD and safety of SS1P in combination with chemotherapy with secondary endpoints of tumor response, progression free survival (PFS) and overall survival (OS). Enrolled patients receive six cycles of standard care chemotherapy consisting of pemetrexed 500 mg/m² and cisplatin 75 mg/m² on day 1. During the first two cycles, patients receive SS1P on days 1, 3, and 5 at escalating dose cohorts of 25 mcg/kg, 35 mcg/kg, 45 mcg/kg, and 55 mcg/kg. Results from the first 19 patients have recently been reported. Five patients were treated at the SS1P dose of 25 mcg/kg, 3 patients at 35 mcg/kg, 10 patients at 45 mcg/kg and 1 patient at the 55 mcg/kg. Of the 14 evaluable patients treated at all dose levels, 7 had a partial response, 3 had stable disease and 4 had progressive disease (18). However, of the 7 evaluable patients treated at the MTD of SS1P in combination with pemetrexed and cisplatin 5 had partial response, 1 had stable disease and 1 had progressive disease. Of the 10 patients whose serum mesothelin levels were evaluated before and at completion of treatment all 5 patients with partial response had a significant decrease in serum mesothelin (63–83%). Two of 3 patients with progressive disease had increased mesothelin levels (16%–34%), while 1 patient had a slight decrease (7%). Two patients with stable disease had discordant responses (17% increase and 60% decrease). Overall the combination of SS1P, cisplatin and pemetrexed was well tolerated with hypoalbuminemia, edema and fatigue being the main side-effects. The trial is now in the expansion phase.

A Phase I trial evaluating SS1P in combination with bevacizumab, carboplatin and paclitaxel for treatment of patients with mesothelin expressing stage IV lung adenocarcinoma was recently closed because of poor accrual. The main reason for closing the trial was that the incidence of mesothelin expression in the patients screened for this study was lower than the expected 50% positivity in lung adenocarcinoma. In addition, similar to the findings in preclinical immunohistochemistry studies the predominant pattern of tumor mesothelin staining was cytoplasmic rather than membranous reactivity (Hassan R, unpublished data, 9).

MORAb-009, a Chimeric Anti-mesothelin mAb

MORAb-009 consists of the heavy and light chain variable regions of a mouse anti-mesothelin single chain Fv grafted to human IgG1 and κ constant regions. MORAb-009 has high affinity for mesothelin with an affinity (K_D) of 1.5 nM. In vitro MORAb-009 inhibits the adhesion between cell lines expressing mesothelin and MUC16, as well as mediates ADCC against mesothelin positive ovarian cancer, pancreatic cancer and mesothelioma cell lines. In a tumor xenograft model using the mesothelin positive cell line A431-K5, treatment with MORAb-009 in combination with gemcitabine or paclitaxel resulted in significant tumor shrinkage compared to MORAb-009 or chemotherapy alone (37). In cynomolgus monkey toxicology studies repeated doses of MORAb-009 of 15 mg/kg were well tolerated. Based on these preclinical data MORAb-009 was evaluated in clinical trials for patients whose tumors were mesothelin positive.

Phase I dose-escalation study of MORAb-009

A Phase I clinical trial of MORAb-009 was performed in patients with mesothelin positive cancers who had failed standard treatment options for their disease, to determine the safety and MTD of this antibody. MORAb-009 was administered as an i.v. infusion at doses ranging from 12.5 to 400 mg/m² on days 1, 8, 15 and 22. Patients with stable disease at day

35 could receive the next cycle of MORAb-009. A total of 24 patients were enrolled on this study, including 13 with mesothelioma, 7 with pancreatic cancer and 4 with ovarian cancer (19). The MTD of MORAb-009 was 200 mg/m² with two patients at the 400 mg/m² having DLT (grade 4 transaminitis and a grade 3 serum sickness). No objective response was noted although 11 patients had stable disease.

An interesting observation during the Phase I clinical trial was the effect of MORAb-009 on serum CA-125 kinetics. As previously described, laboratory studies suggest that mesothelin may be involved in tumor spread by binding to CA-125. In this Phase I trial 8 patients with mesothelioma had serum CA-125 levels measured at baseline and at different time points of MORAb-009 therapy (38). In all of these patients, treatment with MORAb-009 led to a marked increase in serum CA-125 levels even in patients who had normal CA-125 levels at baseline. The increase in CA-125 levels was not due to inflammation or disease progression since the CA-125 levels returned to baseline value once MORAb-009 treatment was stopped. These results suggest that MORAb-009 interferes with mesothelin CA-125 interaction in patients and could have potential utility for prevention of tumor metastasis in patients with mesothelioma and ovarian cancer.

Phase II clinical trials of MORAb-009 with chemotherapy

Given the favorable safety profile and the possibility of clinical benefit two Phase II clinical trials were initiated, which focused on evaluating the efficacy of MORAb-009 in combination with chemotherapy. In the first, MORAb-009 was administered with gemcitabine in patients with unresectable pancreatic cancer (20). This was a double-blind, placebo-controlled trial in patients with advanced pancreatic cancer who were not candidates for surgical resection. The patients were randomized to gemcitabine alone or gemcitabine plus MORAb-009. The primary outcome is overall survival, with progression-free survival being a secondary endpoint. This trial has completed patient enrollment but the results have not yet been published.

The second study is evaluating MORAb-009 as frontline therapy in combination with pemetrexed and cisplatin in patients with unresectable epitheloid malignant pleural mesothelioma. Patients receive pemetrexed 500 mg/m² and cisplatin 75 mg/m² on day 1 with MORAb-009 5 mg/kg on days 1 and 8 of a 3-week cycle. Patients with tumor response or stable disease after 6 cycles of therapy continue on MORAb-009 till disease progression (21). The primary endpoint of this study is to evaluate if combination treatment with MORAb-009 plus pemetrexed and cisplatin improves PFS compared to PFS observed in the pivotal Phase III clinical trial of pemetrexed and cisplatin. No preliminary data has yet been reported and the study is ongoing.

CRS-207, a Mesothelin Tumor Vaccine

CRS-207 is a mesothelin vaccine that utilizes a live attenuated strain of the bacterium *Listeria monocytogenes* (*Lm*), a facultative intracellular bacterium, as the vector (39). CRS-207 is based on CRS-100, which is an engineered vector that has deletions of the two genes that encode the virulence determinants, actA and internalin B. These deletions result in a 1000-fold decrease in virulence compared to the wild-type *Lm*. Preclinical studies show that CRS-207 elicits human mesothelin-specific CD4⁺/CD8⁺ immunity in mice and in cynomolgus monkeys and exhibits therapeutic efficacy in tumor bearing mice (39).

Phase I clinical trial of CRS-207

A Phase I clinical trial of CRS-207 for the treatment of patients with mesothelin expressing cancers was conducted to determine the safety and MTD of CRS-207. The results of this trial were recently reported (22). In this Phase I study CRS-207 was administered i.v. every

3 weeks for a total of 4 doses ranging from 1×10^8 to 1×10^{10} cfu. Seventeen patients including 7 pancreatic cancer, 5 mesothelioma, 3 lung cancer and 2 ovarian cancer were treated. The MTD of CRS-207 was 1×10^9 cfu, with a DLT of hypotension observed at 1×10^{10} cfu dose level. In this group of heavily pretreated patients no objective anti-tumor response was observed but a mesothelin-specific T cell response was observed in 5 of 10 patients tested. Based on the tolerability and immune activation, CRS-207 may be an attractive agent to treat mesothelin-expressing cancers either alone or in combination with other agents.

Phase II clinical trial of CRS-207 with GVAX pancreatic vaccine

A phase II clinical trial of CRS-207 for the treatment of patients with metastatic pancreatic cancer has just opened for enrollment (23). In this Phase II study previously treated patients with metastatic pancreatic cancer will be randomized to either GVAX (irradiated pancreatic cancer cell lines that have been genetically-modified to secrete GM-CSF) alone every 3 weeks \times 6 doses or sequential administration of 2 doses of GVAX vaccine and 4 doses of CRS-207. The primary endpoint of this clinical trial is to determine if administration of CRS-207 will improve overall survival compared to GVAX alone. Secondary endpoints include safety of combining CRS-207 with GVAX and to monitor CRS-207 induced Listeria and mesothelin specific immune response.

Adoptive T Cell Immunotherapy Using Mesothelin Specific Chimeric Antigen Receptors

Since tumor associated antigens are poorly immunogenic, generation of a T cell response to these antigens is limited in patients (40). Over the last several years different approaches have been developed to exploit T cells for cancer therapy such as the administration of tumor infiltrating T cells and adoptive transfer of T cells expressing T-cell receptors against tumor antigens, which have resulted in clinical activity in some tumors (41). An attractive approach to increase the specificity of these T cells for the tumor is the use of chimeric antigen receptors (CARs), which can kill these cells in a non-HLA restricted manner. A CAR consists of an antigen-specific portion of a mAb, such as Fv and the signaling component of the T cells usually the ζ chain of the TCR/CD3 complex. T cells expressing the CARs are specifically directed to tumor cells and then the activation of T cells kills the tumor (42). Adoptive transfer of T cells expressing CARs has shown promise in several cancers (43). Mesothelin is an attractive candidate for T cell adoptive therapy using CARs, and preclinical studies have shown its efficacy in animal models. A CAR that recognizes mesothelin was generated using anti-mesothelin single-chain Fv fused to T cell receptor zeta signal transduction domain, as well as CD28 and CD137(4-1BB) domains. T cells transduced with these CARs using lentiviral vectors were then evaluated for efficacy against mesothelin expressing tumor xenografts in NOD/*scid*/*IL2ry*^{-/-} mice. Both intratumoral, as well as i.v. administration of these transduced T cells resulted in significant shrinkage of these large established mesothelin expressing tumors. T cells expressing CARs that contained the CD28 and CD137 domains had greater efficacy than CARs expressing only the T cell receptor zeta signaling domain (44). Recently, the same group has described an alternative method, which may be safer than the retroviral or lentiviral vector transduction method. Using RNA encoding an anti-mesothelin CAR for T cell transduction shows robust activity against mesothelin expressing xenografts (45). Some of the advantages of using RNA CAR T cell therapy are that it avoids the potential risk of malignant transformation from insertional mutagenesis using retroviral or lentiviral constructs. In addition, in case of toxicity due to CAR T cell therapy, it can be reduced by stopping the administration of RNA CAR T cells, unlike in the case of stably transduced T cells using viral vectors. A phase I

clinical trial of adoptive T cell therapy using mesothelin directed CARs has been initiated for treatment of patients with pleural mesothelioma (24).

Antibody Drug Conjugates (ADC) Targeting Mesothelin

The limited expression of mesothelin on essential human tissues makes it a good target for antibody drug conjugates. Although no mesothelin targeted ADC is yet in the clinic, some of these compounds have undergone preclinical development and include MDX-1204 and BAY 94-9343. The anti-mesothelin ADC MDX-1204 consists of the human anti-mesothelin mAb MDX-1382 conjugated to Duocarmycin, a DNA alkylating agent (46). This compound showed anti-tumor efficacy against mesothelin expressing xenografts in mice and at clinically relevant doses was well tolerated by cynomolgus monkeys without any clinical toxicity due to the antibody binding to mesothelin. Another anti-mesothelin ADC is BAY 94-9343, which consists of the fully human anti-mesothelin IgG1 linked to DM4, a potent tubulin-binding drug (47). This drug conjugate demonstrated cytotoxicity against mesothelin positive cell lines with IC₅₀ in the nanomolar range, as well as anti-tumor activity against mesothelin expressing ovarian, pancreatic and mesothelioma tumor xenografts. A Phase I clinical trial of BAY 94-9343 has recently opened for the treatment of patients with advanced solid tumors. The primary objectives of this trial are to determine the safety and maximum tolerated dose of BAY 94-9343, as well as study its pharmacokinetics (25).

Mesothelin Directed Therapies in Preclinical Development

Some of the different approaches to target mesothelin for cancer therapy that are in preclinical development are discussed below.

Mesothelin Cancer Vaccines

Several lines of work support the utility of mesothelin vaccination for cancer therapy. In patients with advanced mesothelin expressing mesothelioma and ovarian cancer anti-mesothelin IgG antibodies were present in 39% and 41% of patients, respectively (48). This suggests that at least in some patients there is a breakdown of tolerance to mesothelin. In addition, in a clinical trial of patients with advanced pancreatic cancer receiving vaccination with granulocyte macrophage colony-stimulating-secreting pancreatic cancer cell lines (CG8020/CG2505) a strong mesothelin specific CD8⁺ T cell response was observed in 3 of 14 patients who developed a delayed-type hypersensitivity response following vaccination (49). In a subsequent clinical trial of this whole cell vaccine for metastatic pancreatic cancer the investigators observed mesothelin-specific CD8⁺ T cells in most of the patients at baseline but this response was augmented after vaccination (50).

Several preclinical studies have evaluated the utility of mesothelin as a tumor vaccine using different strategies. The study by Yokokawa et al. identified two novel HLA-2 mesothelin epitopes recognized by cytotoxic T lymphocyte. T cells generated from these native or agonist mesothelin epitopes lysed mesothelin expressing pancreatic cancer, ovarian cancer cell and mesothelioma cell lines (51). In addition, studies using a DNA vaccine encoding human mesothelin showed efficacy in a mouse model by generating both CD4⁺ and CD8⁺ T cell as well antibody mediated immune response to mesothelin (52). Other vaccine strategies such as targeting mesothelin directly to dendritic cells using mAbs or using dendritic cells transduced with full-length mesothelin have also shown enhanced induction of mesothelin specific CD4⁺ and CD8⁺ T cell immunity (53, 54). In a study that used vaccination with chimeric virus-like particles that contain human mesothelin substantially inhibited tumor progression in mice, with increases in mesothelin-specific antibodies and cytotoxic T-lymphocyte activity (14). It is likely that some of these vaccination strategies will be evaluated in clinical trials in the near future.

Mesothelin-directed Gene Therapy

Given the potential toxicity to non-target tissues, such as the liver using systemic gene therapy, the use of gene therapy that specifically targets the tumor cells is clinically desirable. Breidenbach et al. conducted studies to evaluate mesothelin as a target for adenoviral vectors for gene therapy of ovarian cancer. Using established ovarian cancer cell lines and purified tumor cells obtained from patients, they demonstrated high mesothelin gene and protein expression (55). Adenoviruses that contained the mesothelin promoter driving reporter gene expression showed that the mesothelin promoter was activated in ovarian cancer cell lines but not in normal control cells. In addition, in an in vivo study in mice, the adenovirus construct containing the mesothelin promoter had a low expression in the liver making it potentially useful for the clinic. To further improve on targeting of adenovirus for gene therapy of ovarian cancer an adenovirus vector containing an Fc-binding domain was conjugated to a mouse anti-human mesothelin mAb. This transductional targeting approach resulted in increased transgene expression in ovarian cancer cells and showed the utility of mesothelin targeting for ovarian cancer gene therapy.

The use of conditionally replicative adenoviruses (CRAd) that contain tumor-specific promoters restricts virus replication to cancer cells. A mesothelin promoter based CRAd with a chimeric Ad5/3 fiber (AdMSLNCRA5/3) was utilized to test the efficacy against a human ovarian cancer xenograft model in immunodeficient mice (56). AdMSLNCRA5/3 treatment resulted in significant tumor growth inhibition and improved overall survival of these mice compared to no virus administration or wild-type adenovirus administration (56). Gene therapy clearly shows some promise in preclinical models. Clinical testing will ultimately determine the efficacy of mesothelin-directed gene therapy and future studies are planned.

Conclusion

Mesothelin is a tumor differentiation antigen with limited expression on normal human tissues that is highly expressed in several malignancies making it a good candidate for cancer therapy. Thus far clinical trials of an anti-mesothelin immunotoxin and mAb, as well as a mesothelin vaccine, have shown encouraging activity without any untoward toxicity due to targeting of normal mesothelial cells that express mesothelin. In addition to the mesothelin targeted therapies (SS1P, MORAb-009, BAY 94-9343, CRS-207 and adoptive T cell immunotherapy) currently in the clinic, there are several other agents in various stages of preclinical development. Although, preclinical studies and clinical trials conducted thus far have validated mesothelin as a drug target, the ongoing clinical trials-some of which have just been initiated, will determine whether mesothelin targeted therapies benefit patients.

Abbreviations

ADC	antibody drug conjugates
CARs	chimeric antigen receptors
CRAd	conditionally replicative adenoviruses
CRS-207	a mesothelin vaccine
DLT	dose limiting toxicities
<i>Lm</i>	<i>Listeria monocytogenes</i>
MORAb-009	a high affinity chimeric (mouse/human) mAb (IgG/κ)

MPF	megakaryocyte-potentiating factor
MTD	maximum tolerated dose
OS	overall survival
PE38	<i>Pseudomonas</i> exotoxin
PFS	progression free survival
SS1P	(SS1 (dsFv)-PE38) is a high affinity recombinant immunotoxin that targets mesothelin

Acknowledgments

Financial Support: This research was supported by the Intramural Research Program of the NIH, National Cancer Institute, Center for Cancer Research

REFERENCES

1. Chang K, Pastan I. Molecular cloning of mesothelin, a differentiation antigen present on mesothelium, mesotheliomas, and ovarian cancers. *Proc Natl Acad Sci U S A*. 1996; 93:136–140. [PubMed: 8552591]
2. Chang K, Pastan I, Willingham MC. Isolation and characterization of a monoclonal antibody, K1, reactive with ovarian cancers and normal mesothelium. *Int J Cancer*. 1992; 50:373–381. [PubMed: 1735605]
3. Hassan R, Bera T, Pastan I. Mesothelin: a new target for immunotherapy. *Clin Cancer Res*. 2004; 10:3937–3942. [PubMed: 15217923]
4. Yamaguchi N, Hattori K, Oh-eda M, Kojima T, Imai N, Ochi N. A novel cytokine exhibiting megakaryocyte potentiating activity from a human pancreatic tumor cell line HPC-Y5. *J Biol Chem*. 1994; 269:805–808. [PubMed: 8288629]
5. Ordonez NG. Application of mesothelin immunostaining in tumor diagnosis. *Am J Surg Pathol*. 2003; 27:1418–1428. [PubMed: 14576474]
6. Ordonez NG. Value of mesothelin immunostaining in the diagnosis of mesothelioma. *Mod Pathol*. 2003; 16:192–197. [PubMed: 12640097]
7. Argani P, Iacobuzio-Donahue C, Ryu B, Rosty C, Goggins M, Wilentz RE, et al. Mesothelin is overexpressed in the vast majority of ductal adenocarcinomas of the pancreas: identification of a new pancreatic cancer marker by serial analysis of gene expression (SAGE). *Clin Cancer Res*. 2001; 7:3862–3868. [PubMed: 11751476]
8. Hassan R, Kreitman RJ, Pastan I, Willingham MC. Localization of mesothelin in epithelial ovarian cancer. *Appl Immunohistochem Mol Morphol*. 2005; 13:243–247. [PubMed: 16082249]
9. Miettinen M, Sarlomo-Rikala M. Expression of calretinin, thrombomodulin, keratin 5, and mesothelin in lung carcinomas of different types: an immunohistochemical analysis of 596 tumors in comparison with epithelioid mesotheliomas of the pleura. *Am J Surg Pathol*. 2003; 27:150–158. [PubMed: 12548160]
10. Hassan R, Ho M. Mesothelin targeted cancer immunotherapy. *Eur J Cancer*. 2008; 44:46–53. [PubMed: 17945478]
11. Bera TK, Pastan I. Mesothelin is not required for normal mouse development or reproduction. *Mol Cell Biol*. 2000; 20:2902–2906. [PubMed: 10733593]
12. Rump A, Morikawa Y, Tanaka M, Minami S, Umesaki N, Takeuchi M, et al. Binding of ovarian cancer antigen CA125/MUC16 to mesothelin mediates cell adhesion. *J Biol Chem*. 2004; 279:9190–9198. [PubMed: 14676194]
13. Gubbels JA, Belisle J, Onda M, Rancourt C, Migneault M, Ho M, et al. Mesothelin-MUC16 binding is a high affinity, N-glycan dependent interaction that facilitates peritoneal metastasis of ovarian tumors. *Mol Cancer*. 2006; 5:50–65. [PubMed: 17067392]

14. Li M, Bharadwaj U, Zhang R, Zhang S, Mu H, Fisher WE, et al. Mesothelin is a malignant factor and therapeutic vaccine target for pancreatic cancer. *Mol Cancer Ther.* 2008; 7:286–296. [PubMed: 18281514]
15. Wang T, Kajino K, Abe M, Tan K, Maruo M, Sun G, et al. Suppression of cell death by the secretory form of N-terminal ERC/mesothelin. *Int J Mol Med.* 2010; 26:185–191. [PubMed: 20596597]
16. Hassan R, Bullock S, Premkumar A, Kreitman RJ, Kindler H, Willingham M, et al. Phase I study of SS1P, a recombinant anti-mesothelin immunotoxin given as a bolus I.V. infusion to patients with mesothelin-expressing mesothelioma, ovarian, and pancreatic cancers. *Clin Cancer Res.* 2007; 13:5144–5149. [PubMed: 17785569]
17. Kreitman RJ, Hassan R, Fitzgerald DJ, Pastan I. Phase I trial of continuous infusion anti-mesothelin recombinant immunotoxin SS1P. *Clin Cancer Res.* 2009; 15:5274–5279. [PubMed: 19671873]
18. Hassan R, Sharon E, Schuler B, Mallory Y, Zhang J, Ling A, et al. Antitumor activity of SS1P with pemetrexed and cisplatin for front-line treatment of pleural mesothelioma and utility of serum mesothelin as a marker of tumor response. *J Clin Oncol.* 2011; 29 suppl abstract 7026.
19. Hassan R, Cohen SJ, Phillips M, Pastan I, Sharon E, Kelly RJ, et al. Phase I clinical trial of the chimeric anti-mesothelin monoclonal antibody MORAb-009 in patients with mesothelin-expressing cancers. *Clin Cancer Res.* 2010; 16:6132–6138. [PubMed: 21037025]
20. Clinicaltrials.gov Identifier: NCT00570713
21. Clinicaltrials.gov Identifier: NCT00738582
22. Le, DT.; Nir-Paz, R.; Hamp, J.; Mathur, S.; Neumanitis, J.; Steiman, D., et al. Results of phase I studies testing two live-attenuated *Listeria* vaccines, ANZ-100 and CrS-207, for the treatment of cancer. AACR Conference, Tumor Immunology: Basic and Clinical Advances; Nov 30–Dec 3, 2010; Miami Beach, FL. (Abstract).
23. Clinicaltrials.gov Identifier: NCT01417000.
24. Clinicaltrials.gov Identifier: NCT01355965.
25. Clinicaltrials.gov Identifier: NCT01439152.
26. Pastan I, Hassan R, FitzGerald DJ, Kreitman RJ. Immunotoxin treatment of cancer. *Annu Rev Med.* 2007; 58:221–237. [PubMed: 17059365]
27. Chowdhury PS, Viner JL, Beers R, Pastan I. Isolation of a high-affinity stable single-chain Fv specific for mesothelin from DNA-immunized mice by phage display and construction of a recombinant immunotoxin with anti-tumor activity. *Proc Natl Acad Sci U S A.* 1998; 95:669–674. [PubMed: 9435250]
28. Chowdhury PS, Pastan I. Improving antibody affinity by mimicking somatic hypermutation in vitro. *Nat Biotechnol.* 1999; 17:568–572. [PubMed: 10385321]
29. Hassan R, Lerner MR, Benbrook D, Lightfoot SA, Brackett DJ, Wang QC, et al. Antitumor activity of SS(dsFv)PE38 and SS1(dsFv)PE38, recombinant antimmesothelin immunotoxins against human gynecologic cancers grown in organotypic culture in vitro. *Clin Cancer Res.* 2002; 8:3520–3526. [PubMed: 12429643]
30. Li Q, Verschraegen CF, Mendoza J, Hassan R. Cytotoxic activity of the recombinant anti-mesothelin immunotoxin, SS1(dsFv)PE38, towards tumor cell lines established from ascites of patients with peritoneal mesotheliomas. *Anticancer Res.* 2004; 24:1327–1335. [PubMed: 15274292]
31. Zhang Y, Xiang L, Hassan R, Chang PH, Carrasquillo JA, Jang B, et al. Synergistic antitumor activity of taxol and immunotoxin SS1P in tumor-bearing mice. *Clin Cancer Res.* 2006; 12:4695–4701. [PubMed: 16899620]
32. Hassan R, Broaddus VC, Wilson S, Liewehr DJ, Zhang J. Anti-mesothelin immunotoxin SS1P in combination with gemcitabine results in increased activity against mesothelin-expressing tumor xenografts. *Clin Cancer Res.* 2007; 13:7166–7171. [PubMed: 18056197]
33. Zhang Y, Xiang L, Hassan R, Pastan I. Immunotoxin and Taxol synergy results from a decrease in shed mesothelin levels in the extracellular space of tumors. *Proc Natl Acad Sci U S A.* 2007; 104:17099–17104. [PubMed: 17940013]

34. Hassan R, Williams-Gould J, Watson T, Pai-Scherf L, Pastan I. Pretreatment with rituximab does not inhibit the human immune response against the immunogenic protein LMB-1. *Clin Cancer Res.* 2004; 10:16–18. [PubMed: 14734446]
35. Mossoba ME, Onda M, Taylor J, Massey PR, Treadwell S, Sharon E, et al. Pentostatin plus cyclophosphamide safely and effectively prevents immunotoxin immunogenicity in murine hosts. *Clin Cancer Res.* 2011; 17:3697–3705. [PubMed: 21521777]
36. Clinicaltrials.gov Identifier: NCT01362790.
37. Hassan R, Ebel W, Routhier EL, Patel R, Kline JB, Zhang J, et al. Preclinical evaluation of MORAb-009, a chimeric antibody targeting tumor-associated mesothelin. *Cancer Immun.* 2007; 7:20. [PubMed: 18088084]
38. Hassan R, Schweizer C, Lu KF, Schuler B, Remaley AT, Weil SC, et al. Inhibition of mesothelin-CA-125 interaction in patients with mesothelioma by the anti-mesothelin monoclonal antibody MORAb-009: Implications for cancer therapy. *Lung Cancer.* 2010; 68:455–459. [PubMed: 19744744]
39. Brockstedt DG, Giedlin MA, Leong ML, Bajhat KS, Gao Y, Lockett W, et al. Listeria-based cancer vaccines that segregate immunogenicity from toxicity. *Proc Natl Acad Sci U S A.* 2004; 101:13832–13837. [PubMed: 15365184]
40. Drake CG, Jaffee E, Pardoll DM. Mechanisms of immune evasion by tumors. *Adv Immunol.* 2006; 90:51–81. [PubMed: 16730261]
41. Morgan RA, Dudley ME, Wunderlich JR, Hughes MS, Yang JC, Sherry RM, et al. Cancer regression in patients after transfer of genetically engineered lymphocytes. *Science.* 2006; 314:126–129. [PubMed: 16946036]
42. Ramos CA, Dotti G. Chimeric antigen receptor (CAR)-engineered lymphocytes for cancer therapy. *Expert Opin Biol Ther.* [PMID21463133] E-pub ahead of print.
43. Till BG, Jensen MC, Wang J, Chen EY, Wood BL, Greisman HA, et al. Adoptive immunotherapy for indolent non-Hodgkin lymphoma and mantle cell lymphoma using genetically modified autologous CD20-specific T cells. *Blood.* 2008; 112:2261–2271. [PubMed: 18509084]
44. Carpenito C, Milone MC, Hassan R, Simonet JC, Lakhai M, Suhoski MM, et al. Control of large, established tumor xenografts with genetically retargeted human T cells containing CD28 and CD137 domains. *Proc Natl Acad Sci U S A.* 2009; 106:3360–3365. [PubMed: 19211796]
45. Zhao Y, Moon E, Carpenito C, Paulos CM, Liu X, Brennan AL, et al. Multiple injections of electroporated autologous T cells expressing a chimeric antigen receptor mediate regression of human disseminated tumor. *Cancer Res.* 70:9053–9061. 15. [PubMed: 20926399]
46. Rao, C.; Huber, M.; Vemuri, K.; Zhang, Q.; Chen, B.; Phillips, J., et al. Efficacy and toxicity of an anti-mesothelin antibody drug conjugate. *Proc Am Assoc Cancer Res*; 2009 Apr 18–22; Denver, CO. Philadelphia (PA): AACR; 2009. Abstract nr 3235.
47. Golfier, S.; Kahnert, A.; Heisler, I.; Kopitz, C.; Berhirster, K.; Stelte-Ludwig, B., et al. Identification of BAY 94-9343, a mesothelin antibody-drug conjugate (ADC): Characterization and anti-tumor activity in mesothelin-positive preclinical tumor models. *Proc, 102nd Annual AACR Mtg*; 2011 Apr 2–6; Orlando, FL. Philadelphia (PA): AACR; 2011. Abstract nr 1754.
48. Ho M, Hassan R, Zhang J, Wang QC, Onda M, Bera T, et al. Humoral immune response to mesothelin in mesothelioma and ovarian cancer patients. *Clin Cancer Res.* 2005; 11:3814–3820. [PubMed: 15897581]
49. Thomas AM, Santarsiero LM, Lutz ER, Armstrong TD, Chen YC, Huang LQ, et al. Mesothelin-specific CD8(+) T cell responses provide evidence of in vivo cross-priming by antigen-presenting cells in vaccinated pancreatic cancer patients. *J Exp Med.* 2004; 200:297–306. [PubMed: 15289501]
50. Laheru D, Lutz E, Burke J, Biedrzycki B, Solt S, Onners B, et al. Allogeneic granulocyte macrophage colony-stimulating factor-secreting tumor immunotherapy alone or in sequence with cyclophosphamide for metastatic pancreatic cancer: a pilot study of safety, feasibility, and immune activation. *Clin Cancer Res.* 2008; 14:1455–1463. [PubMed: 18316569]
51. Yokokawa J, Palena C, Arlen P, Hassan R, Ho M, Pastan I, et al. Identification of novel human CTL epitopes and their agonist epitopes of mesothelin. *Clin Cancer Res.* 2005; 11:6342–6351. [PubMed: 16144939]

52. Chang CL, Wu TC, Hung CF. Control of human mesothelin-expressing tumors by DNA vaccines. *Gene Ther.* 2007; 14:1189–1198. [PubMed: 17581599]
53. Miyazawa M, Iwahashi M, Ojima T, Katsuda M, Nakamura M, Nakamori M, et al. Dendritic cells adenovirally-transduced with full-length mesothelin cDNA elicit mesothelin-specific cytotoxicity against pancreatic cancer cell lines in vitro. *Cancer Lett.* 2011; 305:32–39. [PubMed: 21397388]
54. Wang B, Kuroiwa JM, He LZ, Charalambous A, Keler T, Steinman RM. The human cancer antigen mesothelin is more efficiently presented to the mouse immune system when targeted to the DEC-205/CD205 receptor on dendritic cells. *Ann N Y Acad Sci.* 2009; 1174:6–17. [PubMed: 19769731]
55. Breidenbach M, Rein DT, Everts M, Glasgow JN, Wang M, Passineau MJ, et al. Mesothelin-mediated targeting of adenoviral vectors for ovarian cancer gene therapy. *Gene Ther.* 2005; 12:187–193. [PubMed: 15526007]
56. Tsuruta Y, Pereboeva L, Breidenbach M, Rein DT, Wang M, Alvarez RD, et al. A fiber-modified mesothelin promoter-based conditionally replicating adenovirus for treatment of ovarian cancer. *Clin Cancer Res.* 2008; 14:3582–3588. [PubMed: 18519792]

Table 1

Phase I/II clinical trials of mesothelin targeted agents

Agent	Type of clinical trial	Dose and schedule	Patient population	No. of patients	Outcome	Refs.
SSIP (Immunotoxin)	Phase I single agent bolus study	Dose-escalation of SSIP given as i.v. infusion QOD	Mesothelin positive mesotheliomas, ovarian and pancreatic cancer	34	MTD of SSIP was 45 µg/kg i.v. QOD × 3 doses. DLT was pleuritis. 4 MR and 18 SD out of 33 pts.	16
	Phase I single agent c.i. study	Dose escalation of SSIP given as c.i. over 10 days	Mesothelin positive mesotheliomas, ovarian and pancreatic cancers	24	MTD was 25 µg/kg/d given as c.i. over 10 days. No CR 1 PR.	17
	Phase I SSIP plus pemetrexed and cisplatin	SSIP dose escalation with fixed standard doses of pemetrexed and cisplatin	Chemotherapy naïve malignant mesothelioma patients who are not candidates for curative surgical resection	19	Study open to patient accrual. At the MTD 5 out of 7 evaluable patients have had partial response.	18
MORAb-009 (Chimeric mAb)	SSIP plus paclitaxel, carboplatin & bevacizumab	SSIP dose escalation with fixed dose of chemotherapy and bevacizumab	Newly diagnosed stage IV lung adenocarcinoma that are mesothelin positive	2	Study closed since the incidence of mesothelin positivity was less than the expected 50% positivity in lung adenocarcinoma and the mesothelin staining was predominantly cytoplasmic (Hassan R, unpublished data).	N/R
	Phase I single agent study	Dose-escalation study	Mesothelioma, pancreatic cancer and mesothelin positive lung and ovarian cancer	24	MTD 200 mg/m ² in patients with mesothelioma treatment with MORAb-009 led to increase in serum CA-125 levels	19
	Phase II MORAb-009 plus gemcitabine	Patients randomized to either gemcitabine alone or gemcitabine plus MORAb-009	Locally advanced and metastatic pancreatic cancer	N/R	Study closed. Data not yet available.	20
	Phase II MORAb-009 plus pemetrexed and cisplatin	Single arm study	Newly diagnosed unresectable pleural mesothelioma	N/R	Study is ongoing but closed to new patient accrual.	21
	Phase I single agent	Dose-escalation study	Patients with mesothelin expressing cancers	17	MTD 1 × 10 ⁹ cfu Mesothelin specific immune response observed in 5 out of 10 evaluable patients.	22
CRS-207 (tumor vaccine)	Phase II CRS-207 plus GVAX	Patients randomized to GVAX pancreas vaccine versus GVAX pancreas vaccine plus CRS-207	Previously treated metastatic pancreatic cancer	-	Clinical trial opened August 2011	23
	Phase I	Patients will receive 1 to 3 doses of autologous CIR T cells	Progressive malignant pleural mesothelioma	-	Clinical trial opened May 2011	24
	Phase I	BAY 94-9343 given i.v. every 3 weeks	Patients with advanced solid tumors	-	Clinical trial opened September 2011	25

Abbreviations: c.i., continuous infusion; CIR, Chimeric Immune Receptor; CR, complete response; MR, minor response; MTD, maximum tolerated dose; N/R, not reported; PFS, progression free survival; PR, partial response; QOD, every other day; SD, stable disease .