

RESEARCH ARTICLE

Open Access

# Functional alleles of the flowering time regulator *FRIGIDA* in the *Brassica oleracea* genome

Judith A Irwin<sup>1</sup>, Clare Lister<sup>1</sup>, Eleni Soumpourou<sup>1</sup>, Yanwen Zhang<sup>1</sup>, Elaine C Howell<sup>2</sup>, Graham Teakle<sup>3</sup> and Caroline Dean<sup>1\*</sup>

## Abstract

**Background:** Plants adopt different reproductive strategies as an adaptation to growth in a range of climates. In *Arabidopsis thaliana* *FRIGIDA* (*FRI*) confers a vernalization requirement and thus winter annual habit by increasing the expression of the MADS box transcriptional repressor *FLOWERING LOCUS C* (*FLC*). Variation at *FRI* plays a major role in *A. thaliana* life history strategy, as independent loss-of-function alleles that result in a rapid-cycling habit in different accessions, appear to have evolved many times. The aim of this study was to identify and characterize orthologues of *FRI* in *Brassica oleracea*.

**Results:** We describe the characterization of *FRI* from *Brassica oleracea* and identify the two *B. oleracea* *FRI* orthologues (*BolC.FRI.a* and *BolC.FRI.b*). These show extensive amino acid conservation in the central and C-terminal regions to *FRI* from other Brassicaceae, including *A. thaliana*, but have a diverged N-terminus. The genes map to two of the three regions of *B. oleracea* chromosomes syntenic to part of *A. thaliana* chromosome 5 suggesting that one of the *FRI* copies has been lost since the ancient triplication event that formed the *B. oleracea* genome. This genomic position is not syntenic with *FRI* in *A. thaliana* and comparative analysis revealed a recombination event within the *A. thaliana* *FRI* promoter. This relocated *A. thaliana* *FRI* to chromosome 4, very close to the nucleolar organizer region, leaving a fragment of *FRI* in the syntenic location on *A. thaliana* chromosome 5. Our data show this rearrangement occurred after the divergence from *A. lyrata*. We explored the allelic variation at *BolC.FRI.a* within cultivated *B. oleracea* germplasm and identified two major alleles, which appear equally functional both to each other and *A. thaliana* *FRI*, when expressed as fusions in *A. thaliana*.

**Conclusions:** We identify the two *Brassica oleracea* *FRI* genes, one of which we show through *A. thaliana* complementation experiments is functional, and show their genomic location is not syntenic with *A. thaliana* *FRI* due to an ancient recombination event. This has complicated previous association analyses of *FRI* with variation in life history strategy in the *Brassica* genus.

**Keywords:** *FRIGIDA*, Flowering time, vernalization, synteny, *Brassica oleracea*, *Arabidopsis thaliana*

## Background

The switch to reproductive development is a fundamental process in the plant life cycle. The molecular mechanisms underlying this developmental transition have been extensively studied in *Arabidopsis thaliana*. An integrated network of environmentally responsive genetic pathways converge on a common set of targets to quantitatively regulate the genes required to switch the apical

meristem from a vegetative to a floral state [1-3]. One important environmental cue is prolonged cold, which accelerates flowering in a process termed vernalization and aligns pollination and seed set with the favourable conditions of spring. Variation in requirement for vernalization exists in many plant species and this influences life history strategy with plants requiring vernalization adopting a perennial, biennial or winter annual habit in contrast to summer annuals, which flower in the first growing season. This is in contrast to other species that are more reliant on photoperiodic signals or endogenous cues e.g. rice [4]. The significant fitness consequences of

\* Correspondence: caroline.dean@jic.ac.uk

<sup>1</sup>Department of Cell and Developmental Biology, John Innes Centre, Norwich Research Park, Norwich NR4 7UH, UK

Full list of author information is available at the end of the article

flowering time variation, demonstrated in annual [5,6] and perennial plants [7], have most likely contributed to the evolution of the extensive variability in flowering time control. Flowering also influences the pattern of growth throughout the seasons and affects many agronomic characters including the quantity and quality of crop production. This is particularly apparent in cultivated brassicas, where variation in the flowering process has been selected to produce a diverse array of economically important morphological forms.

A major determinant in the variation of vernalization requirement in *A. thaliana* is allelic variation at *FRIGIDA* (*FRI*) [8-11]. *FRI* represses flowering by promoting the expression of the floral repressor *FLOWERING LOCUS C* (*FLC*) [12,13]. Vernalization acts antagonistically to *FRI* and accelerates flowering by down-regulating *FLC*. A number of rapid-cycling variants of *A. thaliana* that do not need vernalization were found to have arisen through loss of function of *FRI*, an evolutionary step that has occurred multiple times [8,9,11,14]. Parallel evolution through allelic variation at a common target has been found in other organisms [15]. It was therefore interesting to ask whether a similar evolutionary step has occurred in other plant species. Many other species do show variation in vernalization requirement and it is an important agronomic trait in many major crops. For example, in *B. oleracea* (horticultural brassicas) vernalization-requiring biennials are represented by cabbage and Brussels sprouts, with summer annual crops including some calabrese and cauliflower cultivars. Orthologues of *FRI* have been identified in *A. lyrata* [16], *Capsella* species [17] and the halophyte *Thellungiella halophila* [18] within the Brassicaceae, and more broadly in *Medicago truncatula*, *Lotus japonicus*, *Vitis vinifera* [19], *Populus balsamifera* [20] and *Oryza sativa* [21]. To date natural variation in vernalization requirement has been associated with *FRI* polymorphism in *A. lyrata* [8] and allelic variation in one orthologue in *Brassica napus* (*BnaA.FRI.a*) has been associated with flowering time variation [22].

We are interested in understanding the molecular basis of variation in flowering time and vernalization requirement in horticultural brassicas. Genetic information from *A. thaliana* can generally be applied to Brassica species because of their evolutionary relatedness. The *Arabidopsis* and *Brassica* genera are in the same family (Brassicaceae) with *B. oleracea* thought to have arisen from a triplication of an ancestral genome similar to that of *Arabidopsis* [23-26]. Genetic information on the control of flowering in *Arabidopsis* can be applied to Brassica species because of the colinearity of the *Arabidopsis* and *Brassica* genomes [23,27,28]. This has been used to infer candidate genes that might account for QTL underlying flowering time and other variation [22,29,30]; however, in some instances it can be misleading [31]. Here, we


identify the two *FRI* genes in the *B. oleracea* genome and map their genomic locations. We also explored allelic variation at one of the *FRI* loci in cultivated *B. oleracea* germplasm. These new data will provide the necessary information to elucidate how general a role *FRI* plays in life history variation in the Brassicaceae.

## Results and discussion

### Two *FRI* genes are present in the *Brassica oleracea* genome

The *BoFRI* genes were isolated from the JBo BAC library of the *B. oleracea* Chinese kale genotype A12DHd [32] through hybridization with an *A. thaliana* *FRI* genomic clone. From seven positive BAC clones two that showed distinct *FRI* hybridization patterns (JBo72I23 and JBo88G16, Figure 1a, b) were selected for sub-cloning. Analysis of these confirmed they carried different *Brassica* paralogues designated *BoI.C.FRI.a* and *BoI.C.FRI.b* [33] and referred to hereafter as *BoFRIa* and *BoFRIb* [GenBank JN191450 and JN191449]. As in other species, *BoFRIa* and *BoFRIb* contain three exons encoding predicted open reading frames (ORFs) of 594 and 585 residues respectively (Figure 1c). *BoFRIa* contains two coiled-coil domains, typically involved in protein oligomerisation (as predicted by COILS [http://www.ch.embnet.org/software/COILS\\_form.html](http://www.ch.embnet.org/software/COILS_form.html)[34]), very similar to the predicted structure of the *A. thaliana* *FRI* (*AtFRI*) [8,35]. In contrast, *BoFRIb* is predicted to contain only one coiled-coil domain in the C-terminal region as was found to be the case for two of the four *FRI* identified in *B. napus* [22].

*AtFRI* is the original member of a family of seven proteins in *A. thaliana* which, apart from the two predicted coiled-coil domains, show no homology with any other proteins and whose function has yet to be determined. Recent analysis of the *FRI* protein family [19] identified a conserved core central domain. Outside of this domain significant variation is observed that allows the *FRI* family to be subdivided into five distinct groups. *AtFRI* and its orthologues in other species are defined by a conserved region of 37 amino acids in the N-terminal region of the protein. The *BoFRI* proteins we describe here contain this conserved 37 amino acid region reinforcing the view they are *FRI* orthologues; however, the amino acids either side of this region show lower homology to *AtFRI* (Figure 1c). This region includes much of the first predicted coiled-coil in *BoFRIa*. Variations in this domain in *BoFRIb* result in the loss of a predicted coiled-coil, emphasising a possible functional significance for the amino acid polymorphisms in this region. A similar degree of divergence from *AtFRI* is found in the N-terminal region of an orthologue of *FRI* isolated from the halophyte *T. halophila* and in four orthologues of *FRI* identified in *B. napus* [18,22]. By contrast, there is extensive amino acid conservation




**Figure 1 Cloning *BoFRIa* and *BoFRIb*.** (a) Southern hybridisation of 12 BACs identified by colony hybridisation and probed with *AtFRI*. Lanes 1, 4, 5, 8, 11, 12 and 13 contain clones that show homology to *FRIGIDA*. BACs in lanes 2, 3, 6, 7, and 10 do not cross hybridise (b) HindIII digest of six BACs probed with conserved region from exons 2 and 3 of *BoFRIa*. Lane 5 contains JBo72123 from which *BoFRIa* was sequenced. Lane 7 contains JBo88G16. Lanes 1, 2, 3 and 6 contain four further BACs showing the same hybridization pattern as JBo88G16. Note the intensity of the hybridization is indicative of the sequence divergence between *BoFRIa* and *BoFRIb* (See 1c) (c) Comparison of the protein sequences of *BoFRIa* and *BoFRIb* with other members of the FRI sub-family. From top to bottom they are *Brassica oleracea* *BoFRIa*, *Brassica oleracea* *BoFRIb*, *Thellungiella halophila* *ThFRI*, *Arabidopsis thaliana* *AtFRI*, *Arabidopsis lyrata* ssp *lyrata* *AlIFRI* and *Arabidopsis lyrata* ssp *petraea* *AlpFRI*. The N-terminal domain containing the conserved region of 37 amino acids (indicated by solid bar) that defines copies of *FRIGIDA* from other members of the FRI superfamily [17]. The coiled-coil domains are indicated by the red lines.

between *BoFRIa*, *BoFRIb* and *AtFRI* in the central and C-terminal regions (Figure 1c). Transgenic analysis of the functional domains of *AtFRI* in *A. thaliana* where the N or C terminus was deleted revealed that the N-terminal region was less important for function [19], perhaps explaining the high degree of divergence observed.

#### The *BoFRI* genes map to regions that are non-syntenic with *A. thaliana FRI*

A genomic fragment including exon 2, intron 2 and exon 3 of *BoFRIa* (and showing a high level of conservation in *BoFRIb* and *AtFRI*) was hybridized to mapping filters from two *B. oleracea* mapping populations: Chinese kale × calabrese (var. *alboglabra* × var. *italica*; A12DHdxGDDH33, [36]; Figure 2a) and cauliflower × Brussels sprout (var. *botrytis* × var. *gemmifera*; N × G [37]). RFLPs for one of the two *BoFRI* loci segregated in the A12DHdxGDDH33 mapping population that allowed this locus to be mapped to 39.5 cM on linkage group C3 of the *B. oleracea* genome. The locus mapping to C3 was identified as *BoFRIa* by fluorescence in situ hybridization (FISH) with BAC JBo72I23, from which *BoFRIa* was originally sequenced (Figure 2b). JBo88G16 was located on the short arm of chromosome C9 by FISH (Figure 2c, d). Therefore, the second locus, *BoFRIb*, was on linkage group C9. Two further BACs showing the same restriction pattern as JBo88G16 (Figure 1b) hybridized to the same location on C9 (data not shown). These results confirm that the *B. oleracea* genome contains two orthologues.

Comparative analysis of the *Brassica* and *A. thaliana* genomes has shown that the chromosomal regions of C3 and C9 to which the two *BoFRI* loci have been mapped are syntenic to a region of *A. thaliana* chromosome 5 and not to the top of *A. thaliana* chromosome 4, where *AtFRI* (At4g00650) is located [23]. This region of chromosome 5 includes a number of genes known to be involved in the control of flowering including *FLC*, *FY* and *CONSTANS* (*CO*). Several QTL studies have found loci for flowering time variation mapping to this genomic region in a number of *Brassica* populations including *B. oleracea* [38-41], *B. rapa* [42-46]; *B. nigra* [47,48] and *B. napus* [29,49]. The mapping we have undertaken reveals the proximity of *BoFRI* not only to *BoFLC*, but also *BoFY* and *BoCO*; other flowering time genes that have been mapped previously. The sequences of *BoFRIa* and *BoFRIb* further allow us to identify which of the four orthologues of *AtFRI* recently identified in *B. napus* [22] are the two C genome copies. The four copies of *FRI* were designated *BnaA.FRI.a*, and *BnaX.FRI.b-d*. Comparison of the amino acid sequences of these proteins with *BoFRIa* and *BoFRIb* suggest that *BnaX.FRI.d* is the orthologue of *BoFRIa* and the C genome homoeologue of *BnaA.FRI.a*. This conclusion is further supported by the



fact that *BnaA.FRI.a* was mapped to a region of A3 homoeologous to the region of C3 where we have mapped *BoFRIa*. Comparison of the amino acid sequence of *BnaX.FRI.c* shows it to be identical to that of *BoFRIb*. *BnaX.FRI.c* appears most similar to *BnaX.FRI.b* and is


therefore likely to be the A genome homoeologue of *BoFR1b* in *B. napus*.

**A recombination event specific to the *A. thaliana* lineage has relocated the *FRIGIDA* gene to the top of chromosome 4**

In *A. thaliana*, the *AtFRI* locus is located at the top of chromosome 4. However, it has previously been reported that the orthologue of *FRI* in *A. lyrata* maps to linkage group 8 [50,51]. This linkage group is orthologous to the lower arm of *A. thaliana* chromosome 5 [50,52,53]. Interestingly, an annotated gene model in

this region of *A. thaliana* chromosome 5 (*At5g51090*) shows a high degree of homology to *AtFRI*, containing parts of intron 1 and exon 3 but lacking other parts of the coding region, thus it may be a pseudogene [50]. Genevestigator data suggest *At5g51090* is expressed at very low levels, supporting this hypothesis [54]. Downstream, in the opposite orientation, is *At5g51100*, encoding an iron superoxide dismutase and the *BoFRIa* BAC clone contains 3' sequence showing homology to exons 3-9 of this *A. thaliana* chromosome 5 gene (Figure 3a).

Synteny has been studied extensively in the Brassicaceae genomes due to its potential for gene identification



**Figure 3** *BoFRIa* and *BoFR1b* map to regions of the *B. oleracea* genome that are non-syntenic with *A. thaliana*. **(a)** Comparison of *BoFRIa* genomic clone with annotated regions of *A. thaliana* chromosomes 4 and 5 represented in the 5' to 3' orientation. Solid black rectangles correspond to the three exons that make up *FRI*. Arrows indicate the orientation of other genes. (i) *BoFRIa*, genomic clone. (ii) Region of *A. thaliana* chromosome 5 containing an iron superoxide dismutase (*FSD2*, *At5g51100*) and *At51090* showing homology to *FRI* intron 1 and exon 3. (iii) Region of *A. thaliana* chromosome 4 containing *AtFRI* and *At4g400651* showing homology to exon 9 of *FSD2*. **(b)** Proposed derivation of *A. thaliana* and *B. oleracea* linkage groups from ancestral karyotype (AK6 and AK8) as suggested in [53] showing the position of *FRI* and *FLC*. The rearrangement of ancestral blocks W and R results in orthologues of *FRI* and *FLC* being brought into close proximity in the ancestral Brassica genome. This region is present on chromosomes C2, C3 and C9 of *B. oleracea* and is located in regions of the *B. oleracea* genome showing synteny to *A. thaliana* chromosome 5. *BoFRIa* is located on C3 and *BoFR1b* on C9. The copy of *FRI* on C2 has been lost.

and marker development. *Arabidopsis* and *Brassica* are thought to have diverged about 43 Mya with a triplication of an ancestral genome (similar to that of *Arabidopsis*) occurring approximately 23 Mya and giving rise to modern day diploid Brassica genomes [55]. *A. thaliana* and *A. lyrata* are thought to have diverged around 13 Mya, with a reduction in chromosome number, from the ancestral karyotype of  $n = 8$  (as represented in *A. lyrata*) to the derived state in *A. thaliana* of  $n = 5$  [24,55,56]. The ancestral karyotype of the Brassicaceae is proposed to be eight chromosomes composed of 24 conserved chromosomal blocks [57]. These blocks can be rearranged to model the genome structure of *A. thaliana*, *A. lyrata* and the modern day diploid Brassicas [24]. Thus the genomic composition of the nine chromosomes (C1-C9) of *B. oleracea* and ten chromosomes (A1-A10) of *B. rapa* can be related to both the ancestral karyotype and the *A. thaliana* genome.

The ancestral genomic blocks QR and WX from chromosomes 6 and 8 respectively of the ancestral karyotype, and today represented by *A. lyrata* [24], have been recombined in the ancestral *Brassica* genome prior to triplication, leading to the block WR being represented three times in the *B. rapa* genome on A2, A3 and A10 [25,26]. The paralogous regions of *B. oleracea* are on C2, C3 and C9 (Figure 3b). This rearrangement brings orthologues of *FLC* (block R) and *FRI* (block W) together on these chromosomes. Thus in *A. lyrata* and *B. oleracea*, *FRI* maps quite closely to *VIN3* (also in block W and required for vernalization [58]), as well as its major target *FLC* (block R; Figure 3b). *BoFRIa* mapping to C3 and *BoFRIb* to C9 thus represent two of the three syntenic regions. The third paralogue of *FRI* appears to have been lost from C2 during *B. oleracea* evolution; such gene loss is not uncommon [59]. This is in contrast to the current location of *FRI* at top of chromosome 4 in *A. thaliana* that shows homology to block O from chromosome 6 of the ancestral karyotype.

The data we present here suggest that the chromosomal rearrangements that occurred during the evolution of the ancestral Brassicaceae genome into *A. thaliana* included a recombination/rearrangement event that relocated a genomic region containing *AtFRI* to a position near the distal end of the short arm of chromosome 4, close to the nucleolar organiser region, leaving a non-functional remnant in the genomic position on chromosome 5 that is syntenic with *FRI* in the other Brassicaceae (*A. lyrata*, *B. oleracea*; [25,52]).

#### Two common alleles exist for *BoFRIa* in diverse genotypes of *B. oleracea*

The original sequences of *BoFRIa* and *BoFRIb* were obtained from A12DHD, one of the parents of the mapping population used in [38,39]. These studies mapped

QTL for flowering time on C3 in the region where we have mapped *BoFRIa*. We therefore sequenced 650 bp from exon 1 of *BoFRIa* from the other parent of this population, GDDH33 (data not shown). The GDDH33 sequence showed two amino acid substitutions (A118V and Q125E) compared to A12DHD. Thus the parents of this population are carrying different alleles of *BoFRIa* and it is possible that variation at *BoFRIa* is contributing to variation in flowering time in this population. Single amino acid substitutions have also been identified in alleles of *BnaA.FRIa* sequenced from the parents of the Tapidor  $\times$  Ningyou7 (TN) mapping population from *B. napus* and mapped to a region underlying a QTL for flowering time variation [22].

We sequenced *BoFRIa* and *BoFRIb* for two additional genotypes of *B. oleracea italica*. The A12DHD reference sequence is derived from a BAC clone of a Chinese kale, *B. oleracea alboglabra*, which flowers after 8 weeks [38] and can be considered a rapid-cycling type, not requiring vernalization. We therefore selected two additional genotypes of broccoli (*Brassica oleracea italica*); E1, which has a facultative vernalization response, flowering earlier following a period of cold, but which matures in October/November (Autumn) of the year of planting and E8 which has an obligate vernalization requirement and matures in April/May (Spring) of the following year. *BoFRIb* is highly conserved between the three genotypes with only 5 amino acid substitutions (D6G, K20Q, Q372K and R532W in E1 and N105T in E8). The sequencing of *BoFRIa* in these genotypes identified a polymorphic region in exon 1 that included two deletions of seven and three amino acids in E8 relative to E1, either side of the conserved block of 37 amino acids that defines the *FRI* proteins, (Figure 1c). Thirteen non-synonymous and 12 synonymous substitutions differentiate the A12DHD and E1 *BoFRIa* alleles from the E8 allele. We therefore designated the E1 and E8 *BoFRIa* alleles as *BoFRIa-1* and *BoFRIa-4* respectively [GenBank JN191393, JN191392].

We focused our subsequent analysis on *BoFRIa* as this showed most polymorphism and extended our analysis to include 55 genotypes from the cultivated *B. oleracea* Diversity Foundation Set developed at the University of Warwick (BoLDFS, King et al. [http://www.brassica.info/resource/plants/diversity\\_sets.php](http://www.brassica.info/resource/plants/diversity_sets.php); [60] Table 1). A 650 bp region of *BoFRIa* covering the exon 1 polymorphic region containing the two deletions was sequenced (Table 1, GenBank JN191394-191448). We identified six *BoFRIa* alleles within this subset of 55 genotypes from the BoLDFS. These can be divided into two groups; *BoFRIa 1-3* and *BoFRIa 4-6* where alleles in the second group include the seven amino acid and three amino acid deletions. The *BoFRIa-1* and *BoFRIa-4* alleles were the most common within the 55 genotypes studied. In

**Table 1 Amino acid polymorphisms in BoFR1a from cultivated genotypes of *Brassica oleracea***

Line	Accession name	Crop type	Crop group	Ref	origin	cult/landrace/DH	32	36-42	97	102	103	106	107	108	109	110-112	118	119	125	172	BoFR1a Allele
SIR5a	Siria DH line 5a	Cauliflower	Cauliflower	E03	N/A	DH	T	V	V	S	S	P	K	D	A	A	A	C	Q	F	1
HRIGRU002838	ALGROMAIO NO 2	Cauliflower	Cauliflower	E05	NL	cultivar	T	V	V	S	S	P	K	D	A	A	A	C	Q	F	1
HRIGRU004832	TOSCANO	Autumn cauliflower	Cauliflower	D10	ITA	cultivar	T	V	V	S	S	P	K	D	A	A	V	C	E	F	2
HRIGRU005458	DI JESI	Autumn cauliflower	Cauliflower	E02	ITA	cultivar	T	V	V	S	S	P	K	D	A	A	V	C	E	F	2
HRIGRU006254	TASMAN	Autumn cauliflower	Cauliflower	D11	AUS	cultivar	T	V	V	S	S	P	K	D	A	A	A	C	Q	F	1
HRIGRU004239	CANBERRA	Autumn cauliflower	Cauliflower	D08	AUS	cultivar	T	V	V	S	S	P	K	D	A	A	A	C	Q	F	1
HRIGRU008558	GIGANTE DI NAPOLI NATALINO *	Autumn cauliflower	Cauliflower	D12	ITA	N/A	T	V	V	S	S	P	K	D	A	A	A	C	Q	F	1
HRIGRU006797	SOFIA	Autumn cauliflower	Cauliflower	E01	SP	cultivar	T	V	V	S	S	P	K	D	A	A	A	C	Q	F	1
HRIGRU004814	BIANCO NAPOLETANE NATALINO	Summer cauliflower	Cauliflower	E07	ITA	cultivar	T	V	V	S	S	P	K	D	A	A	A	C	Q	F	1
HRIGRU004991	ALL THE YEAR ROUND	Summer cauliflower	Cauliflower	E08	UK	N/A	T	V	V	S	S	P	K	D	A	A	A	C	Q	F	1
HRIGRU004847	VERDE DI MACERATA	Green cauliflower	Cauliflower	F05	ITA	N/A	T	V	V	S	S	P	K	D	A	A	V	C	E	F	2
HRIGRU004850	DI ALBENGA	Green cauliflower	Cauliflower	F06	ITA	cultivar	T	V	V	S	S	P	K	D	A	A	A	C	Q	F	1
HRIGRU004861	ROMANESCO NATALINO	Romanesco cauliflower	Cauliflower	F07	ITA	cultivar	T	V	A	P	P	P	K	D	A	A	V	C	E	V	3
HRIGRU002891	ST MALO HALF HATIF	Winter cauliflower	Cauliflower	E11	FRA	cultivar	T	V	V	S	S	P	K	D	A	A	A	C	Q	F	1
HRIGRU004492	WINTER ROSCOFF	Winter cauliflower	Cauliflower	E12	IRE	landrace	T	V	V	S	S	P	K	D	A	A	A	C	Q	F	1
HRIGRU006230	LATE QUEEN	Winter cauliflower	Cauliflower	F02	IND	cultivar	T	V	V	S	S	P	K	D	A	A	A	C	Q	F	1
ROS152b	Roscoff type F1 DJ1356 DH line 152b	Roscoff winter cauliflower	Cauliflower	E06	N/A	DH	T	V	V	S	S	P	K	D	A	A	V	C	E	F	2
Cor12b	Corvette DH line 12b	broccoli	Broccoli	A07	N/A	DH	T	V	V	S	S	P	K	D	A	A	V	C	E	F	2
CAL 18b	DH line Royal Sluis F1 R571343 (DJ6546)	broccoli	Broccoli	A11	N/A	DH	T	V	V	S	S	P	K	D	A	A	V	C	E	F	2
HRIGRU002398	PICOLINI DI PALERMO	broccoli	Broccoli	B01	ITA	cultivar	T	V	V	S	S	P	K	D	A	A	A	C	Q	F	1
HRIGRU011802	MUGNOLI	broccoli?	Broccoli	B02	ITA	landrace	T	V	V	S	S	P	K	D	A	A	A	C	Q	F	1
HRIGRU005276	CIMA VIOLETTA NATALINO	Purple head broccoli	Broccoli	B08	ITA	cultivar	T	V	V	S	S	P	K	D	A	A	V	C	E	F	2

**Table 1 Amino acid polymorphisms in BoFR1a from cultivated genotypes of *Brassica oleracea* (Continued)**

HRIGRU003543	PURPLE SPROUTING LATE IMPROVED	Broccoli	B09	UK	cultivar	T	V	V	S	P	K	D	A	A	C	Q	F	1		
HRIGRU005416	CAVOLO CAVOLINA RIZZA	Broccoli	B06	ITA	cultivar	T	V	V	S	P	K	D	A	A	C	Q	F	1		
HRIGRU004705	RAMOSO CALABRESE PRECOCE	Broccoli	B03	ITA	cultivar	S	DEL	I	A	S	S	P	N	K	DEL	A	Y	E	F	4
HRIGRU005425	CAVOLO BROCCOLO NATALINO	Broccoli	B05	ITA	cultivar	T	V	V	S	P	K	D	A	A	C	Q	F	1		
BOH 85c	Bohmerwaldkohl DH line 85c	Cabbage	C07	N/A	DH	S	DEL	I	A	S	S	P	N	K	DEL	A	Y	E	F	4
HRIGRU005652	SHETLAND CABBAGE	Cabbage	C11	UK	N/A	S	DEL	I	A	S	S	P	N	K	DEL	A	Y	E	F	4
HRIGRU007833	LARGE BLOOD RED	Cabbage	D01	IND	cultivar	S	DEL	I	A	S	S	P	N	K	DEL	A	Y	E	F	4
HA 84a	Hawke DH line Ha84a	Cabbage	C08	N/A	DH	T	V	V	S	P	K	D	A	A	C	Q	F	1		
HRIGRU004773	CAVOLO VERZA SAN GIOVANNI	Cabbage	D05	ITA	cultivar	T	DEL	I	A	S	S	P	N	K	DEL	A	Y	E	F	5
HRIGRU011490	COUVE REPOLHO BACALAN	Cabbage	D07	PORT	landrace	T	V	V	S	P	K	D	A	A	C	Q	F	1		
HRIGRU004771	CAVOLO CARPUCCIO MEDIO NAPOLETANE	Cabbage	D06	ITA	cultivar	S	DEL	I	A	S	S	P	N	K	DEL	A	Y	E	F	4
HRIGRU002574	CATTLE (EARLY DRUMHEAD)	Cabbage	D02	UK	cultivar	S	DEL	I	A	S	S	P	N	K	DEL	A	Y	E	F	4
AC582	DH ex. Nym	Brussels sprout	B11	N/A	DH	S	DEL	I	A	S	S	P	N	K	DEL	A	Y	E	F	4
HRIGRU000342	EVESHAM GIANT	Brussels sprout	B12	UK	cultivar	S	DEL	I	A	S	S	P	N	K	DEL	A	Y	E	F	4
HRIGRU000605	WILHELMSBURGER	Brussels sprout	C01	DEN	cultivar	S	DEL	I	A	S	S	P	N	K	DEL	A	Y	E	F	4
HRIGRU002227	SANDA ROEM VAN CASTRICUM	Brussels sprout	C02	UK	cultivar	S	DEL	I	A	S	S	P	N	K	DEL	A	Y	E	F	4
HRIGRU002787	GROENENBOOM LATE SELECTION	Brussels sprout	C03	NL	cultivar	S	DEL	I	A	S	S	P	N	K	DEL	A	Y	E	F	4
HRIGRU005086	OLD BEDFORDSHIRE STOCK	Brussels sprout	C04	UK	cultivar	S	DEL	I	A	S	S	P	N	K	DEL	A	Y	E	F	4
HRIGRU006212	CAVOLO DI BRUXELLES MEZZO NANO	Brussels sprout	C05	ITA	cultivar	S	DEL	I	A	S	S	P	N	K	DEL	A	Y	E	F	4
HRIGRU008226	LOCAL SELECTION	Brussels sprout	C06	BHUTAN	landrace	S	DEL	I	A	S	S	P	N	K	DEL	A	Y	E	F	4
CGN14111	Butzo	Kale	G03	N/A	N/A	S	DEL	I	A	S	S	P	N	K	DEL	A	Y	E	F	4
HRIGRU006226	GIANT JERSEY KALE	Kale	G04	UK	cultivar	T	V	V	S	P	K	D	A	A	C	Q	F	1		
HRIGRU009846	RED ON GREEN	Ornamental kale	G08	JPN	cultivar	S	DEL	I	A	S	S	P	N	K	DEL	A	Y	E	F	4
HRIGRU003598	WESTLAND WINTER VERDURA	Borecole kale	F10	UK	cultivar	S	DEL	I	A	S	S	P	N	K	DEL	A	Y	E	F	4



**Table 1 Amino acid polymorphisms in BoFR1a from cultivated genotypes of *Brassica oleracea* (Continued)**

HRIGRU006210	CAVOLO NERO DI TOSCANA O *	Fodder, black kale	Kale	F12	ITA	cultivar	S	DEL	I	A	S	S	P	N	K	DEL	A	Y	E	F	4
HRIGRU011183	PURPLE VIENNA	Kohl rabi	Kohl rabi	G10	USA	cultivar	S	DEL	I	A	S	S	P	N	K	DEL	A	Y	E	F	4
HRIGRU008267	WHITE VIENNA	Kohl rabi	Kohl rabi	G11	IS	cultivar	S	DEL	I	A	S	S	P	N	K	DEL	A	Y	E	F	4
HRIGRU005443	CAVOLO FORTE	Purple kohlrabi	Kohl rabi	G12	ITA	cultivar	S	DEL	I	A	S	S	P	N	K	DEL	A	C	E	F	6
HRIGRU007543	CHINESE KALE	Chinese kale	Alboglaba	A02	CHINA	landrace	T		V	V	S	P	K	D	A		A	C	Q	F	1
Senna (GK95186)		Chinese white kale	Alboglaba	A05	N/A	DH	S	DEL	I	A	S	S	P	N	K	DEL	A	Y	E	F	4
HRIGRU009490	COUVE CORTE	Tronchuda cabbage	Tronchuda cabbage	H02	PORT	landrace	S	DEL	I	A	S	S	P	N	K	DEL	A	Y	E	F	4
HRIGRU009574	COUVE PENCA DE GONDOMAR	Tronchuda cabbage	Tronchuda cabbage	H05	PORT	landrace	T		V	V	S	P	K	D	A		A	C	Q	F	1
HRIGRU007796		Wild cabbage	Wild cabbage	H08	UK	N/A	S	DEL	I	A	S	S	P	N	K	DEL	A	Y	E	F	4

Details of amino acid polymorphisms in the 55 *B. oleracea* genotypes from the BoDFS genotypes [60] for the first 650 bp of BoFR1a. The number of genotypes in a BoFR1a allele type within each vegetable type is also listed

addition the two deletions identified in *BoFRIa-4* always co-occurred and were found at high frequency together with a small number of non-synonymous nucleotide polymorphisms. The two deletions, which had been found in the late-flowering broccoli, were over-represented in *B. oleracea* vegetable types such as Brussels sprouts and kohlrabi with a winter annual or biennial habit, usually grown for consumption of their vegetative rather than floral forms (Table 1, Figure 4). Interestingly *BnaX.FRI.d* from the *B. napus* winter variety Express [22], which we have identified here as the C genome homologue of *BoFRIa* in *B. napus* has both of the deletions identified in the *BoFRI 4-6* class of alleles that are overrepresented in brassica vegetable types with a winter annual or biennial habit. On closer examination *BnaX.FRI.d* was found to have the same amino acid sequence as *BoFRIa-5* and is also present in the European winter type and Chinese semi-winter type parental lines of the TN mapping population [22].

#### Functional analysis of *BoFRIa* alleles in *A. thaliana*

To ascertain if the two most common *BoFRIa* alleles conferred any functional differences we undertook transformation experiments. The coding and 3'UTR sequences from the *BoFRIa-1* and *BoFRIa-4* alleles were used to replace the *AtFRI* coding and 3' UTR sequences in an *A. thaliana* genomic clone. By retaining common regulatory sequences in the 5' region from the *AtFRI* gene we hoped to normalise expression and thus focus on the structural differences between the two *Brassica* proteins. These constructs were transformed into the rapid-cycling *A. thaliana* accession Columbia (Col-0). Col-0 carries a loss-of-function mutation within *AtFRI*, but has a functional *FLC* so these experiments would determine if *BoFRIa* could complement the *fri* mutation in Col-0, and induce late flowering. Both *BoFRIa* alleles complemented the loss-of-function mutation with > 100 primary (T1) transformed plants containing each of the *BoFRIa* alleles flowering very late compared to Col-0 plants and surprisingly also later than Col-0 transformed with a functional *AtFRI* (Figure 5).

To investigate the functionality of *BoFRIa-1* and *BoFRIa-4* alleles under different environmental conditions five transformants carrying each allele were analysed in the next (T2) generation. Flowering time was analysed as days-to-flower and total leaf number in plants that had no vernalization, or had experienced two or four weeks vernalization, at either 5°C or 10°C (Figure 6). In all treatments, except two weeks at 10°C (2W10°C), plants with either *BoFRIa* allele flowered as late as those carrying *AtFRI*. At 2W10°C plants carrying either of the *BoFRIa* alleles flowered later than *AtFRI*. Figure 6 also shows that plants undergoing a vernalization treatment at 10°C compared to 5°C continue to grow and initiate leaves at a faster rate.

Thus, when considering total leaf number as a measure of flowering time it appears that only 4W5°C was an effective vernalization treatment (Figure 6).

Expression of the coding sequences of the two *BoFRIa* alleles under the *AtFRI* 5' regulatory sequences showed that both alleles can produce functionally equivalent proteins that may, under some environmental conditions, be even stronger with respect to flowering time effects than the endogenous *A. thaliana* protein (Figure 6). In contrast, two *A. lyrata FRI* alleles conferred a quantitative difference in flowering time by both association and transgenic studies [16]. The maintenance of both *A. lyrata* alleles at intermediate frequencies in natural populations suggests they are differentially selected in different environments. If the *BoFRIa* alleles do underlie flowering time QTL then there must be expression differences between the two genes to account for the difference in flowering time. Both these genes could be expressed in a very different pattern to *AtFRI* as the rearrangement that moved it to chromosome 4 resulted in completely different 5' sequences less than 1 kb upstream of the transcription start site and places it in a very different chromatin context since it is now 200-300 kb downstream of the heterochromatic nucleolar organizer region NOR4 [61].


#### Conclusions

Knowledge of *B. oleracea FRI* gene number, functionality and map position now puts us in a strong position to undertake an extensive investigation into the contribution of allelic variation at *FRI* to flowering, vernalization and life history behaviours. Differences in life history between *A. lyrata* and *A. thaliana* such as outcrossing versus selfing and a perennial compared to annual habit may result in a requirement for some level of *FRI* functionality in *A. lyrata* that is optional in *A. thaliana* [16]. *B. oleracea*, like *A. lyrata*, is a largely outcrossing species and some wild *B. oleracea*, thought to be the progenitor of the modern crop plants, have been reported to keep flowering for up to 20 years [62]. Our analysis of *BoFRIa* suggests that only a small number of functional *BoFRIa* alleles are captured within the cultivated *B. oleracea* germplasm. To date we have found no evidence for loss-of-function mutations that are frequent in *AtFRI*. Further analysis of the 5' and 3' regulatory regions of *BoFRI* is now underway. The proximity of *BoFRI* to *BoFLC*, *BoFY* and *BoCO* opens up new questions of how this may influence flowering behaviour. It will be particularly important to be in a position to select specific alleles in breeding programmes to allow us to enhance robustness against increasing climate variability.


#### Methods

##### Cloning *BoFRI* genes

The JBo BAC library was hybridised with the *AtFRI* genomic clone, originally from the accession Stockholm



**Figure 4 Comparison of first 160 amino acids of BoFRLi from 55 BoLDFs genotypes.** Protein comparison of BoFRLi from cultivated genotypes of *B. oleracea* listed by crop type. The plate co-ordinates refer to those listed in Table 1. The conserved region of 37 amino acids that defines FRI from other members of the FRI superfamily is delineated by the horizontal black bar.



[8] and seven BACs identified, six having identical restriction patterns and one different. Purified genomic DNA was prepared (Qiagen Maxi Prep Kit) from two of these BACs (72I23 and 88G16) and used to generate shotgun libraries (TOPOor Shotgun Kit) of 1-2 kb fragments, in the pCR<sup>R</sup>4Blunt-TOPO<sup>R</sup> vector, giving 6-fold coverage. Colonies from these libraries were gridded onto nylon membrane (HyBond-N+) and hybridised to three probes generated from *AtFRI* (the 5' region, exon 1, and the 3' end of exon 3 and 3' UTR). BAC subclones were identified with each of these probes. Sequence analysis confirmed that the two BACs carried different *Brassica* paralogues.


### Mapping *BoFRI* loci in *B. oleracea* mapping populations

#### Genetic mapping

Mapping filters of the A12DHdxGDDH33 mapping population were produced and hybridized with a conserved *BoFRI* probe as described in [36,63]. A 900 bp conserved region from exons 2 and 3 from *BoFRIb* was amplified from A12Dhd genomic DNA with primers J2NG\_F1 (5' AAGTATCAAGCGTGGAAAGCA 3') and J2NG\_R1 (5' GTTACGAGGAGACCTGTGATT 3') and used to probe both the A12DHdxGDDH33 and NxG mapping filters (supplied by Graham Teakle, WHRI). Linkage analysis to map the *BoFRIa* locus was performed using Joinmap 3.0 [64] with the mapping data provided at BrassicaDB <http://brassica.bbsrc.ac.uk/BrassicaDB/>.

#### Fluorescence in situ hybridisation (FISH)

FISH was performed on chromosome spreads from the A12Dhd genotype of *B. oleracea* using methods described in [65]. The chromosomes are now named according to their corresponding linkage group. JBo72I23 was applied to meiotic pachytene spreads together with JBo62M08, a BAC which is associated with the RFLP marker pN22 on C3 at 42 cM and previously assigned to chromosome C3 by FISH. JBo88G16 (*BoFRIb*) was applied to mitotic metaphase spreads together with BAC BoB061G14, which



hybridizes to pericentromeric heterochromatin of six pairs of chromosomes, and a 45S rDNA probe from clone pTa71 [66], EMBLX07841. The chromosome pair to which JBo88G16 hybridized lacked signals from the other probes and had morphology suggestive of C9. Therefore, slides were reprobed with JBo32J18, a BAC associated with *BoFLC1* which has been mapped to a region between pN47E4NM (87 cM) and pN3E1 (103 cM) on C9 [31,67] and confirmed to be on C9 by FISH (unpublished). Two further BACs showing the same restriction pattern as JBo88G16 were applied separately with JBo88G16 to pachytene spreads.

#### Sequencing *BoFRIa* in BolDFS

The *B. oleracea* diversity foundation set (BolDFS) is a core collection of lines that represent the genetic variation across the morphologically diverse crops of this species [http://www.brassica.info/resource/plants/diversity\\_sets.php](http://www.brassica.info/resource/plants/diversity_sets.php). DNA was isolated using the DNeasy 96 Plant Kit

(Qiagen) and amplified using the GenomiPhi whole genome amplification kit (GE healthcare). A 650 bp fragment of *BoFR1a* was amplified from genomified DNA of 55 genotypes of the BolDFS by PCR with primers YWFRI\_F (5'CGCACATCGTCCATCAACAAG 3') and FRIJ1\_R2 (5'ATCCTTACCCACCAGCCT 3') using AMPLITAQ GOLD TAQ DNA Polymerase (Life Technologies Ltd (Invitrogen Division)). Sequence analysis was conducted using AlignX in Vector NTI (Invitrogen).

### Functional analysis of *BoFR1a* alleles

Plasmid pFRIg (in pBluescript-KS+, Stratagene) was mutagenised to introduce a *Bam*HI site immediately 5' of the ATG (plasmid pFRIg-B). Digestion of pFRIg-B with *Bam*HI plus *Cla*I allowed removal of the *AtFRI* coding sequences, leaving the 5' region of *AtFRI*. A 4.3 kb fragment containing *BoFR1a* was isolated from genomic DNA of lines E1 and E8 by PCR with primers BoFRI1\_ Bam\_ATG (5'CTTCCGCGGATCC-CATGGCCGTCCGTAAC3') and BoFRI1\_R2\_ ClaI (5'CAGAGATCGATC TCGAGAAAGGTAGCTGTTT 3'), using *PfuUltra* II Fusion HS DNA Polymerase (Agilent Technologies) and sequenced. PCR products were digested with *Bam*HI plus *Cla*I and the purified fragments ligated into *Bam*HI plus *Cla*I-digested pFRIg-B to give final constructs containing the 5'UTR of *AtFRI* with the coding and 3' UTR sequences of *BoFR1a-1* (in BoFRIa-1) or *BoFR1a-4* (in BoFRIa-4). pFRIg-B was used as the *A. thaliana* *FRI* control. The final constructs were ligated into binary vector pSLJ75516 (a gift from Prof. Jonathan Jones, <http://www.tsl.ac.uk/research/jonathan-jones/plasmids.htm>), on an *Eco*RI plus *Xho*I fragment (pFRIg-B) or *Eco*RI plus *Cla*I fragments (from BoFRIa-1 and BoFRIa-4). The constructs were transferred into *Agrobacterium* by triparental mating [68] and transformed into *A. thaliana* accession Col-0 by a floral dipping method (modified from [69]). T1 transformants were isolated by selection for Basta™ resistance. T2 seed were collected and flowering time determined by days-to-flower excluding the period of vernalization treatment and final leaf number at flowering.

### Plant growth

T2 *A. thaliana* seeds were sown on 'Arabidopsis mix' (Scotts® Levington F2 8.75 l bags, 100 l of grit, 200 g of Imidasect® 5 gr.) in plastic pots (7 cm × 7 cm) and stratified in a vernalization chamber at 5°C with an 8 h photoperiod and constant humidity for 3 days. Pots were moved to a naturally lit long day glasshouse for 7 days in May 2010 to allow germination and pre-growth. Seedlings not receiving a vernalization treatment remained in the glasshouse; seedlings to be vernalization treated were transferred back to a vernalization room or controlled environment cabinet (Snijder Economic Deluxe) for a

treatment of either two or four weeks at 5°C or 10°C. After vernalization, 20 plants per line were transplanted into trays with 40 cells of 2 cm × 2 cm and returned to the glasshouse. Trays were moved regularly to random positions to prevent any positional effects on plant growth. Flowering time was recorded as either total leaf number (rosette leaves plus cauline leaves at flowering) or bolting time; bolting time was scored as the number of days-to-flowering determined when the inflorescence stem was 3 cm tall.

### Abbreviations

QTL: Quantitative trait loci; BAC: Bacterial artificial chromosome; ORF: open reading frame; FISH: fluorescence in situ hybridization.

### Acknowledgements

We thank David Laurie, Lars Østergaard and Martin Trick for critical reading of the manuscript, Theresa Townsend for help with initial BAC identification, David Turner (JIC genome lab), for BAC library construction and initial sequencing of *BoFR1a* and *BoFR1b*, John Walshaw for advice on coiled-coil analysis and Sue Kennedy at Elsoms Seeds Ltd for seeds of E1 and E8. We also thank Andreas Mueller for early access to the *B. napus* *FRI* sequences following electronic publication. This work was funded by Defra grant HH37085FV, Defra feasibility Horticulture LINK grant HL0186, and BBSRC Strategic Grant to The John Innes Centre. The *B. oleracea* Diversity Foundation Set was produced at the University of Warwick with funding from Defra projects IF0157 and HH3723XS.

### Author details

<sup>1</sup>Department of Cell and Developmental Biology, John Innes Centre, Norwich Research Park, Norwich NR4 7UH, UK. <sup>2</sup>School of Biosciences, University of Birmingham, Birmingham B15 2TT, UK. <sup>3</sup>School of Life Sciences, University of Warwick, Wellesbourne CV35 9EF, UK.

### Authors' contributions

Jl and CD conceived and designed the experiments, supervised the work and wrote the paper. CL, YZ and Jl analyzed the *BoFR1a* alleles. ES and Jl analyzed the BolDFS and GT contributed DNA from the BolDFS. EH performed the FISH experiments. All authors approved and read the final manuscript.

Received: 2 November 2011 Accepted: 14 February 2012

Published: 14 February 2012

### References

1. Simpson GG, Dean C: Flowering - Arabidopsis, the Rosetta stone of flowering time? *Science* 2002, **296**:285-289.
2. Baurle I, Dean C: The timing of developmental transitions in plants. *Cell* 2006, **125**:655-664.
3. Amasino R: Seasonal and developmental timing of flowering. *Plant J* 2010, **61**:1001-1013.
4. Hayama R, Yokoi S, Tamaki S, Yano M, Shimamoto K: Adaptation of photoperiodic control pathways produced short-day flowering in rice. *Nature* 2003, **422**:719-722.
5. Franks SJ, Sim S, Weis AE: Rapid evolution of flowering time by an annual plant in response to a climate fluctuation. *Proc Natl Acad Sci (USA)* 2007, **104**:1278-1282.
6. Wilczek AM, Roe JL, Knapp MC, Cooper MD, Lopez-Gallego C, Martin LJ, Muir CD, Sim S, Walker A, Anderson J, Egan JF, Moyers BT, Petipas R, Giakountis A, Charbit E, Coupland G, Welch SM, Schmitt J: Effects of Genetic Perturbation on Seasonal Life History Plasticity. *Science* 2009, **323**:930-934.
7. Sandring S, Riihimäki MA, Savolainen O, Agren J: Selection on flowering time and floral display in an alpine and lowland population of *Arabidopsis lyrata*. *J Evolution Biol* 2007, **20**:558-567.
8. Johanson U, West J, Lister C, Michaels S, Amasino R, Dean C: Molecular analysis of *FRIGIDA*, a major determinant of natural variation in *Arabidopsis* flowering time. *Science* 2000, **290**:344-347.

9. Gazzani S, Gendall AR, Lister C, Dean C: **Analysis of the molecular basis of flowering time variation in Arabidopsis accessions.** *Plant Physiol* 2003, **132**:1107-1114.
10. Lempe J, Balasubramanian S, Sureshkumar S, Singh A, Schmid M, Weigel D: **Diversity of flowering responses in wild Arabidopsis thaliana strains.** *PLoS Genetics* 2005, **1**(1):e6.
11. Shindo C, Aranzana MJ, Lister C, Baxter C, Nicholls C, Nordborg M, Dean C: **Role of FRIGIDA and FLOWERING LOCUS C in determining variation in flowering time of Arabidopsis.** *Plant Physiol* 2005, **138**:1163-1173.
12. Sheldon CC, Rouse DT, Finnegan EJ, Peacock WJ, Dennis ES: **The molecular basis of vernalization: The central role of FLOWERING LOCUS C (FLC).** *Proc Natl Acad Sci (USA)* 2000, **97**:3753-3758.
13. Michaels SD, Amasino RM: **Loss of FLOWERING LOCUS C activity eliminates the late-flowering phenotype of FRIGIDA and autonomous pathway mutations but not responsiveness to vernalization.** *Plant Cell* 2001, **13**:935-941.
14. Le Corre V, Roux F, Rebound X: **DNA polymorphism at the FRIGIDA gene in Arabidopsis thaliana: extensive nonsynonymous variation is consistent with local selection for flowering time.** *Mol Biol Evol* 2002, **19**:1261-1271.
15. Colosimo PF, Hosemann KE, Balabhadra S, Villarreal G Jr, Dickson M, Grimwood J, Schmutz J, Myers RM, Schluter D, Kingsley DM: **Widespread parallel evolution in sticklebacks by repeated fixation of Ectodysplasin alleles.** *Science* 2005, **307**:1928-1933.
16. Kuitinen H, Niittyvuopio A, Rinne P, Savolainen O: **Natural variation in Arabidopsis lyrata vernalization requirement conferred by a FRIGIDA indel polymorphism.** *Mol Biol Evol* 2008, **25**:319-329.
17. Slotte T, Huang H, Lascoux M, Cepelitis A: **Polyploid speciation did not confer instant reproductive isolation in Capsella (Brassicaceae).** *Mol Biol Evol* 2008, **25**:1472-1481.
18. Fang Q, Liu J, Xu Z, Song R: **Cloning and characterization of a flowering time gene from Thellungiella halophila.** *Acta Bioch Bioph Sin* 2008, **40**:747-753.
19. Risk JM, Laurie RE, Macknight RC, Day CL: **FRIGIDA and related proteins have a conserved central domain and family specific N- and C- terminal regions that are functionally important.** *Plant Mol Biol* 2010, **73**:493-505.
20. Keller SR, Levensen N, Ingvarsson PK, Olson MS, Tiffin P: **Local selection across a latitudinal gradient shapes nucleotide diversity in Balsam Poplar, Populus balsamifera L.** *Genetics* 2011, **188**:941-952.
21. Goff SA, Ricke D, Lan TH, Presting G, Wang R, Dunn M, Glazebrook J, Sessions A, Oeller P, Varma H, Hadley D, Hutchison D, Martin C, Katagiri F, Lange BM, Moughamer T, Xia Y, Budworth P, Zhong J, Miguel T, Paszkowski U, Zhang S, Colbert M, Sun WL, Chen L, Cooper B, Park S, Wood TC, Mao L, Quail P, et al: **A draft sequence of the rice genome (Oryza sativa L. ssp japonica).** *Science* 2002, **296**:92-100.
22. Wang N, Qian W, Suppanz I, Wei L, Mao B, Long Y, Meng J, Müller AE, Jung C: **Flowering time variation in oilseed rape (Brassica napus L.) is associated with allelic variation in the FRIGIDA homologue BnaA.FRI.a.** *J Exp Bot* 2011, **62**:5641-5658.
23. Parkin IAP, Gulden SM, Sharpe AG, Lukens L, Trick M, Osborn TC, Lydiate DJ: **Segmental structure of the Brassica napus genome based on comparative analysis with Arabidopsis thaliana.** *Genetics* 2005, **171**:765-781.
24. Schranz ME, Lysak MA, Mitchell-Olds T: **The ABC's of comparative genomics in the Brassicaceae: building blocks of crucifer genomes.** *Trends Plant Sci* 2006, **11**:535-542.
25. Trick M, Kwon SJ, Choi SR, Fraser F, Soumpourou E, Drou N, Wang Z, Lee SY, Yang TJ, Mun JH: **Complexity of genome evolution by segmental rearrangement in Brassica rapa revealed by sequence-level analysis.** *BMC Genomics* 2009, **10**:539.
26. Mun JH, Kwon SJ, Kim JA, Jin M, Kim JS, Lim MH, Lee SI, Hong JK, Park TH, Lee SC, Kim BJ, Seo MS, Baek S, Lee MJ, Shin JY, Hahn JH, Hwang YJ, Lim KB, Park JY, Lee J, Yang TJ, Yu HJ, Choi IY, Choi BS, Choi SR, Ramchairy N, Lim YP, Fraser F, Drou N, Soumpourou E, et al: **Sequence and structure of Brassica rapa chromosome A3.** *Genome Biol* 2010, **11**:R94.
27. Suwabe K, Tsukazaki H, Iketani H, Hatakeyama K, Kondo M, Fujimura M, Nunome T, Fukuoka H, Hirai M, Matsumoto S: **Simple sequence repeat-based comparative genomics between Brassica rapa and Arabidopsis thaliana: the genetic origin of clubroot resistance.** *Genetics* 2006, **173**:309-319.
28. Gao M, Li G, Yang B, Qiu D, Farnham M, Quiros C: **High-density Brassica oleracea linkage map: identification of useful new linkages.** *Theor Appl Genet* 2007, **115**:277-287.
29. Long Y, Shi J, Qiu D, Li R, Zhang C, Wang J, Hou J, Zhao J, Shi L, Park BS: **Flowering time quantitative trait loci analysis of oilseed Brassica in multiple environments and genome wide alignment with Arabidopsis.** *Genetics* 2007, **177**:2433.
30. Smooker AM, Wells R, Morgan C, Beaudoin F, Cho K, Fraser F, Bancroft I: **The identification and mapping of candidate genes and QTL involved in the fatty acid desaturation pathway in Brassica napus.** *Theor Appl Genet* 2011, **122**:1075-1090.
31. Razi H, Howell EC, Newbury HJ, Kearsley MJ: **Does sequence polymorphism of FLC paralogs underlie flowering time QTL in Brassica oleracea?** *Theor Appl Genet* 2008, **116**:179-192.
32. O'Neill CM, Bancroft I: **Comparative physical mapping of segments of the genome of Brassica oleracea var. alboglabra that are homoeologous to sequenced regions of chromosomes 4 and 5 of Arabidopsis thaliana.** *Plant J* 2000, **23**:233-243.
33. Østergaard L, King GK: **Standardized gene nomenclature for the Brassica genus.** *Plant Methods* 2008, **4**:10-13.
34. Lupas A, Van Dyke M, Stock J: **Predicting Coiled Coils from Protein Sequences.** *Science* 1991, **252**:1162-1164.
35. Michaels SD, Bezerra IC, Amasino RM: **FRIGIDA-related genes are required for the winter-annual habit in Arabidopsis.** *Proc Natl Acad Sci USA* 2004, **101**:3281-3285.
36. Bohuon EJR, Keith DJ, Parkin IAP, Sharpe AG, Lydiate DJ: **Alignment of the conserved C genomes of Brassica oleracea and Brassica napus.** *Theor Appl Genet* 1996, **93**:833-839.
37. Sebastian RL, Kearsley MJ, King GJ: **Identification of quantitative trait loci controlling developmental characteristics of Brassica oleracea L.** *Theor Appl Genet* 2002, **104**:601-609.
38. Bohuon EJR, Ramsay LD, Craft JA, Arthur AE, Marshall DF, Lydiate DJ, Kearsley MJ: **The association of flowering time quantitative trait loci with duplicated regions and candidate loci in Brassica oleracea.** *Genetics* 1998, **150**:393-401.
39. Rae AM, Howell EC, Kearsley MJ: **More QTL for flowering time revealed by substitution lines in Brassica oleracea.** *Heredity* 1999, **83**:586-596.
40. Okazaki K, Sakamoto K, Kikuchi R, Saito A, Togashi E, Kuginuki Y, Matsumoto S, Hirai M: **Mapping and characterization of FLC homologs and QTL analysis of flowering time in Brassica oleracea.** *Theor Appl Genet* 2007, **114**:595-608.
41. Uptmoor R, Schrag T, Stützel H, Esch E: **Crop model based QTL analysis across environments and QTL based estimation of time to floral induction and flowering in Brassica oleracea.** *Mol Breeding* 2008, **21**:205-216.
42. Schranz ME, Quijada P, Sung SB, Lukens L, Amasino R, Osborn TC: **Characterization and effects of the replicated flowering time gene FLC in Brassica rapa.** *Genetics* 2002, **162**:1457-1468.
43. Nishioka M, Tamura K, Hayashi M, Fujimori Y, Ohkawa Y, Kuginuki Y, Harada K: **Mapping of QTLs for bolting time in Brassica rapa (syn. campestris) under different environmental conditions.** *Breeding Sci* 2005, **55**:127-133.
44. Lou P, Zhao JJ, Kim JS, Shen SX, Del Carpio DP, Song XF, Jin MN, Vreugdenhil D, Wang XW, Koornneef M, Bonnema G: **Quantitative trait loci for flowering time and morphological traits in multiple populations of Brassica rapa.** *J Exp Bot* 2007, **58**:4005-4016.
45. Yang X, Yu YJ, Zhang FL, Zou ZR, Zhao XY, Zhang DS, Xu JB: **Linkage Map Construction and Quantitative Trait Loci Analysis for Bolting Based on a Double Haploid Population of Brassica rapa.** *J Integrative Plant Biol* 2007, **49**:664-671.
46. Zhao J, Kulkarni V, Liu N, Del Carpio DP, Bucher J, Bonnema G: **BrFLC2 (FLOWERING LOCUS C) as a candidate gene for a vernalization response QTL in Brassica rapa.** *J Exp Bot* 2010, **61**:1817-1825.
47. Axelsson T, Shavorskaya O, Lagercrantz U: **Multiple flowering time QTLs within several Brassica species could be the result of duplicated copies of one ancestral gene.** *Genome* 2001, **44**:856-864.
48. Osterberg MK, Shavorskaya O, Lascoux M, Lagercrantz U: **Naturally occurring indel variation in the Brassica nigra COL1 gene is associated with variation in flowering time.** *Genetics* 2002, **161**:299-306.

49. Pires JC, Zhao JW, Schranz ME, Leon EJ, Quijada PA, Lukens LN, Osborn TC: **Flowering time divergence and genomic rearrangements in resynthesized Brassica polyploids (*Brassicaceae*)**. *Biol J Linn Soc* 2004, **82**:675-688.
50. Kuitinen H, de Haan AA, Vogl C, Oikarinen S, Leppala J, Koch M, Mitchell-Olds T, Langley CH, Savolainen O: **Comparing the linkage maps of the close relatives *Arabidopsis lyrata* and *A. thaliana***. *Genetics* 2004, **168**:1575-1584.
51. Beaulieu J, Belzile F, Jean M: **Linkage maps for *Arabidopsis lyrata* subsp. *lyrata* and *Arabidopsis lyrata* subsp. *petraea* combining anonymous and *Arabidopsis thaliana*-derived markers**. *Genome* 2007, **50**:142-150.
52. Yogeewaran K, Frary A, York TL, Amenta A, Lesser AH, Nasrallah JB, Tanksley SD, Nasrallah ME: **Comparative genome analyses of *Arabidopsis* spp.: inferring chromosomal rearrangement events in the evolutionary history of *A. thaliana***. *Genome Res* 2005, **15**:505-515.
53. Hu TT, Pattyn P, Bakker EG, Cao J, Cheng JF, Clark RM, Fahlgren N, Fawcett JA, Grimwood J, Gundlach H, Haberer G, Hollister JD, Ossowski S, Ottillar RP, Salamov AA, Schneeberger K, Spannagl M, Wang X, Yang L, Nasrallah ME, Bergelson J, Carrington JC, Gaut BS, Schmutz J, Mayer KFX, Van De Peer Y, Grigoriev IV, Nordborg M, Weigel D, Guo YL: **The *Arabidopsis lyrata* genome sequence and the basis of rapid genome size change**. *Nat Genet* 2011, **43**:476-481.
54. Hruz T, Laule O, Szabo G, Wessendorp F, Bleuler S, Oertle L, Wdimayer P, Gruissem W, Zimmermann P: **Genevestigator V3: a reference expression database for the meta-analysis of transcriptomes**. *Advances in Bioinformatics* 2008, 420747.q.
55. Beilstein MA, Nagalingum NS, Clements MD, Manchester SR, Mathews S: **Dated molecular phylogenies indicate a Miocene origin for *Arabidopsis thaliana***. *Proc Natl Acad Sci (USA)* 2010, **107**:18724-18728.
56. Koch MA, Matschinger M: **Evolution and genetic differentiation among relatives of *Arabidopsis thaliana***. *Proc Natl Acad Sci (USA)* 2007, **104**:6272-6277.
57. Lysak MA, Berr A, Pecinka A, Schmidt R, McBreen K, Schubert I: **Mechanisms of chromosome number reduction in *Arabidopsis thaliana* and related *Brassicaceae* species**. *Proc Natl Acad Sci (USA)* 2006, **103**:5224-5229.
58. Sung SB, Amasino RM: **Vernalization in *Arabidopsis thaliana* is mediated by the PHD finger protein VIN3**. *Nature* 2004, **427**:159-164.
59. Town CD, Cheung F, Maiti R, Crabtree J, Haas BJ, Wortman JR, Hine EE, Althoff R, Arbogast TS, Tallon LJ, Vigouroux M, Trick M, Bancroft I: **Comparative genomics of *Brassica oleracea* and *Arabidopsis thaliana* reveal gene loss, fragmentation, and dispersal after polyploidy**. *Plant Cell* 2006, **18**:1348-1359.
60. Allender CJ, Allainguillaume J, Lynn J, King GJ: **Simple sequence repeats reveal uneven distribution of genetic diversity in chloroplast genomes of *Brassica oleracea* L. and (n = 9) wild relatives**. *Theor Appl Genet* 2007, **114**:609-618.
61. Schmidt R, West J, Cnops G, Love K, Balestrazzi A, Dean C: **Detailed description of four YAC contigs representing 17 Mb of chromosome 4 of *Arabidopsis thaliana* ecotype Columbia**. *Plant J* 1996, **9**:755-765.
62. Raybould AF, Mogg RJ, Clarke RT, Gliddon CJ, Gray AJ: **Variation and population structure at microsatellite and isozyme loci in wild cabbage (*Brassica oleracea* L.) in Dorset (UK)**. *Genet Resour Crop Ev* 1999, **46**:351-360.
63. Sharpe AG, Parkin IAP, Keith DJ, Lydiate DJ: **Frequent non-reciprocal translocations in the amphidiploid genome of oilseed rape (*Brassica napus*)**. *Genome* 1995, **38**:1112-1121.
64. van Ooijen JW, Voorrips RE: **JoinMap version 3.0: Software for the calculation of genetic linkage map**. Wageningen: Plant Research International; 2001.
65. Howell EC, Barker GC, Jones GH, Kearsey MJ, King GJ, Kop EP, Ryder CD, Teakle GR, Vicente JG, Armstrong SA: **Integration of the cytogenetic and genetic linkage maps of *Brassica oleracea***. *Genetics* 2002, **161**:1225-1234.
66. Gerlach WL, Bedbrook JR: **Cloning and characterisation of ribosomal genes from wheat and barley**. *Nucleic Acids Res* 1979, **7**:1869-1885.
67. Salathia NS: **Regulation of Biological clocks in *Brassica oleracea* and *Arabidopsis thaliana***. University of Warwick: PhD Dissertation; 2003.
68. Figsurki DH, Helinski DR: **Replication of an origin-containing derivative of plasmid RK2 dependent on a plasmid function provided in trans**. *Proc Natl Acad Sci (USA)* 1979, **76**:1648-1652.
69. Bechtold N, Ellis J, Pelletier G: **In planta *Agrobacterium* mediated gene transfer by infiltration of adult *Arabidopsis thaliana* plants**. *C R Acad Sci Paris Life Sci* 1993, **316**:1194-1199.

doi:10.1186/1471-2229-12-21

**Cite this article as:** Irwin et al.: Functional alleles of the flowering time regulator *FRIGIDA* in the *Brassica oleracea* genome. *BMC Plant Biology* 2012 **12**:21.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

