

Complete Genome Sequence of the Dairy Isolate *Streptococcus macedonicus* ACA-DC 198

Konstantinos Papadimitriou,^a Stéphanie Ferreira,^b Nikolaos C. Papandreou,^c Eleni Mavrogonatou,^d Philip Supply,^{b,e,f,g,h} Bruno Pot,^{e,f,g,h} and Effie Tsakalidou^a

Laboratory of Dairy Research, Department of Food Science and Technology, Agricultural University of Athens, Athens, Greece^a; Genoscreen, Genomic Platform and R&D, Campus de l'Institut Pasteur, Lille, France^b; Department of Cell Biology and Biophysics, Faculty of Biology, University of Athens, Panepistimiopolis, Athens, Greece^c; Laboratory of Cell Proliferation and Ageing, Institute of Biology, National Center for Scientific Research Demokritos, Athens, Greece^d; Institut Pasteur de Lille, Center for Infection and Immunity of Lille, Lille, France^e; INSERM U1019, Lille, France^f; CNRS UMR8204, Lille, France^g; and Université Lille Nord de France, Lille, France^h

The species *Streptococcus macedonicus* is associated with the food environment, especially with fermented dairy products. Here we present the complete 2.1-Mb genome sequence of strain ACA-DC 198, which was isolated from naturally fermented Greek kasseri cheese.

Streptococcus macedonicus is an intriguing streptococcal species, since its most frequent source of isolation to date is fermented foods, mainly of dairy origin (5). Within the genus *Streptococcus*, only *Streptococcus thermophilus* is considered nonpathogenic, due to its adaptation to the milk environment (3, 8). Even though *S. macedonicus* has been shown to possess important (bio)technological features similar to those of *S. thermophilus* (5), it belongs to the *Streptococcus bovis/equinus* complex (13). It has been proposed that *S. macedonicus* is a subspecies of *S. gallolyticus*, along with *S. gallolyticus* subsp. *gallolyticus* (formerly *S. bovis* biotype I) and *Streptococcus pasteurianus* (formerly *S. bovis* biotype II.2) (13). This classification scheme is not universally accepted (15); however, there is no doubt that *S. macedonicus* is phylogenetically related to streptococci associated with cases of endocarditis, colorectal cancer, bacteremia, and meningitis (1, 6). Accordingly, the pathogenicity status of *S. macedonicus* is ambivalent, raising concerns about the safety of its use as a starter or adjunct culture in food fermentations.

Sequencing of *S. macedonicus* ACA-DC 198 genome was performed using the 454 GS-FLX (Roche Diagnostics, Basel, Switzerland) and the HiSeq 2000 (Illumina, San Diego, CA) technologies at Genoscreen (Lille, France) and Fidelity Systems, Inc. (Gaithersburg, MD), respectively. Following an initial round of shotgun pyrosequencing, contigs were assembled using Newbler Assembler software (454 Life Sciences, Branford, CT) and further combined with 3-kb paired-end reads down to 7 scaffolds. An additional round of Illumina sequencing was necessary for complete gap closure and finishing. The hybrid assembly between 454 and HiSeq 2000 data (>200× coverage) after analysis with Velvet (16), Newbler, and Fidelity Systems' in-house finishing software resulted in one continuous genomic scaffold of 2,130,034 bp and a plasmid of 12,728 bp. The genome assembly was validated against an NheI optical map of the *S. macedonicus* genome that was produced at OpGen Technologies, Inc. (Madison, WI).

Sequences were annotated with the BaSys (14) and the RAST (2) pipelines and manually curated using Kodon (Applied Maths N.V., Sint-Martens-Latem, Belgium). Final corrections were made based on the quality assessment of the annotation using GenePRIMP (11). We found 2,192 protein-coding genes on the chromosome, 192 of which were identified as potential pseudogenes, indicating an ongoing genome decay process. This hypoth-

esis is also supported by the fact that the *S. macedonicus* genome is approximately 220 kb smaller than the *S. gallolyticus* genome (7, 9, 12), despite the high level of gene synteny between the two species. Such a reductive evolutionary process is common among lactic acid bacteria adapted to the food environment (10) and in the case of *S. thermophilus* was also accompanied by the loss of pathogenicity traits (3). Interestingly, *S. macedonicus* ACA-DC 198 does not carry the *pil1* pilus locus, which is involved in infectious endocarditis caused by *S. gallolyticus* (4). These findings illustrate the usefulness of and the need for comprehensive comparative genomic analysis of *S. macedonicus* against its related streptococcal pathogens in order to assess the safety of the species for its use in foods.

Nucleotide sequence accession numbers. The *S. macedonicus* ACA-DC 198 chromosome and plasmid pSMA198 sequences have been deposited in EMBL under accession numbers [HE613569](#) and [HE613570](#).

ACKNOWLEDGMENTS

The present work was cofinanced by the European Social Fund and the National resources EPEAEK and YPEPTH through the Thales project.

REFERENCES

1. Abdulmir AS, Hafidh RR, Abu Bakar F. 2011. The association of *Streptococcus bovis/gallolyticus* with colorectal tumors: the nature and the underlying mechanisms of its etiological role. *J. Exp. Clin. Cancer Res.* 30:11.
2. Aziz RK, et al. 2008. The RAST Server: rapid annotations using subsystems technology. *BMC Genomics* 9:75.
3. Bolotin A, et al. 2004. Complete sequence and comparative genome analysis of the dairy bacterium *Streptococcus thermophilus*. *Nat. Biotechnol.* 22:1554–1558.
4. Danne C, et al. 2011. Molecular characterization of a *Streptococcus gallolyticus* genomic island encoding a pilus involved in endocarditis. *J. Infect. Dis.* 204:1960–1970.
5. De Vuyst L, Tsakalidou E. 2008. *Streptococcus macedonicus*, a multi-functional and promising species for dairy fermentations. *Int. Dairy J.* 18:476–485.

Received 22 December 2011 Accepted 19 January 2012

Address correspondence to Konstantinos Papadimitriou, kpapadimitriou@aua.gr, or Effie Tsakalidou, et@aua.gr.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JB.06804-11

6. Hensler ME. 2011. *Streptococcus gallolyticus*, infective endocarditis, and colon carcinoma: new light on an intriguing coincidence. *J. Infect. Dis.* **203**:1040–1042.
7. Hinse D, et al. 2011. Complete genome and comparative analysis of *Streptococcus gallolyticus* subsp. *gallolyticus*, an emerging pathogen of infective endocarditis. *BMC Genomics* **12**:400.
8. Hols P, et al. 2005. New insights in the molecular biology and physiology of *Streptococcus thermophilus* revealed by comparative genomics. *FEMS Microbiol. Rev.* **29**:435–463.
9. Lin IH, et al. 2011. Sequencing and comparative genome analysis of two pathogenic *Streptococcus gallolyticus* subspecies: genome plasticity, adaptation and virulence. *PLoS One* **6**:e20519.
10. Makarova KS, Koonin EV. 2007. Evolutionary genomics of lactic acid bacteria. *J. Bacteriol.* **189**:1199–1208.
11. Pati A, et al. 2010. GenePRIMP: a gene prediction improvement pipeline for prokaryotic genomes. *Nat. Methods* **7**:455–457.
12. Rusniok C, et al. 2010. Genome sequence of *Streptococcus gallolyticus*: insights into its adaptation to the bovine rumen and its ability to cause endocarditis. *J. Bacteriol.* **192**:2266–2276.
13. Schlegel L, Grimont F, Ageron E, Grimont PA, Bouvet A. 2003. Reappraisal of the taxonomy of the *Streptococcus bovis*/*Streptococcus equinus* complex and related species: description of *Streptococcus gallolyticus* subsp. *gallolyticus* subsp. nov., *S. gallolyticus* subsp. *macedonicus* subsp. nov. and *S. gallolyticus* subsp. *pasteurianus* subsp. nov. *Int. J. Syst. Evol. Microbiol.* **53**:631–645.
14. Van Domselaar GH, et al. 2005. BASys: a web server for automated bacterial genome annotation. *Nucleic Acids Res.* **33**:W455–W459.
15. Whiley RA, Kilian M. 2003. International Committee on Systematics of Prokaryotes Subcommittee on the taxonomy of staphylococci and streptococci. *Int. J. Syst. Evol. Microbiol.* **53**:915–917.
16. Zerbino DR, Birney E. 2008. Velvet: algorithms for *de novo* short read assembly using de Bruijn graphs. *Genome Res.* **18**:821–829.