

Sequence of the rat α c large chain of the clathrin associated protein complex AP-2

Kerry Lee Tucker, Kate Nathanson and Tomas Kirchhausen*

Department of Anatomy and Cellular Biology, Harvard Medical School, Boston, MA 02115, USA

Submitted July 6, 1990

EMBL accession no. X53773

We have isolated cDNA clones for the large αC chain of the rat brain Clathrin Associated Protein complex AP-2. A rat brain λ -ZAP-II cDNA library (Stratagene) was screened with the synthetic oligonucleotide probe 5'-C(C, T) TCCAT (A, G) TGNCC (A, G) AA (A, G) TC (G, T) ATGTCGTG, corresponding to the stretch of amino acids between His-70 and Ala-79 that are shared by both mouse αA and αC chains (1). Two cDNA clones were sequenced by the dideoxy termination method (2) using internal 17-mer sequencing primers. The combined DNA sequence is shown in Fig. 1. Sequence comparison with the mouse αA and αC homologues demonstrates that the rat protein is 99% identical to the mouse αC chain. It contains 938 amino acids, with no gaps, and two conservative changes corresponding

to Glu-8 instead of Asp-8 and Ile-788 instead of Val-788, respectively.

ACKNOWLEDGEMENTS

We acknowledge the support provided by NIH, March of Dimes, and the M. Peckett Foundation. TK is an Established Investigator of the American Heart Association.

REFERENCES

1. Robinson,M. (1989) *J. Cell Biol.* **108**, 833–842.
 2. Sanger,F., Nicklen,S. and Coulson,A.R. (1977) *Proc. Natl. Acad. Sci. USA* **74**, 5463–5467.

Figure 1. Nucleotide sequence of the rat brain large α c chain. The sequence comparison between the rat and mouse α c chains shows 95% identity at the DNA level and 99% identity at the protein level. Sequence differences and the initiation codon are indicated.

* To whom correspondence should be addressed