

Published in final edited form as:

Mem Inst Oswaldo Cruz. 2011 December ; 106(8): 1017–1023.

Molecular evidence for a single taxon, *Anopheles nuneztovari* s.l., from two endemic malaria regions in Colombia

Luz Marina Jaramillo¹, Lina A Gutiérrez¹, Shirley Luckhart², Jan E Conn^{3,4}, and Margarita M Correa^{1,+}

¹Grupo de Microbiología Molecular, Escuela de Microbiología, Universidad de Antioquia, Calle 67 No 53-108, Lab. 5-430, Medellín, Colombia

²Department of Medical Microbiology and Immunology, University of California at Davis, CA, USA

³Griffin Laboratory, Wadsworth Center, New York State Department of Health, NY, USA

⁴Department of Biomedical Sciences, School of Public Health, State University of New York at Albany, Albany, NY, USA

Abstract

To elucidate the *Anopheles nuneztovari* s.l. taxonomic status at a microgeographic level in four malaria endemic localities from Antioquia and Córdoba, Colombia, fragments of the Cytochrome oxidase subunit I (*COI*) and the *white* gene were used. The *COI* analysis showed low genetic differentiation with F_{ST} levels between -0.02 and 0.137 and Nm values between 3 and infinity, indicating the presence of high gene flow among *An. nuneztovari* s.l. populations from the four localities. The *COI* network showed a single most common haplotype, 1 ($n=55$), present in all localities, as the likely ancestral haplotype. Analysis of the *white* gene showed that *An. nuneztovari* s.l. populations from both departments grouped with haplotypes 19 and 20, which are part of lineage 3 previously reported. The results of the present study suggest that *An. nuneztovari* s.l. is a single taxon in the area of the present study.

Keywords

Malaria; *Anopheles nuneztovari*; *white* gene; Cytochrome oxidase subunit I; Colombia

Introduction

Malaria is an important public health problem in Colombia, as demonstrated by the fact that, after Brazil, Colombia has the second highest number of malaria cases in Latin America (OPS 2009). In the past ten years, the departments of Antioquia and Córdoba, where this investigation was carried out, have reported nearly 60% of the total cases of the country (Padilla *et al.* 2011). There are three main vectors, all in subgenus *Nyssorhynchus*, reported for Colombia present in endemic areas of these departments, including *Anopheles albimanus* Wiedemann, *Anopheles darlingi* -Root-, and *Anopheles nuneztovari* Gabaldón (Olano *et al.* 2001).

Anopheles nuneztovari s.l. is distributed from eastern Panamá to northern South America and Amazonia (Mirabello & Conn 2008) where it presents differences in behaviour (Montoya-Lerma *et al.* 2011). Brazilian collections of this species have detected natural

⁺Corresponding author: mcorrea@quimbaya.udea.edu.co, margaritcorrea@gmail.com .

infection with *Plasmodium* (Arruda *et al.* 1986; Tadei *et al.* 1998), but the role of *An. nuneztovari* s.l. in Brazil seems to be limited to local malaria transmission (Galardo *et al.* 2007). Indeed, populations from Brazil are mainly zoophilic and exophagic with a biting peak near sunset (Tadei & Thatcher 2000). In contrast, in Colombia and Venezuela, *An. nuneztovari* s.l. is a primary malaria vector (Lounibos & Conn 2000). Recent studies in northeastern Colombia showed that *An. nuneztovari* s.l. exhibits anthropophilic and endophagic behavior, with a biting peak between 20:00 and 23:00 hours, and can be naturally infected with *Plasmodium vivax* VK247 (Gutiérrez *et al.* 2009a). In Venezuela, the highest biting activity is close to midnight indoors and outdoors (Rubio-Palis & Curtis 1992).

Based on morphology, cytology, ecology, behavioral and molecular differences, several authors have suggested that *An. nuneztovari* s.l. is a species complex (summarized in Calado *et al.* 2008, Montoya-Lerma *et al.* 2011). Recent *white* gene analyses described five lineages across its range, with one (lineage 3) present in Colombia (Mirabello & Conn 2008). *An. nuneztovari* s.l. is characterized by high intraspecific morphological variability and is thus easily misidentified or confounded with other members of the Oswaldoi Group (Calle *et al.* 2008, Cienfuegos *et al.* 2008, Gómez *et al.* 2010). A deeper dissection of this challenging variability resulted in the description of a new morphological form in the Brazilian Amazon (Bergo *et al.* 2007), later confirmed as *Anopheles goeldii* Rozeboom and Gabaldón by sequence analysis of ITS2, *COI* and *white* genes (Calado *et al.* 2008). Scarpassa & Conn (2011, in press) recently determined that, based on a phylogenetic analysis of *COI* gene sequences, samples from the Amazon Basin could be assigned to two well-supported clades. They hypothesize that one of these clades represents *An. goeldii*, and the second clade represents a previously undescribed lineage or taxon in *An. nuneztovari* s.l.

Few studies have attempted to document the taxonomic status and population structure of *An. nuneztovari* s.l. in Colombia. One study evaluated phenotypic variation in specimens from western and eastern Colombia and detected high levels of wing spot variation, a key character that is often difficult to assess for proper identification (Fajardo *et al.* 2008). Similarly, morphometric analysis of female specimens of *An. nuneztovari* s.l. from Montelíbano, Córdoba, found high intraspecific variability (Gómez *et al.* 2010).

Studies using molecular markers have also attempted to clarify the taxonomic status and population structure of *An. nuneztovari* s.l. from northern and western Colombia. Using random amplified polymorphic DNA (RAPD), Posso *et al.* (2003) found low levels of genetic differentiation among three populations. Analysis of ITS2 sequences concluded that in Colombia, *An. nuneztovari* s.l. from both sides of the Andes mountains comprised a single genetic species (Sierra *et al.* 2004). Given the wide distribution of *An. nuneztovari* s.l. in Antioquia and Córdoba, this study was conducted to test the hypothesis that a single taxon is present in this region, to assess the genetic variability at a microgeographic level, and to identify the *white* gene lineage(s) present.

Materials and methods

Mosquito collection and processing

Adult female specimens of *An. nuneztovari* s.l. were collected between November 2007 and February 2010 in El Bagre (BAG), Zaragoza (ZAR), and San Pedro de Urabá (SPU), in the department of Antioquia, and in Montelíbano (MTL) and Puerto Libertador (PTL), in Córdoba (Figure 1, Table I). The collection protocols and informed consent were reviewed and approved by a Bioethics Committee for Human Research at the University of Antioquia (SIU-UdeA). Mosquitoes were identified by morphological keys (González & Carrejo 2009, Faran 1980, Faran & Linthicum 1981). A subset of specimens collected from each field trip,

is maintained in the University collection. For morphological reference, we retain the wings and legs fixed on slides of a percentage of the specimens. We also maintain photographic records of most specimens that have been processed for molecular analyses.

Molecular analyses

Total DNA was extracted from individual mosquito abdomens by a standard protocol (Rosero *et al.* 2010), used to confirm species identification following a PCR-RFLP-ITS2 assay developed by our group (Zapata *et al.* 2007, Cienfuegos *et al.* 2008, Cienfuegos *et al.* 2011), and then amplified for the *COI* and *white* gene fragments as outlined below. A 1,300 bp fragment of the mtDNA *COI* gene was amplified by PCR using 10 μ M primers UEA3/UEA10 (Lunt *et al.* 1996), in a 25 μ l final reaction volume as previously described (Gutiérrez *et al.* 2010). PCR products were purified using Wizard SV Gel and PCR Cleanup System (Promega, Madison, WI) following the protocol recommended by the manufacturer and sequenced in both directions.

An 800 bp region of the nuclear *white* gene was amplified using primers WF and W2R in a 25 μ l PCR reaction with conditions previously described by Mirabello & Conn (2008). Because of allelic variation (Mirabello & Conn 2008), *white* gene PCR products were cloned as described in Onyabe & Conn (1999). Five clones corresponding to five specimens from each locality were sequenced. Sequences were edited, assembled by pairwise alignment and then multiple alignments with Clustal W were performed using the Geneious software 5.3.4 (Drummond *et al.* 2011).

Descriptive statistics and population differentiation test

Haplotype and nucleotide diversities of mtDNA *COI* sequences were obtained using DnaSP version 5.00.07 (Rozas *et al.* 2003) and Arlequin version 3.11 software (Excoffier *et al.* 2005). Analysis of molecular variance (AMOVA) was used to evaluate within and among population variation using Arlequin software. The *COI* haplotype distribution was evaluated by a statistical parsimony network (TCS) with 95% support (Clement *et al.* 2000).

Phylogenetic relatedness of *An. nuneztovari* s.l. populations from Antioquia and Córdoba with populations from South America using *white* gene sequences

Split networks are successful tools for visualizing incompatible and ambiguous signals and have been used previously for phylogenetics studies (Gutiérrez *et al.* 2009b, Gutiérrez *et al.* 2010). Therefore, a neighbor-net network was estimated using SplitsTree4 version 4.10 (Huson & Bryant 2006, Huson & Steel 2004), for *An. nuneztovari* s.l. *white* gene sequences obtained in this study and for those in GenBank corresponding to specimens from Brazil, Bolivia, Surinam, Venezuela and Colombia (Accession numbers EU500727-EU500758). Bayesian analysis was carried out using Mr Bayes v3.1 (Huelsenbeck & Ronquist 2001), available in Geneious software 5.3.4 (Drummond *et al.* 2011), using the default HKY85 model and gamma distributions. The settings for the Markov Chain Monte Carlo (MCMC) were four chains running every 100,000 generations.

Results

An. nuneztovari s.l. genetic variation at the intra-population level

Out of 2,490 *Anopheles* mosquitoes collected in Antioquia and Córdoba, Colombia, 147 *An. nuneztovari* s.l. specimens were chosen (Table I, Figure 1). A 1,180 bp sequence of the *COI* gene was analyzed. The aligned sequences showed 23 variable sites and 11 that were parsimony informative. The highest nucleotide diversity was detected in San Pedro de Urabá ($P_i = 0.00249 \pm 0.001315$) and moderate to lowest diversity was detected in Puerto Libertador (0.00172 ± 0.001120). The haplotype diversity was highest for San Pedro de

Urabá (0.8134/−0.0418) and lowest for Puerto Libertador (0.709+/-0.0578; Table II). To compare the population pairs using the *COI* gene, samples from BAG and ZAR were combined. These localities are 11 km apart and Nm detected infinite migrants between them. There were 20 unique haplotypes (GenBank accession numbers JN255752 to JN255771), all of which connected parsimoniously. Four haplotypes were shared among all localities. The most widespread haplotype (type 1), showed a frequency of 38% (55/147), and was found in all localities. Three other haplotypes were distributed among all localities: type 2 (21%), type 3 (12%) and type 4 (10%). Haplotype 3 was not found in SPU (Table II). There were few private haplotypes (Figure 2).

Genetic structure and demographic inference

The F_{ST} values estimated for *COI* sequences from localities in Antioquia and Córdoba indicated low to moderate genetic differentiation, with values ranging between −0.02 and 0.137. The number of migrants per generation, Nm , was between 3 and infinity, indicating gene flow among all *An. nuneztovari* s.l. tested (Table III). Analysis of molecular variance, AMOVA, based on *COI* haplotype frequencies and conducted with all localities (populations) as one group showed that 12.49% ($p=0.18$) of the variance could be attributed to among localities (populations) and 87.51% within (Table IV).

The *COI* network showed a single most common haplotype (type 1, $n=55$), which was present in all the localities. There were only 16 private haplotypes, which demonstrates high demographic stability and the existence of gene flow among populations (Figure 2). After removing the intron sequence, a 630 bp fragment of the nuclear *white* gene was analyzed. From 20 sequences obtained, the consensus sequence was submitted to GenBank with accession number JN255772. Phylogenetic neighbor-net network based on *white* gene sequences showed that *An. nuneztovari* s.l. specimens from Antioquia and Córdoba grouped with haplotypes 19 and 20, which are part of lineage 3 (Figure 3) as reported by Mirabello & Conn (2008). However, the Bayesian analysis was completely unresolved and uninformative; whereas the neighbor-net network had an area of boxes, which indicated the need for larger sample sizes from these regions.

Discussion

The results obtained with *COI* sequences of specimens *An. nuneztovari* s.l. collected in localities of high malaria transmission of Antioquia and Córdoba showed low genetic differentiation and gene flow among these localities. In general, levels of *COI* nucleotide diversity were higher for the populations of Antioquia than for populations of Córdoba. Further, most haplotypes were shared between both regions, suggesting that these populations have a common demographic history (Castelloe & Templeton 1994) (Figure 2). Avise (2000) proposed that populations with high nucleotide and haplotype diversity are more stable, constant in size and have higher effective population sizes. In this study, *An. nuneztovari* s.l. from San Pedro de Urabá had the highest nucleotide and haplotype diversity, indicating that this population is the most stable among the populations studied. F_{ST} and Nm values obtained suggest that *An. nuneztovari* s.l. from Antioquia and Córdoba may actually comprise a single population. Nm values in malaria vectors are of relevance because they can be used to predict gene spread, e.g. of those genes important for insecticide resistance or parasite refractoriness. An understanding of gene flow is essential for the introduction of transgenic mosquitoes carrying genes for *Plasmodium* resistance.

White gene phylogenetic analyses of samples from Antioquia and Córdoba together with other South American populations produced mixed results. Our sequences were identical to haplotypes 19 and 20 reported by Mirabello & Conn (2008), which corresponded to their lineage 3. The area of boxes in the neighbor-net network combined with the unresolved

Bayesian analysis, suggest that a broader sampling of this species complex with analyses of *white* gene and other nuclear genes should be undertaken to more formally test the five lineages proposed in Mirabello & Conn (2008). Nevertheless, our results are consistent with previous studies with various markers that detected differentiation between *An. nuneztovari* s.l. populations from Colombia and Venezuela west of the Andes with those in other regions of South America (Elliott 1972, Kitzmiller *et al.* 1973, Conn *et al.* 1993, Scarpassa *et al.* 1996, 1999, 2000, Fritz *et al.* 1994, Mirabello & Conn 2008, Scarpassa & Conn in press 2011).

In comparison with the results presented here, a study using the ribosomal DNA internal transcribed spacer (ITS2) suggested low differentiation between *An. nuneztovari* s.l. populations located on opposite sides of the Andes mountains (NW and NE Colombia), indicating that *An. nuneztovari* s.l. is conspecific in Colombia (Sierra *et al.* 2004). Further, a study using RAPD markers showed low levels of genetic differentiation for *An. nuneztovari* s.l. populations from NW and NE Colombia, with higher levels of gene flow among western populations (Posso *et al.* 2003). Although Posso *et al.* (2003) used RAPD for discrimination of populations in western Colombia, their results are in agreement with the ones in the present study with *COI*, also in western Colombia. Collectively, results of these studies with *An. nuneztovari* s.l. populations from NW or NE Colombia are consistent in the detection of low genetic differentiation among the populations evaluated.

Localities in the present study are placed in the Magdalena biogeographic province (Morrone 2006), which encompasses western Venezuela and NW Colombia. The Magdalena province is characterized by temperatures above 24°C, plains with altitudes no greater than 500 meters, fragmented forest, the presence of livestock and agriculture, and active mining (IGAC 2002). The concurrence of similar ecologic and geographic conditions in this biogeographic region may influence *An. nuneztovari* s.l. populations, favoring the low genetic differentiation observed. The location of *An. nuneztovari* s.l. Cytotypes B and C coincide with the Magdalena biogeographic province, while Cytotype A was reported in the Brazilian Amazon, south of the country. Given that *An. nuneztovari* s.l. is broadly distributed in Colombia, including regions in the south in close proximity to the Brazilian Amazon, genetic analyses of *An. nuneztovari* s.l. should be expanded to include specimens from the southern regions such as the Orinoquía and the Amazonia and other markers such as microsatellites to detect recent mutational events.

Previous work by our group evaluated microgeographic genetic variation of *An. darlingi*, another important Colombian malaria vector, in Antioquia and Córdoba departments. Those results were based on microsatellites and *COI* analysis and also showed that *An. darlingi* populations were highly homogeneous and presented high levels of gene flow (Gutierrez *et al.* 2010). In the report presented here, microsatellite analysis was not performed because at the time of our collections and analyses, these markers had not been available for *An. nuneztovari* (Cardoza *et al.* <http://tomato.bio.trinity.edu/manuscripts/11-4/mer-11-0032.pdf>). It is possible that microsatellite analysis of *An. nuneztovari*, as with *An. darlingi*, would have revealed low genetic differentiation and high gene flow. In particular, analyses of *An. darlingi* in Brazil at the microgeographic level (152 km apart) showed low levels of genetic differentiation ($F_{ST}=0.001-0.095$) and Nm values between 4.7 and 363.8, indicating that at this distance and within similar ecological and geographic conditions, anopheline differentiation may be low (Scarpassa & Conn 2007). However, differentiation at the microgeographic level has been documented for other important malaria vectors such as *Anopheles gambiae* (McLain *et al.* 1989). Specifically, in seven Kenyan localities less than 10 km apart, the authors found significant differences in the genetic distribution of this important African vector indicating the absence of gene flow, probably due to poor dispersal abilities of these populations (McLain *et al.* 1989).

In summary, the results obtained in the present study provide critical data for understanding the primary malaria vector *An. nuneztovari* s.l. in localities with the highest annual reports of malaria cases in Colombia (Padilla *et al.* 2011). In particular, the presence of low genetic differentiation and high gene flow for important malaria vector species such as *An. darlingi* and *An. nuneztovari* s.l., indicate that malaria control programs could apply the same strategies, for example insecticide spraying, in these localities of Antioquia and Córdoba, as has been previously suggested (Gutiérrez *et al.* 2010).

Acknowledgments

We thank G. Gómez, M. Altamiranda, N. Naranjo, and A. Yates who cooperated in specimen collection and identification. We also are grateful to Sara A. Bickersmith (Wadsworth Center, New York State Department of Health, Albany, NY) for preliminary manuscript evaluation and critical editorial comments. LMJ received support for her MSc training from “Estrategia para la Sostenibilidad-Universidad de Antioquia 2007-2008”, code BUPP E01335 to MMC.

Financial support: This study received support from Comité para el Desarrollo de la Investigación, CODI-Universidad de Antioquia, Grant number 8700-074 to MMC, the United States National Institutes of Health grants R03AI076710-02 to MMC and AI R01 54139-02 to JEC.

REFERENCES

- Arruda M, Carvalho MB, Nussenzweig RS, Maracic M, Ferreira AW, Cochrane AH. Potential vectors of malaria and their different susceptibility to *Plasmodium falciparum* and *Plasmodium vivax* in northern Brazil identified by immunoassay. *Am J Trop Med Hyg.* 1986; 35:873–881. [PubMed: 3532844]
- Avice, J. *Phylogeography: The history and formation of species.* Harvard University Press; London: 2000. p. 447
- Bergo ES, Souto RN, Galardo AK, Nagaki SS, Calado DC, Sallum MA. Systematic notes on *Anopheles meigen* (Diptera: Culicidae) species in the state of Amapá, Brazil. *Mem Inst Oswaldo Cruz.* 2007; 102:373–376. [PubMed: 17568944]
- Calado D, Foster P, Bergo E, Santos C, Galardo A, Sallum M. Resurrection of *Anopheles goeldii* from synonymy with *Anopheles nuneztovari* (Diptera, Culicidae) and a new record for *Anopheles dunhami* in the Brazilian Amazon. *Mem Inst Oswaldo Cruz.* 2008; 103:791–799. [PubMed: 19148419]
- Calle DA, Quiñones ML, Erazo HF, Jaramillo N. Discriminación por morfometría geométrica de once especies de *Anopheles* (Nyssorhynchus) presentes en Colombia. *Biomedica.* 2008; 28:371–385. [PubMed: 19034360]
- Cardoza TB, Barbosa De, Sousa AC, Prado Lima M, Pereira De Souza A, Scarpassa VM. Microsatellite markers for the neotropical malaria vector *Anopheles nuneztovari* sensu lato: isolation, development and characterization of eighteen loci. *Mol Ecol Resources.* Available at: <http://tomato.bio.trinity.edu/manuscripts/11-4/mer-11-0032.pdf>.
- Castelloe J, Templeton A. Root probabilities for intraspecific gene trees under neutral coalescent theory. *Mol Phylogenet Evol.* 1994; 3:102–113. [PubMed: 8075830]
- Cienfuegos A, Gómez G, Córdoba L, Luckhart S, Conn J, Correa M. Diseño y evaluación de metodologías basadas en PCR-RFLP de ITS2 para la identificación molecular de mosquitos *Anopheles* spp. (Diptera: Culicidae) de la Costa Pacífica de Colombia. *Rev Biomed.* 2008; 19:35–44.
- Cienfuegos AV, Rosero DA, Naranjo N, Luckhart S, Conn JE, Correa MM. Evaluation of a PCR-RFLP-ITS2 assay for discrimination of *Anopheles* species in northern and western Colombia. *Acta Trop.* 2011; 118:128–135. [PubMed: 21345325]
- Clement M, Posada D, Crandall KA. TCS: a computer program to estimate gene genealogies. *Mol Ecol.* 2000; 9:1657–1659. [PubMed: 11050560]
- Conn J, Puertas Y, Seawright J. A new cytotype of *Anopheles nuneztovari* from western Venezuela and Colombia. *J Am Mosq Control Assoc.* 1993; 9:294–301. [PubMed: 8245938]

- Drummond AJ, Ashton B, Buxton S, Cheung M, Cooper A, Duran C, Field M, Heled J, Kearse M, Markowitz S, Moir R, Stones-Havas S, Sturrock S, Thierer T, Wilson A. Research software for biologists, not computer scientists. 2011 Available from: <http://www.geneious.com>.
- Elliott R. The Influence of vector behavior on malaria transmission. *Am J Trop Med Hyg.* 1972; 21:755–763. [PubMed: 4561523]
- Excoffier L, Laval G, Schneider S. Arlequin ver. 3.0: An integrated software package for population genetics data analysis. *Evol Bioinform Online.* 2005; 1:47–50. [PubMed: 19325852]
- Fajardo Ramos M, González Obando R, Suárez M, López David, Wilkerson R, Sallum M. Morphological analysis of three populations of *Anopheles* (*Nyssorhynchus*) *nuneztovari* Gabaldón (Diptera: Culicidae) from Colombia. *Mem Inst Oswaldo Cruz.* 2008; 103:85–92. [PubMed: 18368239]
- Faran ME. Mosquito studies (Diptera: Culicidae) XXXIV. A revision of the *Albimanus* section of the subgenus *Anopheles Nyssorhynchus* (Diptera: Culicidae). *Am Ent Inst.* 1980; 15:1–215.
- Faran M, Linthicum K. A handbook of the Amazonian species of *Anopheles* (*Nyssorhynchus*) (Diptera: Culicidae). *Mosq Syst.* 1981; 13:1–81.
- Fritz G, Conn J, Cockburn A, Seawright J. Sequence analysis of the ribosomal DNA internal transcribed spacer 2 from populations of *Anopheles nuneztovari* (Diptera: Culicidae). *Mol Biol Evol.* 1994; 11:406–416. [PubMed: 8015435]
- Galardo AK, Arruda M, D'Almeida Couto AA, Wirtz R, Lounibos LP, Zimmerman RH. Malaria vector incrimination in three rural riverine villages in the Brazilian Amazon. *Am J Trop Med Hyg.* 2007; 76:461–469. [PubMed: 17360868]
- Gómez G, Cienfuegos A, Gutiérrez L, Conn J, Correa M. Análisis morfológico y molecular evidencia problemas al identificar *Anopheles nuneztovari* (Diptera: Culicidae) por claves dicotómicas. *Rev Col Entomol.* 2010; 36:68–75.
- Gonzalez, R.; Carrejo, N. Introducción al estudio taxonómico de *Anopheles* de Colombia: claves y notas de distribución. 2nd ed. Programa Editorial Universidad de Valle; Cali: 2009. p. 260
- Gutiérrez LA, Gomez GF, Gonzalez JJ, Castro MI, Luckhart S, Conn JE, Correa MM. Microgeographic genetic variation of the malaria vector *Anopheles darlingi* Root (Diptera: Culicidae) from Córdoba and Antioquia, Colombia. *Am J Trop Med Hyg.* 2010; 83:38–47.
- Gutiérrez LA, Gonzalez JJ, Gomez GF, Castro MI, Rosero DA, Luckhart S, Conn JE, Correa MM. Species composition and natural infectivity of anthropophilic *Anopheles* (Diptera: Culicidae) in the states of Córdoba and Antioquia, Northwestern Colombia. *Mem Inst Oswaldo Cruz.* 2009a; 104:1117–11124.
- Gutiérrez LA, Naranjo NJ, Cienfuegos AV, Muskus CE, Luckhart S, Conn JE, Correa MM. Population structure analyses and demographic history of the malaria vector *Anopheles albimanus* from the Caribbean and the Pacific regions of Colombia. *Malar J.* 2009b; 8:259.
- Huelsenbeck JP, Ronquist F. MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics.* 2001; 17:754–755. [PubMed: 11524383]
- Huson DH, Bryant D. Application of phylogenetic networks in evolutionary studies. *Mol Biol Evol.* 2006; 23:254–267. [PubMed: 16221896]
- Huson DH, Steel M. Phylogenetic trees based on gene content. *Bioinformatics.* 2004; 20:2044–2049. [PubMed: 15044248]
- IGAC, Instituto Geográfico Agustín Codazzi. Atlas de Colombia. 5a ed. Imprenta Nacional de Colombia; Bogotá DC: 2002. p. 342
- Kitzmilller J, Kreutzer R, Tallaferro E. Chromosomal differences in populations of *Anopheles nuneztovari*. *Bull World Health Org.* 1973; 48:435–445. [PubMed: 4543549]
- Lounibos L, Conn J. Malaria vector heterogeneity in South America. *Am Entomol.* 2000; 46:238–249.
- Lunt DH, Zhang DX, Szymura JM, Hewitt GM. The insect cytochrome oxidase I gene: evolutionary patterns and conserved primers for phylogenetic studies. *Insect Mol Biol.* 1996; 5:153–165. [PubMed: 8799733]
- McLain DK, Collins FH, Brandling-Bennett AD, Were JB. Microgeographic variation in rDNA intergenic spacers of *Anopheles gambiae* in western Kenya. *Heredity.* 1989; 62:257–264. [PubMed: 2566589]

- Mirabello L, Conn JE. Population analysis using the nuclear white gene detects Pliocene/Pleistocene lineage divergence within *Anopheles nuneztovari* in South America. *Med Vet Entomol.* 2008; 22:109–119. [PubMed: 18498609]
- Montoya-Lerma J, Solarte Y, Giraldo-Calderón G, Quiñones M, Ruiz-López F, Wilkerson R, González R. Malaria vector species in Colombia - A review. *Mem Inst Oswaldo Cruz.* 2011; 106(Suppl.1):114–122. [PubMed: 21881765]
- Morrone JJ. Biogeographic areas and transition zones of Latin America and the Caribbean islands based on panbiogeographic and cladistic analyses of the entomofauna. *Annu Rev Entomol.* 2006; 51:467–494. [PubMed: 16332220]
- Olano VA, Brochero HL, Sáenz R, Quiñones ML, A MJ. Mapas preliminares de la distribución de especies de *Anopheles* vectores de malaria en Colombia. *Biomédica.* 2001; 21:402–408.
- Onyabe D, Conn J. Intragenomic heterogeneity of a ribosomal DNA spacer (ITS2) varies regionally in the neotropical malaria vector *Anopheles nuneztovari* (Diptera: Culicidae). *Insect Mol Biol.* 1999; 8:435–442. [PubMed: 10634969]
- Padilla JC, Álvarez G, Montoya R, Chaparro P, Herrera S. Epidemiology and control of malaria in Colombia. *Mem Inst Oswaldo Cruz.* 2011; 106(Suppl.1):114–122. [PubMed: 21881765]
- OPS. Informe de la situación del paludismo en las Américas. Organización Panamericana de la Salud; Washington DC: 2009. p. 9
- Posso CE, González R, Cárdenas H, Gallego G, Duque M, Suarez M. Random Amplified Polymorphic DNA analysis of *Anopheles nuneztovari* (Diptera: Culicidae) from western and northeastern Colombia. *Mem Inst Oswaldo Cruz.* 2003; 98:469–476. [PubMed: 12937756]
- Rosero D, Gutiérrez L, Cienfuegos A, Jaramillo L, Correa M. Optimización de un procedimiento de extracción de ADN para mosquitos anofelinos. *Rev Col Entomol.* 2010; 36:260–263.
- Rozas J, Sanchez-DelBarrio J, Messeguer X, Rozas R. DnaSP, DNA polymorphism analyses by the coalescent and other methods. *Bioinformatics.* 2003; 19:2496–2497. [PubMed: 14668244]
- Rubio-Palis Y, Curtis C. Biting and resting behaviour of anophelines in western Venezuela and implications for control of malaria transmission. *Med Vet Entomol.* 1992; 4:325–334. [PubMed: 1463897]
- Scarpassa VM, Conn JE. Tracking of Complex Evolutionary History and Pleistocene Epoch Divergence for Neotropical Malaria Vector *Anopheles nuneztovari* Sensu Lato with Mitochondrial DNA. *Am J Trop Med Hyg.* 2011 In Press.
- Scarpassa V, Conn J. Population genetic structure of the major malaria vector *Anopheles darlingi* (Diptera: Culicidae) from the Brazilian Amazon, using microsatellite markers. *Mem Inst Oswaldo Cruz.* 2007; 102:319–327. [PubMed: 17568937]
- Scarpassa VM, Geurgas S, Azeredo-Espin AM, Tadei WP. Genetic divergence in mitochondrial DNA of *Anopheles nuneztovari* (Diptera: Culicidae) from Brazil and Colombia. *Genet Mol Biol.* 2000; 23:71–78.
- Scarpassa VM, Tadei WP, Suarez MF. Population structure and genetic divergence in *Anopheles nuneztovari* (Diptera: Culicidae) from Brazil and Colombia. *Am J Trop Med Hyg.* 1999; 60:1010–1018. [PubMed: 10403335]
- Scarpassa V, Tadei W, Suarez M. Alloenzyme differentiation among allopatric populations of *Anopheles nuneztovari* (Diptera: Culicidae). *Braz J Genet.* 1996; 19:265–269.
- Sierra DM, Velez ID, Linton YM. Malaria vector *Anopheles* (*Nyssorhynchus*) *nuneztovari* comprises one genetic species in Colombia based on homogeneity of nuclear ITS2 rDNA. *J Med Entomol.* 2004; 41:302–307. [PubMed: 15185929]
- Tadei WP, Thatcher B, Santos JM, Scarpassa VM, Rodrigues IB, Rafael MS. Ecologic observations on anopheline vectors of malaria in the Brazilian Amazon. *Am J Trop Med Hyg.* 1998; 59:325–335. [PubMed: 9715956]
- Tadei WP, Thatcher DB. Malaria vectors in the Brazilian amazon: *Anopheles* of the subgenus *Nyssorhynchus*. *Rev Inst Med Trop Sao Paulo.* 2000; 42:87–94. [PubMed: 10810323]
- Zapata MA, Cienfuegos AV, Quiros OI, Quinones ML, Luckhart S, Correa MM. Discrimination of seven *Anopheles* species from San Pedro de Uraba, Antioquia, Colombia, by polymerase chain reaction-restriction fragment length polymorphism analysis of its sequences. *Am J Trop Med Hyg.* 2007; 77:67, 72. [PubMed: 17620632]




Figure 1. Distribution of collection sites for *An. nuneztovari* s.l. in Antioquia and Córdoba, Colombia. San Pedro de Urabá (SPU), El Bagre (BAG), Zaragoza (ZAR), Montelíbano (MTL), Puerto Libertador (PTL).




Figure 2. Parsimony based haplotype network of 20 *COI* haplotypes of 147 *An. nuneztovari* s.l. specimens. Each locality is represented by a distinct shade. Numbers out of the parentheses represent the haplotype designation and those inside the parentheses indicate the number of sequences included in each haplotype. The size of the circle is proportional to the frequency of detected haplotype. San Pedro de Urabá (SPU), El Bagre (BAG), Zaragoza (ZAR), Montelíbano (MTL), Puerto Libertador (PTL).




Figure 3. Neighbor-Net network of *Anopheles nuneztovari* s.l. *white* gene lineages from Antioquia and Córdoba compared with other *Anopheles nuneztovari* s.l from South America.

Table I
Study sites, dates of collection, coordinates and sample sizes (*n*) of *An. nuneztovari* s.l. collected from Antioquia and Córdoba, Colombia

Municipality (Abbreviation)	Year	Month (number of days)	Coordinates	<i>n</i>
Montelíbano (MTL)	2007	July (5) August (2)	7°59' N, 75°25' W	29
Puerto Libertador (PLB)	2007	January (6)	07°43'N, 75°51'W	32
El Bagre (BAG)	2009	March (3) May (3)	07° 35'N, 74°49'W	41
Zaragoza (ZAR)	2008	January (1) March (3) Jun (1)	07°29'N, 74°51'W	8
San Pedro de Urabá (SPU)	2010	February (5)	06°28'N, 75°33'W	37
Total <i>n</i> =				147

Table II
Shared COI haplotypes and genetic polymorphism statistics for *An. nuneztovari* s.l. collected in localities of Antioquia and Córdoba, Colombia

Locality	<i>n</i>	COI Haplotype	<i>S</i>	<i>h</i>	<i>Hd</i> (DE)	<i>Pi</i> (DE)
MTL	29	<u>1(11)</u> , <u>2(5)</u> , <u>3(2)</u> , <u>4(5)</u> , <u>6(2)</u> , <u>8(2)</u>	9	8	0.807+/-0.5240	0.00189+/-0.001199
PTL	32	<u>1(15)</u> , <u>2(8)</u> , <u>3(1)</u> , <u>4(5)</u> , <u>6(2)</u> , <u>8(2)</u>	7	7	0.709+/-0.0578	0.00172+/-0.001120
Total Córdoba	61	<u>1(26)</u> , <u>2(13)</u> , <u>3(13)</u> , <u>4(10)</u> , <u>6(4)</u>	10	10	0.758+/-0.0397	0.00180+/-0.001128
SPU	37	<u>1(10)</u> , <u>2(13)</u> , <u>4(2)</u> , <u>5(6)</u> , <u>6(2)</u> , <u>9(2)</u>	15	10	0.8134/-0.0418	0.00249+/-0.001315
BAG+ZAR	49	<u>1(21)</u> , <u>2(5)</u> , <u>3(15)</u> , <u>4(3)</u> , <u>7(2)</u>	9	8	0.720+/-0.0441	0.00237+/-0.001417
Total Antioquia	86	<u>1(29)</u> , <u>2(17)</u> , <u>3(15)</u> , <u>4(2)</u> , <u>5(6)</u>	19	13	0.773+/-0.0280	0.00271+/-0.001445
Overall Total	147	<u>1(55)</u> , <u>2(23)</u> , <u>3(28)</u> , <u>4(12)</u> , <u>5(6)</u> , <u>6(4)</u>	23	20	0.781+/-0.0302	0.00234+/-0.001288

S: number of segregating sites, *h*: number of haplotypes, *Hd*: haplotypic diversity, *Pi*: nucleic diversity, *SD*: standard deviation. Numbers inside parenthesis are the frequency of haplotypes by locality. Underline haplotypes are share among populations of Antioquia and Córdoba. Montelbano (MTL), San Pedro de Urabá (SPU), El Bagre (BAG), Zaragoza (ZAR).

Table III
Pairwise genetic differentiation - F_{ST} and gene flow - N_m among *An. nuneztovari* s.l. collected from Antioquia and Córdoba, Colombia

Localities	<i>COI</i>			
	PTL	MTL	BAG+ZAR	SPU
PTL	—	α	6.02	4.09
MTL	-0.02020	—	6.88	3.13
BAG+ZAR	0.07657	0.06767	—	3.12
SPU	0.10872	0.13743	0.13778	—

Above diagonal: N_m value, below diagonal: F_{ST} values, * $p < 0.05$, α : infinity. San Pedro de Urabá (SPU), El Bagre (BAG), Zaragoza (ZAR), Montelíbano (MTL), Puerto Libertador (PTL).

Table IV
Analysis of molecular variance (AMOVA) of *An. nuneztovari* s.l. based on *COI* haplotypes

Source of variation	Variance components	Percentage of variation	Fixation Index
Among populations	0.1896	12.49	$F_{ST} = 0.12^*$
Within populations	1.3234	87.51	

* not significant ($p > 0.05$)