

PUBLISHER'S NOTE: Based on the information available to BioMed Central, this article was apparently handled by Waseem Jerjes, first author of the article and Editor-in-Chief of the journal at that time. The manuscript was reviewed by one recent co-author of the authors and was accepted without revision. BioMed Central asked for the ethics committee reference number. UCL clarified that the ethical approval reference related to data published in this article is LREC reference 07/Q0405/1.

RESEARCH

Open Access

The effect of tobacco and alcohol and their reduction/cessation on mortality in oral cancer patients: short communication

Waseem Jerjes^{1,2,3*}, Tahwinder Upile^{4,5}, Hani Radhi^{2*}, Aviva Petrie⁶, Jesuloba Abiola¹, Aidan Adams³, Panagiotis Kafas⁷, Jacqueline Callear³, Ramin Carbiner⁵, Kartic Rajaram³ and Colin Hopper^{1,5}

Abstract

Background: The use of tobacco is known to increase the incidence of developing oral cancer by 6 times, while the additive effect of drinking alcohol further increases the risk leading to higher rate of morbidity and mortality. In this short communication, we prospectively assessed the effect of tobacco smoking and alcohol drinking in oral cancer patients on the overall mortality from the disease, as well as the effect of smoking and drinking reduction/cessation at time of diagnosis on mortality in the same group.

Materials and methods: A cohort, involved 67 male patients who were diagnosed with oral squamous cell carcinoma, was included in this study. The smoking and drinking habits of this group were recorded, in addition to reduction/cessation after diagnosis with the disease. Comparisons were made to disease mortality at 3 and 5 years.

Results: Follow-up resulted in a 3-year survival of 46.8% and a 5-year survival of 40.4%. Reduction of tobacco smoking and smoking cessation led to a significant reduction in mortality at 3 ($P < 0.001$) and 5 ($P < 0.001$) years. Reduction in drinking alcohol and drinking cessation led to a significant reduction in mortality at 3 ($P < 0.001$) and 5 ($P < 0.001$) years.

Conclusion: Chronic smoking and drinking does have an adverse effect on patients with oral cancer leading to increased mortality from cancer-related causes. Reduction/cessation of these habits tends to significantly reduce mortality in this group of patients. Smoking and drinking cessation counseling should be provided to all newly diagnosed oral cancer patients.

Introduction

Cancers of the oral cavity and the oro-pharyngeal region, a subtype of head and neck, continue to be the 6th most common cancer in the world. They may arise as a primary disease or as a secondary disease by extension from regional or distant areas. The most affected oral sites include the oral tongue, floor of mouth and buccal mucosa [1-5].

This disease is known to affect more males than females, and most commonly reported in the elderly population. The effect of human papilloma virus (HPV) has led to increase in the reports of oral cancer in

younger population. There are a variety of known histological subtypes, but squamous cell carcinomas (SCC) arises in up to 90% of the cases [6-8].

The exact aetiology behind oral cancer is not fully understood, but mutations at the DNA level usually lead to activation of certain oncogenes causing abnormal growth. The most predominant risk factors include tobacco use and alcohol consumption. These risk factors may cause or contribute to the formation of oral leukoplakia, erythroplakia, speckled leukoplakia and other abnormal premalignant oral disorders including submucous fibrosis. Nowadays, the current trends in the spread of HPV16 have led in more oral cancer being diagnosed in non-smokers and non-drinkers [9-11].

The literature is full of case-control and cohort analytic studies establishing a fundamental relationship

* Correspondence: waseem_wk1@yahoo.co.uk; hani_hadi79@hotmail.com

¹UCL Department of Surgery, University College London Medical School, London, UK

²Oral and Maxillofacial Surgery Unit, AL-Mustansiriya University, Baghdad, Iraq
Full list of author information is available at the end of the article

between patients using tobacco and alcohol and developing oral cancer. Smoking tobacco is known to increase the incidence of developing oral cancer by 6 times, while the additive effect of drinking alcohol further increases the risk leading to higher rate of morbidity and mortality [12-23].

Chronic smokers and drinkers are known to succumb from this disease much faster than non-smokers, non-drinkers oral cancer patients. Also, chewing betel, paan and Areca is known to be risk factor as well. Smokeless tobacco contains nicotine and nitrosamines and is usually marketed for oral or nasal use; several studies have provided evidence linking it to oral cancer [12-23].

In this short communication, a small cohort of patients with oral squamous cell carcinoma (SCC) that underwent surgical resection for their pathologies were followed up for a minimum period of 5 years. The aim of this study was to prospectively assess the effect tobacco and alcohol consumption on mortality from this disease; also the effect of smoking and alcohol reduction/cessation at time of diagnosis on survival.

Materials and methods

In this prospective study, identical treatment protocols were used to treat 67 patients with oral SCC referred to the Department of Oral and Maxillofacial Surgery, University College Hospital, London between 1998 and 2003.

All applications were accompanied by multidisciplinary team recommendation, ethical approval, and informed patient consent. The patients' data were entered onto proformas, which were validated and checked by interval sampling. The patients' data included a range of clinical and operative variables. Data collection also included smoking and drinking status, smoking and drinking reduction/cessation at time of diagnosis, comorbidities, recurrence, cause of death, date of death, and last clinic review. The wording "chronic" was used for smokers and drinkers with an ongoing habit for > 20 years.

Demographics of the patients included in the study are highlighted in Table 1. The patient population comprised 67 males. Their mean age at the first diagnosis of Oral SCC was 62.2 years (SD 15.6, min. 25, max. 96 years). Most of the patients were Caucasians (82.1%); other prominent racial groups included Indians (10.4%) and Middle-Easterns (4.5%).

Clinical presentation was mostly an ulcer (74.6%) or a papule (20.9%). Primary sites were mainly identified in the tongue (40.3%), floor of mouth (FOM, 20.9%), lower alveolus (11.9%) and buccal mucosa (7.5%).

Tobacco smoking status was categorised into non-smokers (17.9%), ex-smokers (9.0%), and 5 categories of chronic smokers (i) < 5 cig/day (5.5%), (ii) 5-9 cig/day (10.9%), (iii) 10-14 cig/day (1.8%), (iv) 15-9 cig/day (1.8%) and (v) \geq 20 cig/day (67.3%). Betel nut chewing was

reported by 6 (9.0%) patients. Alcohol drinking status was categorised into non-drinkers (29.9%), ex-drinkers (1.5%), and 3 categories of chronic drinkers (i) < 10 units/week (4.3%), (ii) 11-20 units/week (19.1%) and (iii) > 20 units/week (74.5%). Advice on smoking and drinking and betel nut chewing reduction/cessation was given to all chronic users prior to surgery and their smoking and drinking status was followed-up to assess outcome.

Prior to admission, patients were fully staged. Current staging protocols in our unit requires each patient to undergo magnetic resonance imaging (MRI) of the head and neck, ultrasonographic (US) assessment of the neck nodes, and computed tomography (CT) of the chest and upper abdomen. Clinical staging showed that 13 patients had T1N0 disease, while 10 patients had T2N0 disease; Stage IV tumour was reported in 44 patients (65.7%). Tumour grading was also carried out and reported prior to treatment. Pathological analysis revealed that 13 patients had well differentiated SCC, 4 patients with well-moderately differentiated SCC, 40 patients had moderately differentiated disease, 5 had moderate-poorly differentiated carcinoma and another 5 patients had poorly differentiated disease. Nearly 60% of the cohort received postoperative radiotherapy. Each of the patients was followed-up for a minimum of 5 years.

Statistical analysis

The outcomes of the categorical clinico-pathological variables were summarized as frequencies and percentages for the whole group of patients and the recurrence group. The numerical variables: 'age at 1st diagnosis of SCC', 'depth of invasion (mm)', 'days in ICU post-treatment' and 'days in hospital post-treatment' were summarized by the median, minimal, and maximal values. Fisher's exact test were used to test for statistical significance of the findings, especially the effect of smoking and drinking reduction/cessation on survival at 3 and 5 years.

Results

Tumour clearance was definitively achieved in 39 (58.2%) patients; unfortunately, tumour recurred in 26 patients and was treated by further resection and/or radiotherapy. The status of the surgical margins showed non-cohesive invasion in 34 (50.7%) patients, dysplasia at margin in 19 (28.4%) patients, and presence of severe dysplasia in 33 (49.3%) patients with mean depth of tumour invasion of 10.0 ± 6.5 mm (Min. 1.2, Max. 35.0). Vascular invasion was evident in 3 (4.5%) patients, while nerve invasion was identified only in a patient (Table 1).

The median days of stay in ICU was 1 day while the median days of stay in hospital was 12 days. Follow-up resulted in a 3-year survival of 46.8% and a 5-year survival of 40.4%. The causes of death were either tumour related (i.e. loco-regional or distant metastasis) or non-tumour

Table 1 Demographics of the patients included in the study

	No. of patients (%)	Secondary site (Cont.)	No. of patients (%)	Tumour clearance	No. of patients (%)
Age at 1st presentation					
Mean	62.2	Maxillary tuberosity	1 (1.5)	Positive margins	28 (41.8)
Std. Deviation	15.6	Retromolar area	1 (1.5)	Negative margins	39 (58.2)
Minimum	24.0	Hard palate	1 (1.5)		
Maximum	96.0	Buccal mucosa	5 (7.5)	Depth of invasion (mm)	
		Tonsil	2 (3.0)	Median	10.0
ASA		Smoking status		Minimum	1.2
I	24 (35.8)	Non-smoker	12 (17.9)	Maximum	35.0
II	31 (46.3)	Ex-smoker	6 (9.0)	Dysplasia at margin	19 (28.4)
III	12 (17.9)	Chronic Smoker (< 5 cig/day)	3 (5.5)	Presence of severe dysplasia	33 (49.3)
Ethnicity		Chronic Smoker 9 cig/day)	(5- 6 (10.9)	Non-cohesive Invasive front	34 (50.7)
Caucasian	55 (82.1)	Chronic Smoker (10-14 cig/day)	1 (1.8)	Vascular invasion	3 (4.5)
Indian	7 (10.4)	Chronic Smoker (15-19 cig/day)	1 (1.8)	Nerve invasion	1 (1.5)
Middleeastern	3 (4.5)	Chronic Smoker (≥ 20 cig/day)	37 (67.3)	Bone/cartilage invasion	0 (0.0)
African	2 (3.0)	Advice on smoking cessation		Days in ICU	
Clinical presentation		Reduction of smoking	12 (21.8)	Median	1
Ulcer	50 (74.6)	Cessation of Smoking	13 (23.6)	Minimum	0
Macule	2 (3.0)	Betel nut chewing	6 (9.0)	Maximum	11
Papule	14 (20.9)	Drinking status		Days in Hospital	
Nodule	1 (1.5)	Non-drinker	20 (29.9)	Median	12
Primary site		Ex-drinker	1 (1.5)	Minimum	1
Lateral tongue	25 (37.3)	Chronic drinker (< 10 u/week)	2 (4.3)	Maximum	90
FOM	14 (20.9)	Chronic drinker (10-20 u/week)	9 (19.1)	Post-surgical radiotherapy	40 (59.7)
Lower alveolus	8 (11.9)	Chronic drinker (> 20 u/week)	35 (74.5)	Recurrence	26 (38.8)
Retromolar area	4 (6.0)	Advice on drinking cessation		Survived 3 years	22 (46.8)
Upper alveolus	4 (6.0)	Alcohol reduction	15 (31.9)	Survived 5 years	19 (40.4)
Dorsal tongue	1 (1.5)	Alcohol cessation	9 (19.1)		
Buccal mucosa	5 (7.5)	Tumour staging			
Ventral tongue	1 (1.5)	I	13 (19.4)		
Lower lip	2 (3.0)	II	10 (14.9)		
Soft palate	1 (1.5)	III	3 (4.5)		
Hard palate	1 (1.5)	IV	41 (61.2)		
Tonsil	1 (1.5)	Differentiation			
Secondary site		Well	13 (19.4)		
No site	44 (65.7)	Well-moderate	4 (6.0)		
Lower alveolus	5 (7.5)	Moderate	40 (59.7)		
Ventral tongue	2 (3.0)	Moderate-poor	5 (7.5)		
Soft palate	1 (1.5)	Poor	5 (7.5)		
FOM	5 (7.5)				

Table 2 Tobacco smoking, betel nut chewing and alcohol drinking vs. mortality at 3 and 5 years.

Category	3-years survival	5-years survival	Cause of death
Non-smoker	9/12	8/12	Regional met
Ex-smoker	4/6	3/6	Non-cancer related
Chronic Smoker (< 5 cig/day)	3/3	2/3	Regional met
Chronic Smoker (5-9 cig/day)	2/6	2/6	Regional met
Chronic Smoker (10-14 cig/day)	0/1	0/1	Regional met
Chronic Smoker (15-19 cig/day)	0/1	0/1	Regional met
Chronic Smoker (≥ 20 cig/day)	4/37 (P < 0.001)	4/37 (P < 0.001)	Regional and distant met
Reduction of smoking	10/12 (P < 0.001)	9/12 (P < 0.001)	Regional and distant met
Cessation of smoking	11/13 (P < 0.001)	10/13 (P < 0.001)	Regional and distant met
Betel chewing	2/6	1/6	Regional and distant met
Non-drinker	9/20	9/20	Cancer and non-cancer
Ex-drinker	1/1	1/1	Cancer and non-cancer
Chronic drinker (< 10 u/week)	2/2	2/2	Cancer and non-cancer
Chronic drinker (10-20 u/week)	3/9	4/9	Non-cancer related
Chronic drinker (> 20 u/week)	7/35	3/35	Regional and distant met
Alcohol reduction	10/15 (P < 0.001)	9/15 (P < 0.001)	Non-cancer related
Alcohol cessation	8/9 (P < 0.001)	7/9 (P < 0.001)	Regional met

Also the effect of smoking and drinking reduction/cessation on the mortality. Causes of death have been identified

related (e.g. pneumonia or any other cause that led ultimately to cardiorespiratory failure).

Advice on tobacco smoking reduction and cessation showed that 12 chronic smokers reduced their smoking habits to less than 5 cig/day and 13 chronic smokers stopped smoking immediately after diagnosis. All six patients with betel nut chewing habits stopped after being diagnosed with the disease. Advice on alcohol drinking reduction and cessation showed that 15 chronic drinkers reduced their alcohol intake to less than 10 units/week and 9 patients stopped completely after being diagnosed.

The causes of death of patients according to the smoking, betel nut chewing and drinking status are detailed in Table 2. The reduction of smoking and/or smoking cessation lead to significant reduction in mortality at 3 (P < 0.001) and 5 (P < 0.001) years. Also, the reduction in drinking alcohol and/or drinking cessation lead to significant reduction of mortality at 3 (P < 0.001) and 5 (P < 0.001) years.

Discussion and conclusion

At least three quarters of oral cancers could be prevented by the elimination of tobacco smoking and reduction in alcohol consumption. The removal of these two risk factors also reduces the risk of recurrence or second primary in people with oral cancer. Smoking cessation is associated with a rapid decline in the risk of oral cancers, with a 50% reduction in risk within 3 to 5 years [24].

One of the most effective ways of helping patients prevent or reduce the morbidity and mortality effects of oral cancer is by promoting tobacco and alcohol abandonment.

A need for further promotion of smoking cessation activities by the health professionals was identified [25]. In particular, smoking cessation programs are needed to prevent the increase in mortality from these cancers in many low- and middle-income countries [26].

Dental and medical practitioners have a unique opportunity to lead the advancement in this field aiming to reduce the global effect of oral and oro-pharyngeal and laryngeal cancers.

Reduction or cessation of smoking soon after diagnosis reduced the rate of death significantly (P < 0.001). The same principle applies for cessation of alcohol drinking. Chewing betel can be associated with very poor prognosis and death from loco-regional or distant metastasis most likely due to a wide spread field effect.

Author details

¹UCL Department of Surgery, University College London Medical School, London, UK. ²Oral and Maxillofacial Surgery Unit, AL-Mustansiriya University, Baghdad, Iraq. ³Leeds Institute of Molecular Medicine, School of Medicine, University of Leeds, Leeds, UK. ⁴Department of Otolaryngology/Head and Neck Surgery, Chase Farm & Barnet NHS Trust, Enfield, UK. ⁵Head & Neck Unit, University College London Hospital, London, UK. ⁶Biostatistics Unit, UCL Eastman Dental Institute, London, UK. ⁷Department of Oral Surgery and Radiology, School of Dentistry, Aristotle University, Thessalonica, Greece.

Authors' contributions

All authors have contributed intellectually and to the writing of this manuscript. AP: contributed to the primary analysis of this study and the results are highlighted in Table 1 of this study. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 11 January 2012 Accepted: 12 March 2012
 Published: 12 March 2012

References

1. Bagan JV, Scully C: Recent advances in Oral Oncology 2007: epidemiology, aetiopathogenesis, diagnosis and prognostication. *Oral Oncol* 2008, **44**(2):103-108.
2. Warnakulasuriya S: Global epidemiology of oral and oropharyngeal cancer. *Oral Oncol* 2009, **45**(4-5):309-316.
3. Garzino-Demo P, Dell'Acqua A, Dalmaso P, Fasolis M, La Terra Maggiore GM, Ramieri G, Berrone S, Rampino M, Schemi M: Clinicopathological parameters and outcome of 245 patients operated for oral squamous cell carcinoma. *J Craniomaxillofac Surg* 2006, **34**(6):344-350.
4. Parkin DM, Bray F, Ferlay J, Pisani P: Estimating the world cancer burden: Globocan 2000. *Int J Cancer* 2001, **94**(2):153-156.
5. Llewellyn CD, Johnson NW, Warnakulasuriya KAAS: Risk factors for squamous cell carcinoma of the oral cavity in young people - a comprehensive literature review. *Oral Oncol* 2001, **37**:401-418.
6. Garavello W, Spreafico R, Gaini RM: Oral tongue cancer in young patients: a matched analysis. *Oral Oncol* 2007, **43**(9):894-897.
7. Warnakulasuriya S, Mak V, Möller H: Oral cancer survival in young people in South East England. *Oral Oncol* 2007, **43**(10):982-986.
8. Scully C, Bagan JV: Recent advances in oral oncology 2008; squamous cell carcinoma imaging, treatment, prognostication and treatment outcomes. *Oral Oncol* 2009, **45**(6):e25-e30.
9. Scully C, Bagan J: Oral squamous cell carcinoma overview. *Oral Oncol* 2009, **45**(4-5):301-308.
10. Woolgar JA: Histopathological prognosticators in oral and oropharyngeal squamous cell carcinoma. *Oral Oncol* 2006, **42**(3):229-239.
11. Platz H, Fries R, Hudec M, Min Tjoa A, Wagner RR: The prognostic relevance of various factors at the time of the first admission of the patient. Retrospective DOSAK study on carcinoma of the oral cavity. *J Maxillofac Surg* 1983, **11**(1):3-12.
12. Rothman K, Keller A: The effect of joint exposure to alcohol and tobacco on the risk of cancer of the mouth and pharynx. *J Chron Dis* 1972, **25**:711-716.
13. Wynder EL, Mushinski MH, Spivak JC: Tobacco and alcohol consumption in relation to the development of multiple primary cancers. *Cancer* 1977, **40**:1872-1878.
14. Wigle DT, Mao Y, Grace M: Relative importance of smoking as a risk for selected cancers. *Can J Pub Health* 1980, **71**:269-275.
15. Winn DM, Blot WJ, Shy CM, Pickle LW, Toledo A, Fraumeni JF Jr: Snuff dipping and oral cancer among women in the southern United States. *N Engl J Med* 1981, **304**:745-749.
16. Winn DM: Smokeless tobacco and cancer: the epidemiologic evidence. *CA Cancer J Clin* 1988, **38**:236-243.
17. Zheng TZ, Boyle P, Hu HF, Duan J, Jiang PJ, Ma DQ, Shui LP, Niu SR, MacMahon B: Tobacco smoking, alcohol consumption and risk of oral cancer: a case-control study in Beijing, People's Republic of China. *Cancer Causes Control* 1990, **1**:173-179.
18. Choi SY, Kahyo H: Effect of cigarette smoking and alcohol consumption in the etiology of cancer of the oral cavity, pharynx, and larynx. *Int J Epidemiol* 1991, **20**:878-885.
19. Kato I, Nomura AM, Stemmermann GN, Chyou PH: Prospective study of the association of alcohol with cancer of the upper aerodigestive tract and other sites. *Cancer Causes Control* 1992, **3**:145-151.
20. Johnson NW, Bain CA: Tobacco and oral disease. EU-Working Group on Tobacco and Oral Health. *Br Dent J* 2000, **189**(4):200-6.
21. Sanderson RJ, Ironside JA: Squamous cell carcinomas of the head and neck. *BMJ* 2002, **325**(7368):822-827.
22. Znaor A, Brennan P, Gajalakshmi V, Mathew A, Shanta V, Varghese C, Boffetta P: Independent and combined effects of tobacco smoking, chewing and alcohol drinking on the risk of oral, pharyngeal and esophageal cancers in Indian men. *Int J Cancer* 2003, **105**(5):681-686.
23. Cogliano V, Straif K, Baan R, Grosse Y, Secretan B, El Ghissassi F: Smokeless tobacco and tobacco-related nitrosamines. *Lancet Oncol* 2004, **5**(12):708.
24. Some JM: The health benefits of smoking cessation. *Med Clin North Am* 1992, **76**(2):399-414.
25. Saito A, Nishina M, Murai K, Mizuno A, Ueshima F, Makiishi T, Ichinohe T: Health professional's perceptions of and potential barriers to smoking cessation care: a survey study at a dental school hospital in Japan. *BMC Res Notes* 2010, **7**(3):329.
26. Adair T, Hoy D, Dettrick Z, Lopez AD: Trends in oral, pharyngeal and oesophageal cancer mortality in Australia: the comparative importance of tobacco, alcohol and other risk factors. *Aust N Z J Public Health* 2011, **35**(3):212-219.

doi:10.1186/1758-3284-4-6

Cite this article as: Jerjes et al.: The effect of tobacco and alcohol and their reduction/cessation on mortality in oral cancer patients: short communication. *Head & Neck Oncology* 2012 **4**:6.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

