

Binding of Collagen $\alpha 1$ Chains to Human Platelets

THOMAS M. CHIANG, EDWIN H. BEACHEY, and ANDREW H. KANG

From the Veterans Administration Hospital and the Departments of Biochemistry, Medicine, and Microbiology, University of Tennessee Center for the Health Sciences, Memphis, Tennessee 38104

ABSTRACT We previously reported that purified $\alpha 1$ chains of type I chick skin collagen induce platelet aggregation. We now describe immunological and biochemical evidence that the peptide binds to intact platelets as an early event in the induction of platelet aggregation and the release reaction.

Antibody against $\alpha 1$ (I) was obtained by immunizing rabbits with complete Freund's adjuvant mixed with purified $\alpha 1$. Immunofluorescence studies showed that $\alpha 1$ (I)-treated platelets exhibited strong immunofluorescence. The intensity of fluorescence was markedly decreased by the pretreatment of platelets with $\alpha 1$ -CB5 and glucosylgalactosylhydroxylysine.

Dose-response curves of platelet aggregation induced by $\alpha 1$ and the binding of $\alpha 1$ by washed intact platelets are correlated. The biochemical studies showed that the binding of the $\alpha 1$ chain to washed intact platelets was platelet concentration and temperature dependent, and that it reached a maximum in 10 min. The process was reversible and specific, with an association constant of 1.7 μ M. The inhibitor of $\alpha 1$ -induced platelet aggregation, glucosylgalactosylhydroxylysine, inhibited the $\alpha 1$ binding. These results suggest that $\alpha 1$ (I) chains bind to specific receptor site(s) on platelet membranes to trigger aggregation and the release reaction.

INTRODUCTION

Platelets play an important role in hemostasis and in pathologic thromboses. After blood vessel injury, platelets adhere to the exposed subendothelial connective tissue, collagen in particular (1-5), and aggregate, releasing several biologically active substances. Collagen and other agents, such as ADP, thrombin, epinephrine, and antigen-antibody complexes, have been shown to induce platelets to aggregate and

undergo the "physiologic" release reaction in preparations of platelet-rich plasma (6-13).

It has been suggested that these aggregating agents may initiate these phenomena by acting on the platelet surface. Thus, it has been shown that ADP binds to isolated platelet membranes (14), and that thrombin binds to the membrane of intact platelets (15, 16). The mechanisms whereby collagen may mediate platelet aggregation and the structural features responsible for the interaction have not been completely clarified. Previously, we reported that the purified, denatured $\alpha 1$ (I) chain of chick skin collagen is capable of causing platelet aggregation and the release reaction, and that the carbohydrate moiety located in one of the CNBr peptides, $\alpha 1$ (I)-CB5, is involved in the interaction (17-19). Inhibition of adenylate cyclase of platelet membranes by collagen and the $\alpha 1$ chain (19) suggested the possibility that these substances act directly on the platelet surface. In the present paper, we report evidence to indicate that $\alpha 1$ chains bind specifically and reversibly to the intact platelet surface, and that the sugar moiety, glucosylgalactosylhydroxylysine (Glc-Gal-Lys[OH]),¹ is involved in the interaction. The close correlation between the degree of binding and aggregation supports the hypothesis that the binding of $\alpha 1$ to membrane receptor(s) may be an initial event which culminates in platelet aggregation.

METHODS

Preparation of washed human platelets. Blood was collected from normal volunteers after they had fasted overnight and placed in polypropylene tubes containing 0.1 vol of 3.8% sodium citrate. Platelet-rich plasma was prepared by centrifuging the citrated blood at room temperature for 15 min at 226 g (18). The contaminating erythrocytes were removed by centrifugation at 1,500 g for 1 min. The erythrocyte-free platelet-rich plasma was then spun at 2,000 g for 15 min in the presence of 0.001 M EDTA. The platelets

Dr. Kang is a Medical Investigator of the Veterans Administration.

Received for publication 20 August 1976 and in revised form 22 November 1976.

¹Abbreviations used in this paper: Glc-Gal-Lys(OH), glucosylgalactosylhydroxylysine; PBS, 0.02 M phosphate/0.15 M, NaCl, pH 7.4; Tris-EDTA, 0.01 M Tris/0.001 M EDTA per 0.15 M NaCl, pH 7.5.




FIGURE 1 Human platelets treated with $\alpha 1(I)$ chain (A) or soluble tropocollagen (B) prepared from chick skin followed by rabbit anti- $\alpha 1$ or antitropocollagen, respectively, and then by fluorescein-conjugated goat antirabbit IgG. Preincubation of platelets with Glc-Gal-Lys(OH) before incubation with $\alpha 1$ chain markedly reduced immunofluorescence (C) which was comparable to control platelets incubated with PBS rather than $\alpha 1$ or tropocollagen.

were washed twice with 0.01 M Tris/0.001 M EDTA per 0.15 M NaCl, pH 7.5 (Tris-EDTA), and then resuspended in the same buffer at the concentration of 300,000–400,000/mm³. The washed platelets were kept in an ice bath until used. The experiments were performed within 3 h after collection.

Preparation of chick skin collagen, $\alpha 1$ chains, and CNBr peptides. Neutral salt-soluble collagen was extracted from the skin of 3-wk-old, White Leghorn chicks which had been rendered lathyritic by administration of β -aminopropionitrile (Aldrich Chemical Co. Inc., Milwaukee, Wis.) for 2 wk.

Extracted collagen was purified by repeated differential precipitation with NaCl as previously described (20). The $\alpha 1$ chain was prepared from purified lathyritic collagen by chromatography on carboxymethylcellulose (20). For preparation of radiolabeled $\alpha 1$, the animals had been injected intraperitoneally with 100 μ Ci of [¹⁴C]glycine (sp act, 49.73 μ Ci/mmol, New England Nuclear, Boston, Mass.) in 1 ml of sterile 0.9% NaCl once a day for 3 days before they were killed.

The CNBr peptides of the $\alpha 1$ chain were prepared by a combination of ion-exchange and molecular-sieve chroma-




FIGURE 2 Time-dependent binding of ^{14}C -labeled chick skin $\alpha 1$ by platelets. $25 \mu\text{g}$ of ^{14}C -labeled $\alpha 1$ was incubated with the washed platelets (1×10^6) at 37°C . At the time indicated, platelet-bound ^{14}C -labeled $\alpha 1$ was isolated by Millipore (Millipore Corp.) filtration. Each point is the mean of duplicates. All values were corrected for nonspecific binding.

tography of CNBr digests of the $\alpha 1$ chain on various resins. The details of these procedures as well as the criteria of their purity have been described in detail elsewhere (21).

Preparation of antiserum against $\alpha 1$ chain and collagen. Antisera were raised in rabbits immunized with 1 mg of collagen or $\alpha 1$ chain emulsified in 1.0 ml of complete Freund's adjuvant (Difco Laboratories, Detroit, Mich.). 4 wk later, the animals were boosted with 1 mg of alum-precipitated (22) collagen or $\alpha 1$ chain injected intraperitoneally at weekly intervals for 3 wk and again on wk 10, followed by an intravenous injection of the same dose on wk 11. The rabbits were bled 7 days after the last injection. Sera were stored at 4°C . Antibody titers were assayed by passive hemagglutination of tanned human group O erythro-

cytes coated with collagen or $\alpha 1$ chain (23). The antibodies were partially purified by quarternary-aminoethyl-Sephadex (Pharmacia Fine Chemicals, Inc., Piscataway, N. J.) column chromatography as previously described (24).

Immunofluorescence studies. Smears of washed platelets were prepared on glass microscope slides. The dried smears were washed with 0.02 M phosphate/0.15 M NaCl, pH 7.4 (PBS), and fixed with 1% paraformaldehyde for 1 min. The washed, fixed smears were treated with 1-mg/ml solutions of collagen or $\alpha 1$ chain in PBS for 20 min at room temperature, washed with PBS twice, incubated with antiserum either to collagen or $\alpha 1$ chain for 20 min at room temperature, washed twice again with PBS, and finally treated with a 0.1 dilution of fluorescein-conjugated, goat anti-rabbit IgG for 20 min. At the end of incubation, the smears were washed with PBS and mounted with glass coverslips using one drop of Gelvatol (Monsanto Co., St. Louis, Mo.). The smears were examined with a fluorescence microscope (Carl Zeiss, N. Y.).

Binding experiments. Assays for the binding of the ^{14}C -labeled $\alpha 1$ chain to washed platelets were performed in Tris-EDTA at various temperatures. The assay mixtures consisted of $25 \mu\text{g}$ of [^{14}C]glycine-labeled $\alpha 1$ chain in 0.1 ml of Tris-EDTA and various amounts of washed platelets for various periods of time as indicated in the text. In some experiments, various other proteins or peptides were added to the mixtures as specified. After specified intervals, incubations were terminated by passing the mixtures through a $0.45\text{-}\mu\text{m}$ Millipore filter (Millipore Corp., Bedford, Mass.) under reduced pressure. The platelets were washed with 10 ml of 0.05 M Tris, pH 7.5, containing 0.5% bovine serum albumin, and counted in a Nuclear-Chicago scintillation counter (Nuclear-Chicago, Des Plaines, Ill.) using Aquasol (New England Nuclear) as solvent.

Identification of platelet-bound radioactivity. The bound ^{14}C -labeled $\alpha 1$ chain was dissociated by soaking the filter in 20 ml of 0.05 M Tris, pH 7.5, containing 1.25 mg/ml of unlabeled $\alpha 1$ at 37°C for 1 h with constant shaking. Approximately 80% of the total radioactivity was recovered in the extract.

A portion of the lyophilized extract was dissolved in 2 ml




FIGURE 3 Dissociation of platelet-bound $25 \mu\text{g}$ of ^{14}C -labeled $\alpha 1$ was preincubated with the washed platelets (6×10^6) in $100 \mu\text{l}$ of Tris-EDTA buffer for 30 min at 37°C . At 0 time, 1.25 mg of unlabeled $\alpha 1$ dissolved in 0.2 ml of the same buffer was added and incubated for the additional indicated times (●). To the controls was added 0.2 ml of the buffer only (○). Each point represents separate duplicate incubations.




FIGURE 4 The ratio, bound/free ^{14}C -labeled $\alpha 1$, is plotted as a function of bound $\alpha 1$ according to the method of Scatchard (26). B, amount of $\alpha 1$ bound (pmol/ 7×10^7 platelets) to platelets; F, amount of free $\alpha 1$ in micromoles. The intercept with the abscissa, 67 pmol/ 7×10^7 platelets, corresponds to an approximation of the binding capacity of platelets at saturation. From the Avogadro number (6.02×10^{23}), the number of binding sites per platelet was calculated to be approximately 600,000.

of 0.01 M Tris/1 M CaCl_2 , pH 7.4, applied to a 2×110 -cm column of agarose A 1.5 m (Bio-Rad Laboratories, Richmond, Calif.), and eluted with the same buffer. 3-ml fractions were collected. The remaining extract was dissolved in 5 ml of 0.05 M sodium acetate, pH 4.8, together with 10 mg of unlabeled $\alpha 1$ to serve as carrier. The mixture was applied to a 2×10 -cm column of carboxymethylcellulose which had been equilibrated with the same buffer at 43°C . The column was eluted with a linear gradient of NaCl from 0 to 0.1 M over a total volume of 800 ml and 13-ml fractions were collected. The optical density (230 nm) of each of the effluent fractions was determined using a Zeiss spectrophotometer and the radioactivity was assayed in the scintillation counter.

Determination of protein concentration. The concentrations of collagen-derived peptides were determined by hydroxyproline analyses after hydrolysis as described previously (25). The platelet counts were performed by phase-contrast microscopy.

RESULTS

Immunofluorescence studies. Evidence that $\alpha 1$ chain of collagen binds to the intact human platelet was obtained by immunofluorescence studies. The $\alpha 1$ chain-treated platelets fluoresced intensely and uniformly (Fig. 1A) whereas tropocollagen-treated platelets fluoresced unevenly (Fig. 1B) as compared to the absence of fluorescence in control platelets treated with PBS. The intensity of immunofluorescence produced by $\alpha 1$ and collagen was markedly reduced by preincubation of platelets with $\alpha 1$ -CB5 or Glc-Gal-Lys(OH) (Fig. 1C), but not with other CNBr peptides, including CB4, CB7, and CB8, suggesting that Glc-Gal-Lys(OH) is a determinant in the binding of $\alpha 1$ by platelets.

Binding of $\alpha 1$ chain to washed human platelets.

The binding of $\alpha 1$ chain to platelets is time (Fig. 2) and platelet concentration dependent (data not shown). The binding of $\alpha 1$ chain to platelets is also temperature dependent; the total amount bound to platelets at 37°C is three to four times higher than that at room temperature and 0°C (data not shown). In addition, the rate of binding at 37°C exceeded that at room temperature and 0°C . In subsequent studies therefore, the binding experiments were performed at 37°C . The amount bound at various temperatures was corrected for the background using the same amount of ^{14}C -labeled $\alpha 1$ chain at parallel points in the absence of platelets.

The binding of the $\alpha 1$ chain to platelets was shown to be a reversible process (Fig. 3). The bound $\alpha 1$ chain was rapidly displaced from platelets by adding a 50-fold excess of unlabeled $\alpha 1$ chain. The undisplaceable radioactivity was considered to represent non-specific binding. In subsequent experiments, therefore, duplicate incubations were performed with and without the addition of 50-fold excess of unlabeled $\alpha 1$ chain. Each value of binding was corrected by subtracting the value obtained in a duplicate incubation mixture containing excess unlabeled $\alpha 1$.

Determination of association constant. To determine the average number of $\alpha 1$ -chain receptor sites per platelet as well as their affinity, the amount of $\alpha 1$

TABLE I
Effects of $\alpha 1$ CNBr Peptides, Glc-Gal-Lys(OH), and Hydroxylysine on ^{14}C -Labeled $\alpha 1$ Binding by Platelets*

Test substances	Control
	%
Control	100
$\alpha 1$, 125 μg	43
$\alpha 2$, 125 μg	40
$\beta 12$, 125 μg	61
CB4, 125 μg	106
CB5, 125 μg	65
CB5, 250 μg	41
CB7, 125 μg	97
CB8, 125 μg	84
BSA \ddagger , 125 μg	110
Glc-Gal-Lys(OH), 125 μg	79
Glc-Gal-Lys(OH), 250 μg	52
Hydroxylysine, 125 μg	115
Hydroxylysine, 250 μg	104

* Washed platelets (6×10^8) were incubated with the CNBr peptides or Glc-Gal-Lys(OH), or hydroxylysine at 37°C for 20 min. After the addition of 25 μg of ^{14}C -labeled $\alpha 1$, the mixtures were incubated at 37°C for an additional 20-min period. The bound $\alpha 1$ was isolated by Millipore filtration. Values are expressed as the percent of control experiments.

\ddagger BSA, bovine serum albumin.




FIGURE 5 Agarose A 1.5 m column chromatography of $\alpha 1$ chains displaced from the washed platelets which had been incubated with ^{14}C -labeled $\alpha 1$. The radioactivity was displaced with a 50-fold excess of unlabeled $\alpha 1$. The dissociation material was lyophilized and redissolved in 0.01 M Tris-1 M CaCl_2 , pH 7.4, applied to a 2×110 -cm column, and eluted with the same buffer. Aliquots of the column effluent fractions were analyzed for radioactivity (○) and optical density at 230 nm (●).

bound to platelets was determined as a function of free $\alpha 1$ added to the system. The resulting data were analyzed by a Scatchard plot (26) (Fig. 4). The number of molecules bound to platelets at saturation was calculated from the intercept with the abscissa. From a number of experiments using different platelet sources, it was determined that there was a single population of approximately 600,000 binding sites per platelet with an association constant of $1.7 \mu\text{M}$.

Inhibition studies. In an attempt to identify the interaction site(s) between $\alpha 1$ chain and platelets,

Glc-Gal-Lys(OH) and $\alpha 1$ -CB5 were tested for their ability to inhibit binding of $\alpha 1$ to intact platelets. The results summarized in Table I demonstrate that these two substances inhibit binding of the $\alpha 1$ chain to platelets. In contrast, the other collagen peptides, CB4, CB7, CB8, as well as hydroxylysine and bovine serum albumin had no effect on the binding of $\alpha 1$.

Identification of the displaced $\alpha 1$ chain. To determine whether $\alpha 1$ was altered or whether it remained unaltered after it became bound to the intact platelet, ^{14}C - $\alpha 1$ chains were first reacted with platelets, and then displaced with a 50-fold excess of unlabeled $\alpha 1$. The dissociated material was mixed with unlabeled, carrier $\alpha 1$ and chromatographed on a column of agarose A 1.5 m. Only one radioactive peak which coeluted with the unlabeled $\alpha 1$ chain was observed (Fig. 5). Recovery of the radioactivity from the chromatogram was 90%. No evidence of heterogeneity in radioactivity in radioactive material was observed. Similar results were obtained by carboxymethylcellulose chromatography of a separate aliquot (figure not shown). These results indicate that the $\alpha 1$ chain remained intact during the binding and displacement reactions.

Relationship of the $\alpha 1$ binding with platelet aggregation. The various concentrations of $\alpha 1$ chain used to induce platelet aggregation (17-19) together with the amounts of $\alpha 1$ bound to washed intact platelets at each concentration of $\alpha 1$ are depicted in Fig. 6. The dose-response curve of $\alpha 1$ chain-induced platelet aggregation demonstrates a direct relationship with $\alpha 1$ chain binding to platelets. A relatively small amount of binding is associated with the functional




FIGURE 6 Comparison of platelet aggregation induced by $\alpha 1$ (●) and the binding of ^{14}C -labeled $\alpha 1$ by washed platelets (○).

response, aggregation. A similar relationship has been observed between hormone binding such as insulin and the biological response of the target cells (27).

DISCUSSION

The immunofluorescence studies reported here indicate that collagen and the $\alpha 1(I)$ chains of chick skin bind to the surface of platelets, and that the binding is inhibited by Glc-Gal-Lys(OH), a known inhibitor of $\alpha 1$ chain-induced platelet aggregation (17, 18). These observations were confirmed by biochemical studies which indicate that binding of $\alpha 1(I)$ to intact platelets is reversible and specific as evidenced by the fact that ^{14}C -labeled $\alpha 1$ chain can be displaced from intact washed platelets by unlabeled $\alpha 1$. Scatchard plot analyses suggest that platelets possess a single population of binding site(s) with an association constant of $1.7 \mu M$ for the interaction with $\alpha 1$, and that at saturation an average of 600,000 molecules of $\alpha 1$ are bound per platelet. The observed association constant is consistent with the amount of $\alpha 1$ necessary to induce platelet aggregation and the release reaction (19). However, it is still possible that there may have been an additional population(s) present in platelets as has been shown in thrombin-binding experiments (15). Because of the limitations in obtaining $\alpha 1$ chain of sufficiently high specific activity by in vivo labeling, it was impossible to examine binding at much lower concentrations of $\alpha 1$ than used in the present study.

Inhibition studies using various collagen-derived peptides indicate that Glc-Gal-Lys(OH) and the glycopeptide, $\alpha 1$ -CB5, interfere with the $\alpha 1$ binding. In contrast, free hydroxylysine, and other CNBr peptides not containing the hydroxylysineglycoside, have no effect. These results are consistent with previously reported findings that Glc-Gal-Lys(OH) inhibits $\alpha 1$ -induced platelet aggregation (17, 19) but not aggregation induced by other agents such as ADP (unpublished observations).

Platelet aggregation occurred after binding of relatively small quantities of $\alpha 1$ to platelets. This is consistent with the relationship reported previously for insulin binding and the biological response of fat cells (27). These observations strongly support the idea that the collagen $\alpha 1$ chain induces platelet aggregation by acting on the platelet membrane. It should be noted, however, that these studies have been carried out using radiolabeled chick skin $\alpha 1$ chains for cogent technical reasons, and that the reactivity of human platelet membrane to collagens of other sources may differ (17, 18). Further studies to identify the receptor sites on platelet membranes may provide useful information both in regard to platelet variations and in

regard to collagen abnormalities in various disease states manifested by abnormal collagen-platelet interactions.

ACKNOWLEDGMENTS

The authors gratefully acknowledge the expert technical assistance of Mr. Bobby J. Curtis.

These studies were conducted under Research Projects 0805-02, 4826-01, and 4826-02 from the Veterans Administration and supported in part by research grants AI-10085 and AM-16506 from the U. S. Public Health Service.

REFERENCES

1. Hughes, J. 1960. Accolement des plaquettes au collagène. *C. R. Seances Soc. Biol. Fil.* **154**: 866-868.
2. Zucker, M. B., and J. Borelli. 1962. Platelet clumping produced by connective tissue suspensions and by collagen. *Proc. Soc. Exp. Biol. Med.* **109**: 779-787.
3. Hovig, T. 1963. Aggregation of rabbit blood platelets produced in vitro by saline "extract" of tendons. *Thromb. Diath. Haemorrh.* **9**: 248-263.
4. Spaet, T. H., J. Cintron, and M. Spivack. 1962. Some properties of the platelet-connective tissue mixed agglutination reaction. *Proc. Soc. Exp. Biol. Med.* **111**: 292-295.
5. Jaffe, R., and D. Deykin. 1974. Evidence for a structural requirement for the aggregation of platelets by collagen. *J. Clin. Invest.* **53**: 875-883.
6. Wilner, G. D., H. L. Nossel, and E. C. LeRoy. 1968. Aggregation of platelets by collagen. *J. Clin. Invest.* **47**: 2616-2621.
7. Holmsen, H., H. J. Day, and H. Stormorken. 1969. The blood platelet release reaction. *Scand. J. Haematol. Suppl.* **8**: 1-26.
8. Ardlie, N. G., M. A. Packham, and J. F. Mustard. 1970. Adenosine diphosphate-induced platelet aggregation in suspensions of washed rabbit platelets. *Br. J. Haematol.* **19**: 7-17.
9. Born, G. V. R. 1962. Aggregation of blood platelets by adenosine diphosphate and its reversal. *Nature (Lond.)* **194**: 927-929.
10. Spaet, T. H., and M. B. Zucker. 1964. Mechanism of platelet plug formation and role of adenosine diphosphate. *Am. J. Physiol.* **206**: 1267-1274.
11. Haslam, R. J. 1964. Role of adenosine diphosphate in the aggregation of human blood-platelets by thrombin and fatty acids. *Nature (Lond.)* **202**: 765-768.
12. Mills, D. C. B., and G. C. K. Roberts. 1967. Effects of adrenaline on human blood platelets. *J. Physiol. (Lond.)* **193**: 443-453.
13. Siqueira, M., and R. A. Nelson, Jr. 1961. Platelet agglutination by immune complexes and its possible role in hypersensitivity. *J. Immunol.* **86**: 516-525.
14. Nachman, R. L., and B. Ferris. 1974. Binding of adenosine diphosphate by isolated membranes from human platelets. *J. Biol. Chem.* **249**: 704-710.
15. Tollefsen, D. M., J. R. Feagler, and P. W. Majerus. 1974. The binding of thrombin to the surface of human platelets. *J. Biol. Chem.* **249**: 2646-2651.
16. Tollefsen, D. M., C. M. Jackson, and P. W. Majerus. 1975. Binding of the products of prothrombin activation to human platelets. *J. Clin. Invest.* **56**: 241-245.
17. Katzman, R. L., A. H. Kang, and E. H. Beachey. 1973. Collagen-induced platelet aggregation: Involvement

- ment of an active glycopeptide fragment ($\alpha 1$ -CB5). *Science (Wash. D. C.)*. **181**: 670-672.
18. Kang, A. H., E. H. Beachey, and R. L. Katzman. 1974. Interaction of an active glycopeptide from chick skin collagen ($\alpha 1$ -CB5) with human platelets. *J. Biol. Chem.* **249**: 1054-1059.
 19. Chiang, T. M., E. H. Beachey, and A. H. Kang. 1975. Interaction of a chick skin collagen fragment ($\alpha 1$ -CB5) with human platelets. Biochemical Studies during the aggregation and release reaction. *J. Biol. Chem.* **250**: 6916-6922.
 20. Kang, A. H., K. A. Piez, and J. Gross. 1969. Characterization of the α chains of chick skin collagen and the nature of the NH₂-terminal cross-link region. *Biochemistry*. **8**: 3648-3655.
 21. Kang, A. H., K. A. Piez, and J. Gross. 1969. Characterization of the cyanogen bromide peptides from the $\alpha 1$ chain of chick skin collagen. *Biochemistry*. **8**: 1506-1514.
 22. Chase, M. W., E. A. Kabat, and A. A. Benedict. 1967. Production of antiserum. *Methods Immunol. Immunochem.* **1**: 197-306.
 23. Avrameas, S., B. Taudou, and S. Chuilon. 1969. Glutaraldehyde, cyanuric chloride and tetraazotized o-dianisidine as coupling reagents in the passive hemagglutination test. *Immunochemistry*. **6**: 67-76.
 24. Bisno, A. L., I. Ofek, E. H. Beachey, R. W. Chandler, and J. W. Curran. 1975. Human immunity to *Neisseria gonorrhoeae*: Acquired opsonic antibodies. *J. Lab. Clin. Med.* **86**: 221-229.
 25. Bergman, I., and R. Loxley. 1963. Two improved and simplified methods for the spectrophotometric determination of hydroxyproline. *Anal. Chem.* **35**: 1961-1965.
 26. Scatchard, G. 1949. The attractions of proteins for small molecules and ions. *Ann. N. Y. Acad. Sci.* **51**: 660-672.
 27. Olefsky, J. M. 1976. The effects of spontaneous obesity on insulin binding, glucose transport, and glucose oxidation of isolated rat adipocytes. *J. Clin. Invest.* **57**: 842-851.