

RESEARCH

Open Access

Loss of HLTF function promotes intestinal carcinogenesis

Sumit Sandhu¹, Xiaoli Wu¹, Zinnatun Nabi¹, Mojgan Rastegar^{1,3,4}, Sam Kung³, Sabine Mai^{1,2} and Hao Ding^{1*}

Abstract

Background: HLTF (Helicase-like Transcription Factor) is a DNA helicase protein homologous to the SWI/SNF family involved in the maintenance of genomic stability and the regulation of gene expression. HLTF has also been found to be frequently inactivated by promoter hypermethylation in human colon cancers. Whether this epigenetic event is required for intestinal carcinogenesis is unknown.

Results: To address the role of loss of HLTF function in the development of intestinal cancer, we generated *Hltf* deficient mice. These mutant mice showed normal development, and did not develop intestinal tumors, indicating that loss of *Hltf* function by itself is insufficient to induce the formation of intestinal cancer. On the *Apc*^{min/+} mutant background, *Hltf* deficiency was found to significantly increase the formation of intestinal adenocarcinoma and colon cancers. Cytogenetic analysis of colon tumor cells from *Hltf*^{-/-}/*Apc*^{min/+} mice revealed a high incidence of gross chromosomal instabilities, including Robertsonian fusions, chromosomal fragments and aneuploidy. None of these genetic alterations were observed in the colon tumor cells derived from *Apc*^{min/+} mice. Increased tumor growth and genomic instability was also demonstrated in HCT116 human colon cancer cells in which HLTF expression was significantly decreased.

Conclusion: Taken together, our results demonstrate that loss of HLTF function promotes the malignant transformation of intestinal or colonic adenomas to carcinomas by inducing genomic instability. Our findings highly suggest that epigenetic inactivation of HLTF, as found in most human colon cancers, could play an important role in the progression of colon tumors to malignant cancer.

Keywords: HLTF, Mouse gene-targeting, Adenomatous polyposis coli (*Apc*), Intestinal adenocarcinoma, Colonic tumor or cancer, Chromosomal instability, HCT116 cells

Background

Human colon cancer is the second leading cause of cancer-related death in developed countries. About 50% of the Western population develops adenomatous polyps (a benign colon tumor) by the age of 70, and the lifetime risk for colon cancer is estimated to be 5% [1]. The formation of colon cancer involves a multiple-step process, starting from a small adenomatous polyp and followed by the development of a large adenoma with dysplasia that ultimately leads to the formation of invasive carcinoma (see the recent review by Fearon ER [2]). It is widely accepted that most human colon cancers are initiated by the inactivation of the Adenomatous

Polyposis Coli (APC)/Wnt signaling pathway and then progress as the result of a series of mutational activation of oncogenes coupled with the inactivation of tumor-suppressor genes [2,3]. Apart from genetic mutations, epigenetic alterations, particularly aberrant CpG island methylation, have been demonstrated as a major alternative mechanism for suppressing gene function during the development of colon cancer [4-6]. To date, many genes that are epigenetically silenced in human colon cancers as well as in colonic adenomas have been identified. However, the function of many of these genes in colon carcinogenesis is still largely unknown. In this study, we have characterized the role of one of these methylated genes, termed Helicase-like Transcription Factor (HLTF), in intestinal carcinogenesis.

HLTF (SMARCA3 in OMIM) is homologous to the SWI/SNF family of chromatin remodelers [7-10].

* Correspondence: dingh@cc.umanitoba.ca

¹Department of Biochemistry and Medical Genetics, University of Manitoba, 745 Bannatyne Avenue, Winnipeg MB R3E 0J9, Canada
Full list of author information is available at the end of the article

Although HLTF was originally identified as a DNA-binding protein that could interact with several gene promoters and enhancers [7-11], recent studies indicate that this DNA helicase is more involved in the DNA-damage repair pathway. First, HLTF has been shown to exhibit an E3 ubiquitin ligase activity for the polyubiquitination of proliferating cell nuclear antigen (PCNA), which is required for the initiation of an error-free replication through DNA damage lesions [12,13]. Second, HLTF has also been found to display a double-stranded DNA translocase activity, which promotes the resolution of stalled replication forks at DNA damage lesions [14,15]. Third, a recent study indicates that HLTF also possesses a chromatin remodeling activity, which leads to the displacement of DNA-bound proteins on stalled replication forks and facilitates DNA-damage repair [16]. These findings demonstrate that HLTF may be a functional homologue of yeast rad5 and that it plays an important role in an error-free post-replicative repair pathway. The requirement of HLTF for repair of damaged DNA may also implicate a tumor suppression role in human colon cancers, where HLTF has been identified as a common target for methylation and epigenetic gene silencing.

Epigenetic inactivation of HLTF gene expression by promoter hypermethylation has been reported in more than 40% of human colon cancers [17-20]. The frequency of *HLTF* promoter methylation was found to increase drastically between early stage of adenomas and advanced adenomas, suggesting that this epigenetic alteration could be a later event in colon carcinogenesis [17,19]. In addition, *HLTF* methylation was demonstrated to be significantly correlated with poor prognosis in human colon cancer patients [21,22]. Besides colon tumors, *HLTF* methylation was also commonly detected in human gastric tumors or cancers, but not in other human cancers, such as lung and breast cancers [17,23-25], suggesting that this epigenetic alteration is unique to the tumors of the gastrointestinal tract. However, although *HLTF* methylation has been demonstrated as a common event in human colon cancer and it has been suggested as a prognostic biomarker [22,26], it is largely unclear whether this epigenetic event is important for the development of this cancer. In the present study, we generated an *Hltf* deficient mouse allele, and applied this mouse model to characterize the role of loss of HLTF function in the development of intestinal cancer. Our data demonstrates that loss of *Hltf* has a promoting effect on the transition of intestinal adenomas to invasive cancers, which highly suggests that aberrant HLTF methylation, as found in most human colon cancers, may have an important pathogenic role in the development of this cancer.

Results


Generation of the *Hltf* null mouse allele

To study the loss of function of *Hltf* *in vivo*, we mutated *Hltf* in mice by homologous recombination. A gene-targeting vector was generated to replace the exons 1-5 coding for residues 2-208 of *Hltf* with a nuclear localized *LacZ* (*nls-LacZ*) cDNA that was fused in-frame with the endogenous *Hltf* starting codon (Figure 1A). Correct homologous recombination in the targeted ES cells was confirmed by Southern blot analysis using both 5' and 3' probes located upstream and downstream of the targeted *Hltf* sequence (Figure 1B and 1C). Two independent ES clones were used to produce germline-transmitting chimeras that were further bred with a ubiquitous Cre transgenic line (*Elia-Cre*) to delete the *loxP* flanked *pGK-neo* selection marker from the targeted locus (Figure 1A). The resultant *Hltf*^{+/-} mouse, which was maintained on either a C57BL/6 or C57BL/6/129S1 background, was used to produce *Hltf* homozygous mutant mice for phenotypic characterization.

To determine whether *Hltf* expression is completely inactivated in *Hltf*^{-/-} mice, we performed Northern blot analysis of total RNA prepared from control and mutant E10.5 embryos. Whereas *Hltf* transcripts were identified in control (wild-type and *Hltf*^{+/-}) embryos using the 5' and 3' *Hltf* cDNA probes, transcripts were not detected in *Hltf*^{-/-} samples (Figure 1D). To further demonstrate the absence of *Hltf* expression in *Hltf*^{-/-} mice, we also applied a more sensitive real time RT-PCR assay to examine the expression of *Hltf* in derived *Hltf*^{-/-} mouse embryonic stem (ES) cells or mouse embryonic fibroblast (MEF) cells as well as in *Hltf*^{-/-} E10.5 embryos. With the primer set located on the 3' of *Hltf* cDNA, a strong amplification of *Hltf* was detected in the control group, but no signal was found in *Hltf*^{-/-} samples (Figure 1E). These data clearly indicate that the targeted mutation that we introduced into the *Hltf* mouse genomic locus gave rise to a null allele.

Normal development and lack of intestinal tumors in *Hltf* null mice


With the established *Hltf* null allele, we then asked whether loss of *Hltf* function could affect mouse development and induce the formation of intestinal tumors. As a first approach to address this, we determined the expression pattern of *Hltf* during development. We took advantage of the inserted *nls-LacZ* reporter gene in *Hltf* knockout mice, whose expression is controlled by the endogenous *Hltf* regulatory elements. Both *Hltf*^{+/-} and *Hltf*^{-/-} mice showed the same *LacZ* expression pattern. By determining *LacZ* activity in these mice, we found that *Hltf* was specifically expressed in the heart at an early developmental stage (E8.5 to E9.5) (Figure 2A).



Hltf exhibited a broader expression pattern at E10.5, with LacZ signals detected in somites, branchial arches, limb bud and brain (Figure 2B). At later embryonic developmental stages, such as E16.5, *Hltf* showed wide and strong expression in many tissues, including heart, lung, liver, kidney, spleen and pancreas (Figure 2C). This wide-spread expression of *Hltf* was also observed in adult mice (data not shown). In both adult intestine

and colon, *Hltf* expression was mainly detected in the crypts and in the intestinal epithelial cells (Figure 2D).

This broad expression pattern of *Hltf* during mouse development might suggest that *Hltf* is important for development. However, *Hltf*^{-/-} mice were found to have normal embryonic and postnatal development. The *Hltf* null mice were fertile, and exhibited similar a body weight and life span as their wild-type (wt) littermates.



In addition, *Hltf*^{-/-} mice did not display obvious abnormalities of intestinal proliferation or differentiation as determined by the BrdU incorporation assay as well as by the analyses with different intestinal cell lineage markers (Additional file 1 and Additional file 2). To determine whether loss of *Hltf* function could induce

the formation of intestinal tumors, we monitored a cohort of 60 mice fed with a normal diet for a two-year period (30 *Hltf*^{-/-} mice on C57BL/6 background and 30 wt mice matched for age and genetic background). Similar to the wt control group, 10% of *Hltf*^{-/-} mice (3 out of 30) developed liver tumors or lymphomas between 16

and 24 months. However, none of *Hltf*^{-/-} mice formed intestinal tumors or gastric tumors.

Taken together, our results demonstrate that *Hltf* is dispensable for normal development. Our work also indicates that loss of *Hltf* function by itself is insufficient to drive the oncogenic process in the gastrointestinal tract, which implicates that the epigenetic inactivation of HLTF, as commonly found in human colon cancers, could be involved in the late stage of intestinal tumor development.

Development of intestinal cancers in *Hltf*^{-/-}/*Apc*^{min/+} mutant mice

To determine whether the loss of HLTF function could have a role in the progression of intestinal tumors, we introduced the *Hltf* null mutation into *Apc*^{min/+} mice. The *Apc*^{min/+} mice harbor a premature stop codon in one allele of the *Apc* tumor suppressor gene, and develop multiple intestinal adenomas that mimic human familial adenomatous polyposis [27]. The adenomas in *Apc*^{min/+} mice rarely progress to invasive intestinal or colon cancers [28], making this mouse model an excellent genetic tool for studying the later events involved in intestinal carcinogenesis.

To avoid strain-specific modification of the *Apc*^{min/+} intestinal tumor phenotype [29,30], we generated *Hltf*^{-/-}/*Apc*^{min/+} mice on the C57BL/6 background. *Hltf*^{-/-}/*Apc*^{min/+} mice became moribund at a similar age as *Apc*^{min/+} mice. Post-mortem examination revealed that *Hltf*^{-/-}/*Apc*^{min/+} and *Apc*^{min/+} mice developed similar numbers of tumors in the small intestine (average of 65 macroscopic tumors/per mouse in both groups). However, upon histological characterization, most *Hltf*^{-/-}/*Apc*^{min/+} mice (24 out of 30) were found to form invasive intestinal adenocarcinomas characterized by deeper invasion of tumor cells into the muscularis propria (Figure 3B and Additional file 3). The invaded neoplastic glandular cells exhibited high β -catenin activity as reflected by the nuclear accumulation of β -catenin detected by immuno-staining with anti- β -catenin antibody (Figure 3B). In contrast, most *Apc*^{min/+} mice developed only well-defined tubular adenomas or adenomas within the lamina propria (Figure 3A).

Hltf^{-/-}/*Apc*^{min/+} mice were also found to develop a similar number of neoplastic lesions in the colon as *Apc*^{min/+} mice (average of 1.5 macroscopic tumors/per mouse). Histologically, although both *Hltf*^{-/-}/*Apc*^{min/+} and *Apc*^{min/+} colon tumors showed tumor growth with a pedunculated morphology protruding into the colonic lumen, the majority of *Hltf*^{-/-}/*Apc*^{min/+} colon tumors displayed a more dramatic glandular atypia that resulted in the formation of numerous mucin-filled cysts (Figure 4A, B), a characteristic pathological feature for human


colon cancers [31]. In addition, most *Hltf*^{-/-}/*Apc*^{min/+} colon tumors were also found to have a strong desmoplastic stromal reaction, suggesting invasion of the lamina propria (Figure 4B). Furthermore, *Hltf*^{-/-}/*Apc*^{min/+} colon tumors were also found to have stronger and more ubiquitous nuclear-stained β -catenin signals than the tumor cells in *Apc*^{min/+} mice (Figure 4C). These data indicate that colon tumors from *Hltf*^{-/-}/*Apc*^{min/+} mice had progressed to much higher grades than those from *Apc*^{min/+} mice. To further demonstrate the malignancy of colon tumors developed in *Hltf*^{-/-}/*Apc*^{min/+} mice, we derived cells from these tumors and then subcutaneously injected them into *Rag1*^{-/-}/*IL2*^{-/-} immunodeficient mice. Within 30 days after injection, these cells formed subcutaneous tumors that maintained the morphological and immunohistochemical features of the parental colon tumors (Figure 4D). This capability of propagating tumor cells into another host, which was not observed with cells derived *Apc*^{min/+} colon tumors, further supports that many colon tumors developed in *Hltf*^{-/-}/*Apc*^{min/+} mice could be malignant cancers.

Collectively, our results demonstrate that loss of *Hltf* function is able to promote intestinal or colonic carcinogenesis in mice with a mutant *Apc* background. This finding, together with human epigenetic evidence that HLTF is frequently silenced in the advanced human colon adenomas or cancers but uncommonly in early adenomas [17,19], indicates that epigenetic inactivation of HLTF could be an important event involved in the transition of benign adenomas to malignant colon cancer.

Chromosomal instability in *Hltf*^{-/-}/*Apc*^{min/+} colon tumor cells

Chromosomal instability (CIN) is a genetic hallmark of most human colon cancers, and it has been demonstrated to be important for the progression of colon tumors by increasing the rate of genetic aberration [2,32,33]. Since HLTF has been shown to be required for the maintenance of genomic stability [12-16], it is likely that intestinal or colonic tumors developed in *Hltf*^{-/-}/*Apc*^{min/+} mice would have increased genomic instability. To examine this, we analyzed metaphase spreads generated from tumor cells (passage 2) derived from colon tumors developed in *Apc*^{min/+} and *Hltf*^{-/-}/*Apc*^{min/+} mice. These tumors all showed pronounced cribriform tumor growth without invasive neoplastic glands and were histologically characterized as late stage colonic adenomas.

Consistent with previous findings [34,35], colon tumor cells from *Apc*^{min/+} mice showed a near-diploid karyotype and did not harbor obvious chromosomal abnormalities (Figure 5A). In contrast, three colon tumor cell lines derived from *Hltf*^{-/-}/*Apc*^{min/+} mice displayed an



aneuploid phenotype (Figure 5B and 5C). Spectral karyotype (SKY) analysis on these tumor cells further revealed a high incidence of genetic alterations in the forms of nonreciprocal translocations, chromosomal fusions (which include both dicentric chromosomes and centromeric fusions), and chromosomal fragments and breaks (Figure 5D-G and Table 1). In addition, many

chromosomal fusions in *Hltf*^{-/-}/*Apc*^{min/+} colon tumor cells were found to lack telomere signals (Figure 5I), indicating the presence of telomere dysfunction in these tumor cells. Collectively, these results demonstrate that *Hltf*^{-/-}/*Apc*^{min/+} colon tumors had gross chromosomal abnormalities, similar to the CIN phenotype as described in most human colorectal cancers.





Figure 4 *Hltf*^{-/-}/*Apc*^{min/+} mice develop malignant colon tumors that are transplantable into another host. (A) Haematoxylin-eosin staining of representative colon tumors collected from *Apc*^{min/+} mice, showing pronounced tumor growth with a pedunculated morphology protruding into the colonic lumen (indicated by arrow). (B) Representative colon tumors collected from *Hltf*^{-/-}/*Apc*^{min/+} mice, demonstrating the formation of numerous mucin-filled cysts within tumors (white arrowheads) and strong desmoplastic stromal reaction. (C) Immuno-staining with anti- β -catenin antibody. *Hltf*^{-/-}/*Apc*^{min/+} colon tumors had stronger and more ubiquitous nuclear-stained β -catenin signals than the tumor cells in *Apc*^{min/+} mice. Arrowheads indicate the tumor cells with nuclear-stained β -catenin. (D) The formation of subcutaneous tumors from derived *Hltf*^{-/-}/*Apc*^{min/+} colon tumor cells in *Rag1*^{-/-}/*IL2*^{-/-} immunodeficient mice. The subcutaneous tumors displayed neoplastic glandular structures, which contained strong nuclear-stained β -catenin signals and were also positive for Villin, a marker for intestinal epithelial cells. Arrows indicate the tumor, whereas arrowheads mark the subcutaneous muscle layer.

Increased tumor growth and chromosomal instability in HCT116 human colon cancer cells with HLTF knockdown

To determine whether loss of HLTF function would also lead to a similar tumor-promoting effect in human colon cancer as we demonstrated in the *Hltf*^{-/-}/*Apc*^{min/+} mouse model, we applied a lentiviral based shRNA knockdown approach to down-regulate HLTF expression in HCT116 human colon cancer cells. HCT116 cells are one of a few colon cancer cell lines that lack *HLTF* promoter methylation [17], and express high

levels of HLTF (Figure 6A). In addition, these cells are near-diploid with a stable karyotype [36], which would also allow us to determine whether loss of HLTF function in human colon cancer cells could induce chromosomal abnormalities in a manner similar to the *Hltf*-deficient mouse colon tumor cells.

As shown in Figure 6B, HLTF expression was significantly reduced to almost undetectable protein level in the transduced HCT116 cells that were stably integrated with HLTF shRNAs (HCT116^{HLTFshRNA1} and



HCT116^{HLTFshRNA2}) as compared to the scramble shRNA control (HCT116^{Scramble}). To determine whether this down-regulation of HLTF expression was able to modulate HCT116 tumor growth, HCT116 cells with or without HLTF knockdown were injected subcutaneously

into *Rag1^{-/-}/IL2^{-/-}* immunodeficient mice. All the mice with injected tumor cells (10⁵ cells/per mouse) developed subcutaneous tumors within 20 days. However, the tumors formed by HCT116^{HLTFshRNA1} or HCT116^{HLTFshRNA2} cells showed significantly increased

Table 1 Summary of cytogenetic aberrations observed in colon tumors from *Hltf*^{-/-}*Apc*^{min/+} mice

Genotype of tumors	Metaphase Analyzed	Aneuploidy (%) ^a	Translocations (%)	End-to-End Fusions (%) ^b	Fragments (%)	Breaks (%)
<i>Apc</i> ^{min/+}	20	3 (15%)	0	0	0	0
<i>Hltf</i> ^{-/-} / <i>Apc</i> ^{min/+}	20	17 (85%)	7 (35%)	9 (45%)	16 (80%)	3 (15%)
<i>Hltf</i> ^{-/-} / <i>Apc</i> ^{min/+}	33	31 (93%)	10 (32%)	21 (63%)	15 (50%)	4 (20%)
<i>Hltf</i> ^{-/-} / <i>Apc</i> ^{min/+}	20	20 (100%)	3 (15%)	4 (20%)	1 (5%)	4 (20%)

^a Expressed as percentage of cells with other than 40 chromosomes (each Robertsonian fusion counted as two chromosomes).

^b Expressed as total number of end-to-end associated chromosomes per metaphase, including dicentric chromosomes, telomere associations and Robertsonian fusions




Figure 6 Down-regulation of HLTf in human HCT116 colon cancer cells promotes tumor growth. (A) Western blot analysis with anti-HLTf antibody on the cell lysates prepared from *HLTF* methylated human colon cancer cells (RKO and SW480) and *HLTF* unmethylated human colon cancer cells (HCT116). (B) Western blot analysis on the cell lysates prepared from HCT116 cells that were stably integrated with scramble control shRNA and *HLTF* shRNAs. *HLTF* expression was efficiently inhibited by two independent *HLTF* shRNAs. (C) Representative image of subcutaneous tumors formed from HCT116^{Scramble}, HCT116^{HLTFshRNA1} and HCT116^{HLTFshRNA2} cells in *Rag1*^{-/-}/*IL2*^{-/-} immunodeficient mice. (D) Haematoxylin-eosin staining demonstrates that both *HLTF* knockdown tumors displayed tumor cell invasion into the muscle layer (indicated by arrows) which was not typically observed in the scramble control tumors. *HLTF* knockdown tumors also showed low expression of *HLTF* as detected by immunostaining with anti-*HLTF* antibody.

tumor size as compared with that of the scramble control (Figure 6C). Moreover, both HCT116^{HLTFshRNA1} and HCT116^{HLTFshRNA2} subcutaneous tumors also frequently exhibited the invasion of tumor cells into the subcutaneous muscle layer that was uncommonly observed in the control tumors (Figure 6D). To confirm that HLTF expression was indeed down-regulated in these tumors, we performed immunohistochemistry with an anti-HLTF antibody. Both HCT116^{HLTFshRNA1} and HCT116^{HLTFshRNA2} subcutaneous tumors showed significantly lower expression of HLTF as compared to the scramble control tumors (Figure 6D). Taken together, our data indicate that down-regulation of HLTF in human colon cancer cells may promote tumor growth and malignancy, similar to what we found in the *Hltf*^{-/-}/*Apc*^{min/+} mouse model.

We then performed cytogenetic analysis to determine whether down-regulation of HLTF expression in human colon cancer cells induced chromosomal abnormalities. Detailed SKY analysis of metaphase spreads from both short term (10 days) cultured HCT116 parental cells and HCT116^{scramble} cells revealed the presence of near-diploid chromosomes with a few highly conserved chromosomal translocations (Figure 7A), consistent with the documented karyotype for HCT116 parental cells [37]. This near-diploid karyotype was also observed in the second generation of HCT116 cells and HCT116^{scramble} cells that were derived from the subcutaneous tumors formed by these cells (Figure 7C), further indicating that HCT116 cells have a stable chromosome karyotype as demonstrated previously [38,39]. In contrast, both short-term cultured and the second generation of HCT116^{HLTFshRNA1} and HCT116^{HLTFshRNA2} tumor cells exhibited significantly increased chromosomal abnormalities in the forms of chromosomal loss and chromosomal fusions and breaks as compared to the control groups ($p < 0.001$) (Figure 7B-C). In addition, approximately 15% of metaphases from HCT116^{HLTFshRNA1} and HCT116^{HLTFshRNA2} tumor cells showed high numbers of trisomy (Figure 7D). These results clearly indicate that down-regulation of HLTF in human colon cancer cells is able to induce chromosomal abnormalities.

Discussion

There is now strong evidence that a series of genetic alterations are required for the pathogenesis of human colon cancer. Specific gene mutations, such as in the *APC* gene, initiate the formation of colonic adenomas and others (e.g. *TP53* mutations as well as other alterations, including *KRAS* activation) drive the malignant transformation of the adenomas in a multistep progression model [2,3]. More recently, epigenetic alterations, specifically aberrant DNA methylation, have been found to occur commonly in colon cancers [4,5]. Although

there is controversy regarding the significance of these alterations in the pathogenesis of colon cancer, more and more data indicate that the aberrant methylation of at least some of these genes, such as *MLH1* (MutL homolog 1), *MGMT* (O-6-methylguanine-DNA methyltransferase) and *HIC1* (hypermethylated in cancer 1) can be pathogenetic in colorectal carcinogenesis [2,6]. Here, we have added another example to demonstrate the importance of epigenetic alterations in this carcinogenesis.

In this study, we focused on determining the role of loss of HLTF function in the development of colon cancer. The *HLTF* promoter has been found to be hypermethylated in more than 40% of human colon cancers [17-20], suggesting that *HLTF* is a common target for methylation in this cancer. All colon cancer cells that lacked HLTF expression had the methylation of CpG islands within the *HLTF* promoter, while methylation was not detected in the HLTF-expressing cells [17] (Figure 6A), further indicating that this epigenetic event leads to a complete inactivation of HLTF expression in colon cancer cells. To determine the role of loss of HLTF function, in this study, we generated *Hltf* deficient mice (Figure 1). Although these mice did not develop intestinal or colonic tumors, the majority of them were found to develop invasive intestinal and colonic cancers upon *Apc* mutation-mediated tumor initiation (Figures 4 and 5). Our results strongly suggest that aberrant methylation of HLTF, which leads to the loss of this gene's function as found in most human colon cancers, could have a pathogenetic role rather than being a consequence of colorectal carcinogenesis. This epigenetic event could be important for driving the transition of benign adenomas to the malignant adenocarcinomas, which is consistent with human tumor data showing that *HLTF* promoter methylation is mainly detected in the advanced colonic adenomas and cancers, but not in the early stage of adenomatous polyp [17,19]. The high frequency of *HLTF* promoter hypermethylation observed in these tumors indicates that the silencing of this gene could confer a selective advantage for the progression of these tumors to the malignant cancers.

Although the mechanism by which loss of HLTF function results in this malignant transformation still needs to be defined, our data demonstrating gross chromosomal abnormalities in *Hltf* deficient colon tumor cells may suggest that genomic instability induced in these tumors is a driving force for this carcinogenesis. Genomic instability is a characteristic of human colon cancers, and has been implicated in the initiation and progression of these cancers by increasing the rate of genetic alterations [2]. Two types of genetic instability have been found in human colon cancer: microsatellite instability (MIN) and chromosomal instability (CIN).




Figure 7 HLTf knockdown induces chromosomal abnormalities in HCT116 colon cancer cells. (A) Representative SKY image of HCT116^{scramble} cells, showing a stable karyotype with 45 total chromosomes, der(10)t(10;6) (white arrow), der(16)t(8;16) (white arrowhead) and der(18)t(17;18) (yellow arrow), a documented signature for HCT116 parental cells. (B) Representative SKY images of HLTf knockdown HCT116 cells. Several additional chromosomal abnormalities were identified, which include chromosomal fragment (yellow arrowhead), monosomy chromosome 3 (white arrow), der(15)(15;11) (white arrowhead) and der(16)(16;22) (yellow arrow). (C) The number of structural chromosomal abnormalities in HCT116 cells with and without HLTf knockdown. Two groups of cells were analyzed, one from a short term *in vitro* culture (10 days) and another from subcutaneous tumors that were generated by these cells. The horizontal bars indicate the mean numbers of chromosomal abnormalities in each population of cells. Each data point represents the total number of structural chromosomal rearrangements per metaphase spread after subtraction of the rearrangements (i.e. stable present translocations) found in HCT116 parental cells. (D) Representative SKY images of HLTf knockdown HCT116 cells, demonstrating the presence of high number of trisomies in the tumor cells (indicated by arrows)

Tumors with MIN display normal diploid with no obvious chromosomal defects but have high incidence of instability in microsatellite repeats linked to impaired mismatch repair (MMR) [40]. The predominant mechanism identified to inactivate MMR in these tumors is epigenetic silencing through promoter methylation [41], although somatic mutations in MMR genes also occur [42]. CIN, which accounts for 85% of colorectal cancers, exhibits aneuploid, allelic losses and other chromosomal abnormalities, such as translocation, fusion and breaks [32]. Colon cancers with CIN phenotype have been found to behave more aggressively in terms of invasiveness and metastasis [32]. Although its molecular basis remains largely elusive, CIN is thought to arise from structural defects involving centromere or centrosome, microtubule dysfunction, telomere defects, chromosomal fragility and cell cycle checkpoint failure [2,6]. In this work, we have provided solid evidence that *Hltf* deficient mouse colon tumors display severe chromosomal abnormalities, similar to CIN as described in human colon cancers (Figure 5). We have also demonstrated that down-regulation of HLTF increased chromosomal abnormalities in human colon cancer cells (Figure 7). Furthermore, similar to human colon cancers with a CIN phenotype, we have also found that *Hltf* deficient mouse colon tumors or *HLTF* knockdown human colon cancer cells developed more malignant features, such as invasiveness, as compared to the tumor cells that contain functional HLTF (Figures 4, 5 and 6). All these findings indicate that the genomic instability induced in HLTF deficient colon tumors likely plays an important role in the malignant transformation of these tumors. Future studies using our established *Hltf* knockout mouse colon tumor model to identify the genetic mutations involved in this pathogenesis have the potential to provide a deeper understanding of colon cancer development.

The presence of a CIN phenotype in *Hltf* knockout or knockdown tumor cells also implicates loss of HLTF function as an additional mechanism to induce CIN phenotypes in human colon cancer. The involvement of HLTF in the post-replicative DNA damage-repair pathway has been demonstrated by several recent studies [12-16]. In these work, HLTF was shown to display a similar function as yeast rad5 in the polyubiquitination of PCNA, which is required for initiating an error-free post-replicative repair activity [12,13]. HLTF was also found to have both DNA translocase and chromatin remodeling activities to facilitate the repair of damaged DNA on stalled replication forks [14-16]. Therefore, HLTF could be an important factor in the post-replicative repair pathway. Without HLTF, it is likely that the progression of the replication fork would be blocked at DNA damage lesions, resulting in an accumulation of

DNA intermediates that trigger recombination and genomic instability [43]. Our cytogenetic data demonstrating the gross chromosomal abnormalities present in *Hltf* deficient mouse tumor or in HLTF knockdown HCT116 colon cancer cells strongly support this notion. Our data also agree with many other studies indicating that aberrant post-replicative repair activity is a major source of the mutations and chromosome rearrangements that could be important for tumorigenesis. Interestingly, mice deficient in *Hltf* alone were found to develop normally and did not display any pathological disorders associated with genetic instability. This may be due to the robust replication checkpoints present in normal mouse cells, which, upon DNA damage, could boost a DNA damage response that arrests cells that harbor unrepaired DNA damage [44]. These unrepaired cells could undergo apoptosis and then be replenished by cells differentiated from a pool of progenitor and stem cells during development. However, when these checkpoint pathways are not functional, the cells that contain defective or unrepaired DNA lesions could still progress through the cell cycle, resulting in broken chromosomes, genome aberrations and an accumulation of mutations [44,45]. A recent study indicates that APC could play an important role in the DNA replication checkpoints by stabilizing the association of DNA replication complexes at stalled DNA replication forks [46]. Consistent with this finding, tumor cells that have lost APC function, such as human SW480 colon cancer cells, have been found to lose the capacity to arrest cells with damaged DNA [46]. Therefore, *Hltf*^{-/-}/*Apc*^{min/+} colon tumor cells may have impaired replication checkpoints, which could facilitate cell cycle progression to enhance the genomic instability in these tumor cells.

HLTF has also been demonstrated to interact with several different gene promoters and enhancers [7-11], suggesting its potential role in the regulation of gene expression. We recently performed microarray-based gene expression analyses and did not find significant changes of gene expression between *Hltf*^{+/+} and *Hltf*^{-/-} mouse ES cells or in HCT116 cells with and without HLTF knockdown. In both cases, a very few genes showed 2-fold change with the statistical significance (Additional file 4). None of these genes have been implicated in tumorigenesis or in the maintenance of genomic stability. Therefore, the loss of HLTF function is unlikely to mis-regulate the expression of specific genes, leading to the development of colon cancers.

Conclusions

We have provided direct genetic evidence that loss of HLTF function promotes the malignant transformation of intestinal or colonic adenomas to carcinomas. We have also demonstrated that this pathogenetic outcome

could be associated with the role of HLTf in the maintenance of genomic stability. Given the high frequency of epigenetic inactivation by hypermethylation of HLTf in human colon cancers, our studies strongly suggest that this epigenetic alteration could be directly involved in the development of colon cancer rather than a consequence of this carcinogenesis.

Methods

Construction of *Hltf* gene-targeting vector

The *Hltf* gene-targeting vector was generated based on a PCR-based cloning strategy [47]. Briefly, the mouse *Hltf* genomic fragments required for the 5' and 3' arms of homology were PCR-amplified from the genomic DNA of R1 ES cells (on 129S1 background) with a high-fidelity polymerase (TaKaRa). The primer sequences are listed in Additional file 5. After validation by DNA sequencing, the *Hltf* 5' arm DNA fragment was ligated upstream of a *nlsLacZpA-loxP-pGKneo-loxP* cassette. This vector was further inserted by a DNA fragment containing *Hltf* 3' arm and *pGK-DTA* cassette to generate the final gene-targeting vector that contains *Hltf* 5' arm-*nlsLacZpA-loxP-pGKneo-loxP-Hltf* 3' arm-*pGK-DTA* (Figure 1A).

Generation of *Hltf* deficient mice

The *Hltf* gene-targeting vector was linearized with *SacII* and then electroporated into R1 ES cells. The transfected ES cells were selected with G418 (250 µg/ml), and G418-resistant ES clones were screened by Southern blot analysis for the correctly targeted allele using *BamHI* (for the 5' external probe) and *EcoRI* (for the 3' external probe) digestions. Both the 5' and 3' external probes were PCR amplified with primers listed in Additional file 5. Two independently targeted ES cell clones were used to generate chimeric mice by ES cell-diploid embryo aggregation. Transmitting chimeric males were crossed with an *EIIa-Cre* line (Jackson laboratories, Bar Harbor, Maine) to delete the *loxP* flanked *neo* cassette from the targeted allele. The resultant mouse allele was further bred with 129S1 females to produce hemizygous transgenic offspring on 129S1/C57BL/6 background. *Hltf*^{+/-} mice were also backcrossed to C57BL/6 for eight generations. All mouse experiments were performed in accordance with procedures approved by the University of Manitoba Animal Care and Use Committee.

Genotyping

A PCR based genotyping method was applied to genotype *Hltf* knockout mice. Primers to amplify the targeted allele were the sense primer (P1) (5'-GGAGCTTTAT-CAGGCTGTCTGGGA-3') and antisense primer (P2) specific for the *LacZpA* cassette (5'-AGGAAGATCG-CACTCCAGCCAGCT-3'). To detect the wild-type

HLTF allele, wt P1 primer (5'-GTCCATGTCCTAGC-CATGAGTA-3') and an antisense primer (P3) locating in exon 1 (5'-GCTTGGTAAGGACTACAAAGCA-3') were used for PCR.

Mouse intestinal and colonic tumor analysis on *Hltf/Apc* mutant mice

The *Apc*^{Min/+} mice were purchased from The Jackson Laboratory and maintained on a C57/B6 genetic background. These mice were bred with *Hltf*^{+/-} (C57/B6 background) to generate a cohort of *Hltf*^{-/-/Apc}^{min/+}, *Apc*^{min/+} and *Hltf*^{+/-/Apc}^{min/+} mice for intestinal tumor analysis. Mice were sacrificed by cervical dislocation, and the entire small intestine and colon were flushed with cold PBS and dissected longitudinally. The number, location and size of tumors in the intestine or colon were examined and recorded. Small intestinal and colonic tumors were further histologically analyzed by hematoxylin/eosin staining and immunohistochemistry.

Establishment of colon tumor cells from *Hltf/Apc* mutant mice and growth in *Rag1*^{-/-/IL2}^{-/-} immunocompromised mice

Tumor cells from *Hltf*^{-/-/Apc}^{min/+} and *Apc*^{min/+} colon tumors were generated as described [35]. The cells were grown in 6 cm culture dishes plated with mitomycin-treated mouse embryonic fibroblast (MEF) cells using the culture conditions as described [35]. The early passage (p2) cells were collected either for tumorigenicity analysis by injecting 10⁵ cells subcutaneously to *Rag1*^{-/-/IL2}^{-/-} mice (purchased from Jackson Laboratory), or for cytogenetic characterization.

Immunohistochemistry (IHC)

IHC was performed as described previously [48]. Briefly, dissected intestinal and colonic tumors were fixed in 10% formalin in PBS overnight. The fixed tissues were embedded in paraffin, and 5 µm paraffin-embedded tissue sections were pretreated with Retrieval solution (Dako) and blocked with either mouse IgG blocking reagent (Vector Laboratory) or serum free blocking reagent (Dako), and then incubated with primary antibodies overnight at 4°. Primary antibodies used were mouse anti-BrdU (Roche, dilution 1:100), mouse anti-Ki67 (BD Pharmingen, dilution 1:100), mouse anti-β-catenin (BD Pharmingen, dilution 1:200), rabbit anti-HLTf (Sigma, dilution 1:100) rabbit anti-lysozyme (Dako, dilution 1:200), and rabbit anti-chromogranin A (ImmunoStar, dilution 1:4000). After washing with TBS containing 0.1% Tween20 buffer, sections were incubated with biotinylated anti-rabbit or anti-mouse antibody for 60 min at room temperature. The sections were washed 3 times with TBS-0.1% Tween20 buffer and incubated with avidin-biotin-peroxidase complex

(Vector Laboratory), according to the manufacturer's protocol. Color was developed using metal 3,3'-diaminobenzidine tetrachloride.

Lentiviral based shRNA knockdown of HLTF expression in HCT116 cells

Lentiviral vectors expressing shRNA against human *HLTF* (NM 003071, V2LHS 254799, _153142, _153143, _153144, _153146) or scrambled control were obtained from the GIPZ lentiviral shRNAmir library (Open Biosystems, Thermo Scientific) acquired at the University of Manitoba. The pGIPZ lentiviral vector expresses a short hairpin RNA modeled by human microRNA-30 (miR30) primary transcript for increased Drosha and Dicer processing. TurboGFP reporter gene and puromycin drug resistance marker are expressed as part of a bicistronic transcript. Preparation and infection of lentiviral particles into HCT116 cells (from ATCC) were performed based on an established procedure as described previously [49]. After infection, HCT116 cells were treated with 0.75 µg/ml puromycin for four days. The puromycin-resistant clones were pooled to determine HLTF expression by Western blot analysis with anti-HLTF antibody (Sigma). 10⁵ cells from these established stable cell lines were subcutaneously injected to *Rag1*^{-/-}/*IL2*^{-/-} mice to determine the effect of HLTF knockdown on tumor growth. These cells, as well as cells derived from subcutaneous tumors, were also utilized for cytogenetic analysis.

Cytogenetic analysis

Metaphase spreads for SKY analysis were performed using the SKT kit (ASI) according to the manufacturer's instructions and previously published methods [50]. In each cell line, at least 20 metaphase spreads were studied. Q-FISH analysis was performed as previously described [51].

Northern blot analysis

Total RNA from freshly dissected mouse E10.5 embryos was extracted using TRIzol (Life Technologies, Inc.). 20 µg of total RNA was separated on a 1% agarose-formaldehyde gel and transferred to Hybond nylon membrane (Amersham). Hybridization was carried out in PerfectHyb (Sigma) with 1.5 × 10⁶ cpm/ml of 5' and 3' *Hltf* cDNA probes. Both 5' and 3' cDNA probes were PCR-amplified with the primers listed in Additional file 5.

X-gal staining

Dissected mouse embryos or tissues were fixed for 30 min (for E8.5 to E9.5) or 60 min (for E10.5 embryos and tissues) with 4% paraformaldehyde (PFA), 0.2% glutaraldehyde and 0.02% NP-40 in PBS. Subsequently, fixed samples were washed three times with PBS containing 0.02% NP-40, and stained at 37°C overnight with a

staining solution containing 4 mM K₄Fe(CN)₆, 4 mM K₃Fe(CN)₆, 2 mM MgCl₂, and 0.2% X-gal in PBS.

Western blot analysis

For Western blots, cells were lysed in RIPA buffer. 30 µg lysate was separated by 8% SDS-PAGE and transferred to nitrocellulose membrane, which was blocked with 5% non-fat milk and incubated with rabbit anti-HLTF antibody (Sigma, dilution 1:500) at 4° overnight. Protein was detected using an enhanced chemiluminescence system (Amersham Life Science).

Statistical analysis

A paired *t*-test (two tailed) was used to analyze the statistical significance of chromosomal abnormalities in tumor cells. *P* < 0.05 was considered indicative of statistical significance.

Additional material

Additional file 1: Loss of *Hltf* function does not affect the cellular proliferation in small intestine and colon. The small intestines and colons collected from 2-month old *Hltf*^{+/+} and *Hltf*^{-/-} mice were stained with Haematoxylin-eosin and anti-BrdU antibody. Both *Hltf*^{+/+} and *Hltf*^{-/-} intestine and colon displayed normal morphology and a similar number of BrdU positive cells within the crypt (2 h after BrdU injection) and in crypt-villus axis (24 h after BrdU injection).

Additional file 2: Loss of *Hltf* function does not affect the differentiation of epithelial cells in small intestine and colon. The small intestines and colons from 2-month old *Hltf*^{+/+} and *Hltf*^{-/-} mice were analyzed using several intestinal cell-lineage markers. The Goblet cells were determined by staining with Alcian blue and periodic acid-Schiff (PAS). The enteroendocrine cells were analyzed by immunostaining with anti-chromogranin A, and the Paneth cells in small intestine were detected by anti-lysozyme antibody. Both *Hltf*^{+/+} and *Hltf*^{-/-} intestines or colons showed very similar staining patterns for these markers.

Additional file 3: *Hltf*^{-/-}/*Apc*^{min/+} mice frequently developed invasive intestinal adenocarcinomas. Several additional Haematoxylin-eosin stained images demonstrate the formation of invasive intestinal adenocarcinomas (indicated by arrows) in *Hltf*^{-/-}/*Apc*^{min/+} mice.

Additional file 4: List of genes that showed expression changes between *Hltf*^{+/+} and *Hltf*^{-/-} mouse ES cells and in HCT116 cells with and without HLTF knockdown. The microarray expression assays were carried out by the Center for Applied Genomics at the Toronto Hospital for Sick Children. Three independent cell lines from each group were analyzed. The changes with the statistical significance (*p* < 0.05) are highlighted by yellow. Red indicates the change of gene expression cannot be validated by real-time PCR.

Additional file 5: List of PCR primers that were applied in this study.

Abbreviations

HLTF: Helicase-like Transcription Factor; APC: Adenomatous polyposis coli; PCNA: Proliferating cell nuclear antigen; wt: Wild type; CIN: Chromosomal instability; SKY: Spectral karyotype; MLH1: MutL homolog 1; MGMT: O-6-methylguanine-DNA methyltransferase; HIC1: Hypermethylated in cancer 1; MIN: Microsatellite instability; MMR: Mismatch repair; ES: Embryonic stem cells; MEF: Mouse embryonic stem cell; IHC: Immunohistochemistry; PFA: Paraformaldehyde.

Acknowledgements

We thank Drs. Mark Nachtigal and Mary Lynn Duckworth for their critical comments. We also thank Mr. Wenjun Liu for the analysis of microarray data. This work was in part supported by grants from the Canada Research Chair Program (#950-201825 to HD), the Canadian Institute of Health Research (MOP 77783 to HD) and the Manitoba Health Research Council (G00302513 to HD). SS and ZN are supported by the fellowships from NSERC (Natural Sciences and Engineering Research Council of Canada) (to SS) and the Manitoba Health Research Council (MHRC) (to ZN). HD is the holder of a Canada Research Chair in Genetic Modeling.

Author details

¹Department of Biochemistry and Medical Genetics, University of Manitoba, 745 Bannatyne Avenue, Winnipeg MB R3E 0J9, Canada. ²Manitoba Institute of Cell Biology, 675 McDermot Avenue, Winnipeg MB R3E 0V9, Canada. ³Department of Immunology, University of Manitoba, 750 McDermot Avenue, Winnipeg MB R3E 0T5, Canada. ⁴Regenerative Medicine Program, University of Manitoba, 745 Bannatyne Avenue, Winnipeg MB R3E 0J9, Canada.

Authors' contributions

SS, XW, SM and HD conceived and designed the experiments. SS, XW and HD generated the gene-targeting vector and produced the knockout mice and characterized the tumor phenotypes. SS and SM performed cytogenetic analysis. ZN, MR and SK provided reagents and technical supports for this study. SS and HD wrote the manuscript with subsequent contributions from all authors. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 16 January 2012 Accepted: 27 March 2012

Published: 27 March 2012

References

- Edwards BK, Ward E, Kohler BA, Ehemann C, Zauber AG, Anderson RN, Jemal A, Schymura MJ, Lansdorf-Vogelaar I, Seeff LC, et al: **Annual report to the nation on the status of cancer, 1975-2006, featuring colorectal cancer trends and impact of interventions (risk factors, screening, and treatment) to reduce future rates.** *Cancer* 2010, **116**:544-573.
- Fearon ER: **Molecular genetics of colorectal cancer.** *Annu Rev Pathol* 2011, **6**:479-507.
- Fearon ER, Vogelstein B: **A genetic model for colorectal tumorigenesis.** *Cell* 1990, **61**:759-767.
- Issa JP: **CpG island methylator phenotype in cancer.** *Nat Rev Cancer* 2004, **4**:988-993.
- Kim MS, Lee J, Sidransky D: **DNA methylation markers in colorectal cancer.** *Cancer Metastasis Rev* 2010, **29**:181-206.
- Grady WM, Carethers JM: **Genomic and epigenetic instability in colorectal cancer pathogenesis.** *Gastroenterology* 2008, **135**:1079-1099.
- Sheridan PL, Schorpp M, Voz ML, Jones KA: **Cloning of an SNF2/SWI2-related protein that binds specifically to the SPH motifs of the SV40 enhancer and to the HIV-1 promoter.** *J Biol Chem* 1995, **270**:4575-4587.
- Ding H, Descheemaeker K, Marynen P, Nelles L, Carvalho T, Carmo-Fonseca M, Collen D, Belayew A: **Characterization of a helicase-like transcription factor involved in the expression of the human plasminogen activator inhibitor-1 gene.** *DNA Cell Biol* 1996, **15**:429-442.
- Hayward-Lester A, Hewetson A, Beale EG, Oefner PJ, Doris PA, Chilton BS: **Cloning, characterization, and steroid-dependent posttranscriptional processing of RUSH-1 alpha and beta, two uteroglobin promoter-binding proteins.** *Mol Endocrinol* 1996, **10**:1335-1349.
- Gong X, Kaushal S, Ceccarelli E, Bogdanova N, Neville C, Nguyen T, Clark H, Khatib ZA, Valentine M, Look AT, Rosenthal N: **Developmental regulation of Zbu1, a DNA-binding member of the SWI2/SNF2 family.** *Dev Biol* 1997, **183**:166-182.
- Mahajan MC, Weissman SM: **DNA-dependent adenosine triphosphatase (helicase-like transcription factor) activates beta-globin transcription in K562 cells.** *Blood* 2002, **99**:348-356.
- Unk I, Hajdu I, Fatyol K, Hurwitz J, Yoon JH, Prakash L, Prakash S, Haracska L: **Human HLTf functions as a ubiquitin ligase for proliferating cell nuclear antigen polyubiquitination.** *Proc Natl Acad Sci USA* 2008, **105**:3768-3773.
- Motegi A, Liaw HJ, Lee KY, Roest HP, Maas A, Wu X, Moinova H, Markowitz SD, Ding H, Hoeijmakers JH, Myung K: **Polyubiquitination of proliferating cell nuclear antigen by HLTf and SHPRH prevents genomic instability from stalled replication forks.** *Proc Natl Acad Sci USA* 2008, **105**:12411-12416.
- Blastyak A, Hajdu I, Unk I, Haracska L: **Role of double-stranded DNA translocase activity of human HLTf in replication of damaged DNA.** *Mol Cell Biol* 2010, **30**:684-693.
- Lin JR, Zeman MK, Chen JY, Yee MC, Cimprich KA: **SHPRH and HLTf act in a damage-specific manner to coordinate different forms of postreplication repair and prevent mutagenesis.** *Mol Cell* 2011, **42**:237-249.
- Achar YJ, Balogh D, Haracska L: **Coordinated protein and DNA remodeling by human HLTf on stalled replication fork.** *Proc Natl Acad Sci USA* 2011, **108**:14073-14078.
- Moinova HR, Chen WD, Shen L, Smiraglia D, Olechnowicz J, Ravi L, Kasturi L, Myeroff L, Plass C, Parsons R, et al: **HLTf gene silencing in human colon cancer.** *Proc Natl Acad Sci USA* 2002, **99**:4562-4567.
- Bai AH, Tong JH, To KF, Chan MW, Man EP, Lo KW, Lee JF, Sung JJ, Leung WK: **Promoter hypermethylation of tumor-related genes in the progression of colorectal neoplasia.** *Int J Cancer* 2004, **112**:846-853.
- Kim YH, Petko Z, Dzieciatkowski S, Lin L, Ghiassi M, Stain S, Chapman WC, Washington MK, Willis J, Markowitz SD, Grady WM: **CpG island methylation of genes accumulates during the adenoma progression step of the multistep pathogenesis of colorectal cancer.** *Genes Chromosomes Cancer* 2006, **45**:781-789.
- Leung WK, To KF, Man EP, Chan MW, Bai AH, Hui AJ, Chan FK, Sung JJ: **Quantitative detection of promoter hypermethylation in multiple genes in the serum of patients with colorectal cancer.** *Am J Gastroenterol* 2005, **100**:2274-2279.
- Hibi K, Nakao A: **Highly-methylated colorectal cancers show poorly-differentiated phenotype.** *Anticancer Res* 2006, **26**:4263-4266.
- Wallner M, Herbst A, Behrens A, Crispin A, Stieber P, Goke B, Lamerz R, Kolligs FT: **Methylation of serum DNA is an independent prognostic marker in colorectal cancer.** *Clin Cancer Res* 2006, **12**:7347-7352.
- Leung WK, Yu J, Bai AH, Chan MW, Chan KK, To KF, Chan FK, Ng EK, Chung SC, Sung JJ: **Inactivation of helicase-like transcription factor by promoter hypermethylation in human gastric cancer.** *Mol Carcinog* 2003, **37**:91-97.
- Hamai Y, Oue N, Mitani Y, Nakayama H, Ito R, Matsusaki K, Yoshida K, Toge T, Yasui W: **DNA hypermethylation and histone hypoacetylation of the HLTf gene are associated with reduced expression in gastric carcinoma.** *Cancer Sci* 2003, **94**:692-698.
- Kim JJ, Chung SW, Kim JH, Kim JW, Oh JS, Kim S, Song SY, Park J, Kim DH: **Promoter methylation of helicase-like transcription factor is associated with the early stages of gastric cancer with family history.** *Ann Oncol* 2006, **17**:657-662.
- Herbst A, Wallner M, Rahmig K, Stieber P, Crispin A, Lamerz R, Kolligs FT: **Methylation of helicase-like transcription factor in serum of patients with colorectal cancer is an independent predictor of disease recurrence.** *Eur J Gastroenterol Hepatol* 2009, **21**:565-569.
- Su LK, Kinzler KW, Vogelstein B, Preisinger AC, Moser AR, Luongo C, Gould KA, Dove WF: **Multiple intestinal neoplasia caused by a mutation in the murine homolog of the APC gene.** *Science* 1992, **256**:668-670.
- Boivin GP, Washington K, Yang K, Ward JM, Pretlow TP, Russell R, Besselsen DG, Godfrey VL, Doetschman T, Dove WF, et al: **Pathology of mouse models of intestinal cancer: consensus report and recommendations.** *Gastroenterology* 2003, **124**:762-777.
- Dietrich WF, Lander ES, Smith JS, Moser AR, Gould KA, Luongo C, Borenstein N, Dove W: **Genetic identification of Mom-1, a major modifier locus affecting Min-induced intestinal neoplasia in the mouse.** *Cell* 1993, **75**:631-639.
- Shoemaker AR, Moser AR, Midgley CA, Clipson L, Newton MA, Dove WF: **A resistant genetic background leading to incomplete penetrance of intestinal neoplasia and reduced loss of heterozygosity in ApcMin/+ mice.** *Proc Natl Acad Sci USA* 1998, **95**:10826-10831.
- Bresalier R, Kim Y: *Malignant neoplasms of the large intestine* Philadelphia: Saunders; 1998.
- Rajagopalan H, Nowak MA, Vogelstein B, Lengauer C: **The significance of unstable chromosomes in colorectal cancer.** *Nat Rev Cancer* 2003, **3**:695-701.

33. Nowak MA, Komarova NL, Sengupta A, Jallepalli PV, Shih Ie M, Vogelstein B, Lengauer C: **The role of chromosomal instability in tumor initiation.** *Proc Natl Acad Sci USA* 2002, **99**:16226-16231.
34. Haigis KM, Caya JG, Reichelderfer M, Dove WF: **Intestinal adenomas can develop with a stable karyotype and stable microsatellites.** *Proc Natl Acad Sci USA* 2002, **99**:8927-8931.
35. Hinoi T, Akyol A, Theisen BK, Ferguson DO, Greenson JK, Williams BO, Cho KR, Fearon ER: **Mouse model of colonic adenoma-carcinoma progression based on somatic Apc inactivation.** *Cancer Res* 2007, **67**:9721-9730.
36. Lengauer C, Kinzler KW, Vogelstein B: **Genetic instability in colorectal cancers.** *Nature* 1997, **386**:623-627.
37. Masramon L, Ribas M, Cifuentes P, Arribas R, Garcia F, Egozcue J, Peinado MA, Miro R: **Cytogenetic characterization of two colon cell lines by using conventional G-banding, comparative genomic hybridization, and whole chromosome painting.** *Cancer Genet Cytogenet* 2000, **121**:17-21.
38. Roschke AV, Stover K, Tonon G, Schaffer AA, Kirsch IR: **Stable karyotypes in epithelial cancer cell lines despite high rates of ongoing structural and numerical chromosomal instability.** *Neoplasia* 2002, **4**:19-31.
39. Ribas M, Masramon L, Aiza G, Capella G, Miro R, Peinado MA: **The structural nature of chromosomal instability in colon cancer cells.** *FASEB J* 2003, **17**:289-291.
40. Thibodeau SN, Bren G, Schaid D: **Microsatellite instability in cancer of the proximal colon.** *Science* 1993, **260**:816-819.
41. Veigl ML, Kasturi L, Olechnowicz J, Ma AH, Lutterbaugh JD, Periyasamy S, Li GM, Drummond J, Modrich PL, Sedwick WD, Markowitz SD: **Biallelic inactivation of hMLH1 by epigenetic gene silencing, a novel mechanism causing human MSI cancers.** *Proc Natl Acad Sci USA* 1998, **95**:8698-8702.
42. Liu B, Nicolaidis NC, Markowitz S, Willson JK, Parsons RE, Jen J, Papadopolous N, Peltomaki P, de la Chapelle A, Hamilton SR, *et al*: **Mismatch repair gene defects in sporadic colorectal cancers with microsatellite instability.** *Nat Genet* 1995, **9**:48-55.
43. Unk I, Hajdu I, Blastyak A, Haracska L: **Role of yeast Rad5 and its human orthologs, HLTF and SHPRH in DNA damage tolerance.** *DNA Repair (Amst)* 2010, **9**:257-267.
44. Branzei D, Foiani M: **Maintaining genome stability at the replication fork.** *Nat Rev Mol Cell Biol* 2010, **11**:208-219.
45. Branzei D, Foiani M: **The checkpoint response to replication stress.** *DNA Repair (Amst)* 2009, **8**:1038-1046.
46. Brocardo MG, Borowiec JA, Henderson BR: **Adenomatous polyposis coli protein regulates the cellular response to DNA replication stress.** *Int J Biochem Cell Biol* 2011, **43**:1354-1364.
47. Wu X, Ding H: **Generation of conditional knockout alleles for PDGF-C.** *Genesis* 2007, **45**:653-657.
48. Wu X, Sandhu S, Nabi Z, Ding H: **Generation of a mouse model for studying the role of upregulated RTTEL1 activity in tumorigenesis.** *Transgenic Res* 2012, DOI 10.1007/s11248-011-9586-7.
49. Rastegar M, Hotta A, Pasceri P, Makarem M, Cheung AY, Elliott S, Park KJ, Adachi M, Jones FS, Clarke ID, *et al*: **MECP2 isoform-specific vectors with regulated expression for Rett syndrome gene therapy.** *PLoS One* 2009, **4**: e6810.
50. Ding H, Roncari L, Shannon P, Wu X, Lau N, Karaskova J, Gutmann DH, Squire JA, Nagy A, Guha A: **Astrocyte-specific expression of activated p21-ras results in malignant astrocytoma formation in a transgenic mouse model of human gliomas.** *Cancer Res* 2001, **61**:3826-3836.
51. Ding H, Schertzer M, Wu X, Gertsenstein M, Selig S, Kammori M, Pourvali R, Poon S, Vulto I, Chavez E, *et al*: **Regulation of murine telomere length by Rtel: an essential gene encoding a helicase-like protein.** *Cell* 2004, **117**:873-886.

doi:10.1186/1476-4598-11-18

Cite this article as: Sandhu *et al*: Loss of HLTF function promotes intestinal carcinogenesis. *Molecular Cancer* 2012 **11**:18.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

