

RESEARCH

Open Access

Identification of single nucleotide polymorphisms in the bovine Toll-like receptor 1 gene and association with health traits in cattle

Christopher D Russell^{1,2}, Stephanie Widdison^{1,2}, James A Leigh² and Tracey J Coffey^{1,2*}

Abstract

Bovine mastitis remains the most common and costly disease of dairy cattle worldwide. A complementary control measure to herd hygiene and vaccine development would be to selectively breed cattle with greater resistance to mammary infection. Toll-like receptor 1 (TLR1) has an integral role for the initiation and regulation of the immune response to microbial pathogens, and has been linked to numerous inflammatory diseases. The objective of this study was to investigate whether single nucleotide polymorphisms (SNPs) within the bovine TLR1 gene (*boTLR1*) are associated with clinical mastitis (CM).

Selected *boTLR1* SNPs were analysed within a Holstein Friesian herd. Significant associations were found for the tagging SNP -79 T > G and the 3'UTR SNP +2463 C > T. We observed favourable linkage of reduced CM with increased milk fat and protein, indicating selection for these markers would not be detrimental to milk quality. Furthermore, we present evidence that some of these *boTLR1* SNPs underpin functional variation in bovine TLR1. Animals with the GG genotype (from the tag SNP -79 T > G) had significantly lower *boTLR1* expression in milk somatic cells when compared with TT or TG animals. In addition, stimulation of leucocytes from GG animals with the TLR1-ligand Pam3csk4 resulted in significantly lower levels of CXCL8 mRNA and protein. SNPs in *boTLR1* were significantly associated with CM. In addition we have identified a bovine population with impaired *boTLR1* expression and function. This may have additional implications for animal health and warrants further investigation to determine the suitability of identified SNPs as markers for disease susceptibility.

Introduction

Bovine mastitis is an inflammatory udder disease of great economic importance to the dairy industry. Mastitis is caused mainly by intramammary bacterial infection hence effective disease control measures rely upon farm management practices to limit the duration of infection and to restrict contagious spread of pathogens through the herd. As a result of these control measures opportunistic pathogens such as *Escherichia coli* and *Streptococcus uberis* predominate, and are now the leading cause of mastitis in the UK and common causes of mastitis worldwide [1]. Vaccination programmes may be required to further control mastitis [2], however no effective vaccines are currently available. An additional long term and complementary strategy is the genetic

selection of cattle that are less prone to mastitis. A lack of phenotypic data on clinical mastitis (CM) limits many breeding programmes, which select on the basis of a lowered somatic cell count (SCC) as a surrogate of CM [3]. The correlation between basal SCC and CM continues to be questioned [4,5] and the elucidation of more accurate genetic risk factors for CM is required. The identification of single nucleotide polymorphisms (SNPs) within genes involved with the mammary innate immune response are receiving particular interest as DNA markers for CM susceptibility [6-9].

The innate immune system provides an early defence to microbial pathogens [10] and involves cell surface proteins from the Toll-like receptor (TLR) family. TLRs are a structurally conserved type I membrane-bound class of pathogen recognition receptor (PRR), that can differentiate between an array of pathogen-associated molecular patterns (PAMPs) [11,12]. Functional expression of TLRs on resident mammary immune cells and

* Correspondence: tracey.coffey@nottingham.ac.uk

¹Bovine Genomics Group, Institute for Animal Health, Compton, Berkshire, RG20 7NN, UK

Full list of author information is available at the end of the article

epithelia has been shown to be important in the initiation and control of innate immune responses towards mastitis-causing bacteria [13-16]. Ten functional bovine (bo) TLRs have been characterised (Coffey et al. unpublished) [17], revealing close identity to their human orthologues [18]. Each individual TLR possesses its own ligand recognition repertoire, and upon stimulation activate internal adaptor proteins to initiate signalling for the NF κ B-mediated production of pro-inflammatory cytokines including the chemokine CXCL8 [19]. During mastitis, CXCL8 production increases rapidly [20] providing a potent chemotactic gradient to promote neutrophil movement into the mammary gland, a vital early defence mechanism against infection.

The presence of SNPs within mammalian TLRs has been shown to predispose susceptibility to a number of inflammatory diseases [21-23]. TLR1 variants have been postulated to underlie the altered immune responses to a wide spectrum of bacterial pathogens. In humans, a non-synonymous *TLR1* SNP located within the transmembrane domain, which had an impact on TLR1 trafficking, NF κ B activity and cytokine output, was found to have strong associations with susceptibility to and progression of infection by Gram positive bacteria and *Mycobacterium leprae* [24,25]. Polymorphisms affecting bo*TLR1* may therefore predispose excessive or poor immune responses to pathogens and hence could confer an increased risk of mammary infection.

Mammalian TLR1 is a member of a unique TLR subfamily that includes TLR6 and TLR10. All have the ability to form heterodimers with TLR2 to expand the repertoire of recognised ligands. TLR1/TLR2 complexes mediate cellular responses to natural triacylated lipoprotein structures [26,27], which are cell wall constituents of Gram-positive and Gram-negative bacteria [28]. TLR6/TLR2 complexes expand ligand recognition to diacylated lipoproteins, although in cattle the boTLR1/TLR2 complex may be further activated by diacylated lipoproteins [29]. Whilst more investigation is required to uncover their function, TLR10/TLR2 complexes may extend recognition of TLR1 agonists via an unknown signalling mechanism [30]. Phylogenetic analysis of bo*TLR1*, 6 and 10 reveals common ancestry with all three genes located in tandem to form a ~69 kb *TLR6-TLR1-TLR10* gene cluster on *Bos taurus* chromosome 6 (bta6) [31]. Genome-wide quantitative trait locus (QTL) studies reveal the bo*TLR6-TLR1-TLR10* gene cluster to be situated within a dense QTL region for a number of milk production traits and CM [32,33]. Furthermore, bo*TLR1* has been highlighted as a strong candidate for underlying QTL regions for disease resistance such as mastitis [34].

Despite strong QTL evidence, associative studies between bo*TLR1* and mastitis are limited. The bo*TLR1*

coding sequence (CDS) is highly polymorphic, containing many SNPs across several cattle breeds [35]. Four SNPs within the CDS of bo*TLR1* were identified in Holstein cows [36], but only one non-synonymous SNP, located within the transmembrane domain, was reported to result in highly significant differences in SCC between all genotype populations. More investigation is required to verify such studies, and to establish associations with novel bo*TLR1* SNPs/haplotypes. More importantly, the ability to use data indicating actual CM incidence, rather than a surrogate such as SCC, and associate this with specific genetic markers has the potential to yield more accurate information.

We report the identification of bo*TLR1* SNPs within coding and non-coding regions, and their potential as genetic markers for mastitis. We further demonstrate evidence of variable receptor response between defined bo*TLR1* SNP genotypes.

Materials and methods

Ethics statement

All experiments conformed to local and national guidelines on the use of experimental animals.

Sample population

The herd of Holstein Friesian cows used during this project (Mayfield Dairy, Institute for Animal Health, Compton, UK) contributes to the National Milk Records (NMR) [37] which submits production records for each animal to a database accessed through the InterHerd software programme (University of Reading/PAN Live-stock services team, UK). This, along with health data held locally, provided comprehensive information for each cow within the herd. All animal data were recorded within the years 2001-2010, during which, the mean CM incidence for the milking herd was 60 cases per 100 cows per year.

Isolation and extraction of genomic DNA from milk somatic cells

Milk samples (50 mL) were held on ice prior to centrifugation (3000 g; 5 min; 4°C). Supernatants were carefully removed along with any residual fat layer. Remaining casein micelles were removed using an adaptation of a procedure devised previously [38]. In brief, cell pellets were re-suspended in 1.5 mL PBS/EDTA solution (1 mL PBS, 300 μ L 0.5 M EDTA and 200 μ L TE buffer - 10 mM-Tris-HCl-1 EDTA pH 7.6), mixed well and allowed to stand at room temperature (RT) for 10 min. Cells were re-pelleted by centrifugation (7000 g; 1 min; RT), the supernatant discarded and cells re-suspended in 1.5 mL of TE buffer. Cells were pelleted again (7000 g; 1 min) and resuspended in 200 μ L PBS.

Milk somatic cells were processed using the Qiagen DNeasy blood and tissue kit for genomic DNA isolation (Qiagen Ltd, Sussex, UK). Briefly, 20 μ L proteinase K solution (Qiagen Ltd.) was added to resuspended cells and mixed well before being incubated overnight at 4°C. To ensure RNA-free genomic DNA, 4 μ L RNase A (100 mg/mL) was added following overnight incubation and samples left for 2 min at RT. DNA was extracted using the column protocol according to manufacturer's instructions (Qiagen Ltd). The quantity and quality of DNA was analysed using a NanoDrop 8000 spectrophotometer (Thermo Scientific, USA) and agarose gel electrophoresis.

Identification and sequencing of sample population for boTLR1 SNPs

BoTLR1-specific primers for PCR amplification of coding and non-coding genomic regions were designed based on the *Bos taurus* genome annotation build: Btau_4.0 (NC_007304.4). Sequencing of PCR amplicons from a DNA pool comprising animals with high (greater than six lifetime CM cases) or no mastitis case histories was initially chosen as a method of accelerating detection of disease-associated SNPs. Selected animals were unrelated and shared similar ages and lactation histories. PCR was performed using DreamTaq™ DNA polymerase (Fermentas UK, York, UK) following the manufacturer's protocol. Sequencing of purified PCR products was carried out using BigDye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, Warrington, UK). Samples were sequenced by capillary electrophoresis on an ABI PRISM 310 Genetic Analyser (Applied Biosystems), and analysed using the Sequencher™ package version 4.1.4 (Gene Codes Corporation, Ann Arbor, MI, USA). All primers (Additional file 1) were designed using Primer3 software [39] and synthesised by Sigma-Genosys Ltd. (Haverhill, UK). All boTLR1 SNP offsets are given relative to their position (bp) from the A nucleotide of the ATG codon. Sequencing of PCR amplicons was extensively used to genotype the herd population.

Restriction fragment length polymorphism (RFLP) - PCR analysis

In addition to sequencing of PCR amplicons, SNPs +798 C > T and +1762A > G were genotyped using RFLP analysis of PCR products; whereby restriction enzyme MboII cuts in the presence of the C allele for SNP +798 C > T, and restriction enzyme BclI cuts only in the presence of the A allele (Isoleucine) for the non-synonymous SNP +1762A > G; as described previously [36]. Restriction digests were performed in a final volume of 20 μ L containing 10 μ L PCR product and 2U of appropriate restriction enzyme. Digests were incubated at the recommended temperature for 2 h, with a subsequent

heat inactivation of enzyme at 65°C for 20 min. Homozygous and heterozygous samples, confirmed by sequencing, were included as controls.

Identifying associations between SNPs and clinical mastitis and milk quality

Identified SNPs were correlated to health or milk productivity trait data available through the InterHerd software programme and NMR data. These included CM case histories, milk SCC, mean 305 d milk yields and mean milk protein/fat concentrations. All Animal data were recorded within and up to, the first three lactations only. Median SCC was transformed logarithmically as $NEWSCC = \text{LOG}_2(\text{SCC}/100\,000) + 3$. This model accounts for the skewness of data distribution as used previously [3]. CM was analysed for presence or absence of cases across the entire milking tenure of their first three lactations. Mean incidence of CM is presented as cases per cow per year ($[\text{total number of cases}/\text{total number of milking days}] * 365$) within the first three lactations. CM was defined as an individual clinically apparent event, coupled with a rise in milk SCC and/or isolation of pathogen, recorded on a single day within any/all quarters during a lactation period. Repeated cases within the same quarter were noted but only regarded as a new case after seven days return to milking.

Statistical analysis

Genotypic frequencies were tested for deviation from Hardy-Weinberg equilibrium according to Chi² test and level of significance at 1 degree of freedom. Haplotypes and linkage disequilibrium (LD) scores (based on the r^2 statistic), were analysed using the Haploview programme [40]. Owing to substantial positive skewness (due to zero values) of data distribution, mean CM rate was transformed: $NEWCM = \text{LOG}_{10}(\text{CM} + 1)$. A genotypic model was used for separate genotype pair-wise comparisons. One-way analysis of variance (ANOVA) and Tukey's multiple comparisons for CM rate, milking traits and functional data means, were performed using Prism[®] 5.04 software (Graph Pad Software, Inc, CA, USA). Presence of any CM event across all lactations (CM recorded = 1, no recorded CM = 0) between genotypes were pair-wise tested for significance by a Chi² odds ratio test using Minitab[®] 16 software (Minitab Statistical Software, Inc, PA, USA). Calculation of Pearson correlation coefficients between factors and analysis of CM response in a General linear model (GLM) was performed using Minitab[®] 16 software.

Isolation of polymorphonuclear leukocytes from bovine blood

For stimulation assays, polymorphonuclear leukocytes (PMNs) were isolated from 50 mL blood samples of healthy, age-matched, genotyped cattle that had recorded

no intramammary infection in the two weeks prior to sample collection. Briefly, blood was collected into EDTA (final concentration of ~1.5%w/v), centrifuged at RT (15 min; 1000 g) and the plasma and buffy coat phases carefully removed. For every 2.5 g of lower blood phase, 10 mL of H₂O was added. The mixture was agitated and left for 40 s before adding half the water volume of NaCl/PO₄ reagent (2.7% NaCl/PO₄ reagent: 2.7 g NaCl into 10 mL 0.132 M phosphate buffer (18.74 g Na₂HPO₄ and 17.96 g KH₂PO₄ into 1 L H₂O, pH 6.8) and made up to 100 mL with H₂O) to restore the isotonic strength. PMNs were pelleted by centrifugation (125 g; 10 min; RT), the supernatant discarded and the cells re-suspended in 1.5 mL of PBS. This was repeated twice to wash the cells. Typically 1 × 10⁸ PMN cells were obtained per 50 mL sample. FACS analysis of polymorphic PMN samples suggested little difference in cell populations and purity obtained from the different genotypes.

Isolated PMNs were seeded to 1 × 10⁶/1 mL in RPMI 1640 + GlutaMAX™ with 10% foetal calf serum (FCS) (both Invitrogen Ltd. UK) and 100 units mL⁻¹/100 µg mL⁻¹ Penicillin/Streptomycin (Sigma, St Louis, MO, USA). Prior to stimulation, seeded cells were incubated at 37°C, in 5% CO₂ for approximately 18 h in 24 well tissue culture plates.

Stimulation of PMNs with TLR ligands

All TLR ligands (InvivoGen, San Diego, CA, USA) were reconstituted in sterile endotoxin-free water according to the manufacturers' instructions, to a concentration of 1 mg mL⁻¹, and stored at 20°C. The synthetic TLR1-specific ligand PAM3 was added to the cells in 100 µL of culture media, at final concentrations of 100 ng/mL, 250 ng/mL and 500 ng/mL. A media control (containing no ligand) was used. LPS (100 ng/mL) was used as a non-TLR1 stimulant. Stimulated cells (4 h and 8 h post stimulation) were lysed and total RNA extracted. Supernatants were taken 24 h post stimulation for detection of cytokine production by ELISA.

RNA isolation and cDNA synthesis

Pelleted cell samples were lysed in RLT buffer for total RNA extraction using the RNeasy mini kit (Qiagen Ltd) according to manufacturer's instructions. Samples were DNase treated (DNAfree, Ambion, Austin, TX, USA) and analysed for quantity and quality. Total RNA was used for cDNA synthesis using SuperScript II Reverse Transcriptase (Invitrogen Ltd. UK) using the recommended protocol. Quantification of cDNA was carried out using the NanoDrop 8000 spectrophotometer.

Q-PCR analysis

Quantitative PCR (Q-PCR) was performed with the ABI Prism 7500 FAST Sequence Detection System (Applied

Biosystems) using FAST Universal Mastermix (Applied Biosystems) according to the manufacturer's protocol. Taqman[®] probe and primers for *GAPDH*, *CXCL8* and *Bcl2A1* have been designed previously [41-43]. Taqman[®] probe and primers for bo*TLR1* were designed from sequenced templates, avoiding any SNPs, using the Primer3 software [39] (Forward: 5'-GCA CCA CAG TGA GTC TGG AA -3', Reverse: 5'-GTA CGC CAA ACC AAC TGG AG -3', Probe: 5'-TGT GTG CTT GAT GAT AAT GGG TGT CCT -3'). All primers and probes were synthesised by Sigma-Genosys Ltd. and Eurogentec Ltd. (Romsey, UK) respectively. Probes were labelled at the 5' end with the reporter dye FAM (6-carboxyfluorescein) and at the 3' end with the quencher dye TAMRA (6-carboxytetramethylrhodamine). 100 ng of cDNA template from each animal were tested in triplicate and quantified by comparison with a standard curve from plasmid DNA of known copy number. Relative target gene expression was then calculated by normalising to the mean expression levels of the housekeeping genes glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*) and ribosomal protein, large, P2 (*RPLP2*). Analysis was performed using Microsoft[®] Excel 2008 (Microsoft Co., Redmond, WA, USA) and Prism[®] 5.04 software. Differences between groups were assessed by one-way analysis of variance (ANOVA) and Tukey's multiple comparisons.

ELISA

Levels of CXCL8 were determined by luminescent ELISA, using a pair of commercial human IL-8 specific antibodies (R&D Systems, Oxon, UK) according to a modified luminescence-based ELISA described previously [44]. Samples were tested in triplicate, with recombinant bovine CXCL8 (Kingfisher Biotech, Inc., MN, USA) as the standard. Levels of bovine IL-6 were determined using a commercial TMB ELISA kit according to the manufacturer's protocol (Thermo Fisher Scientific, Inc. USA). Differences between groups were assessed by one-way analysis of variance (ANOVA) and Tukey's multiple comparisons.

Results

Characterisation of bo*TLR1* SNPs

The genomic organisation of bo*TLR1* was confirmed using internal cDNA sequencing, and analysis of EST data submitted to the NCBI database. As described previously [31], the full length bo*TLR1* mRNA transcript comprises five exons and four introns. The CDS for full length bo*TLR1*; a 2379 bp transcript corresponding to a 793aa chain [45], is located within exon 5. Sequencing of bo*TLR1* within a Holstein Friesian herd (*n* = 246) revealed eleven SNPs; five of which are novel (Table 1). Eight of the eleven SNPs were exonic, six were located

Table 1 Identified SNPs within boTLR1 from Holstein Friesians.

SNP	Genomic (bta6) position	Amino acid position	Amino acid	Restriction site presence	Previously reported	dbSNP ID
-1205 <u>A</u> > G	60440366	-	-	-	No	-
-1151 <u>T</u> > A	60440312	-	-	-	No	-
-79 <u>T</u> > G	60439240	-	-	-	No	-
+798 <u>C</u> > T	60438363	266	Phe-Phe	MbolI	Yes ¹	ss73689409
+1641 <u>A</u> > C	60437520	547	Ser-Ser	Bst X-I	Yes ²	ss73689413
+1716 <u>G</u> > A	60437445	572	Lys-Lys	Bsi Y	Yes ²	ss73689414
+1762 <u>A</u> > G	60437324	587	Ile-Val	Bcl I	Yes ²	ss73689415
+2100 <u>C</u> > T	60436989	700	Phe-Phe	-	Yes ²	ss73689417
+2103 <u>T</u> > C	60436986	701	Val-Val	BsmAI	Yes ²	ss73689418
+2463 <u>C</u> > T	60436698	-	-	MslI	No	-
+2731A > G	60436430	-	-	-	No	-

Abbreviations: bta6 = *Bos taurus* chromosome 6, dbSNP = Single-Nucleotide Polymorphism Database [46]. All boTLR1 sequences were aligned with the published Btau_4.0 sequence NC_007304.4 (underlined allele denotes the published sequence). Light shading - synonymous coding SNPs, dark shading - non-synonymous coding SNP. All boTLR1 SNP offsets are given relative to their position (bp) from the A nucleotide of the first ATG codon. ¹ [31], ² [35].

in the CDS, of which only one was a non-synonymous change (+1762A > G, Iso > Val) positioned within the transmembrane domain, while the remaining two were located in the non-coding 3'UTR. The three non-exonic SNPs were identified within the introns of the 5' UTR (Figure 1).

Genotypes and allelic frequencies


Five boTLR1 SNPs: -79 T > G, +798 C > T, +1762A > G, +2463 C > T, and +2731A > G (Figure 1) were detected within the herd ($n = 156, 246, 125, 225$ and 232 respectively). The genotypic and allelic frequencies for each SNP are similar, with heterozygotes the most abundant genotypes (Table 2). All SNPs were found to be in agreement with Hardy-Weinberg equilibrium. The three SNPs -79 T > G, +798 C > T, and +2731A > G were in strong linkage disequilibrium ($LD = > 0.9$) (Figure 2) and as a result only the SNP -79 T > G was used for association analysis. LD between the tagging SNP -79 T > G and the non-synonymous +1762A > G was moderate ($LD = \sim 0.8$). LD between SNPs -79 T > G and +2463 C > T was lower. The lowest LD was observed between SNPs +1762A > G and +2463 C > T (Figure 2). Consequently, analysis of association with traits for SNPs -79 T > G, +1762A > G and +2463 C > T was performed individually. Preliminary haplotype construction for SNPs 79 T > G, +798 C > T, +1762A > G, +2463 C > T, and +2731A > G, yielded two major haplotypes of G₇₉T₇₉₈A₁₇₆₂C₂₄₆₃G₂₇₃₁ (frequency 0.42) and T₇₉C₇₉₈G₁₇₆₂T₂₄₆₃A₂₇₃₁ (frequency 0.41) (underlined allele denotes change from the published Btau_4.0 sequence NC_007304.4). Owing to the low number of individuals typed for all five markers ($n = 125$), haplotype association analysis was not performed.

Analysis of boTLR1 SNPs with clinical mastitis

The three boTLR1 SNPs -79 T > G, +1762A > G and +2463 C > T were analysed in Holstein Friesian cows for association with CM and milk production traits. Analysis of the tagging SNP -79 T > G revealed the TG and GG genotypes associated with an increased rate of CM (0.75 and 0.70 cases/cow/year, respectively) compared with the TT genotype (0.40 cases/cow/year) (Table 3). The association with CM rate was found to be significantly ($P < 0.05$) higher for the heterozygote (TG) population, the most abundant genotype, when compared with homozygote (TT) animals. The proportion of animals within each genotyped population having any episodes of CM during their first three lactations was highest in the GG genotype (70%) compared with TT and TG genotypes (58% and 60%, respectively), although this was found not to be statistically significant. Analysis of the non-synonymous, transmembrane domain SNP; +1762A > G, showed AA and AG animals had higher rates of CM (0.69 and 0.66 cases/cow/year, respectively) compared with GG animals (0.38 cases/cow/year), although these differences were not significantly different. Analysis of the SNP +2463 C > T located within the 3'UTR revealed that CC and CT animals had higher rates of CM (0.70 cases/cow/year) compared to TT animals (0.39 cases/cow/year). However, only the increase seen in the most abundant genotype, heterozygous CT cows, was shown to be statistically significant ($P < 0.05$).

Association of boTLR1 SNPs with milk productivity traits

Analysis of the productivity traits within the genotyped animals revealed that the TG and GG genotypes of the tagging SNP -79 T > G associated significantly with lowered milk fat (TG = $P < 0.05$, GG = $P < 0.01$) and protein (TG = $P < 0.01$, GG = $P < 0.001$) concentrations



when compared to those observed in the TT genotype (Table 3). A similar trend was detected in the genotypes of the SNP +1762A > G; AA and AG genotypes associated with lower milk fat and protein concentrations; however these differences were not statistically significant. Analysis of genotypes within the SNP +2463 C >

T revealed that CT and CC genotypes produced milk with lower protein concentration ($P < 0.05$) when compared with the TT population. The same genotypes tended to show lower milk fat concentration, but these differences were not significant. No differences in either milk yield or mean SCC were detected between any of

Table 2 Genotype and allele frequencies including significance from Hardy-Weinberg equilibrium for bo*TLR1* SNPs selected for trait association.

SNP	Genotype	n =	Genotype Frequency	HWE	Allele	Allele Frequency
-79 T > G	TT	53	0.34	0.831	T	0.568
	TG	71	0.46		G	0.432
	GG	32	0.20			
+798 C > T ¹	CC	78	0.32	0.885	C	0.543
	CT	122	0.49		T	0.456
	TT	46	0.19			
+1762A > G ¹	AA	33	0.26	0.326	A	0.495
	GA	57	0.46		G	0.505
	GG	35	0.28			
+2463 C > T	CC	64	0.28	0.163	C	0.507
	CT	102	0.46		T	0.493
	TT	59	0.26			
+2731A > G	AA	77	0.33	0.611	A	0.546
	AG	110	0.48		G	0.454
	GG	45	0.19			

¹SNPs genotyped by RFLP-PCR. HWE not significant at 1 degree of freedom.




Figure 2 Linkage disequilibrium (LD) plot for five extensively genotyped SNPs in relation to the *boTLR1* gene model. Open box represents *boTLR1* coding exon, grey boxes represent non-coding untranslated (UTR) exonic regions. The non-synonymous amino acid change is shown in brackets below corresponding coding SNP. The graphical representation, generated by Haploview, of the r^2 LD relationship between each SNP is expressed on a gray-scale gradient; where black - very high LD ($r^2 = 1.0$) and white - very low LD (< 0.1). The r^2 value for each pair-wise comparison is shown, where no value is indicated $r^2 = 1$.

the genotyped populations, for any of the SNPs described.

Multiple comparisons

Correlation coefficient analysis was performed revealing a weak positive correlation between CM and SCC (Pearson coefficient = 0.198) and weak negative correlations between CM and milk fat and milk protein percentages (Pearson coefficient = -0.2 and -0.1 respectively). All significant associations between individual SNP genotypes and CM were found to survive adjustments for each factor using a General linear model (GLM).

Q-PCR analysis of basal *boTLR1* mRNA abundance

Milk somatic cells from cows with representative *boTLR1* -79 T > G SNP genotypes were assessed for differences in basal *boTLR1* expression. All animals were of similar age and stage of lactation and none had any known health issues. Q-PCR assays were performed

using RNA from five animals of each genotype. Cows with the GG genotype demonstrated lower *boTLR1* mRNA abundance ($P < 0.01$) compared with either the TT or TG genotypes (Figure 3). Due to close homology and potential for co-expression, the abundance of *boTLR6* mRNA was also measured and revealed no significant differences.

Variation in immune responses to a TLR1 ligand

Cells from twelve animals (four from each representative *boTLR1* -79 T > G SNP genotype) were stimulated for up to 24 h with the ligand Pam3csk4 (PAM₃), which has an affinity for TLR1/TLR2 heterodimers [49]. *CXCL8* and *IL-6* mRNA abundance and *CXCL8* and *IL-6* production were used as markers for TLR1 stimulation. Q-PCR analysis indicated that animals with the GG genotype had lower levels of *CXCL8* mRNA 4 h post-stimulation with PAM₃ compared to equivalent samples from animals with either of the TT or TG genotypes (Figure 4). Samples from animals that were homozygous TT showed significantly greater ($P < 0.05$) *CXCL8* mRNA abundance 4 h after stimulation with PAM₃ at concentrations of 250 and 500 ng/mL when compared to GG homozygotes (Figure 4a). Homozygous TT animals tended to show greater *IL-6* abundance 4 h after stimulation however no significant differences were detected between genotypes across all ligand concentrations (data not shown). A similar trend in *CXCL8* and *IL-6* data was observed from cells stimulated for 8 h (data not shown). Animals that were homozygous TT displayed significantly greater ($P < 0.05$) *IL-6* mRNA abundance 8 h after stimulation with PAM₃ at 500 ng/mL when compared to GG homozygotes (data not shown).

The mRNA abundance of the anti-apoptotic factor *Bcl2A1* was used as an additional marker to indicate differences in TLR1 stimulation. Following stimulation with PAM₃, Q-PCR analysis showed greater levels of *Bcl2A1* mRNA in animals with the TT genotype compared to those with either the TG or GG genotypes (Figure 4b). Significantly ($P < 0.05$) greater levels of mRNA were detected in samples from animals with the TT genotype compared to those with the TG genotype following stimulation with PAM₃ at 250 ng/mL and 500 ng/mL. In contrast, a statistically significant difference between the TT and GG genotypes was only detected following stimulation at the higher concentration.

Supernatants (24 h post-stimulation) were assayed for *CXCL8* and *IL-6* production by ELISA. *CXCL8* data showed a similar trend to those obtained by Q-PCR, with significantly ($P < 0.05$) lower levels of *CXCL8* produced by cows with the GG genotype than those with the TT genotype following stimulation with 500 ng/mL PAM₃ (Figure 5a). Individual variation was largest within those samples from TG heterozygotes, with


Table 3 Analysis of boTLR1 SNP genotypes by CM and milk productivity traits over the first three lactations.

SNP	Gen.	n =	P% with CM ¹	Mean CM rate ²	P = ₃	Median SCC ⁴	P = ₃	Mean 305 d milk yield ⁴	P = ₃	Mean milk fat (%) ⁴	P = ₃	Mean milk protein (%) ⁴	P = ³
tag-79 T > G	<u>TT</u>	78*	58%	0.40 (± 0.11)	-	1.96 (± 0.63)	-	7868 (± 1181)	-	4.15 (± 0.41)	-	3.28 (± 0.19)	-
	TG	122*	60%	0.75 (± 0.18)	< 0.05	1.85 (± 0.61)	-	8058 (± 1139)	-	4.02 (± 0.38)	< 0.05	3.2 (± 0.17)	< 0.01
	GG	46*	70%	0.70 (± 0.24)	-	1.98 (± 0.67)	-	7983 (± 1273)	-	3.89 (± 0.41)	< 0.01	3.16 (± 0.18)	< 0.001
+1762A > G	AA	33	63%	0.69 (± 0.27)	-	1.96 (± 0.64)	-	8101 (± 1106)	-	3.95 (± 0.41)	-	3.19 (± 0.18)	-
	GA	57	61%	0.66 (± 0.18)	-	1.89 (± 0.69)	-	8067 (± 1330)	-	4.04 (± 0.45)	-	3.21 (± 0.2)	-
	<u>GG</u>	35	59%	0.38 (± 0.16)	-	1.90 (± 0.62)	-	8239 (± 1185)	-	4.05 (± 0.38)	-	3.25 (± 0.16)	-
+2463 C > T	CC	64	67%	0.70 (± 0.24)	-	1.95 (± 0.66)	-	7841 (± 1067)	-	4.02 (± 0.45)	-	3.19 (± 0.18)	< 0.05
	CT	102	61%	0.70 (± 0.14)	< 0.05	1.90 (± 0.63)	-	8137 (± 1206)	-	3.98 (± 0.45)	-	3.18 (± 0.2)	< 0.05
	<u>TT</u>	59	59%	0.39 (± 0.11)	-	1.88 (± 0.63)	-	7940 (± 1262)	-	4.13 (± 0.38)	-	3.27 (± 0.18)	-

Abbreviations: Gen. = genotype, P% = population percentage, CM = clinical mastitis, SCC = somatic cell count. * Numbers from tagged SNP +798 C > T in Table 2. ¹No significance was detected between genotypes by Chi² odds ratio test. ²Mean ± 95% confidence interval given in column parentheses. ³P values analysed by ANOVA and Tukey's multiple comparisons, showing only results tested against the underlined (lower CM rate) genotype; - (dash) denotes no significance. ⁴Standard deviation from the mean given in column parentheses.


higher concentrations of PAM₃ resulting in increased variability in CXCL8 production. IL-6 data showed significantly ($P < 0.05$) lower levels produced by cows with the GG genotype than the TT genotype following stimulation with 500 ng/mL PAM₃ (Figure 5b). Overall

stimulation with TLR1 ligand across all concentrations did not increase cytokine production in GG variants. The TLR4 ligand LPS was used to assess if the detected variation in response to PAM₃ reflected the ability of the cell populations to respond to any ligand. The response to LPS was variable across all genotypes and did not segregate by the defined genotypes (Additional file 2).




Discussion

In this study we analysed five SNPs within exonic and non-exonic regions of boTLR1 for susceptibility to CM. We identified significant associations between the tagging SNP -79 T > G, and the 3'UTR SNP +2463 C > T and susceptibility to CM. Rates of CM for genotypes -79 TT and +2463 TT were much lower in comparison to the homozygous genotypes (-79 GG and +2463 CC) and significantly lower than their respective heterozygous genotypes. The CM rate differences observed between defined genotypes can have a significant impact on both herd welfare and farm expenditure. For example, the -79 TT animals recorded on average 40 CM cases per 100 cows per year, below the historic national average for England and Wales, which has been estimated at 47 cases [1]. GG and TG animals recorded 70 and 75 cases respectively, a substantial increase from the national average. As a simple economic evaluation, if estimated average costs for a clinical case are £175 [50] potential losses per 100 cows when compared to TT animals equate to £6125 and £5250 for TG and GG animals respectively.



Dairy cows are subjected to intense selection pressures for higher milk yields and reduced SCC that may unintentionally be detrimental to CM incidence [5]. However, in this study, no significant differences were observed between boTLR1 SNPs for mean 305 d milk yields or SCC. This contradicts a study of 208 Chinese Holsteins, in which the non-synonymous SNP +1762A > G associated with highly significant differences in milk SCC. Animals with an AA (Isoleucine) at this position had lower SCC and were classed as more resistant to mastitis [36]. In contrast, our results suggest the GG (Valine) at this location is the more favourable genotype in terms of having a lower CM incidence in comparison to AA. Although this discrepancy may reflect breed and herd variation, it may also demonstrate the dichotomy



that the surrogate measure of mastitis is also the presence of the effector cell which can at times eliminate infection in the absence of overt clinical signs.

All boTLR1 SNP genotypes associated with a lowered rate of CM display significantly greater milk fat and protein concentrations when compared to the other two genotypes. It is unclear whether a functional association exists between milk fat and protein concentrations and mastitis susceptibility. Interestingly the casein gene cluster is located ~28 Mb from the TLR6-TLR1-TLR10 gene cluster on bta6 [51]. Protein and fat contents are financially rewarded milk qualities and breeding for these factors could influence boTLR1 genotypes.

In addition to being one of the first studies to examine association between boTLR1 SNPs and CM incidence, we further demonstrate evidence of variable receptor

responses between the defined genotypes. In this study animals segregated by -79 T > G SNP genotypes exhibited significantly lower boTLR1 expression in GG animals when compared with TT or TG. This difference, which appeared to be specific for boTLR1 as it was not seen for boTLR6 a paralogue of boTLR1, may indicate interference with transcription. Although more variable, intermediate basal expression levels observed in TG cows suggests that the presence of a T allele is compensating for the lowered expression originating from the G allele. Preliminary analysis for causal mutations using the programme MOTIF [52] highlighted the -79 T > G SNP as part of an Octamer-1 (Oct-1) motif. Oct-1 is thought to work closely with AP-1 and GATA-1 as important regulatory elements for TLR expression [53]. However the motif was only identified with the T variant (threshold > 80%), and was absent in those with the G substitution. Although this is an attractive hypothesis it is also possible that the SNP may simply be a marker for other unidentified causal mutations.

Currently there are no boTLR1 antibodies available to directly compare levels of protein expression. To ascertain the functional consequence of potentially reduced boTLR1 expression, production of cytokines known to be induced by TLR1 was determined from cells stimulated with the TLR1 ligand PAM₃. Our study demonstrated genotype-specific variation in cytokine responses of bovine PMN to PAM₃. Across all time points and ligand concentrations analysed, PAM₃ stimulation of cells from animals segregated by tagging -79 T > G SNP genotypes revealed lower levels of CXCL8 and IL-6 mRNA and protein from those of a GG genotype and consistently high CXCL8 and IL-6 response from those with TT genotypes. The output of cytokines from cells from the TG animals was the most variable. The NF-κB-transcribed gene [54] encoding the pro-survival protein *Bcl2A1*, which is up-regulated in neutrophils following stimulation with TLR ligands including PAM₃ [55], was used as an additional marker of effective boTLR1 stimulation. Significantly greater levels of the *Bcl2A1* transcript were detected in animals with the TT genotype compared to those with the TG or GG genotypes. All genotypes were also stimulated with LPS, and no differences were detected in levels of CXCL8 or IL6. This further indicates that differences observed following stimulation with PAM₃ were influenced by variation in the response via boTLR1.

PMNs contain a high proportion of neutrophils, whose primary role is the phagocytosis of microbes to contain and eliminate infection. However stimulated neutrophil TLRs further induce the neutrophil to synthesise cytokines that influence their own activity and survival, recruit additional immune cells and modulate the immune response [56-59]. The greater

expression of *Bcl2A1* in animals with the TT genotype may promote PMN survival to complement the increased cytokine output. Reduced boTLR1 stimulation by pathogens in GG cows may equate to a dampened inflammatory response thus enabling establishment and persistence of infection. In terms of influence upon mastitis susceptibility, GG cows have a higher CM rate than TT cows; however heterozygous TG cows have a similar rate to GG cows, despite expressing boTLR1 at a similar level to a TT cow. TG animals showed intermediary cytokine responses to TLR1 ligand, and like GG animals, lower expression of *Bcl2A1*. This may suggest that differences in boTLR1 transcript levels alone may not account for the variation in receptor response, and/or that post-transcriptional control of translation may also be affected by the cumulative presence of SNPs. The animals used for the functional assays conform to observed haplotypes: TT = T₇₉C₇₉₈G₁₇₆₂T₂₄₆₃A₂₇₃₁ and GG = G₇₉T₇₉₈A₁₇₆₂C₂₄₆₃G₂₇₃₁; with TG animals heterozygous for each SNP. In terms of the likelihood of being causative of a variable TLR1 response, the non-synonymous SNP +1762 Iso > Val is an unlikely candidate. Both Isoleucine and Valine are non-polar branched amino acids with hydrophobic properties, and so a substitution within the transmembrane domain should not impact the tertiary structure; although further targeted structure/function studies are required to verify this. Furthermore, influence of polymorphisms within TLR-related or target genes such as *TLR2* and *CXCL8* should be considered.

In conclusion the findings presented here demonstrate an association between SNPs, common to boTLR1, and CM susceptibility. Furthermore, we present evidence that some of these SNPs underpin functional variation in boTLR1. A rapid immune response, conferred by the favourable boTLR1 SNP -79 TT variant, could reduce detrimental clinical manifestations of diseases via an efficient yet controlled influx of inflammatory cells to neutralise pathogens and control infection. The complexity of mastitis infection suggests a polygenic and multi-factorial immune response comprising many different proteins which will in turn interact with one another. The ability to detect significant associations with CM for a single gene indicates the importance of boTLR1 related responses in the control of bacterial disease in cattle. A bovine population with impaired boTLR1 expression may also be more susceptible to other diseases and health issues. Preliminary evidence for this comes from our study where it was observed that animals with the GG variant seemed to be prematurely culled compared to other populations.

The potential of SNPs within boTLR1 to act as genetic markers for altered susceptibility to mastitis warrants further investigation. The favourable linkage of lowered

CM with increased milk fat and protein concentrations in the boTLR1 variants demonstrated here indicates selection for lowered CM using these markers would not be detrimental to milk quality, as has been previously suggested in studies linking SNPs with increased SCC as a marker for CM susceptibility.

Additional material

Additional file 1: Primers used for PCR amplification of boTLR1 genomic regions, RFLP PCR digests, and quantitative PCR (QPCR) assays.

Additional file 2: Table to show expression levels of CXCL8 and Bcl2 A1 and CXCL8 and IL-6 production levels following PMN stimulation with LPS.

Acknowledgements

The authors wish to acknowledge the help of R Parsons (Institute for Animal Health/IAH Compton) and A Bowen (IAH Compton) for contributing valuable tools to this project. We also thank the staff at Mayfield Dairy, IAH Compton, UK, for animal handling and sampling assistance. This study is part of a PhD project funded through a Doctoral Training Partnership from the Biotechnology and Biological Sciences Research Council administered through the IAH.

Author details

¹Bovine Genomics Group, Institute for Animal Health, Compton, Berkshire, RG20 7NN, UK. ²The School of Veterinary Medicine and Science, The University of Nottingham, Sutton Bonington Campus, Sutton Bonington, Leicestershire, LE12 5RD, UK.

Authors' contributions

Conceived and designed the experiments: CDR, SW, JAL, TJC. Performed the experiments: CDR. Analysis and discussion of data: CDR, SW, JAL, TJC. Wrote the manuscript: CDR. Edited the manuscript: SW, JAL, TJC. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 20 December 2011 Accepted: 14 March 2012

Published: 14 March 2012

References

- Bradley AJ, Leach KA, Breen JE, Green LE, Green MJ: Survey of the incidence and aetiology of mastitis on dairy farms in England and Wales. *Vet Rec* 2007, **160**:253-257.
- Leigh JA: Streptococcus uberis: a permanent barrier to the control of bovine mastitis? *Vet J* 1999, **157**:225-238.
- Rupp R, Boichard D: Genetic parameters for clinical mastitis, somatic cell score, production, udder type traits, and milking ease in first lactation Holsteins. *J Dairy Sci* 1999, **82**:2198-2204.
- Barkema HW, Schukken YH, Lam TJ, Beiboer ML, Wilmink H, Benedictus G, Brand A: Incidence of clinical mastitis in dairy herds grouped in three categories by bulk milk somatic cell counts. *J Dairy Sci* 1998, **81**:411-419.
- Suriyasathaporn W, Schukken YH, Nielsen M, Brand A: Low somatic cell count: a risk factor for subsequent clinical mastitis in a dairy herd. *J Dairy Sci* 2000, **83**:1248-1255.
- Skelding A, Schenkel FS, Sharma BS, Verschoor C, Pant S, Biggar G, Boermans H, Karrow N: Identification of single nucleotide polymorphisms in the bovine interleukin-12 and interleukin-23 receptor genes and their associations with health and production traits in Holstein cows. *J Dairy Sci* 2010, **93**:4860-4871.
- Beecher C, Daly M, Childs S, Berry DP, Magee DA, McCarthy TV, Giblin L: Polymorphisms in bovine immune genes and their associations with somatic cell count and milk production in dairy cattle. *BMC Genet* 2010, **11**:99.
- Chen R, Yang Z, Ji D, Mao Y, Chen Y, Li Y, Wu H, Wang X, Chang L: Polymorphisms of the IL8 gene correlate with milking traits, SCS and mRNA level in Chinese Holstein. *Mol Biol Rep* 2011, **36**:4083-4088.
- Sharma BS, Leyva I, Schenkel F, Karrow NA: Association of toll-like receptor 4 polymorphisms with somatic cell score and lactation persistency in Holstein bulls. *J Dairy Sci* 2006, **89**:3626-3635.
- Oviedo-Boyso J, Valdez-Alarcon JJ, Cajero-Juarez M, Ochoa-Zarzosa A, Lopez-Meza JE, Bravo-Patino A, Baizabal-Aguirre VM: Innate immune response of bovine mammary gland to pathogenic bacteria responsible for mastitis. *J Infect* 2007, **54**:399-409.
- Kawai T, Akira S: Pathogen recognition with Toll-like receptors. *Curr Opin Immunol* 2005, **17**:338-344.
- Werling D, Jungi TW: TOLL-like receptors linking innate and adaptive immune response. *Vet Immunol Immunopathol* 2003, **91**:1-12.
- Ibeagha-Awemu EM, Lee JW, Ibeagha AE, Bannerman DD, Paape MJ, Zhao X: Bacterial lipopolysaccharide induces increased expression of toll-like receptor (TLR) 4 and downstream TLR signaling molecules in bovine mammary epithelial cells. *Vet Res* 2008, **39**:11.
- De Schepper S, De Ketelaere A, Bannerman DD, Paape MJ, Peelman L, Burvenich C: The toll-like receptor-4 (TLR-4) pathway and its possible role in the pathogenesis of Escherichia coli mastitis in dairy cattle. *Vet Res* 2008, **39**:5.
- Goldammer T, Zerbe H, Molenaar A, Schuberth HJ, Brunner RM, Kata SR, Seyfert HM: Mastitis increases mammary mRNA abundance of beta-defensin 5, toll-like-receptor 2 (TLR2), and TLR4 but not TLR9 in cattle. *Clin Diagn Lab Immunol* 2004, **11**:174-185.
- Yang W, Zerbe H, Petzl W, Brunner RM, Gunther J, Draing C, von Aulock S, Schuberth HJ, Seyfert HM: Bovine TLR2 and TLR4 properly transduce signals from Staphylococcus aureus and E. coli, but S. aureus fails to both activate NF-kappaB in mammary epithelial cells and to quickly induce TNFalpha and interleukin-8 (CXCL8) expression in the udder. *Mol Immunol* 2008, **45**:1385-1397.
- McGuire K, Jones M, Werling D, Williams JL, Glass EJ, Jann O: Radiation hybrid mapping of all 10 characterized bovine Toll-like receptors. *Anim Genet* 2006, **37**:47-50.
- Menzies M, Ingham A: Identification and expression of Toll-like receptors 1-10 in selected bovine and ovine tissues. *Vet Immunol Immunopathol* 2006, **109**:23-30.
- Farhat K, Sauter KS, Bric M, Frey J, Ulmer AJ, Jungi TW: The response of HEK293 cells transfected with bovine TLR2 to established pathogen-associated molecular patterns and to bacteria causing mastitis in cattle. *Vet Immunol Immunopathol* 2008, **125**:326-336.
- Lee JW, Bannerman DD, Paape MJ, Huang MK, Zhao X: Characterization of cytokine expression in milk somatic cells during intramammary infections with Escherichia coli or Staphylococcus aureus by real-time PCR. *Vet Res* 2006, **37**:219-229.
- Schroder NW, Schumann RR: Single nucleotide polymorphisms of Toll-like receptors and susceptibility to infectious disease. *Lancet Infect Dis* 2005, **5**:156-164.
- Mockenhaupt FP, Cramer JP, Hamann L, Stegemann MS, Eckert J, Oh NR, Otchwemah RN, Dietz E, Ehrhardt S, Schroder NW, Bienzle U, Schumann RR: Toll-like receptor (TLR) polymorphisms in African children: Common TLR-4 variants predispose to severe malaria. *Proc Natl Acad Sci USA* 2006, **103**:177-182.
- Misch EA, Hawn TR: Toll-like receptor polymorphisms and susceptibility to human disease. *Clin Sci (Lond)* 2008, **114**:347-360.
- Misch EA, Macdonald M, Ranjit C, Sapkota BR, Wells RD, Siddiqui MR, Kaplan G, Hawn TR: Human TLR1 deficiency is associated with impaired mycobacterial signaling and protection from leprosy reversal reaction. *PLoS Negl Trop Dis* 2008, **2**:e231.
- Hawn TR, Misch EA, Dunstan SJ, Thwaites GE, Lan NT, Quy HT, Chau TT, Rodrigues S, Nachman A, Janer M, Hien TT, Farrar JJ, Aderem A: A common human TLR1 polymorphism regulates the innate immune response to lipopeptides. *Eur J Immunol* 2007, **37**:2280-2289.

26. Takeuchi O, Sato S, Horiuchi T, Hoshino K, Takeda K, Dong Z, Modlin RL, Akira S: **Cutting edge: role of Toll-like receptor 1 in mediating immune response to microbial lipoproteins.** *J Immunol* 2002, **169**:10-14.
27. Omuetti KO, Beyer JM, Johnson CM, Lyle EA, Tapping RI: **Domain exchange between human toll-like receptors 1 and 6 reveals a region required for lipopeptide discrimination.** *J Biol Chem* 2005, **280**:36616-36625.
28. Babu MM, Priya ML, Selvan AT, Madera M, Gough J, Aravind L, Sankaran K: **A database of bacterial lipoproteins (DOLOP) with functional assignments to predicted lipoproteins.** *J Bacteriol* 2006, **188**:2761-2773.
29. Werling D, Jann OC, Offord V, Glass EJ, Coffey TJ: **Variation matters: TLR structure and species-specific pathogen recognition.** *Trends Immunol* 2009, **30**:124-130.
30. Guan Y, Ranoa DR, Jiang S, Mutha SK, Li X, Baudry J, Tapping RI: **Human TLRs 10 and 1 share common mechanisms of innate immune sensing but not signaling.** *J Immunol* 2010, **184**:5094-5103.
31. Opsal MA, Vage DI, Hayes B, Berget I, Lien S: **Genomic organization and transcript profiling of the bovine toll-like receptor gene cluster TLR6-TLR1-TLR10.** *Gene* 2006, **384**:45-50.
32. Ogorevc J, Kunej T, Razpet A, Dovc P: **Database of cattle candidate genes and genetic markers for milk production and mastitis.** *Anim Genet* 2009, **40**:832-851.
33. Klungland H, Sabry A, Heringstad B, Olsen HG, Gomez-Raya L, Vage DI, Olsaker I, Odegard J, Klemetsdal G, Schulman N, Vilki J, Ruane J, Aasland M, Rønningen K, Lien S: **Quantitative trait loci affecting clinical mastitis and somatic cell count in dairy cattle.** *Mamm Genome* 2001, **12**:837-842.
34. Jann OC, King A, Corrales NL, Anderson SI, Jensen K, Ait-Ali T, Tang H, Wu C, Cockett NE, Archibald AL, Glass EJ: **Comparative genomics of Toll-like receptor signalling in five species.** *BMC Genomics* 2009, **10**:216.
35. Seabury CM, Cargill EJ, Womack JE: **Sequence variability and protein domain architectures for bovine Toll-like receptors 1, 5, and 10.** *Genomics* 2007, **90**:502-515.
36. Li C, Shi W, Chu MX, An Y, Chen H, Di R, Fang L: **Polymorphisms of TLR gene and their relationship with somatic cell Score in Holstein cows.** *Sci Agric Sin* 2009, **42**:2118-2125.
37. **The National Milk Records.** [<http://www.nmr.co.uk>].
38. Murphy MA, Shariflou MR, Moran C: **High quality genomic DNA extraction from large milk samples.** *J Dairy Res* 2002, **69**:645-649.
39. Rozen S, Skaletsky H: **Primer3 on the WWW for general users and for biologist programmers.** *Methods Mol Biol* 2000, **132**:365-386.
40. Barrett JC, Fry B, Maller J, Daly MJ: **Haploview: analysis and visualization of LD and haplotype maps.** *Bioinformatics* 2005, **21**:263-265.
41. Collins RA, Howard CJ, Duggan SE, Werling D: **Bovine interleukin-12 and modulation of IFN γ production.** *Vet Immunol Immunopathol* 1999, **68**:193-207.
42. Widdison S, Watson M, Piercy J, Howard C, Coffey TJ: **Granulocyte chemotactic properties of *M. tuberculosis* versus *M. bovis*-infected bovine alveolar macrophages.** *Mol Immunol* 2008, **45**:740-749.
43. Widdison S, Watson M, Coffey TJ: **Early response of bovine alveolar macrophages to infection with live and heat-killed *Mycobacterium bovis*.** *Dev Comp Immunol* 2011, **35**:580-591.
44. Kwong LS, Hope JC, Thom ML, Sopp P, Duggan S, Bembridge GP, Howard CJ: **Development of an ELISA for bovine IL-10.** *Vet Immunol Immunopathol* 2002, **85**:213-223.
45. Farhat K, Riekenberg S, Jung G, Wiesmuller KH, Jungi TW, Ulmer AJ: **Identification of full length bovine TLR1 and functional characterization of lipopeptide recognition by bovine TLR2/1 heterodimer.** *Vet Res* 2010, **41**:34.
46. **National Center for Biotechnology Information: SNP Database.** [<http://www.ncbi.nlm.nih.gov/projects/SNP>].
47. **SMART: Simple Modular Architecture Research Tool.** [<http://smart.embl-heidelberg.de/>].
48. **Leucine-Rich Repeat Finder (LRRfinder) web-based tool.** [<http://www.lrrfinder.com/>].
49. Jin MS, Kim SE, Heo JY, Lee ME, Kim HM, Paik SG, Lee H, Lee JO: **Crystal structure of the TLR1-TLR2 heterodimer induced by binding of a triacylated lipopeptide.** *Cell* 2007, **130**:1071-1082.
50. Kossaibati MA: **The costs of clinical mastitis in UK dairy herds.** *Cattle Practice* 2000, **8**:323-328.
51. Threadgill DW, Womack JE: **Genomic analysis of the major bovine milk protein genes.** *Nucleic Acids Res* 1990, **18**:6935-6942.
52. **Motif: Searching DNA/Protein Sequence Motifs.** [<http://motif.genome.jp>].
53. Roger T, Miconnet I, Schiesser AL, Kai H, Miyake K, Calandra T: **Critical role for Ets, AP-1 and GATA-like transcription factors in regulating mouse Toll-like receptor 4 (TLR4) gene expression.** *Biochem J* 2005, **387**:355-365.
54. Liang Y, Zhou Y, Shen P: **NF- κ B and its regulation on the immune system.** *Cell Mol Immunol* 2004, **1**:343-350.
55. Francois S, El Benna J, Dang PM, Pedruzzi E, Gougerot-Pocidallo MA, Elbim C: **Inhibition of neutrophil apoptosis by TLR agonists in whole blood: involvement of the phosphoinositide 3-kinase/Akt and NF- κ B signaling pathways, leading to increased levels of Mcl-1, A1, and phosphorylated Bad.** *J Immunol* 2005, **174**:3633-3642.
56. Hayashi F, Means TK, Luster AD: **Toll-like receptors stimulate human neutrophil function.** *Blood* 2003, **102**:2660-2669.
57. Kobayashi Y: **The role of chemokines in neutrophil biology.** *Front Biosci* 2008, **13**:2400-2407.
58. Parker LC, Whyte MK, Dower SK, Sabroe I: **The expression and roles of Toll-like receptors in the biology of the human neutrophil.** *J Leukoc Biol* 2005, **77**:886-892.
59. Paape MJ, Bannerman DD, Zhao X, Lee JW: **The bovine neutrophil: Structure and function in blood and milk.** *Vet Res* 2003, **34**:597-627.

doi:10.1186/1297-9716-43-17

Cite this article as: Russell et al.: Identification of single nucleotide polymorphisms in the bovine Toll-like receptor 1 gene and association with health traits in cattle. *Veterinary Research* 2012 **43**:17.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

