



Published as: *Cell Metab.* 2012 April 4; 15(4): 439–450.

The *C. elegans* microRNA *mir-71* acts in neurons to promote germline-mediated longevity through regulation of DAF-16/FOXO

Konstantinos Boulias and H. Robert Horvitz

Department of Biology and Howard Hughes Medical Institute, Massachusetts Institute of Technology, Cambridge, MA, 02139, USA

Summary

The lifespan of *Caenorhabditis elegans* is controlled by signaling between the germline and the soma. Germ cell removal extends lifespan by triggering the activation of the DAF-16/FOXO transcription factor in the intestine. Here we analyze microRNA function in *C. elegans* aging and show that the microRNA *mir-71* functions to mediate the effects of germ cell loss on lifespan. *mir-71* is required for the lifespan extension caused by germline removal, and overexpression of *mir-71* further extends the lifespan of animals lacking germ cells. *mir-71* functions in the nervous system to facilitate the localization and transcriptional activity of DAF-16 in the intestine. Our findings reveal a novel microRNA-dependent mechanism of lifespan regulation by the germline and indicate that signaling among the gonad, the nervous system and the intestine coordinates the lifespan of the entire organism.

Introduction

Genetic studies of *C. elegans* have identified numerous genes that function in highly conserved pathways to control aging (Kenyon, 2010). For example, in an insulin-like signaling pathway the DAF-2 Insulin/IGF-1 receptor homolog activates a conserved phosphatidylinositol 3-kinase pathway to shorten lifespan by inhibiting the activity of DAF-16, a FOXO family transcription factor (Tatar et al., 2003). DAF-16 promotes longevity by regulating the expression of a number of targets, including antioxidant, antimicrobial and metabolic enzymes (Lee et al., 2003a; Murphy et al., 2003; Oh et al., 2006).

The reproductive systems of worms and possibly also those of flies and mammals regulate lifespan (Kenyon, 2010). For example, when the germline of *C. elegans* is removed either by laser microsurgery or by mutations that block germ-cell proliferation, animals live up to 60% longer than control animals (Arantes-Oliveira et al., 2002; Hsin and Kenyon, 1999). This lifespan extension requires the activities of DAF-16 and of the steroid hormone receptor DAF-12 (Hsin and Kenyon, 1999). In animals lacking germ cells, DAF-16 accumulates specifically in the intestinal nuclei and activates the transcription of stress-related and metabolic genes (Lin et al., 2001; Wang et al., 2008; Yamawaki et al., 2008). Upon germline removal, the somatic gonad promotes longevity by triggering a pathway

© 2012 Elsevier Inc. All rights reserved.

horvitz@mit.edu, phone: 617-253-4671, fax: 617-253-8126.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

involved in the biosynthesis of the endogenous ligands for DAF-12 (Gerisch et al., 2007; Gerisch et al., 2001; Yamawaki et al., 2010).

microRNAs, a class of small non-coding RNAs, have emerged as critical posttranscriptional regulators of gene expression in diverse biological processes (Ambros, 2004; Stefani and Slack, 2008). The first microRNAs discovered, the products of the *C. elegans* genes *lin-4* and *let-7*, control the timing of developmental events (Ambros and Horvitz, 1984; Chalfie et al., 1981; Reinhart et al., 2000). *C. elegans* microRNAs also control cell-fate specification, embryonic development, physiology, behavior, neural synaptic activity and longevity (Alvarez-Saavedra and Horvitz, 2010; Boehm and Slack, 2005; Chang et al., 2004; Simon et al., 2008; Yoo and Greenwald, 2005). In this study we performed a comprehensive search for microRNA genes that regulate *C. elegans* lifespan by determining the lifespans of mutants for most known microRNA genes. We show that the microRNA *mir-71* acts in the nervous system to mediate the effects of the germline on longevity.

Results

mir-71 functions to promote longevity and stress resistance and delay aging

We have previously described the isolation and initial characterization of a large collection of strains that carry deletions in most of the 115 known *C. elegans* microRNA genes (Miska et al., 2007). To identify microRNAs that function in the aging process, we tested these microRNA mutants for defects in aging and longevity. Specifically, we determined the lifespans of 81 mutant strains that carried deletions in 90 microRNA genes. We found that most microRNAs are dispensable for normal lifespan (Supplemental Table 1). Strikingly, two independently isolated deletion mutants of *mir-71* each displayed a severe decrease in lifespan of about 40% (Fig. 1A). Since longevity and resistance to stresses are frequently coupled, we subjected *mir-71* mutant adults to various stresses. We found that *mir-71* mutants showed increased sensitivity to both heat shock and oxidative stress (Fig. S1A,B,C). A transgene that contained a wild-type copy of the *mir-71* genomic locus rescued both the short lifespan (Fig. 1B) and the heat-stress sensitivity of *mir-71* mutants (Fig. S1D). Removing the 22 bp sequence of the mature *mir-71* microRNA from this transgene abolished its rescuing activity (Fig. 1C). Taken together, these results indicate that *mir-71* is required for normal lifespan and normal responses to heat and oxidative stress. In agreement with our findings, a recent study of the temporal patterns of microRNA expression during aging reported that *mir-71* is upregulated in aging adults and promotes longevity and stress resistance (de Lencastre et al., 2010).

To determine whether overexpression of *mir-71* is sufficient to extend *C. elegans* lifespan, we integrated extrachromosomal arrays carrying the *mir-71* locus into the genome and determined the lifespans of multiple resulting transgenic lines. We found that extra copies of *mir-71* modestly extended lifespan by 20% (Fig. 1D), suggesting that *mir-71* functions to promote longevity and that the short lifespan of *mir-71* mutants is not the result of non-specific pathology. Consistent with this hypothesis, aging *mir-71* mutant adults showed a premature reduction in the rates of both locomotion and pharyngeal pumping (Fig. 1E,F), two behaviors that normally decline with age (Huang et al., 2004). These results indicate that loss of *mir-71* causes an accelerated aging phenotype and suggest that *mir-71* acts to delay aging.

mir-71 mediates the effects of germ cell loss on lifespan

To test if *mir-71* functions in one of the pathways known to control *C. elegans* aging, we assayed genetic interactions between *mir-71* and various longevity genes. When temperature-sensitive *glp-1* mutants grow at the restrictive temperature they fail to develop

mature germ cells and as a result live 60% longer than wild-type animals (Fig. 2A) (Arantes-Oliveira et al., 2002). Interestingly, we found that mutations in *mir-71* strongly suppressed the long lifespan of germline-deficient *glp-1* mutants (Fig. 2A). To confirm that *mir-71* is required for the lifespan extension caused by germ cell loss, we used laser microsurgery to ablate the germline precursor cells Z2 and Z3 in wild-type and *mir-71* mutant animals. Whereas germline ablation resulted in a robust lifespan extension of wild-type animals, it failed to extend the lifespan of *mir-71* mutants (Fig. 2B). These results indicate that *mir-71* is required for the increased longevity of germline-less animals and suggest that *mir-71* mediates the effects of germ cell loss on lifespan.

To examine if *mir-71* is specifically required for germ cell loss to extend lifespan, we tested whether deletion of *mir-71* suppresses the long lifespan of animals with compromised insulin/IGF signaling or defective mitochondrial function. Partial loss-of-function mutations of the *daf-2* Insulin/IGF receptor homolog cause animals to live twice as long as the wild type (Fig. 2C) (Kenyon et al., 1993). We observed that loss of *daf-2* function similarly extended the lifespan of *mir-71* mutants more than two-fold (Fig. 2C). RNAi that reduced the levels of *cco-1* or *T02H6.11* or a loss-of-function mutation of *isp-1* (all three of these genes encode enzymes that function in mitochondrial respiration (Dillin et al., 2002; Feng et al., 2001; Lee et al., 2003b), extended the lifespan of *mir-71* mutants to a similar degree as in a wild-type background (Fig. 2D and Fig. S2A,B). In addition, *mir-71* mutants could extend lifespan in response to dietary restriction (Fig. S2C). Taken together, these results indicate that *mir-71* is specifically required for the lifespan extension caused by germline removal.

Extra copies of *mir-71* further extend the lifespan of germline-less animals

To examine the effect of *mir-71* overexpression on the lifespan of germline-deficient animals, we introduced the integrated *mir-71* transgenes into *glp-1* mutants. While extra copies of *mir-71* resulted in a modest lifespan extension in germline-intact animals, overexpression of *mir-71* extended the lifespan of germline-deficient *glp-1* animals by more than 40% (Fig. 2E). In agreement with this observation, ablation of the germline precursor cells Z2 and Z3 further extended the lifespan of *mir-71* overexpressors by about 40%, compared to germline-ablated wild-type controls. Interestingly, extra copies of *mir-71* also robustly extended the lifespan of somatic gonad-ablated animals, indicating that the somatic gonad is not required for *mir-71*-mediated lifespan extension in the absence of the germline (Fig 2F). This effect was specific to germline-deficient animals, as extra copies of *mir-71* caused little or no effect on the long lifespan of either *daf-2* mutants or animals with defective mitochondrial respiration, respectively (Fig. S3). These results indicate that *mir-71* overexpression is sufficient to extend further the lifespan of germline-less animals and suggest that the presence of the germline limits the lifespan-promoting activity of *mir-71*.

mir-71 is broadly expressed during development and adulthood

To identify the tissue(s) in which *mir-71* functions to regulate longevity, we monitored the spatial pattern of *mir-71* expression using a *gfp* transcriptional reporter. We found that a *Pmir-71::gfp* reporter was broadly expressed; stronger signal was detected in the intestine, body wall muscles and neurons during larval development and adulthood and weaker expression was observed in the hypoderm of adult animals (Fig. 3A) (Martinez et al., 2008). Germline removal did not affect the expression pattern of this *Pmir-71::gfp* reporter during development and adulthood (Fig. S4).

mir-71 functions in neurons to promote germline-mediated longevity

To test if *mir-71* functions in the intestine, muscles, neurons or hypoderm to mediate the effects of germ cell loss on lifespan, we generated genetic mosaics that lacked *mir-71* gene

function in a subset of cell lineages. Genetic mosaics were produced by the spontaneous loss of an extrachromosomal array that carries the only wild-type gene copy of *mir-71* as well as *gfp* reporters that serve as cell lineage markers (Yochem and Herman, 2003). We identified two classes of genetic mosaics: *mir-71* AB(-) mosaics that lacked *mir-71* function in the AB lineage, which generates almost the entire nervous system and most of the hypoderm, but retained *mir-71* function in the P1 lineage, which gives rise to the intestine, muscles, gonad and part of the hypoderm; and *mir-71* E(-) mosaics that lacked *mir-71* function in the E lineage (a subset of the P1 lineage), which generates the entire intestine, but retained *mir-71* function in the AB lineage (Fig. 3B and Experimental Procedures). As expected, germline removal in animals that carried the *mir-71*-expressing array in presumably all tissues caused extension of lifespan by 60% (Fig. 3C). We observed that the lifespan of *mir-71* AB(-) mosaics, was extended by only 23% by germline removal, indicating that the activity of *mir-71* in the AB lineage was largely required for the effect of germ cell loss on lifespan (Fig. 3D). On the other hand, we found that the lifespan of *mir-71* E(-) mosaics, was extended by 60% by germline removal, indicating that *mir-71* function in the AB lineage is sufficient to fully restore the effect of germ cell loss on lifespan whereas intestinal *mir-71* activity is dispensable (Fig. 3E). Therefore, our mosaic analysis shows that *mir-71* activity in the AB lineage, which generates almost all neurons and most of the hypoderm, is both necessary and sufficient for germline-mediated longevity.

Since *Pmir-71::gfp* expression was detected in neuronal and hypodermal cells, we hypothesized that *mir-71* likely acts in either the nervous system or the hypoderm to mediate the extension of lifespan that occurs in the absence of the germline. To distinguish between those alternatives, we performed tissue-specific rescue experiments. Expression of *mir-71* under the control of the ubiquitous *rpl-28* promoter strongly rescued the short lifespan of *mir-71*; *gfp-1* mutants (Fig. 4A,B). Similarly, driving expression of *mir-71* in neurons alone, using the pan-neuronal *unc-119* and *rab-3* promoters, resulted in strong rescue of the lifespan defect of *mir-71* mutants; their lifespan was extended by germline removal by more than 40% or 30%, respectively (Fig. 4C and Fig. S5). By contrast, hypodermal-specific expression of *mir-71* failed to restore the effect of germ cell loss on lifespan (Fig. 4D). Taken together, these results indicate that *mir-71* functions in the nervous system to promote germline-mediated longevity.

mir-71*-mediated lifespan extension in germline-less animals depends on *daf-16

The long lifespan of animals that lack germ cells depends on the activities of DAF-16 and a pathway that signals through the DAF-12 steroid hormone receptor (Gerisch et al., 2007; Hsin and Kenyon, 1999). It is thought that signals from the germline regulate the subcellular localization and transcriptional activity of DAF-16, while signals from the somatic gonad control the biosynthesis of DAF-12 ligands (Lin et al., 2001; Yamawaki et al., 2008; Yamawaki et al., 2010). To test whether *mir-71* genetically interacts with either *daf-16* or *daf-12* to regulate lifespan upon germ cell removal, we examined the effect of *mir-71* overexpression in loss-of-function mutants of *daf-16* or *daf-12*. Importantly, we observed that *mir-71*-mediated lifespan extension was fully suppressed by a null allele of *daf-16* but was unaffected by complete loss of *daf-12* function (Fig. 5A,B,C and Fig. S6). That *daf-12* is dispensable for *mir-71* to promote the lifespan of germline-less animals is not surprising, given that the somatic gonad is also not required (Fig. 2F). These results indicate that *mir-71* acts upstream of or in parallel to *daf-16* to promote germline-mediated longevity.

Intestinal *daf-16* function is sufficient for *mir-71* to promote germline-mediated longevity

To identify the tissue(s) in which *daf-16* activity is required for *mir-71*-mediated lifespan extension, we assayed whether tissue-specific expression of DAF-16 is sufficient for *mir-71* overexpression to extend the lifespan of germline-deficient animals. We found that driving

daf-16 expression only in neurons had little or no effect on the lifespan of *daf-16; glp-1* animals overexpressing *mir-71* (Fig. 5D). By contrast, intestinal expression of DAF-16::GFP fully rescued *mir-71*-mediated lifespan extension, indicating that the activity of DAF-16 in the intestine is sufficient for *mir-71* to promote longevity in animals lacking germ cells (Fig. 5E). In short, our results suggest that upon germline removal, *mir-71* functions in the nervous system to promote lifespan extension by facilitating the activation of *daf-16* in the intestine.

***mir-71* promotes the localization and transcriptional activity of DAF-16 in the intestine of germline-less animals**

When the germline is removed DAF-16 accumulates in the nuclei of intestinal cells and promotes the expression of several DAF target genes (Lin et al., 2001; Yamawaki et al., 2008). Our findings raise the intriguing possibility that *mir-71* regulates the expression, subcellular localization or transcriptional activity of DAF-16 in the intestine of germline-deficient animals. To distinguish among these alternatives, we first examined the expression and subcellular localization of a functional DAF-16::GFP fusion protein. Interestingly, we observed that loss of *mir-71* function partially blocked the accumulation of DAF-16::GFP in the intestine of germline-deficient adults without affecting the overall levels of DAF-16 in the presence or absence of germ cells (Fig. 6A,B,C). This effect is specific, since the nuclear translocation of DAF-16 in response to heat shock and in *daf-2* mutants was not dependent on *mir-71* (Fig. S7). These results suggest that *mir-71* specifically facilitates the translocation of DAF-16 to the intestinal nuclei of animals lacking germ cells.

To examine whether the absence of *mir-71* affects the ability of DAF-16 to activate its targets, we examined the expression of *Psod-3::gfp*, a well-characterized and widely used sensor of DAF-16 activity (Libina et al., 2003; Yamawaki et al., 2008). As previously shown, we found that *Psod-3::gfp* expression was upregulated primarily in the intestine of germline-deficient *glp-1* adults (Fig. 6D) (Yamawaki et al., 2008). Importantly, we found that loss of *mir-71* function partially blocked the induction of *Psod-3::gfp*, suggesting that *mir-71* is required for the transcriptional activation of DAF-16-dependent gene targets in the intestine (Fig. 6D). To directly test this, we measured by qRT-PCR the transcript levels of a number of genes known to be upregulated in the intestine of germline-deficient animals in a *daf-16*-dependent manner. The induction of superoxide dismutase *sod-3*, the triglyceride lipase *K04A8.5* and the glyceraldehyde 3-phosphate dehydrogenase *gpd-2* is totally dependent on DAF-16, while the upregulation of the putative steroid dehydrogenase *dod-8* and the nicotinamide nucleotide transhydrogenase *nnt-1* is partially *daf-16*-dependent (Wang et al., 2008; Yamawaki et al., 2008). We found that loss of *mir-71* function in animals lacking germ cells resulted in 50% reduction in the levels of *sod-3* and *K04A8.5*; the levels of *gpd-2*, *dod-8* and *nnt-1* were not significantly affected (Fig. 6F). By contrast, the levels of *sod-3* and *K04A8.5* were not affected by the absence of *mir-71* in germline-intact animals (Fig. 6E). Taken together, our results suggest that *mir-71* promotes germline-mediated longevity by regulating the localization and transcriptional activity of DAF-16.

Discussion

In this study we report a systematic analysis of microRNA genes in the regulation of aging of an entire organism. We have established that the microRNA gene *mir-71* is a critical factor in mediating the effect of germ cell loss on lifespan: *mir-71* is necessary for the lifespan extension caused by germline removal and promotes the longevity of animals lacking germ cells by regulating the localization and transcriptional activity of DAF-16. We propose a model in which *mir-71* functions in the nervous system to mediate the production or modification of a lifespan-extending signal that promotes the intestinal expression of key DAF-16-dependent target genes (Fig. 7). According to this model, *mir-71* could inhibit

posttranscriptionally the expression of neuronal factor(s) that antagonize cell non-autonomously the activity of DAF-16 in the intestine (Fig. 7).

Several approaches using bioinformatics have been developed to help identify direct targets of microRNAs. Although these algorithms have identified a large number of predicted microRNA targets, experimental validation of most of these putative targets is lacking (Hammell et al., 2008; Lall et al., 2006; Lewis et al., 2005). Recently, de Lencastre et al. (2010) implicated *cdc-25.1* as a potential target of *mir-71* function. We have tested if *cdc-25.1* is a biologically significant target of *mir-71* in controlling germline-mediated longevity in genetic epistasis experiments; our preliminary results do not support this hypothesis (K. Boulias and H.R. Horvitz, unpublished observations). Given our finding that *mir-71* activity in neurons is important for germline-mediated longevity, it will be interesting to test experimentally whether predicted *mir-71* targets known to be highly enriched in the nervous system function to mediate the effects of germline on lifespan.

Our results indicate that the germline strongly suppresses the ability of *mir-71* to promote longevity, since extra copies of *mir-71* have only a modest effect on the lifespan of intact animals. When the germline is removed, *mir-71*-mediated lifespan extension requires the intestinal activity of *daf-16*, whereas lifespan extension does not depend on the presence of the somatic gonad or the *daf-12* pathway. Since *mir-71* expression is not regulated by the germline, we postulate that *mir-71*-mediated lifespan extension requires factors that are triggered by germline removal and act on DAF-16 function. Previous studies have shown that germline removal triggers the up-regulation of TCER-1, a transcription elongation factor that acts to promote the transcriptional activity of DAF-16 in the intestine (Ghazi et al., 2009). We found that *mir-71*-mediated lifespan extension is partially dependent on *tcer-1* gene function (Fig. 5F). Thus, *mir-71* might act with TCER-1 and possibly other factors to promote intestinal DAF-16 activity in germline-deficient animals.

Current evidence suggests that the rate of aging at least in *C. elegans* and *Drosophila* is coordinated through communication and signaling among different tissues. For example, tissue-specific manipulations of insulin/IGF-1 signaling in the intestine or the nervous system have been shown to regulate the lifespan of the whole organism, while genetic ablation of a set of neurons can affect the ability of worms to extend lifespan in response to dietary restriction (Bishop and Guarente, 2007; Broughton et al., 2005; Hwangbo et al., 2004; Libina et al., 2003; Wolkow et al., 2000). In addition, a recent study showed that perturbation of mitochondrial function can modulate *C. elegans* aging in a cell non-autonomous fashion (Durieux et al., 2011). Furthermore, the germline of *C. elegans* is thought to send signals that inhibit the lifespan of the entire animal (Kenyon, 2010), while the somatic gonad is thought to be involved in the production of a steroid hormone that promotes the lifespan of animals lacking germ cells (Yamawaki et al., 2010). Tissue-specific rescue experiments suggested that the somatic gonad might act through the hypoderm, the endocrine XXX cells or sensory neurons to promote longevity (Yamawaki et al., 2010). Our finding that *mir-71* activity in neurons is sufficient to promote longevity underscores the importance of the nervous system and neuroendocrine signaling for the control of germline-mediated longevity. Based on our results, we suggest that *mir-71* functions cell non-autonomously in neurons to promote DAF-16 activity in the intestine. In short, our results implicate the nervous system in lifespan control upon germ cell removal and support a model in which signaling among the germline, the somatic gonad, the intestine and the nervous system coordinates the rate of aging of the whole organism.

Experimental Procedures

Strains

Strains were cultured as described (Brenner, 1974) and maintained at 20°C unless specified otherwise. Strains that contained the *glp-1(e2141ts)* allele were maintained at 15 °C. *mir-71(n4115)* was outcrossed 8x, *mir-71(n4105)* 6x and *nIs286*, *nIs287* and *nIs289* 4x to the wild type. A list of the strains used in this study is provided in Supplementary Material.

Rescue Experiments and Transgenic Animals

For rescue experiments and mosaic analysis, we amplified a 3 kb fragment surrounding the *mir-71* locus (2 kb upstream of and 1 kb downstream of the *mir-71* locus) from wild-type genomic DNA using PCR and cloned this fragment into the PCRII-TOPO (Invitrogen) vector. We used site-directed ligase-independent mutagenesis to generate a control plasmid in which the mature microRNA sequence was deleted (Chiu et al., 2004). To generate the *Pmir-71::gfp* reporter, we amplified by PCR the 2 kb upstream region of the *mir-71* locus present in the rescuing construct and cloned this fragment into the pPD96.62 (Adgene) vector.

For tissue-specific rescue experiments we substituted the *gfp* coding sequence of pPD95.75 with the *mir-71* precursor sequence (pPD95.75-*mir-71pr*). Subsequently, we cloned either the *rpl-28* promoter fragment (ubiquitous expression) from pPD129.57 or the *unc-119* promoter fragment (pan-neuronal expression) from *Punc-119::gfp* (Nakano et al., 2010) or the *dpy-7* promoter fragment (hypodermal expression) from *Pdpy-7::2Xnls::yfp* (Myers and Greenwald, 2005) into pPD95.75-*mir-71pr*.

Germline transformation experiments were performed as described (Mello et al., 1991). For rescue experiments, injection mixes contained plasmids at 5 ng/μl (for *mir-71* and for a *mir-71* control plasmid with the *mir-71* mature sequence deleted), 20 ng/μl of pTG96 (*Psur-5::gfp*) as a co-transformation marker and 80 ng/μl of 1 kb DNA ladder (Invitrogen) as carrier DNA. For mosaic analyses, *mir-71(n4115)* hermaphrodites were injected with a mix that contained 5 ng/μl of *mir-71* rescuing construct, 50 ng/μl of *Posm-6::gfp* construct (Collet et al., 1998), 50 ng/μl of *Pges-1::gfp* construct (Bishop and Guarente, 2007) and 80 ng/μl of 1 kb DNA ladder. For tissue-specific rescue experiments, *mir-71(n4115)* hermaphrodites were injected with a mix that contained 20 ng/μl of *Prpl-28::mir-71* construct or 100 ng/ul *Punc-119::mir-71* construct or 100 ng/ul *Prab-3::mir-71* construct or 20 ng/ul *Pdpy-7::mir-71* construct along with 100 ng/ul pRF4 and 80 ng/μl of 1 kb DNA ladder. We generated the *Pmir-71::gfp* transgenic strain by injecting *lin-15AB(n765)* hermaphrodites with 30 ng/μl of *Pmir-71::gfp* construct, 33 ng/μl *plin-15(EK)* and 80 ng/μl of 1 kb DNA ladder.

nIs286, *nIs287* and *nI289*, integrants of the *Ex[mir-71(+)* + pTG96] transgene, and *nIs298*, an integrant of the *Ex[Pmir-71::gfp]* transgene, were isolated after a standard γ -ray integration screen and were backcrossed twice to the wild type before analysis (Mello et al., 1991).

Lifespan Analyses

Unless stated otherwise, lifespan assays were performed by standard methods at 20 °C using NGM plates seeded with OP50 bacteria containing 25 μM FUDR. In lifespan experiments assaying strains that carried the *glp-1(e2141ts)* mutation, animals (*glp-1* and control germline-intact controls) were raised at 25 °C during embryogenesis and were either kept at 25 °C on plates without FUDR throughout the analysis or shifted to 20 °C after the L4 stage and for the rest of the lifespan analysis. Statistical analysis was performed with GraphPad

Prism 4 software, which uses the log-rank (Mantel-Cox) method to calculate p values. RNAi lifespan assays were performed according to the standard feeding protocol (Kamath et al., 2003).

Behavioral Assays

Single animals were maintained on individual NGM plates throughout adulthood and were transferred to fresh plates seeded with OP50 bacteria before locomotion or pumping rates were counted. Locomotion rates were determined by counting body bends per 20 sec of animals moving on a fresh bacterial lawn using a dissecting microscope. Pumping rates were assayed by counting the number of movements per min of the rear bulb of the pharynx of animals within the bacterial lawn using a dissecting microscope.

Laser Ablation Experiments

Laser ablations of germline precursor cells (Z2 and Z3) and somatic gonad precursor cells (Z1 and Z4) of newly hatched L1 larvae were performed as described previously (Avery and Horvitz, 1987). At adulthood the absence of a germline was determined using a dissecting microscope. Intact controls were anesthetized and recovered from the same sodium azide agarose pads as experimental animals.

Mosaic analysis

We used *mir-71(n4115); nEx1717* and *mir-71(n4115); glp-1(e2141); nEx1717* animals to generate *mir-71* genetic mosaics. *nEx1717* is an extrachromosomal array containing a genomic copy of *mir-71* as well as lineage specific markers (see Rescue Experiments and Transgenic Animals). Approximately 200,000 progeny of each of *mir-71(n4115); nEx1717* and *mir-71(n4115); glp-1(e2141); nEx1717* animals were raised at 25 °C until the L4 stage and then were screened using a fluorescence dissecting microscope (Olympus) for mosaic animals in which either the AB-specific marker *Posm-6::gfp* (expressed in ciliated neurons) or the E-specific marker *Pges-1::gfp* (expressed in the intestine generated by the P1 lineage) was absent. AB(-) mosaics (*Posm-6::gfp* negative, *Pges-1::gfp* positive) lost the *mir-71* locus in the AB lineage and presumably retained *mir-71* locus in the P1 lineage. E(-) mosaics (*Posm-6::gfp* positive, *Pges-1::gfp* negative) lost the *mir-71* locus in the E lineage, presumably retained *mir-71* locus in the AB lineage and might or might not have carried the *mir-71* locus in the MS and P2 lineages. Thus E(-) mosaics probably included a mix of E(-), EMS(-) and P1(-) mosaic animals. Mosaics were selected as L4 larvae and were transferred to 20 °C for lifespan analysis. *mir-71(n4115); nEx1717* and *mir-71(n4115); glp-1(e2141); nEx1717* controls (array present in all cells), and *mir-71(n4115)* and *mir-71(n4115); glp-1(e2141)* controls (array lost in all cells) underwent the same procedure. They were selected using the fluorescence dissecting microscope in parallel with the mosaic animals and were exposed to UV radiation for approximately the same time.

GFP Fluorescence Microscopy and Quantification

Animals were anaesthetized on agarose pads containing 20–50 mM NaN₃. Images were taken with a CCD digital camera using a 5X objective on a Zeiss Axioskop microscope. For each trial, exposure time was calibrated to minimize the number of saturated pixels and was kept constant though the experiment. The ImageJ software was used to quantify mean fluorescence intensity per worm as measured by intensity of each pixel in the selected area. No expression of the transgene was visible in embryos prior to egg laying.

Quantitative RT-PCR Analysis

Germline-deficient *glp-1(e2141ts)* and wild-type N2 animals were raised at 25 °C until the L4 stage and then shifted to 20 °C. On day 2 of adulthood, animals were collected for RNA

extraction. RNA extraction, purification, and reverse transcription and qPCR were carried as described (Andersen et al., 2008). Data were generated from four biological replicates. mRNA levels of *snb-1* and *rpl-26* were used for normalization (Curran et al., 2009). Primer sequences are available upon request.

Statistical Analysis

Error bars represent the standard error of the mean (S.E.M). p values were calculated using the unpaired Student's *t* test.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank S. Nakano, D.P. Denning, A. Saffer, E. Alvarez-Saavedra and A. Chalkiadaki for critically reading the manuscript; N.A. Bishop and L. Guarente for plasmids and for the dietary restriction analyses; A. Antebi, R.K. Herman and the *Caenorhabditis* Genetics Center for strains and plasmids and members of the Horvitz laboratory for discussions. This work was supported by a European Molecular Biology Organization fellowship, a grant from the Ellison Medical Foundation and by the Howard Hughes Medical Institute. H.R.H is an Investigator of the Howard Hughes Medical Institute.

References

- Alvarez-Saavedra E, Horvitz HR. Many families of *C. elegans* microRNAs are not essential for development or viability. *Curr Biol.* 2010; 20:367–373. [PubMed: 20096582]
- Ambros V. The functions of animal microRNAs. *Nature.* 2004; 431:350–355. [PubMed: 15372042]
- Ambros V, Horvitz HR. Heterochronic mutants of the nematode *Caenorhabditis elegans*. *Science.* 1984; 226:409–416. [PubMed: 6494891]
- Andersen EC, Saffer AM, Horvitz HR. Multiple levels of redundant processes inhibit *Caenorhabditis elegans* vulval cell fates. *Genetics.* 2008; 179:2001–2012. [PubMed: 18689876]
- Arantes-Oliveira N, Apfeld J, Dillin A, Kenyon C. Regulation of life-span by germ-line stem cells in *Caenorhabditis elegans*. *Science.* 2002; 295:502–505. [PubMed: 11799246]
- Avery L, Horvitz HR. A cell that dies during wild-type *C. elegans* development can function as a neuron in a *ced-3* mutant. *Cell.* 1987; 51:1071–1078. [PubMed: 3690660]
- Bishop NA, Guarente L. Two neurons mediate diet-restriction-induced longevity in *C. elegans*. *Nature.* 2007; 447:545–549. [PubMed: 17538612]
- Boehm M, Slack F. A developmental timing microRNA and its target regulate life span in *C. elegans*. *Science.* 2005; 310:1954–1957. [PubMed: 16373574]
- Brenner S. The genetics of *Caenorhabditis elegans*. *Genetics.* 1974; 77:71–94. [PubMed: 4366476]
- Broughton SJ, Piper MD, Ikeya T, Bass TM, Jacobson J, Driege Y, Martinez P, Hafen E, Withers DJ, Leivers SJ, et al. Longer lifespan, altered metabolism, and stress resistance in *Drosophila* from ablation of cells making insulin-like ligands. *Proc Natl Acad Sci U S A.* 2005; 102:3105–3110. [PubMed: 15708981]
- Chalfie M, Horvitz HR, Sulston JE. Mutations that lead to reiterations in the cell lineages of *C. elegans*. *Cell.* 1981; 24:59–69. [PubMed: 7237544]
- Chang S, Johnston RJ Jr, Frokjaer-Jensen C, Lockery S, Hobert O. MicroRNAs act sequentially and asymmetrically to control chemosensory laterality in the nematode. *Nature.* 2004; 430:785–789. [PubMed: 15306811]
- Chiu J, March PE, Lee R, Tillett D. Site-directed, Ligase-Independent Mutagenesis (SLIM): a single-tube methodology approaching 100% efficiency in 4 h. *Nucleic Acids Res.* 2004; 32:e174. [PubMed: 15585660]

- Collet J, Spike CA, Lundquist EA, Shaw JE, Herman RK. Analysis of *osm-6*, a gene that affects sensory cilium structure and sensory neuron function in *Caenorhabditis elegans*. *Genetics*. 1998; 148:187–200. [PubMed: 9475731]
- Curran SP, Wu X, Riedel CG, Ruvkun G. A soma-to-germline transformation in long-lived *Caenorhabditis elegans* mutants. *Nature*. 2009; 459:1079–1084. [PubMed: 19506556]
- de Lencastre A, Pincus Z, Zhou K, Kato M, Lee SS, Slack FJ. MicroRNAs Both Promote and Antagonize Longevity in *C. elegans*. *Curr Biol*. 2010; 20:2159–2168. [PubMed: 21129974]
- Dillin A, Hsu AL, Arantes-Oliveira N, Lehrer-Graiwer J, Hsin H, Fraser AG, Kamath RS, Ahringer J, Kenyon C. Rates of behavior and aging specified by mitochondrial function during development. *Science*. 2002; 298:2398–2401. [PubMed: 12471266]
- Durieux J, Wolff S, Dillin A. The cell-non-autonomous nature of electron transport chain-mediated longevity. *Cell*. 2011; 144:79–91. [PubMed: 21215371]
- Feng J, Bussiere F, Hekimi S. Mitochondrial electron transport is a key determinant of life span in *Caenorhabditis elegans*. *Dev Cell*. 2001; 1:633–644. [PubMed: 11709184]
- Gerisch B, Rottiers V, Li D, Motola DL, Cummins CL, Lehrach H, Mangelsdorf DJ, Antebi A. A bile acid-like steroid modulates *Caenorhabditis elegans* lifespan through nuclear receptor signaling. *Proc Natl Acad Sci U S A*. 2007; 104:5014–5019. [PubMed: 17360327]
- Gerisch B, Weitzel C, Kober-Eisermann C, Rottiers V, Antebi A. A hormonal signaling pathway influencing *C. elegans* metabolism, reproductive development, and life span. *Dev Cell*. 2001; 1:841–851. [PubMed: 11740945]
- Ghazi A, Henis-Korenblit S, Kenyon C. A transcription elongation factor that links signals from the reproductive system to lifespan extension in *Caenorhabditis elegans*. *PLoS Genet*. 2009; 5:e1000639. [PubMed: 19749979]
- Hammell M, Long D, Zhang L, Lee A, Carmack CS, Han M, Ding Y, Ambros V. mirWIP: microRNA target prediction based on microRNA-containing ribonucleoprotein-enriched transcripts. *Nat Methods*. 2008; 5:813–819. [PubMed: 19160516]
- Hsin H, Kenyon C. Signals from the reproductive system regulate the lifespan of *C. elegans*. *Nature*. 1999; 399:362–366. [PubMed: 10360574]
- Huang C, Xiong C, Kornfeld K. Measurements of age-related changes of physiological processes that predict lifespan of *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A*. 2004; 101:8084–8089. [PubMed: 15141086]
- Hwangbo DS, Gershman B, Tu MP, Palmer M, Tatar M. *Drosophila* dFOXO controls lifespan and regulates insulin signalling in brain and fat body. *Nature*. 2004; 429:562–566. [PubMed: 15175753]
- Kamath RS, Fraser AG, Dong Y, Poulin G, Durbin R, Gotta M, Kanapin A, Le Bot N, Moreno S, Sohrmann M, et al. Systematic functional analysis of the *Caenorhabditis elegans* genome using RNAi. *Nature*. 2003; 421:231–237. [PubMed: 12529635]
- Kenyon C, Chang J, Gensch E, Rudner A, Tabtiang R. A *C. elegans* mutant that lives twice as long as wild type. *Nature*. 1993; 366:461–464. [PubMed: 8247153]
- Kenyon CJ. The genetics of ageing. *Nature*. 2010; 464:504–512. [PubMed: 20336132]
- Lall S, Grun D, Krek A, Chen K, Wang YL, Dewey CN, Sood P, Colombo T, Bray N, Macmenamin P, et al. A Genome-Wide Map of Conserved MicroRNA Targets in *C. elegans*. *Curr Biol*. 2006; 16:461–470.
- Lee SS, Kennedy S, Tolonen AC, Ruvkun G. DAF-16 target genes that control *C. elegans* life-span and metabolism. *Science*. 2003a; 300:644–647. [PubMed: 12690206]
- Lee SS, Lee RY, Fraser AG, Kamath RS, Ahringer J, Ruvkun G. A systematic RNAi screen identifies a critical role for mitochondria in *C. elegans* longevity. *Nat Genet*. 2003b; 33:40–48. [PubMed: 12447374]
- Lewis BP, Burge CB, Bartel DP. Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell*. 2005; 120:15–20. [PubMed: 15652477]
- Libina N, Berman JR, Kenyon C. Tissue-specific activities of *C. elegans* DAF-16 in the regulation of lifespan. *Cell*. 2003; 115:489–502. [PubMed: 14622602]

- Lin K, Hsin H, Libina N, Kenyon C. Regulation of the *Caenorhabditis elegans* longevity protein DAF-16 by insulin/IGF-1 and germline signaling. *Nat Genet.* 2001; 28:139–145. [PubMed: 11381260]
- Martinez NJ, Ow MC, Reece-Hoyes JS, Barrasa MI, Ambros VR, Walhout AJ. Genome-scale spatiotemporal analysis of *Caenorhabditis elegans* microRNA promoter activity. *Genome Res.* 2008; 18:2005–2015. [PubMed: 18981266]
- Mello CC, Kramer JM, Stinchcomb D, Ambros V. Efficient gene transfer in *C. elegans*: extrachromosomal maintenance and integration of transforming sequences. *Embo J.* 1991; 10:3959–3970. [PubMed: 1935914]
- Miska EA, Alvarez-Saavedra E, Abbott AL, Lau NC, Hellman AB, McGonagle SM, Bartel DP, Ambros VR, Horvitz HR. Most *Caenorhabditis elegans* microRNAs are individually not essential for development or viability. *PLoS Genet.* 2007; 3:e215. [PubMed: 18085825]
- Murphy CT, McCarroll SA, Bargmann CI, Fraser A, Kamath RS, Ahringer J, Li H, Kenyon C. Genes that act downstream of DAF-16 to influence the lifespan of *Caenorhabditis elegans*. *Nature.* 2003; 424:277–283. [PubMed: 12845331]
- Myers TR, Greenwald I. *lin-35* Rb acts in the major hypodermis to oppose ras-mediated vulval induction in *C. elegans*. *Dev Cell.* 2005; 8:117–123. [PubMed: 15621535]
- Nakano S, Ellis RE, Horvitz HR. Otx-dependent expression of proneural bHLH genes establishes a neuronal bilateral asymmetry in *C. elegans*. *Development.* 2010; 137:4017–4027. [PubMed: 21041366]
- Oh SW, Mukhopadhyay A, Dixit BL, Raha T, Green MR, Tissenbaum HA. Identification of direct DAF-16 targets controlling longevity, metabolism and diapause by chromatin immunoprecipitation. *Nat Genet.* 2006; 38:251–257. [PubMed: 16380712]
- Reinhart BJ, Slack FJ, Basson M, Pasquinelli AE, Bettinger JC, Rougvie AE, Horvitz HR, Ruvkun G. The 21-nucleotide *let-7* RNA regulates developmental timing in *Caenorhabditis elegans*. *Nature.* 2000; 403:901–906. [PubMed: 10706289]
- Simon DJ, Madison JM, Conery AL, Thompson-Peer KL, Soskis M, Ruvkun GB, Kaplan JM, Kim JK. The microRNA *miR-1* regulates a MEF-2-dependent retrograde signal at neuromuscular junctions. *Cell.* 2008; 133:903–915. [PubMed: 18510933]
- Stefani G, Slack FJ. Small non-coding RNAs in animal development. *Nat Rev Mol Cell Biol.* 2008; 9:219–230. [PubMed: 18270516]
- Tatar M, Bartke A, Antebi A. The endocrine regulation of aging by insulin-like signals. *Science.* 2003; 299:1346–1351. [PubMed: 12610294]
- Wang MC, O'Rourke EJ, Ruvkun G. Fat metabolism links germline stem cells and longevity in *C. elegans*. *Science.* 2008; 322:957–960. [PubMed: 18988854]
- Wolkow CA, Kimura KD, Lee MS, Ruvkun G. Regulation of *C. elegans* lifespan by insulinlike signaling in the nervous system. *Science.* 2000; 290:147–150. [PubMed: 11021802]
- Yamawaki TM, Arantes-Oliveira N, Berman JR, Zhang P, Kenyon C. Distinct activities of the germline and somatic reproductive tissues in the regulation of *Caenorhabditis elegans*' longevity. *Genetics.* 2008; 178:513–526. [PubMed: 18202391]
- Yamawaki TM, Berman JR, Suchanek-Kavipurapu M, McCormick M, Maria Gaglia M, Lee SJ, Kenyon C. The somatic reproductive tissues of *C. elegans* promote longevity through steroid hormone signaling. *PLoS Biol.* 2010; 8:e1000468. [PubMed: 20824162]
- Yochem J, Herman RK. Investigating *C. elegans* development through mosaic analysis. *Development.* 2003; 130:4761–4768. [PubMed: 12952898]
- Yoo AS, Greenwald I. LIN-12/Notch activation leads to microRNA-mediated down-regulation of Vav in *C. elegans*. *Science.* 2005; 310:1330–1333. [PubMed: 16239437]

Highlights

- *C. elegans* lifespan extension caused by germ cell loss depends on the microRNA *mir-71*
- *mir-71* functions in neurons to promote germline-mediated longevity
- *mir-71*-mediated lifespan extension depends on intestinal *daf-16* function
- *mir-71* facilitates the localization and activity of DAF-16 in the intestine




Fig. 1. The microRNA *mir-71* functions to promote longevity and delay aging (A) *mir-71* mutants have a short lifespan ($p < 0.0001$; 40% reduction in mean lifespan). (B) A genomic fragment containing the *mir-71* locus rescued the lifespan defect of *mir-71(n4115)* mutants. ($p < 0.0001$; 45–70% mean lifespan extension compared to *mir-71(n4115)*). (C) A genomic fragment lacking the mature *mir-71* sequence failed to rescue the lifespan defect of *mir-71(n4115)* mutants. ($p > 0.1$ compared to *mir-71(n4115)*). (D) Extra copies of *mir-71* extended lifespan. ($p < 0.0001$; 15–25% mean lifespan extension). (E) Young day-1 *mir-71* adults showed normal levels of locomotion ($p > 0.1$; mean 19.0 body bends/20 sec) compared to wild-type day-1 adults (mean velocity=19.9 body bends/20 sec). At day 7, *mir-71* adults showed a 50% decrease in locomotion ($p < 0.01$; mean 5.6 body bends/20 sec) compared to wild-type adults of the same age (mean 12.1 body bends/20 sec) (wild-type, $n = 11$;

mir-71(n4115), $n=12$). **(F)** Young *mir-71(n4115)* adults showed normal levels of pharyngeal pumping compared to the wild-type. The pumping rate of day-2 *mir-71* adults was only slightly reduced ($p<0.01$; mean pumping rate 237.1 pumps/min) compared to the wild-type (mean pumping rate 250.0 pumps/min), and day-4 *mir-71* adults had a pumping rate ($p>0.1$; mean pumping rate 213.3 pumps/min) indistinguishable from that of the wild type (mean pumping rate 217.3 pumps/min). At day 8 *mir-71* adults showed a large decrease in pumping rate ($p<0.0001$; mean pumping rate 33.2 pumps/min) compared to wild-type adults of the same age (mean pumping rate 170.5 pumps/min) (wild-type, $n=11$; *mir-71(n4115)*, $n=12$). All experiments were repeated at least once with similar effects. Mean lifespan values and statistical analyses of lifespan assays are shown in Supplementary Table 2. p values refer to the experimental strain and corresponding wild-type control animals unless otherwise noted.




Fig. 2. *mir-71* mediates the effects of germ cell loss on longevity

(A) Loss of *mir-71* function suppresses the long lifespan of germline-deficient *glp-1(e2141)* mutants. Germline removal by *glp-1(e2141)* resulted in a 55% extension of mean lifespan in otherwise wild-type animals ($p < 0.0001$) compared to a 15% extension in *mir-71(n4115)* mutants ($p < 0.0001$). (B) Loss of *mir-71* function fully suppresses the increased longevity of germline-ablated animals. Ablation of germline precursor cells Z2 and Z3 resulted in a 60% extension of wild-type mean lifespan ($p < 0.0001$), while it had no effect ($p > 0.5$) on *mir-71(n4115)* mutant lifespan. (C) Loss of *daf-2* function extended both wild-type lifespan ($p < 0.0001$; 100% mean lifespan extension) and the lifespan of *mir-71(n4115)* mutants ($p < 0.0001$; 180% mean lifespan extension) (D) *cco-1* RNAi extended both wild-type lifespan ($p < 0.0001$; 35% mean lifespan extension) and the lifespan of *mir-71(n4115)*

mutants ($p < 0.0001$; 45% mean lifespan extension). **(E)** Extra copies of *mir-71* modestly extended the lifespan of intact animals at 25 °C ($p < 0.0003$, 14–15 % mean lifespan extension), whereas it caused a robust extension on the lifespan of germline-deficient *glp-1(e2141)* animals at 25 °C ($p < 0.0001$, 40–45% mean lifespan extension). **(F)** Extra copies of *mir-71* caused a robust lifespan extension on the lifespan of both germline-ablated (Z2 and Z3) ($p < 0.0001$, 40% mean lifespan extension compared to wild type Z2Z3(-) animals) and somatic gonad-ablated (Z1 and Z4) animals ($p < 0.0001$, 75% mean lifespan extension compared to wild type Z1Z4(-) animals). All experiments were repeated at least once with similar effects. Mean lifespan values and statistical analyses of lifespan assays are shown in Supplementary Table 2.




Fig. 3. *mir-71* functions in the AB lineage to regulate the lifespan of germline-deficient animals. (A) A *Pmir-71::gfp* reporter was strongly expressed in the intestine, body wall muscles and neurons during adulthood. (B) A cell-lineage diagram indicating tissues produced by the early blastomeres of *C. elegans*. (C–E) *mir-71* functions primarily in AB-derived tissue(s) to mediate the effects of germline on lifespan. Germline removal had a minor effect on the lifespan of *mir-71* mutants ($p < 0.0001$; 7% mean lifespan extension), whereas it robustly extended the lifespan of animals that carried the *mir-71*-expressing array presumably in all tissues ($p < 0.0001$; 60% mean lifespan extension). Whereas *mir-71* E(-) mosaics fully responded to germline removal ($p < 0.0001$; 60% mean lifespan extension), the lifespan of *mir-71* AB(-) mosaics was only modestly extended by germ cell loss ($p < 0.0001$; 23% mean lifespan extension). Mean lifespan values and statistical analyses of lifespan assays are shown in Supplementary Table 2.




Fig. 4.

Expression of *mir-71* in neurons alone was sufficient to promote germline-mediated longevity. (A–D) Driving *mir-71* expression either ubiquitously (using the *rpl-28* promoter) or in the nervous system of *mir-71* mutants (using the pan-neuronal *unc-119* promoter) resulted in strong rescue ($p < 0.0001$; 40–45% mean lifespan extension). On the other hand, hypodermal-specific expression of *mir-71* (using the *dpy-7* promoter) failed to rescue the lifespan defect of *mir-71; glp-1* mutants ($p < 0.0001$; 5–10% mean lifespan extension). All experiments were repeated at least once with similar effects. Mean lifespan values and statistical analyses of lifespan assays are shown in Supplementary Table 2.




Fig. 5. *mir-71* functions upstream of or parallel to DAF-16 to promote the longevity of germline-deficient animals. (A) Extra copies of *mir-71* caused a robust extension of the lifespan of germline-deficient *glp-1(e2141)* animals. ($p < 0.0001$, 40–45% mean lifespan extension). (B) Loss of *daf-16* function suppressed *mir-71*-mediated lifespan extension in germline-deficient *glp-1(e2141)* animals. Extra copies of *mir-71* only slightly extended the lifespan of germline-deficient *daf-16; glp-1* animals ($p < 0.0001$, 13–20% mean lifespan extension). (C) Loss of *daf-12* function did not suppress *mir-71*-mediated lifespan extension in germline-deficient *glp-1(e2141)* animals. Extra copies of *mir-71* caused a robust extension on the lifespan of germline-deficient *glp-1; daf-12* animals ($p < 0.0001$, 45–60% mean lifespan extension). (D) Expression of DAF-16 in neurons was not sufficient for *mir-71* overexpression to substantially extend the lifespan of germline-deficient *glp-1(e2141)* animals. Extra copies of *mir-71* caused only a slight extension on the lifespan of germline-deficient *glp-1* animals that express *daf-16* only in neurons ($p < 0.0001$, 15% mean lifespan extension). (E) Expression of DAF-16 in the intestine was sufficient for *mir-71* overexpression to substantially extend the lifespan of germline-deficient *glp-1(e2141)* animals. Extra copies of *mir-71* caused a robust extension on the lifespan of germline-deficient *glp-1* animals that express *daf-16* only in the intestine ($p < 0.0001$, 40% mean lifespan extension). (F) Deletion of *tcer-1* partially suppressed *mir-71*-mediated lifespan extension in germline-deficient *glp-1(e2141)* animals. Extra copies of *mir-71* only modestly extended the lifespan of germline-deficient *tcer-1(tm1452); glp-1* animals ($p < 0.0001$, 15–25% mean lifespan extension). All experiments were repeated at least once with similar effects. Mean lifespan values and statistical analyses of lifespan assays are shown in Supplementary Table 2.




Fig. 6. *mir-71* facilitates the localization and transcriptional activity of DAF-16 in the intestine of animals lacking germ cells. (A–C) Loss of *mir-71* function partially affected the nuclear accumulation of DAF-16::GFP in the intestine of germline-deficient *glp-1(e2141)* animals. (A) Representative images for each genotype. (B) Loss of *mir-71* function does not cause major changes in overall levels of DAF-16::GFP expression in day 2 adults ($n = 20$). Error bar, standard error of the mean (SEM). (C) Intestinal DAF-16::GFP nuclear accumulation was assessed by measuring the number of nuclei from the images of day 2 adults (low accumulation, <10 nuclei; intermediate accumulation, 10–20 nuclei; high accumulation, >20 nuclei) ($n = 78$ for each genotype). Error bar, standard error of two biological replicates. All strains contain the *daf-16(mu86)* allele and the functional *muIs109[Pdaf-16::DAF-16::GFP]* reporter. (D) Loss of *mir-71* function blocked the induction of *Psod-3::gfp* in day-3 germline-deficient *glp-1(e2141)* animals ($p < 0.0001$; 40% reduction in mean GFP fluorescence intensity) (wild-type, $n = 32$; *mir-71(n4115)*, $n = 35$; *glp-1(e2141)*, $n = 40$; *mir-71(n4115); glp-1(e2141)*, $n = 40$). (E) Loss of *mir-71* function did not affect the expression of DAF-16 targets in intact animals. mRNA levels of DAF-16 target genes *sod-3*, *K04A8.5*, *gpd-2*, *dod-8* and *nnt-1* were measured by qRT-PCR in wild-type and *mir-71(n4115)* day-2 germline-intact adults. Loss of *mir-71* function did not affect the levels of *sod-3*, *K04A8.5*, *gpd-2* and *dod-8*, while it resulted in a 50% increase in the levels of *nnt-1* ($p < 0.05$). mRNA levels are relative to wild-type levels. Error bars, standard error of four biological replicates. (F) Loss of *mir-71* function reduced the expression of a subset of DAF-16 targets in germline-less animals. mRNA levels of DAF-16 target genes *sod-3*, *K04A8.5*, *gpd-2*, *dod-8* and *nnt-1* were measured by qRT-PCR in germline-deficient *glp-1* and *mir-71; glp-1* day-2 adults. Loss of *mir-71* function in germline-defective *glp-1* animals resulted in 50% decrease in the levels of *sod-3* ($p < 0.05$) and *K04A8.5* ($p < 0.01$) and a small

reduction in the levels of *gpd-2* ($p < 0.05$). mRNA levels are shown relative to *glp-1(e2141)* levels. Error bars, standard error of four biological replicates.




Figure 7.

A model for the regulation of germline-mediated longevity by the *mir-71* microRNA. We propose that *mir-71* functions in neurons to inhibit posttranscriptionally the expression of factor(s) involved in the production or modification of a signal that controls the localization and activity of DAF-16 in the intestine.