

Substrate Selectivity of YgfU, a Uric Acid Transporter from *Escherichia coli*⁵

Received for publication, February 23, 2012, and in revised form, March 15, 2012. Published, JBC Papers in Press, March 21, 2012, DOI 10.1074/jbc.M112.355818

Konstantinos Papakostas and Stathis Frillingos¹

From the Laboratory of Biological Chemistry, University of Ioannina Medical School, 45110 Ioannina, Greece

Background: YgfU is homologous to nucleobase transporters of the ubiquitous family NCS2.

Results: YgfU transports uric acid. A single-amino acid replacement (T100A) converts YgfU to a dual-selectivity transporter for uric acid and xanthine.

Conclusion: Thr-100 is essential for the uric acid selectivity.

Significance: Dissecting selectivity determinants in family NCS2 is crucial for understanding evolution of purine uptake.

The ubiquitous nucleobase-ascorbate transporter (NAT/NCS2) family includes more than 2,000 members, but only 15 have been characterized experimentally. *Escherichia coli* has 10 members, of which the uracil permease UraA and the xanthine permeases XanQ and XanP are functionally known. Of the remaining members, YgfU is closely related in sequence and genomic locus with XanQ. We analyzed YgfU and showed that it is a proton-gradient dependent, low-affinity (K_m 0.5 mM), and high-capacity transporter for uric acid. It also shows a low capacity for transport of xanthine at 37 °C but not at 25 °C. Based on the set of positions delineated as important from our previous Cys-scanning analysis of permease XanQ, we subjected YgfU to rationally designed site-directed mutagenesis. The results show that the conserved His-37 (TM1), Glu-270 (TM8), Asp-298 (TM9), and Gln-318 and Asn-319 (TM10) are functionally irreplaceable, and Thr-100 (TM3) is essential for the uric acid selectivity because its replacement with Ala allows efficient uptake of xanthine. The key role of these residues is corroborated by the conservation pattern and homology modeling on the recently described x-ray structure of permease UraA. In addition, site-specific replacements at TM8 (S271A, M274D, V282S) impair expression in the membrane, and V320N (TM10) inactivates the permease, whereas R327G (TM10) or S426N (TM14) reduces the affinity for uric acid (4-fold increased K_m). Our study shows that comprehensive analysis of structure-function relationships in a newly characterized transporter can be accomplished with relatively few site-directed replacements, based on the knowledge available from Cys-scanning mutagenesis of a prototypic homolog.

Conservation of sequence and overall structure in many evolutionarily broad families of transporters often correlates with high functional variations between homologs. In most cases,

however, it is common that few of the members have been studied for structure-function relationships, whereas high-resolution evidence from crystallography is scarce. To explore the spectrum of substrate selectivity determinants in such families, it is important to introduce effective mutagenesis designs for the comprehensive *ab initio* study of new homologs, based on existing evidence from known members. Capitalization on data from Cys-scanning or other systematic analyses of a well studied homolog, if available, offers a powerful approach to this end.

The above considerations apply promptly to the nucleobase-ascorbate transporter (NAT)² or nucleobase-cation symporter-2 (NCS2) family, an interesting example of a conserved but functionally diverse group of transporters present in all major taxa of organisms. NAT transporters model on the template of the uracil permease UraA, the first and only x-ray structure to be described recently for a member of this family, which represents a novel fold (1). With respect to function, only 15 of more than 2,000 predicted members have been characterized in detail; these are specific for the cellular uptake of uracil, xanthine, or uric acid (microbial, plant, and nonprimate mammalian genomes) or vitamin C (mammalian genomes) (1–3). Two of them, the xanthine permease XanQ of *Escherichia coli* (4–10) and the uric acid/xanthine permease UapA of *Aspergillus nidulans* (11–15), have been studied extensively with Cys-scanning mutagenesis and reverse and forward genetics, respectively. These studies have shown striking similarities between key NAT determinants of the two transporters, reinforcing the idea that few residues at conserved motifs of the family may be invariably critical for function or underlie specificity differences (for a summary of current knowledge, see supplemental Fig. S1). Based on homology modeling, most of these residues are found at the vicinity or at the periphery of the binding site in transmembrane segments TM1, TM3, TM8, and TM10 (1, 10, 15).

In the current work, we enrich the data set of functionally known NAT members by studying previously uncharacterized homologs from the genome of *E. coli* K-12, and we analyze

* This work was supported by the research project EVOTRANS co-funded by the European Union–European Social Fund (ESF) and National Sources, in the framework of program “Thalis” through the Operational Program “Education and Lifelong Learning” of the Hellenic Ministry of Education, Lifelong Learning and Religious Affairs, in the context of the National Strategic Reference Framework (NSRF).

⁵ This article contains supplemental Tables S1 and S2 and Fig. S1.

¹ To whom correspondence should be addressed. E-mail: efriligo@cc.uoi.gr.

² The abbreviations used are: NAT, nucleobase-ascorbate transporter; TM, transmembrane segment; BAD, biotin acceptor domain; HIU, 5-hydroxyisourate; OHCU, 2-oxo-4-hydroxy-4-carboxy-5-ureidoimidazole; hSVCT, human Na⁺-coupled ascorbic acid transporter-2.

function, substrate selectivity, and the role of key residues in one of them (YgfU), using mutagenesis designs that are based on the well studied homolog XanQ. Strikingly, the genome of *E. coli* K-12 contains 10 predicted members, of which the uracil permease UraA (16) and the xanthine permeases XanQ and XanP (4) are functionally known. Of the remaining members, three cluster together with UraA, XanQ, and XanP in COG2233 (YgfU, RutG, YbbY), and four cluster separately in COG2252 (YgfQ, YjcD, YicO, PurP). Characteristic NAT sequence motifs are retained only by the three COG2233 members (see Fig. 1A). Of them, RutG is related in sequence with the uracil permease UraA and is considered to be part of an unconventional pyrimidine utilization operon (17–19), whereas YgfU is found along with XanQ in a cluster of putative purine catabolic genes (20) and is related in sequence with xanthine or uric acid permeases from Gram-positive bacteria (21) (supplemental Table S1). The COG2252 members are more related in sequence with the fungal and plant AzgA-like adenine-guanine-hypoxanthine transporters (22, 23), and one of them (PurP) has been proposed to be a high-affinity adenine transporter, based on genetic (24) and systems biology evidence (25).

In the context of this work, we cloned and overexpressed YgfU and showed that it is a proton-gradient dependent, low-affinity (K_m , 0.5 mM), and high-capacity transporter for uric acid that also transports xanthine, but with disproportionately low capacity. Subsequently, we subjected YgfU to site-directed mutagenesis, based on data available for the homologous xanthine permease XanQ, and found that residues irreplaceable for the mechanism occur at five highly conserved positions, whereas a single-amino acid replacement (T100A) converts the uric acid-selective YgfU to a dual-selectivity transporter for both uric acid and xanthine. These results are supported with mirror-image replacements made in XanQ and homology modeling on the recently described structure of UraA.

EXPERIMENTAL PROCEDURES

Materials—[8-¹⁴C]Uric acid (51.5 mCi mmol⁻¹), [8-³H]xanthine (28 Ci mmol⁻¹), and [5,6-³H]uracil (59 Ci mmol⁻¹) were purchased from Moravék Biochemicals. Nonradioactive nucleobases were from Sigma. Oligodeoxynucleotides were synthesized from BioSpring GmbH. High-fidelity *Taq* polymerase (Phusion high-fidelity PCR system) was from Finnzymes. Restriction endonucleases used were from Takara. Horseradish peroxidase (HRP)-conjugated avidin was from Amersham Biosciences. All other materials were reagent grade and obtained from commercial sources.

Bacterial Strains and Plasmids—*E. coli* K-12 was transformed according to Inoue *et al.* (26). TOP10F' (Invitrogen) was used for initial propagation of recombinant plasmids. T184 (27) harboring pT7-5/*xanQ* or pT7-5/*ygfU* with given replacements was used for isopropyl-1-thio- β -D-galactopyranoside-inducible expression from the *lacZ* promoter/operator.

DNA Manipulations—Construction of expression plasmids and biotin acceptor domain (BAD)-tagged versions of NAT homologs was essentially as described previously for XanQ and XanP (4). Briefly, the coding sequences of NAT genes were amplified by PCR on the template of genomic DNA prepared from *E. coli* T184 and transferred to plasmid vector pT7-5 by

restriction fragment replacement; BAD-tagged versions were prepared using two-stage (overlap extension) PCR (28) on the templates of pT7-5/*ygfU* (or other NAT) and pT7-5/*xanQ*-BAD. For construction of mutants, two-stage PCR was performed on the template of YgfU-BAD or XanQ-BAD, as indicated. The entire coding sequence of all engineered constructs was verified by double-strand DNA sequencing in an automated DNA sequencer (MWG Biotech) (supplemental Table S2).

Growth of Bacteria—*E. coli* T184 harboring given plasmids was grown aerobically at 37 °C in Luria-Bertani medium containing streptomycin (0.01 mg/ml) and ampicillin (0.1 mg/ml). Fully grown cultures were diluted 10-fold, allowed to grow to mid-logarithmic phase, induced with isopropyl-1-thio- β -D-galactopyranoside (0.5 mM) for an additional 2 h at 37 °C, harvested, and washed with appropriate buffers.

Transport Assays and Kinetic Analysis—*E. coli* T184 were assayed for active transport of [³H]xanthine (0.1–250 μ M), [³H]uracil (1–100 μ M), and [¹⁴C]uric acid (0.01–2 mM), by rapid filtration, at 25 and 37 °C, pH 7.5, as described (4). For kinetic uptake measurements, initial rates were assayed in T184 cells, at 5–20 s, and data were fitted to the Michaelis-Menten equation using Prism4, to determine K_m and V_{max} values. For ligand competition experiments with XanQ mutants, uptake of [³H]xanthine (1 μ M) was assayed in the absence or presence of unlabeled analogues (1 mM) (4–6).

Immunoblot Analysis—Membrane fractions were prepared from 10-ml cultures of *E. coli* T184 harboring given plasmids and subjected to SDS-PAGE (12%), as described (4). Proteins were electroblotted to polyvinylidene difluoride membranes (Immobilon-PVDF; Pall Corp.). The BAD-tagged permeases were probed with avidin-HRP. Signals were developed with enhanced chemiluminescence (ECL).

In Silico Analysis—Comparative sequence analysis of NAT/NCS2 homologs was based on BLAST-p search and ClustalW alignment; the most recent genome annotations were used for retrieving sequence data. Initial analysis of transmembrane topology of YgfU was performed using the program TMHMM (29). Homology threading was performed using the known x-ray structure of UraA as a template (Protein Data Bank (PDB) code 3QE7) on the web-based SWISSPROT modeling server (30); results were displayed and analyzed with PyMOL version 1.4 (Schrödinger, LLC).

RESULTS

YgfU Is a Uric Acid Transporter—The NAT genes *ygfU*, *rutG*, *ybbY*, *ygfQ*, *yjcD*, *yicO*, and *purP* were mobilized from the *E. coli* K-12 genome, transferred to transcriptional control of the *lacZ* promoter/operator in plasmid vector pT7-5, and induced for overexpression in *E. coli* T184 at conditions of negligible endogenous activity of oxidized purine or uracil uptake (4). Our data show that with one exception, all NAT constructs can be expressed in the *E. coli* membrane at high levels, comparable with the ones of XanQ used as a control (Fig. 1B), whereas transport activity at 25 °C is observed only with YgfU, which transports [¹⁴C]uric acid (1 mM) to high levels, and RutG, which has a marginally detectable activity for [³H]xanthine (1 μ M) (Fig. 1C). Kinetic analysis reveals that RutG transports xanthine

Selectivity of Transporter YgfU for Uric Acid




FIGURE 1. Sequence, immunoblot, and nucleobase uptake analysis of NAT homologs from *E. coli* K-12 were aligned using ClustalW, and part of this alignment referring to conserved sequence motifs of the family is shown in *A*. The members include the known nucleobase transporters XanQ (P67444), XanP (P0AGM9), and UraA (P0AGM7) and the unknown YgfU (Q46821), RutG (P75892), YbbY (P77328), YgfQ (Q46817), YjcD (P0AF52), YicO (P31440), and PurP (P31466). The motifs shown are characteristic for functionally known members of the family and have been shown to contain functionally important residues in XanQ (6–10). Sequence identity scores are shown on the *right*. The coding sequences from the unknown homologs were mobilized from the genome, induced for overexpression in *E. coli* T184 under control of the *lacZ* promoter/operator in pT7-5/NAT-BAD, and assayed for expression levels in the membrane and active transport of xanthine, uracil, and uric acid. *B*, samples containing 100 μ g of membrane protein were subjected to SDS-PAGE (12%) and immunoblotting using HRP-conjugated avidin. *C*, cells expressing each homolog were normalized to 0.7 mg of protein/ml of cell suspension and assayed for active transport of [14 C]uric acid (1 mM, 25 $^{\circ}$ C) in the presence or absence of carbonyl cyanide *m*-chlorophenyl hydrazone (CCCP) (10 μ M). Transport assays were also performed for [3 H]xanthine (1 μ M, 25 $^{\circ}$ C), using XanQ as a positive control (4), and for [3 H]uracil (1 μ M, 25 $^{\circ}$ C). Uracil uptake could not be detected with any of the unknown homologs; xanthine uptake was observed only with RutG, albeit to a marginal extent (not shown).

with 50-fold lower affinity (K_m 0.15 mM) in comparison with XanQ or XanP (4), and YgfU transports uric acid with low affinity (K_m 0.5 mM) and high capacity (V_{max} 715 nmol min $^{-1}$ mg $^{-1}$) (Table 1). The steady-state level of uric acid accumulation achieved with YgfU after 2–15 min of exposure to uric acid (1 mM) (Fig. 1C) is 381 ± 4 nmol mg $^{-1}$; taking into account an internal volume for *E. coli* of about 5.8 μ l mg $^{-1}$ (31), this level corresponds to a concentration gradient of 128-fold (in/out). In addition, this uptake activity is abolished in the presence of the protonophore carbonyl cyanide *m*-chlorophenyl hydrazone, implying that transport is dependent on proton symport (32). With respect to xanthine uptake, YgfU displays a marginal xanthine uptake activity at 25 $^{\circ}$ C, which precludes kinetic analysis, but it can transport xanthine at 37 $^{\circ}$ C, albeit with very low affinity (K_m 0.3 mM; 100-fold higher than the K_m values of XanP or XanQ) and capacity (V_{max} 14 nmol min $^{-1}$ mg $^{-1}$) (Table 1).

Delineation of Mutagenesis Targets for Structure-Function Analysis of YgfU—The Cys-scanning and site-directed analysis of the homologous XanQ (5–10) offers a basis to search for

targets of efficient mutagenesis designs in YgfU. Important residues of XanQ have been delineated at positions where a side chain is irreplaceable with respect to function (Glu-272, Asp-304, Gln-324, Asn-325) or expression in the membrane (Gly-83, Pro-318), replaceable with a limited number of other side chains (His-31, Asn-93, Phe-94, Asp-276, Ala-279, Thr-280, Gly-305, Gly-351, Pro-354, Ile-432), or highly sensitive to inactivation of a substituted Cys mutant by *N*-ethylmaleimide (Leu-86, Val-261, Gly-275, Ser-284, Ala-323, Asn-326, Gly-327, Val-328, Ile-329, Thr-332, Gly-333, Ser-336, Asn-430). Most of these 29 residues are highly conserved in the NAT/NCS2 family (Fig. 2), and YgfU differs from XanQ at 9 of the 29 positions (supplemental Fig. S1). To examine whether these residues contribute to the different substrate preference of YgfU, we engineered mutants replacing each one of the 9 residues with the corresponding amino acid found in XanQ. Mutagenesis was also performed at conserved side chains corresponding to the 4 functionally irreplaceable residues of XanQ and His-31, which is essential for substrate affinity (7). Thus, our initial mutagen-

TABLE 1

 K_m and V_{max} values of YgfU, XanQ, and mutants

E. coli T184 expressing the corresponding constructs were assayed for initial rates of [14 C]uric acid (0.01–2 mM) and [3 H]xanthine (0.1–250 μ M) uptake at 5–20 s, at either 25 °C or 37 °C, as given; negative control values obtained from T184 harboring vector pT7–5 alone were subtracted from the sample measurements in all cases. Kinetic parameters were determined from non-linear regression fitting to the Michaelis-Menten equation using Prism4; values represent the means of three independent determinations with standard deviations shown. ND, assays were performed, but kinetic values were not determined due to very low uptake rates. All mutants and the wild-type versions used contained a carboxyl-terminal BAD.

	Kinetics for uric acid uptake (25 °C)		
	K_m	V_{max}	V_{max}/K_m
	mM	nmol min ⁻¹ mg ⁻¹	μ l min ⁻¹ mg ⁻¹
Permease			
YgfU(wt)	0.5 ± 0.1	715 ± 81	1430
H37N	1.4 ± 0.3	23 ± 8	16
H37K/H37L	ND	ND	
T100A	0.9 ± 0.4	700 ± 160	777
T100S	1.9 ± 0.7	296 ± 87	159
S256Q	0.8 ± 0.1	698 ± 50	872
T259V	0.8 ± 0.3	808 ± 150	1049
L278T	0.6 ± 0.3	609 ± 165	1015
E270D/E270N/E270Q	ND	ND	
D298E/D298N	ND	ND	
S317A	0.5 ± 0.2	535 ± 104	1070
Q318N/Q318E	ND	ND	
N319Q/N319D	ND	ND	
V320N	ND	ND	
R327G	2.2 ± 0.9	2064 ± 565	938
S426N	1.9 ± 0.3	808 ± 150	946
	Kinetics for xanthine uptake (37 and 25 °C)		
	K_m	V_{max}	V_{max}/K_m
	μ M	nmol min ⁻¹ mg ⁻¹	μ l min ⁻¹ mg ⁻¹
Permease			
YgfU(wt) (25 °C)	ND	ND	
YgfU(wt) (37 °C)	300 ± 60	14 ± 4	46
T100A (37 °C)	240 ± 5	132 ± 20	550
T100S	ND	ND	
XanQ(wt) (25 °C)	4.1 ± 0.5	7.2 ± 1.1	1756
N93T	11.6 ± 2.9	5.0 ± 0.4	431
N93A	2.4 ± 0.7	5.0 ± 0.5	2083
Q258S	ND	ND	
V261T	7.4 ± 0.7	23.2 ± 0.8	3135
A273S	24.1 ± 0.5	113.1 ± 3.5	4688
D276M	ND	ND	
T280L	4.3 ± 0.3	14.8 ± 0.4	3442
S284V	5.1 ± 0.5	8.8 ± 0.3	1725
A323S	13.9 ± 4.1	37.5 ± 4.5	2698
N326V	11.4 ± 1.4	23.1 ± 1.0	2026
G333R	2.8 ± 0.3	2.4 ± 0.2	857
N430S	12.0 ± 1.3	6.6 ± 0.3	550

esis targets included His-37, Thr-100, Thr-259, Glu-270, Met-274, Leu-278, Val-282, Asp-298, Ser-317, Gln-318, Asn-319, Val-320, Arg-327, and Ser-426 (residues conserved in XanQ are underlined) (Fig. 3).

Site-directed Mutagenesis of YgfU—Based on the above, we constructed and analyzed single-replacement mutants (a) E270D, E270N, E270Q, D298E, D298N, Q318N, Q318E, N319Q, N319D (conservative replacements of residues that are invariant and functionally irreplaceable in XanQ); (b) H37N, H37K, H37L, T100N, T100S, T100A (replacements of side chains that correspond to affinity- or specificity-related residues in XanQ); and (c) T259V, M274D, L278T, V282S, S317A, V320N, R327G, S426N (replacements of nonconserved residues of the important set with the corresponding amino acid found in XanQ). In addition, prompted by our initial data (see below), we engineered replacements at positions of TM8 that correspond to moderately *N*-ethylmaleimide-sensitive mutants (S256Q and S271A). The complete set of YgfU residues sub-

jected to mutagenesis and their counterparts in XanQ is given in supplemental Fig. S1. After verification of the sequence, each YgfU mutant was transferred and expressed in *E. coli* T184 and assayed for its expression levels in the membrane and its ability to catalyze active transport of uric acid and xanthine (Fig. 4). Of the 25 mutants, four are undetectable in the membrane (T100N, S271A, M274D, V282S). Of the remaining 21 mutants, which display high protein levels comparable with wild type, nine show negligible uptake activities for uric acid or xanthine (E270D, E270N, E270Q, D298E, N319Q, N319D, H37K, H37L, V320N). Of the 12 active mutants, four show marginal activity for uric acid uptake (5–12% of wild-type) (Q318N, Q318E, H37N, D298N), two accumulate uric acid at low (T100S, 20%) or intermediate levels (S317A, 40%), and six are highly active (60–120%) (T100A, S256Q, T259V, L278T, R327G, S426N). With respect to xanthine uptake, significant activities above the wild-type threshold are only attained by mutants T100A, T100S, and S317A (Fig. 4C). Mutant T100A, in particular, which is highly active with both uric acid and xanthine, displays very significant xanthine uptake activity (about 50% of wild-type XanQ used as a control in these assays) even at 25 °C, where wild-type YgfU is inactive (Fig. 5). Kinetic transport analysis shows that three mutants (T100S, R327G, S426N) have significantly lower affinity for uric acid than wild type (4-fold higher K_m values), whereas T100A does not differ significantly from wild type in affinity but displays a 10-fold higher V_{max} for uptake of xanthine at 37 °C (Table 1). When comparing uric acid with xanthine uptake activity for wild-type and T100A, it is clear that this replacement converts the uric acid-selective YgfU to a dual-selectivity transporter for both uric acid and xanthine (Fig. 5D).

Mirror-image Replacements in XanQ—To evaluate further the role of nonconserved YgfU residues, we performed reverse site-directed mutagenic analysis in XanQ. Each one of the 11 important residues of XanQ that differ in YgfU was replaced individually with the corresponding amino acid found in YgfU. Thus, we constructed XanQ mutants Q258S, V261T, A273S, T280L, S284V, A323S, N326V, and N430S and assayed them for expression, uptake activity, and substrate selectivity in parallel with mutants N93T (7), N93A (10), D276M (9), and G333R (8) that had been engineered by us in previous studies. As shown in Fig. 6, all these mutants are expressed in the *E. coli* membrane to high levels. Of them, two show negligible uptake activities for xanthine or uric acid (Q258S, D276M), two display very low xanthine uptake activities (10–20% of wild type) (N93T, N430S), seven display intermediate xanthine uptake activities (40–60% of wild type) (N93A, V261T, T280L, S284V, A323S, N326V, G333R), and one is highly active for xanthine uptake (100–120%) (A273S). On the other hand, only N93A shows a low but detectable uptake activity for uric acid (Fig. 6C). Kinetic transport analysis shows that five mutants have significantly lower affinity for xanthine than wild type (3-fold higher K_m values for mutants N93T, A323S, N326V, and N430S and 6-fold higher K_m value for A273S) (Table 1). Finally, the specificity profile of each mutant was investigated by assaying the extent of inhibition of [3 H]xanthine uptake in the presence or absence of a series of purine analogues (Table 2). It is very interesting that despite their inefficiency for uric acid uptake, most

Selectivity of Transporter YgfU for Uric Acid




FIGURE 2. Conservation pattern of important residues of XanQ in NAT family. A logotype of 29 residues delineated as important from the Cys-scanning analysis of XanQ (5–10) is shown on top. Larger sized letters indicate irreplaceable XanQ residues, medium sized letters indicate residues that are replaceable with a limited number of side chains, and smaller sized letters indicate residues where a Cys replacement is highly sensitive to inactivation by *N*-ethylmaleimide ($IC_{50} < 100 \mu\text{M}$). A perpendicular line indicates a residue where replacements distort specificity for the imidazole moiety of substrate (leading to aberrant recognition of uric acid and 8-methylxanthine). Italics indicate residues that are crucial (smaller size) or irreplaceable (larger size) for expression in the membrane. The histogram in the upper panel shows the number and relative abundance of different amino acids found at each position of the important set in the 189 closest homologs (>33% sequence identity with XanQ) (10); the corresponding variants are listed in decreasing order of frequency, whereas variants with very low occurrence ($\leq 1.6\%$) are omitted for clarity. The sequence alignments in the lower panel show the residues found at each position of the important set in 15 characterized NAT/NCS2 transporters, including *E. coli* XanQ (P67444), XanP (POAGM9) and UraA (POAGM7), *B. subtilis* PbuX (P42086), PucK (O32140) and PucJ (O32139), *Lactococcus lactis* PyrP (AAK05701), *A. nidulans* UapA (Q07307) and UapC (P487777), *Aspergillus fumigatus* AfUapA (XP748919), *Candida albicans* Xut1 (AAX2221), *Zea mays* Lpe1 (AAB17501), *Homo sapiens* SVCT1 (SLC23A1) (AAH50261) and SVCT2 (SLC23A2) (Q9UGH3), and *Rattus norvegicus* SNBT1 (SLC23A4) (AB511909). The full-length sequences were aligned using ClustalW, and part of this alignment including the residues of interest is presented. Major substrates of each transporter are shown in parentheses on the left. The abbreviations are: *Xan*, xanthine; *UA*, uric acid; *Ura*, uracil; and *Asc*, ascorbate.

of these mutants allow the recognition of uric acid or other analogues that differ from xanthine at the imidazole moiety and are not recognized by wild-type XanQ. Thus, uric acid inhibits N93A, V261T, or G333R by 40–50%, 8-methylxanthine inhibits N93A or G333R by 80–90% and V261T, A273S, T280L, A323S, or N326V by 50–70%, and 7-methylxanthine inhibits G333R by 80% (Table 2).

DISCUSSION

The results in this study yield important new knowledge in two respects. First, it is shown for the first time that the *E. coli* K-12 genome encodes a uric acid permease activity (YgfU). Second, key determinants of the substrate binding site function and specificity of this permease are delineated, and Thr-100 in TM3 is demonstrated to be essential for the selectivity for uric acid.

The finding that the *ygfU* gene product in *E. coli* is a high-capacity transporter for uric acid raises questions on the putative physiological significance of such an uptake activity because *E. coli* is not currently known to contain enzymes for uric acid catabolism or to have a complete catabolic pathway for purines. Previous studies have shown that *E. coli* K-12 is unable to use purines other than adenine as sole nitrogen sources and, even with adenine, nitrogen assimilation is incomplete (20), but it can grow on allantoin as a sole nitrogen source, albeit only at anaerobic conditions (33). It also contains xanthine dehydrogenase activity (20), which can divert hypoxanthine and xanthine from the purine salvage pathway and produce uric acid. The missing link, however, is in the catabolic oxidation of uric acid to allantoin.

As shown relatively recently (34), the oxidation of uric acid to allantoin is expected to involve three enzymatic steps, the first




FIGURE 3. **Topology model of YgfU permease and mutagenesis targets.** The model is derived from homology threading using the UraA structure (3QE7) as a template and initial topology predictions based on the program TMHMM and analysis of the XanQ homolog for the accessibility of loops to hydrophilic reagents (8). The α -helical segments are indicated in *rectangles with continuous line* (core domain) or *broken line perimeters* (gate domain). Residues corresponding to the ones analyzed with Cys-scanning mutagenesis in XanQ are *circled*. Residues corresponding to the important set (Fig. 2) are shown in a *dark background* (functionally important in XanQ) or in a *bold circle* (crucial for expression in XanQ). Residues corresponding to positions of XanQ mutants that are moderately sensitive to *N*-ethylmaleimide (IC_{50} values from 100 μ M to 1 mM) are in a *gray background*. The residues of YgfU analyzed in this study with site-directed mutagenesis are *numbered* and shown in *blue* (irreplaceable for function), *red* (essential for the counter-selectivity against xanthine), *dark red* (replacements lead to lower affinity for uric acid), *orange* (replacements lead to abolishment of expression in the membrane), or *gray scale* (remaining positions).

one catalyzed by urate oxidase and yielding 5-hydroxy-isourate (HIU), the second one catalyzed by HIU hydrolase, which converts HIU to 2-oxo-4-hydroxy-4-carboxy-5-ureidoimidazole (OHCU), and the third one catalyzed by OHCU decarboxylase, which converts OHCU to dextrorotatory *S*(+)-allantoin, the substrate for (*S*)-allantoin amidohydrolase (AllB) (35). In *E. coli* K-12, no enzymatic activity or any candidate *E. coli* gene has been reported for the step of oxidation of uric acid to HIU, whereas HIU hydrolase, and suggestively, OHCU decarboxylase activity has been assigned experimentally to the product of gene *yedX* (EcoGene database).³ Interestingly, *E. coli* HIU hydrolase and its *Bacillus subtilis* homolog PucM (36) show clear sequence and structural similarity with the vertebral thyroid hormone distributor protein transthyretin, which probably originated from a duplication of an ancestral HIU hydrolase gene (37). In contrast to most bacterial homologs, the HIU hydrolase genes from *E. coli* and other enterobacteria appear not to be associated with purine metabolism operons and have been suggested to encode proteins with periplasmic localization, which might utilize HIU produced by a spontaneous, non-enzymatic reaction of uric acid acting as an antioxidant in the periplasm (38).

³ K. Rudd, personal communication.

On the other hand, a complete gene cluster for purine utilization (*hpx* cluster) has been recently identified in *Klebsiella pneumoniae* (39) and *Klebsiella oxytoca* (40) containing separate genes for all three enzymatic steps of uric acid oxidation, namely *hpxO* (urate hydroxylase), *hpxT* (HIU hydrolase), and *hpxQ* (OHCU decarboxylase). Apart from HpxT (HIU hydrolase), which is homologous to *E. coli* HIU hydrolase and *B. subtilis* PucM, HpxO was found to function as a novel FAD-dependent urate oxidase (41), distinct from the classical urate oxidase (uricase) known for *B. subtilis* (36) and other species with complete purine utilization pathways (42). Parts of the *hpx* gene cluster of *Klebsiella* including the HpxO homolog are also encountered in other γ -proteobacterial genera such as *Xanthomonas* and *Acinetobacter* (40) but not in *E. coli*. Finally, the *hpx* cluster contains a single putative permease gene (*hpxP*), which was suggested to encode a general purine permease (40). HpxP belongs to the NAT/NCS2 family and is closely homologous with the *E. coli* xanthine permease XanQ (45% sequence identity) but shares limited sequence homology with YgfU (25% identity).

Overall, it appears that there is more to learn on the purine utilization pathways in *E. coli* K-12, and our identification of YgfU as a uric acid transporter is an important contribution to this end. Although YgfU is found along with XanQ in a cluster

Selectivity of Transporter YgfU for Uric Acid




FIGURE 4. Expression and uptake activities of YgfU mutants. *E. coli* T184 harboring pT7-5/ygfU(wild-type-BAD) with given mutations were grown aerobically to mid-logarithmic phase, induced with isopropyl-1-thio- β -D-galactopyranoside (0.5 mM) for 2 h, and subjected to immunoblot analysis of membrane fractions to mid-logarithmic phase, induced with isopropyl-1-thio- β -D-galactopyranoside (0.5 mM) for 2 h, and subjected to immunoblot analysis of membrane fractions or assayed for transport of [14 C]uric acid (1 mM, 25 °C) and [3 H]xanthine (0.1 mM, 37 °C). **A**, samples containing ~100 μ g of membrane protein were subjected to SDS-PAGE (12%) and immunoblotting using HRP-conjugated avidin. Prestained molecular mass standards (in kDa; Bio-Rad, low range) are shown on the left. **B**, initial rates of uric acid uptake (closed histogram bars) were measured at 5–15 s, and steady-state levels of uric acid accumulation (open histogram bars) were measured at 2–10 min. Control values obtained from T184 harboring pT7-5 alone were subtracted from the sample measurements in all cases. Results are expressed as a percentage of the activity of wild-type control with standard deviations from three independent determinations. **C**, comparison of initial rates of uric acid and xanthine uptake for the active YgfU mutants. Results are expressed as a percentage of the rate of YgfU and XanQ, respectively. The percentage ratio of the xanthine uptake value (red) to the uric acid uptake value (black) is given for each mutant in a third histogram bar (blue).

of putative purine catabolic genes (20), this cluster does not appear to contain homologs for catabolic utilization of uric acid. In addition, YgfU has a limited array of orthologs in γ -Proteobacteria (for example, only six homologs with >50% identity in other enterobacteria), but it shares up to 50% sequence identity with many homologs from Gram-positive bacteria including the known xanthine permease PbuX and uric acid permeases PucJ and PucK from *B. subtilis* (supplemental Table S1). Apart from uric acid, YgfU can utilize as a substrate xanthine, but with disproportionately low capacity, which is only apparent at 37 °C (Fig. 5 and Table 1). In addition, the affinity of YgfU for xanthine is very low relative to the affinities of the xanthine permeases XanP and XanQ (4). Based on the above, the major substrate transported by YgfU appears to be uric acid, and we would like to suggest UacT (uric acid transporter) as an alternative, function-based designation for YgfU.

The comprehensive mutagenesis study of YgfU (UacT) presented herein reveals the key role of His-37 (TM1), Glu-270 (TM8), Asp-298 (TM9), and Gln-318 and Asn-319 (TM10), which are functionally irreplaceable, and of Thr-100 (TM3),

which is essential for the uric acid selectivity. Structural modeling of YgfU based on the known x-ray structure of the uracil-transporting homolog UraA (1) shows that five of the six important residues fall at transmembrane helices, which form a binding shelter around substrate, and at least three of them (Glu-270, Gln-318, Thr-100) are in the substrate binding pocket (Fig. 7). In other members of the NAT/NCS2 family, these residues are either absolutely conserved (His-37, Glu-270, Asn-319) or conform to a distinct conservation pattern with limited side chain changes (Fig. 2). Most importantly, as well, critical roles have been described for the corresponding residues of other known NAT homologs (1, 7–13, 43), as addressed in more detail below.

The five irreplaceable YgfU residues are conserved in the xanthine-transporting homolog XanQ and the dual-specificity uric acid/xanthine transporter UapA, and four out of five are also irreplaceable for function in XanQ (8, 9) and in UapA (12, 13). The irreplaceable role of the counterparts of Gln-318 and Asn-319, which belong to the conserved NAT signature motif (11), has been linked with defining the function of the binding




FIGURE 5. Analysis of substrate selectivity for mutant T100A. *E. coli* T184 harboring pT7-5/*ygfU* (wild-type-BAD), pT7-5/*ygfU* (T100A-BAD), pT7-5/*xanQ* (wild-type-BAD), or pT7-5 alone were grown, induced, and assayed for transport of [¹⁴C]uric acid (0.01–2 mM, 25 °C) and [³H]xanthine (0.1 μM–0.25 mM, 37 °C or 25 °C), as described in the legends for Fig. 4 and Table 1. A, time courses of xanthine (0.1 mM) uptake at 37 °C. B, time courses of xanthine (0.1 mM) uptake at 25 °C. C, time courses of uric acid (1 mM) uptake at 25 °C. D, comparison of the transport efficiencies (V_{max}/K_m) for uric acid (25 °C) and xanthine (37 °C) displayed by YgfU and T100A (see also Table 1). The abbreviations are: UA, uric acid; and X, xanthine.

site in both XanQ (5, 8) and UapA (12, 13), and site-directed alkylation studies in XanQ have implied that the corresponding residues interact directly with purine substrate (8). In the known x-ray structure of the pyrimidine-specific homolog UraA (1), the position of Gln-318 is occupied by a Glu (Glu-290), which forms two direct hydrogen bonds with substrate (uracil); the counterpart of Asn-319 (Asn-291) stabilizes the structure through an indirect hydrogen-bonding effect involving the counterpart of His-37 (His-24) in TM1. The role of another irreplaceable YgfU residue (Glu-270) might also be linked directly with binding because its counterpart in UraA (Glu-241) also forms two direct hydrogen bonds with substrate in the crystal structure (1). The fourth invariably irreplaceable residue (Asp-298 in YgfU) corresponds to a position of UraA, which is at the same depth in the membrane but distal from the binding site in the crystal structure and might have a distinct conformational role in the purine translocation mechanism, as suggested for permease XanQ (7, 9). In any event, the fact that four conserved residues associated either directly or indirectly with binding are functionally irreplaceable in XanQ, YgfU (UacT), and UapA highlights the functional conservation of the purine binding site in NAT/NCS2 homologs of different selectivity.

His-37 (TM1) is absolutely conserved in the NAT/NCS2 family. Although replaceable in transporters other than YgfU, this histidine appears to be irreplaceable for function in YgfU (Fig. 4). From the other homologs, which include the bacterial XanQ (7), the fungal UapA (15), and the mammalian hSVCT1

(43) and hSVCT2 (44), a conserved hydrogen-bonding role is apparent for this residue, which is crucial for either the overall structural folding or the constitution of a proper affinity and specificity binding site or for both. Namely, the presence of an uncharged hydrogen-bonding residue at this position is essential for high-affinity binding and transport in XanQ (7) and in the hSVCTs (43, 44), whereas in UapA, it has been suggested as crucial for proper folding and targeting to the plasma membrane (15). The recently elucidated crystal structure of the different-specificity homolog UraA (1) as well as homology models built on this structure (10, 15) show that the conserved His of TM1 hydrogen-bonds to the conserved Asn of the NAT signature motif at the binding site periphery, substantiating further the previous findings. Based on the mutagenesis data of Fig. 4, it appears that this hydrogen-bonding interaction between His-37 and Asn-319 (Fig. 7) is more stringent in YgfU than in other homologs, and it may play an irreplaceable role with respect to the binding site integrity in this transporter.

Replacement of Thr-100 (TM3) with Ala changes the uric acid-selective YgfU to a dual-selectivity variant transporting both uric acid and xanthine with comparable efficiencies (Fig. 5), and replacement of the same residue with Ser also allows improved xanthine uptake relative to the uric acid uptake activity (Fig. 4). Thus, the side chain of Thr-100 is linked with the transporter ability to differentiate between uric acid (8-oxy-xanthine) and xanthine. A similar role has been proposed for the corresponding amino acid in permease XanQ (Asn-93) because replacement of Asn-93 with Ala or Ser converts the

Selectivity of Transporter YgfU for Uric Acid




FIGURE 6. Expression and uptake activities of XanQ mutants. *E. coli* T184 harboring pT7-5/*xanQ*(wild-type-BAD) with given mutations were grown, induced, and subjected to immunoblot analysis of membrane fractions (A) or assayed for transport of [³H]xanthine (1 μ M, 25 °C) (B) and [¹⁴C]uric acid (2 mM, 25 °C) (C), exactly as described in the legend for Fig. 4. Closed and open histogram bars represent initial rate and steady-state values, respectively. Results are expressed as a percentage of the activity of wild-type control with standard deviations from three independent determinations.

xanthine-selective XanQ to dual-selectivity variants (7, 10); in the case of XanQ, however, the dual-selectivity variants transport the non-wild-type substrate (uric acid) with very low capacity (10). The selectivity-related role of Thr-100/Asn-93 appears to be unique because no other single-replacement mutant of either XanQ (6–10) or YgfU (this study) has been shown to convert the native transporter to a dual-selectivity one. Interestingly, a similar albeit less prominent effect was observed with mutants replacing the corresponding residue (Ser-154) in the fungal homolog UapA, where introduction of an Ala leads to higher affinity for xanthine relative to uric acid, thus shifting the dual-selectivity profile to the xanthine-selective direction (15).

Although the side chains of Thr-100 or Asn-93 *per se* are poorly conserved in other homologs (10), the polar character of Thr-100/Asn-93 is conserved invariably in the known nucleobase-transporting NAT members (Asn, Thr, or Ser), whereas the ascorbate-transporting members have an Ala (Fig. 2). Furthermore, all the dual-selectivity uric-acid/xanthine transporters (Xut1, UapA, UapC, AfUapA, Lpe1) have a Ser at this position (Fig. 2). To understand the structural relevance of this difference, we have built models of the xanthine-selective XanQ, the uric acid-selective YgfU, and the dual-selectivity fungal homologs on the template of the known structure of UraA (Fig. 7 and data not shown). The models indicate that the relevant position of TM3 is vicinal to the substrate binding site formed between residues of the middle parts of TM3, TM8, and TM10 (1). Strikingly, however, the Thr or Ser replacing Asn-93 in YgfU and all dual-selectivity NATs is distal from the conserved, substrate-relevant glutamate of TM8 (minimal distance between oxygen atoms, 6.0 Å), whereas Asn-93 in XanQ is sig-

nificantly closer (distance between oxygen atoms, 4.5 Å). This difference is most prominent in YgfU (Fig. 7) and UapA (10), which conserve nearly all the other functionally important residues of TM1, TM3, TM8, or TM10, except Asn-93.

In UapA (10, 15), occupation of the Asn-93 position by Ser orients this residue away from the substrate-relevant Glu of TM8, leading to relaxation of a constraint for the recognition of analogues at position 8 and allowing binding and transport of uric acid (8-oxy-xanthine), which shifts the NAT selectivity to a less stringent, dual-substrate profile. Accordingly, the XanQ mutants replacing Asn-93 with Ser or Ala yield efficient recognition of 8-methylxanthine and low but significant uptake of uric acid, mimicking in part the fungal, dual-selectivity NATs (10).

In YgfU, Thr-100 is also oriented away from the carboxyl group of Glu-270 (TM8), leaving more space in the substrate binding pocket but, at the same time, the pK_a of Glu-270 may be distorted significantly relative to the corresponding carboxylic acid in XanQ due to its vicinity with hydrophobic groups from Thr-100 and Met-274 (Fig. 7B). These changes on the substrate binding Glu-272/Glu-270 might account for the selectivity difference between YgfU (uric acid) and XanQ (xanthine). Interestingly, however, replacement of Asn-93 with Thr in XanQ cannot imitate the YgfU profile but leads to low affinity for all xanthine analogues, possibly due to interference of the methyl group of Thr-93 in the vicinity of the essential Glu-272, which is not counterbalanced by other changes (10). An even more dramatic effect is observed in the current study with the mirror-image YgfU replacement T100N, which leads to complete lack of expression in the membrane (Fig. 4). Based on these observations, it is evident that further combinatorial replacements

TABLE 2

Specificity profile of XanQ mutants

E. coli T184 expressing the corresponding constructs were assayed for initial rates of [³H]xanthine (1 μ M) uptake at 5–20 s, in the absence or presence of 1000-fold excess (1 mM) of unlabeled competitors. The uptake value obtained in the absence of competitor was taken as 100%. Values represent the means of three determinations, with standard deviations <20%. Most significant differences from the wild-type profile are highlighted in bold. All mutants and the wild-type XanQ version used contained a carboxyl-terminal BAD. Data on mutants N93T/N93A and G333R are from Refs. 10 and 8, respectively.

	Xanthine analogue used as competitor ^a										
	XanQ	N93T	N93A	V261T	A273S	T280L	S284V	A323S	G333R	N326V	N430S
Imidazole moiety	%	%	%	%	%	%	%	%	%	%	%
Uric acid	96	115	60	55	102	102	98	100	50	100	75
8-Methylxanthine	96	69	10	48	39	37	82	33	18	35	81
7-Methylxanthine	105	121	62	69	75	74	92	61	20	63	69
Oxypurinol	31	13	2	39	17	8	16	14	10	44	45
Pyrimidine moiety											
2-Thioxanthine	14	43	16	34	12	6	32	0	12	24	18
3-Methylxanthine	25	49	41	55	50	18	29	6	1	42	29
6-Thioxanthine	20	84	24	42	15	11	30	5	11	36	42

^a Xanthine uptake rate retained is shown as percentage.




FIGURE 7. **Arrangement of important residues in modeled structure of YgfU.** The sequences of YgfU and XanQ were threaded on the known x-ray structure of UraA (1) using the SWISSPROT modeling server. The structural models were displayed with PyMOL version 1.4. Shown are four TMs that contain binding residues and form a shelter around substrate in UraA, namely TM1 (gray), TM3 (wheat), TM8 (teal), and TM10 (salmon). A, arrangement of the four TMs in the YgfU model. B, the same model (YgfU) excluding the β -strand regions of TM3 and TM10 and indicating the side chains of residues analyzed with mutagenesis in this study. C, arrangement of the corresponding residues and helices in XanQ. The side chains of important residues of YgfU (B) and XanQ (C) are shown as thick sticks (irreplaceable and/or substrate binding-relevant) or spheres (sensitive to alkylation or bulky replacement in XanQ; crucial for the protein expression or affinity in YgfU). The side chain of Thr-100 (or Asn-93), which is associated with the uric acid selectivity, is shown with red label.

are needed to promptly convert XanQ to the YgfU selectivity profile or *vice versa*, and targets for such replacements can be selected from residues of the important set analyzed here (Fig. 2), taking into account their orientation relative to the binding site in the modeled structures and functional properties of the relevant single-replacement mutants.⁴

⁴ E. Karna, K. Papakostas, and S. Frillingos, manuscript in preparation.

Apart from the results discussed above, our initial mutagenic analysis of YgfU permease revealed that site-specific replacements at TM8 (S271A, M274D, V282S) impair expression in the membrane, V320N (TM10) inactivates the permease, S317A (TM10) allows substantial uptake of xanthine, and R327G (TM10) or S426N (TM14) reduces the affinity for uric acid (4-fold increased K_m). The significance of these additional findings will be evaluated further with ongoing mutagenesis

Selectivity of Transporter YgfU for Uric Acid

studies in this and other NAT homologs.⁵ In any event, it is important to note that several mutants of YgfU at positions of TM8, TM10, TM3, and TM1 flanking the binding site in the modeled structure lead to grossly impaired activity (H37N, V320N) or lack of expression in the membrane (T100N, S271A, M274D, V282S), whereas their counterpart or mirror-image replacement mutants in XanQ are active and expressed normally (Fig. 6). This observation implies that the binding site of YgfU is more prone to destabilization induced by peripheral side chain changes, whereas the reciprocal changes in XanQ are permissive for structure and function.

Finally, from a methodological point of view, it should be emphasized that all the results presented here for YgfU permease were based on site-directed mutagenesis designs deduced from Cys-scanning analysis data of a well studied homolog (XanQ). In this respect, we have shown that comprehensive analysis of structure-function relationships in a newly characterized transporter can be accomplished with relatively few site-directed replacements, based on the knowledge available from Cys-scanning mutagenesis of a prototypic homolog.

Acknowledgment—We thank Dr. Kenneth Rudd for helpful discussions on an appropriate function-based designation for YgfU.

REFERENCES

- Lu, F., Li, S., Jiang, Y., Jiang, J., Fan, H., Lu, G., Deng, D., Dang, S., Zhang, X., Wang, J., and Yan, N. (2011) Structure and mechanism of the uracil transporter UraA. *Nature* **472**, 243–246
- Yamamoto, S., Inoue, K., Murata, T., Kamigaso, S., Yasujima, T., Maeda, J. Y., Yoshida, Y., Ohta, K. Y., and Yuasa, H. (2010) Identification and functional characterization of the first nucleobase transporter in mammals: implication in the species difference in the intestinal absorption mechanism of nucleobases and their analogs between higher primates and other mammals. *J. Biol. Chem.* **285**, 6522–6531
- Tsukaguchi, H., Tokui, T., Mackenzie, B., Berger, U. V., Chen, X. Z., Wang, Y., Brubaker, R. F., and Hediger, M. A. (1999) A family of mammalian Na⁺-dependent L-ascorbic acid transporters. *Nature* **399**, 70–75
- Karatza, P., and Frillingos, S. (2005) Cloning and functional characterization of two bacterial members of the NAT/NCS2 family in *Escherichia coli*. *Mol. Membr. Biol.* **22**, 251–261
- Karatza, P., Panos, P., Georgopoulou, E., and Frillingos, S. (2006) Cysteine-scanning analysis of the nucleobase-ascorbate transporter signature motif in YgfO permease of *Escherichia coli*: Gln-324 and Asn-325 are essential, and Ile-329-Val-339 form an α -helix. *J. Biol. Chem.* **281**, 39881–39890
- Papakostas, K., Georgopoulou, E., and Frillingos, S. (2008) Cysteine-scanning analysis of putative helix XII in the YgfO xanthine permease: Ile-432 and Asn-430 are important. *J. Biol. Chem.* **283**, 13666–13678
- Karena, E., and Frillingos, S. (2009) Role of intramembrane polar residues in the YgfO xanthine permease: His-31 and Asn-93 are crucial for affinity and specificity, and Asp-304 and Glu-272 are irreplaceable. *J. Biol. Chem.* **284**, 24257–24268
- Georgopoulou, E., Mermelekas, G., Karena, E., and Frillingos, S. (2010) Purine substrate recognition by the nucleobase-ascorbate transporter signature motif in the YgfO xanthine permease: Asn-325 binds and Ala-323 senses substrate. *J. Biol. Chem.* **285**, 19422–19433
- Mermelekas, G., Georgopoulou, E., Kallis, A., Botou, M., Vlantos, V., and Frillingos, S. (2010) Cysteine-scanning analysis of helices TM8, TM9a, and TM9b and intervening loops in the YgfO xanthine permease: a carboxyl group is essential at Asp-276. *J. Biol. Chem.* **285**, 35011–35020
- Karena, E., and Frillingos, S. (2011) The role of transmembrane segment TM3 in the xanthine permease XanQ of *Escherichia coli*. *J. Biol. Chem.* **286**, 39595–39605
- Diallinas, G., Valdez, J., Sophianopoulou, V., Rosa, A., and Scazzocchio, C. (1998) Chimeric purine transporters of *Aspergillus nidulans* define a domain critical for function and specificity conserved in bacterial, plant, and metazoan homologues. *EMBO J.* **17**, 3827–3837
- Koukaki, M., Vlanti, A., Goudela, S., Pantazopoulou, A., Gioule, H., Tournaviti, S., and Diallinas, G. (2005) The nucleobase-ascorbate transporter (NAT) signature motif in UapA defines the function of the purine translocation pathway. *J. Mol. Biol.* **350**, 499–513
- Papageorgiou, I., Gournas, C., Vlanti, A., Amillis, S., Pantazopoulou, A., and Diallinas, G. (2008) Specific interdomain synergy in the UapA transporter determines its unique specificity for uric acid among NAT carriers. *J. Mol. Biol.* **382**, 1121–1135
- Kosti, V., Papageorgiou, I., and Diallinas, G. (2010) Dynamic elements at both cytoplasmically and extracellularly facing sides of the UapA transporter selectively control the accessibility of substrates to their translocation pathway. *J. Mol. Biol.* **397**, 1132–1143
- Amillis, S., Kosti, V., Pantazopoulou, A., Mikros, E., and Diallinas, G. (2011) Mutational analysis and modeling reveal functionally critical residues in transmembrane segments 1 and 3 of the UapA transporter. *J. Mol. Biol.* **411**, 567–580
- Andersen, P. S., Frees, D., Fast, R., and Mygind, B. (1995) Uracil uptake in *Escherichia coli* K-12: isolation of uraA mutants and cloning of the gene. *J. Bacteriol.* **177**, 2008–2013
- Loh, K. D., Gyaneshwar, P., Markenscoff Papadimitriou, E., Fong, R., Kim, K. S., Parales, R., Zhou, Z., Inwood, W., and Kustu, S. (2006) A previously undescribed pathway for pyrimidine catabolism. *Proc. Natl. Acad. Sci. U.S.A.* **103**, 5114–5119
- Kim, K. S., Pelton, J. G., Inwood, W. B., Andersen, U., Kustu, S., and Wemmer, D. E. (2010) The Rut pathway for pyrimidine degradation: novel chemistry and toxicity problems. *J. Bacteriol.* **192**, 4089–4102
- Parales, R. E., and Ingraham, J. L. (2010) The surprising Rut pathway: an unexpected way to derive nitrogen from pyrimidines. *J. Bacteriol.* **192**, 4086–4088
- Xi, H., Schneider, B. L., and Reitzer, L. (2000) Purine catabolism in *Escherichia coli* and function of xanthine dehydrogenase in purine salvage. *J. Bacteriol.* **182**, 5332–5341
- Schultz, A. C., Nygaard, P., and Saxild, H. H. (2001) Functional analysis of 14 genes that constitute the purine catabolic pathway in *Bacillus subtilis* and evidence for a novel regulon controlled by the PucR transcription activator. *J. Bacteriol.* **183**, 3293–3302
- Cecchetto, G., Amillis, S., Diallinas, G., Scazzocchio, C., and Drevet, C. (2004) The AzgA purine transporter of *Aspergillus nidulans*: characterization of a protein belonging to a new phylogenetic cluster. *J. Biol. Chem.* **279**, 3132–3141
- Mansfield, T. A., Schultes, N. P., and Mourad, G. S. (2009) AtAzg1 and AtAzg2 comprise a novel family of purine transporters in Arabidopsis. *FEBS Lett.* **583**, 481–486
- Burton, K. (1994) Adenine transport in *Escherichia coli*. *Proc. Biol. Sci.* **255**, 153–157
- Cho, B. K., Federowicz, S. A., Embree, M., Park, Y. S., Kim, D., and Palsson, B. O. (2011) The PurR regulon in *Escherichia coli* K-12 MG1655. *Nucleic Acids Res.* **39**, 6456–6464
- Inoue, H., Nojima, H., and Okayama, H. (1990) High-efficiency transformation of *Escherichia coli* with plasmids. *Gene* **96**, 23–28
- Teather, R. M., Bramhall, J., Riede, I., Wright, J. K., Fürst, M., Aichele, G., Wilhelm, U., and Overath, P. (1980) Lactose carrier protein of *Escherichia coli*: structure and expression of plasmids carrying the Y gene of the lac operon. *Eur. J. Biochem.* **108**, 223–231
- Ho, S. N., Hunt, H. D., Horton, R. M., Pullen, J. K., and Pease, L. R. (1989) Site-directed mutagenesis by overlap extension using the polymerase chain reaction. *Gene* **77**, 51–59
- Granseth, E., Daley, D. O., Rapp, M., Melén, K., and von Heijne, G. (2005) Experimentally constrained topology models for 51,208 bacterial inner membrane proteins. *J. Mol. Biol.* **352**, 489–494
- Schwede, T., Kopp, J., Guex, N., and Peitsch, M. C. (2003) SWISS-MODEL: an automated protein homology-modeling server. *Nucleic Acids*

⁵ M. Botou, K. Papakostas, and S. Frillingos, manuscript in preparation.

- Res.* **31**, 3381–3385
31. Felle, H., Porter, J. S., Slayman, C. L., and Kaback, H. R. (1980) Quantitative measurements of membrane potential in *Escherichia coli*. *Biochemistry* **19**, 3585–3590
 32. Kaback, H. R., Reeves, J. P., Short, S. A., and Lombardi, F. J. (1974) Mechanisms of active transport in isolated bacterial membrane vesicles. 18. The mechanism of action of carbonylcyanide *m*-chlorophenylhydrazone. *Arch. Biochem. Biophys.* **160**, 215–222
 33. Cusa, E., Obradors, N., Baldomà, L., Badia, J., and Aguilar, J. (1999) Genetic analysis of a chromosomal region containing genes required for assimilation of allantoin nitrogen and linked glyoxylate metabolism in *Escherichia coli*. *J. Bacteriol.* **181**, 7479–7484
 34. Ramazzina, I., Folli, C., Secchi, A., Berni, R., and Percudani, R. (2006) Completing the uric acid degradation pathway through phylogenetic comparison of whole genomes. *Nat. Chem. Biol.* **2**, 144–148
 35. Rintoul, M. R., Cusa, E., Baldomà, L., Badia, J., Reitzer, L., and Aguilar, J. (2002) Regulation of the *Escherichia coli* allantoin regulon: coordinated function of the repressor AllR and the activator AllS. *J. Mol. Biol.* **324**, 599–610
 36. Lee, Y., Lee, D. H., Kho, C. W., Lee, A. Y., Jang, M., Cho, S., Lee, C. H., Lee, J. S., Myung, P. K., Park, B. C., and Park, S. G. (2005) Transthyretin-related proteins function to facilitate the hydrolysis of 5-hydroxyisourate, the end product of the uricase reaction. *FEBS Lett.* **579**, 4769–4774
 37. Zanotti, G., Cendron, L., Ramazzina, I., Folli, C., Percudani, R., and Berni, R. (2006) Structure of zebra fish HIUase: insights into evolution of an enzyme to a hormone transporter. *J. Mol. Biol.* **363**, 1–9
 38. Henneby, S. C. (2009) Evolutionary changes to transthyretin: structure and function of a transthyretin-like ancestral protein. *FEBS J.* **276**, 5367–5379
 39. de la Riva, L., Badia, J., Aguilar, J., Bender, R. A., and Baldomà, L. (2008) The *hpx* genetic system for hypoxanthine assimilation as a nitrogen source in *Klebsiella pneumoniae*: gene organization and transcriptional regulation. *J. Bacteriol.* **190**, 7892–7903
 40. Pope, S. D., Chen, L. L., and Stewart, V. (2009) Purine utilization by *Klebsiella oxytoca* M5al: genes for ring-oxidizing and -opening enzymes. *J. Bacteriol.* **191**, 1006–1017
 41. O'Leary, S. E., Hicks, K. A., Ealick, S. E., and Begley, T. P. (2009) Biochemical characterization of the HpxO enzyme from *Klebsiella pneumoniae*, a novel FAD-dependent urate oxidase. *Biochemistry* **48**, 3033–3035
 42. Gournas, C., Oestreicher, N., Amillis, S., Diallinas, G., and Scazzocchio, C. (2011) Completing the purine utilization pathway of *Aspergillus nidulans*. *Fungal Genet. Biol.* **48**, 840–848
 43. Ormazabal, V., Zuñiga, F. A., Escobar, E., Aylwin, C., Salas-Burgos, A., Godoy, A., Reyes, A. M., Vera, J. C., and Rivas, C. I. (2010) Histidine residues in the Na⁺-coupled ascorbic acid transporter-2 (SVCT2) are central regulators of SVCT2 function, modulating pH sensitivity, transporter kinetics, Na⁺ cooperativity, conformational stability, and subcellular localization. *J. Biol. Chem.* **285**, 36471–36485
 44. Varma, S., Campbell, C. E., and Kuo, S. M. (2008) Functional role of conserved transmembrane segment 1 residues in human sodium-dependent vitamin C transporters. *Biochemistry* **47**, 2952–2960