



Published in final edited form as:

*Diabetes*. 1997 January ; 46(1): 51–56.

## Oscillations in Oxygen Consumption by Permeabilized Clonal Pancreatic $\beta$ -cells (HIT) Incubated in an Oscillatory Glycolyzing Muscle Extract: Roles of free $\text{Ca}^{2+}$ , substrates, and the ATP/ADP ratio

Vildan N. Civelek, Jude T. Deeney, Glenn E. Fusonie, Barbara E. Corkey, and Keith Tornheim

Diabetes and Metabolism Unit, Evans Department of Medicine, and the Department of Biochemistry, Boston University School of Medicine, Boston, Massachusetts, USA

### Abstract

To determine whether oscillations in glycolysis could underlie the oscillations in  $\text{O}_2$  consumption observed in intact islets, we evaluated the capacity of an islet extract to exhibit spontaneous oscillations in glycolysis. When a cell-free extract obtained from approximately 1,000 islets was supplied with glucose and glycolytic cofactors, oscillations in NADH fluorescence were obtained. After this demonstration of spontaneous oscillations in islet extracts, we bathed permeabilized clonal  $\beta$ -cells in the more plentiful spontaneously oscillating glycolytic muscle extract that generates pulses of alpha-glycerophosphate and pyruvate and induces oscillations in free  $\text{Ca}^{2+}$  and the ATP/ADP ratio. This preparation was used to investigate whether changes in  $\text{Ca}^{2+}$  and possibly alpha-glycerophosphate or pyruvate supply could underlie observed oscillations in  $\text{O}_2$  consumption and explain coordination between cytosolic and mitochondrial metabolism. We found that oscillations of  $\text{O}_2$  consumption and  $\text{Ca}^{2+}$  of a similar period were induced. Removal of medium  $\text{Ca}^{2+}$  with EGTA did not prevent the oscillations in  $\text{O}_2$  consumption nor were they greatly affected by the substantial rise in medium  $\text{Ca}^{2+}$  on treatment with thapsigargin to inhibit sequestration into the endoplasmic reticulum. The  $\text{O}_2$  oscillations were also not eliminated by the addition of relatively high concentrations of pyruvate or alpha-glycerophosphate. However, they were lost on addition of fructose-2,6-P<sub>2</sub> at concentrations that prevent oscillations of glycolysis and the ATP/ADP ratio. Addition of a high concentration of ADP increased  $\text{O}_2$  consumption and also prevented  $\text{O}_2$  oscillations. These results suggest that the changes in respiration reflected in the  $\text{O}_2$  oscillations occur in response to the oscillations in the ATP/ADP ratio or ADP concentration and that this parameter is a primary regulator of  $\text{O}_2$  consumption in the pancreatic  $\beta$ -cell.

### INTRODUCTION

Insulin secretion *in vivo*, in the perfused pancreas, and by groups of perfused islets is pulsatile (1-3). Similar oscillations in intracellular free  $\text{Ca}^{2+}$  and  $\text{O}_2$  consumption have been observed in glucose-stimulated perfused islets (1,4).  $\text{Ca}^{2+}$  oscillations have also been seen in single  $\beta$ -cells stimulated with glucose (5-7). We have suggested that these oscillatory phenomena may derive from oscillatory metabolism of glucose, such as has been

---

Address correspondence and reprint requests to Dr. K. Tornheim, Boston University School of Medicine, 80 E. Concord St., E-211, Boston, MA, 02118, United States of America. vncivelek@gmail.com.

Contributions: Vildan N. Civelek, [M.D., Endocrinologist, Ph.D.; Prof. Dr. of Biochemistry and Medicine, Project Leader], Jude T. Deeney, [Ph.D.; Assoc. Prof., Project Leader], Glenn E. Fusonie [M.D.; Senior Microsurgery Fellow, Project Leader], Barbara E. Corkey, [Ph.D.; Prof. Dr. of Biochemistry and Medicine, Project Leader, President], and Keith Tornheim, [Ph.D.; Prof. Dr. of Biochemistry and Medicine, Project Leader].

demonstrated in extracts of skeletal muscle (8-12), heart (13), and yeast (14,15), as well as in intact yeast (16-18) and ascites cells (19). The mechanism of the oscillations in skeletal muscle extracts involves a rather unusual form of enzyme regulation in which the product activates the enzyme. Autocatalytic, AMP-dependent activation of the key glycolytic enzyme phosphofructokinase (PFK) by its product, fructose-1,6-P<sub>2</sub>, has been well documented, and its role is supported by observations that the oscillations can be blocked by addition of sufficiently high concentrations of fructose-2,6-P<sub>2</sub> or glucose-1,6-P<sub>2</sub> (10,11), activators that are natural analogs of fructose-1,6-P<sub>2</sub> and compete for the same regulatory side on PFK. There are three mammalian isoforms of PFK with differing tissue distribution (20,21). Adult muscle has only the M-type subunit, and liver has mainly the L-type subunit. Brain has C-type as well as M- and L-type subunits. Skeletal muscle PFK is strongly activated by micromolar levels of fructose-1,6-P<sub>2</sub> (22,23), whereas there is little, if any, activation of liver or platelet PFK (mainly C- and L-type subunits) in the presence of near physiological concentrations of ATP (several millimolar) and AMP and fructose-6-phosphate (micromolar) (24). Our recent Western blots of pancreatic islets and clonal  $\beta$ -cells (INS-1) showed the presence of both M- and C-type isoforms, with perhaps a smaller amount of L-, and kinetic studies demonstrated that the activity under near-physiological conditions was dominated by the M-type fructose-1,6-P<sub>2</sub> activatable form (25). Thus, the autocatalytic activation of PFK by micromolar levels of fructose-1,6-P<sub>2</sub>, which is the basis of the glycolytic oscillations in the muscle extract system, should be expected to occur in islets. Because of the limited amount of islet material obtainable, oscillations were examined by NADH fluorescence, a sensitive technique frequently used to monitor glycolytic oscillations in other cell types. We previously used a model experimental system consisting of permeabilized clonal pancreatic  $\beta$ -cells incubated in a synthetic cytosol containing oscillatory glycolyzing muscle extract; under these conditions, oscillations in free Ca<sup>2+</sup> were generated (26). Because such permeabilized cells maintain mitochondrial function, we examined this model system for oscillations in O<sub>2</sub> consumption. The behavior of the oscillatory muscle extract has been well characterized; it also produces pulses of the mitochondrial substrates pyruvate and alpha-glycerophosphate (8,9), as well as oscillations in the ATP/ADP ratio or ADP concentration (9-12). Furthermore, several of the mitochondrial dehydrogenases are activated by Ca<sup>2+</sup> (27-32). We therefore tested which of these factors was involved in generating the observed oscillations in O<sub>2</sub> consumption. Our data demonstrate that an islet extract exhibits spontaneous metabolic oscillations like the muscle extract, and using the muscle extract, oscillations in respiration by the permeabilized cells are indeed driven by the glycolytic oscillations. Removal or elevation of Ca<sup>2+</sup> or the addition of high saturating concentrations of mitochondrial substrate did not dampen the O<sub>2</sub> oscillations. In contrast, inhibition of glycolytic oscillations with fructose-2,6-P<sub>2</sub> or stimulation of maximal rates of respiration with ADP eliminated oscillations in O<sub>2</sub> consumption, suggesting that the ATP/ADP ratio or ADP concentration is probably the dominant regulator of these oscillations.

## RESEARCH DESIGN AND METHODS

### Isolation, extraction, and assessment of islets

Islets were prepared by collagenase digestion as described previously (1). Islet extracts were prepared by placing islets in a microfuge tube, washing with Hanks' buffer, removing as much medium as possible, and freezing until the day of use. Islets suspended in 0.2 ml/1,000 islets of 15 mmol/l potassium phosphate, pH 7.0, 100 mmol/l KCl, and 1 mmol/l dithiothreitol were sonicated for 10 s in ice and centrifuged for 5 min in a microfuge.

Oscillations were induced in a reaction mixture containing ATP, MgCl<sub>2</sub>, NAD, potassium phosphate, KCl, and HEPES buffer and a 50% volume of islet extract in a total volume of 70

$\mu$ l. Hexokinase and apyrase were added as needed to achieve the desired rates of ATP usage and glucose phosphorylation.

Glycolytic oscillations were monitored by pyridine nucleotide fluorescence (9,15,16) using a Farrand fluorometer.

### Growth and incubation of HIT cells

Clonal pancreatic  $\beta$ -cells (HIT-T 15) were cultured in RPMI 1640 medium supplemented with 50 U/ml penicillin, 50  $\mu$ g/ml streptomycin, and 10% fetal calf serum (33). Cells were used between passages 64 and 80, harvested with phosphate-buffered saline containing 0.02% EDTA, and washed in Hanks'  $\text{Ca}^{2+}$ - $\text{Mg}^{2+}$ -free buffer, pH 7.4. Cells were incubated at 30°C in a buffer containing an oscillating glycolytic muscle extract (high speed supernatant fraction, gel filtered to remove endogenous metabolites) (12) in a total volume of 1.7 ml. Although we provide evidence of metabolic oscillations in islet extracts, the muscle extract has two important advantages as a reagent at this point: it is easily prepared in sufficient quantity, and it has been well characterized (8-12) and already used successfully with permeabilized clonal  $\beta$ -cells (26). Saponin (60-80  $\mu$ g/ml) was added to selectively permeabilize the plasma membrane of cells in suspension. The precise concentration used was the minimum amount required to permeabilize the plasma membrane within 5 min without altering the  $\text{Ca}^{2+}$  set point (33).

### Oxygen consumption

$\text{O}_2$  consumption was measured using a Clark type electrode with an amplifier designed and built by the Bio-Instrumentation Group of the University of Pennsylvania in a stirred water-jacketed open chamber. An open chamber was used in these studies, since otherwise the total  $\text{O}_2$  of the chamber was consumed too quickly to allow observation of multiple oscillations. Air was blown over the top surface of the solution at a rate sufficient to balance average  $\text{O}_2$  use such that a concentration of  $\text{O}_2$  in the chamber of  $\sim 150 \mu\text{mol/l}$  was maintained.

Thus, a rise in the  $\text{O}_2$  traces shown indicates a decreased rate of  $\text{O}_2$  consumption, and a fall in the trace indicates an increased rate of  $\text{O}_2$  consumption compared with the average.

### $\text{Ca}^{2+}$ measurements

$\text{Ca}^{2+}$  was measured in permeabilized cells in the presence of 2  $\mu\text{mol/l}$  free fura 2 added to the buffer.  $\text{Ca}^{2+}$  values were calculated from the fura 2 signals at excitation wavelengths of 340 and 380 nm and emission at 510 nm (34) using a time-sharing fluorometer designed and built by the Bio-Instrumentation Group of the University of Pennsylvania (35).

### Metabolite assays

Samples (0.1 ml) of the reaction mixture were deproteinized and assayed by enzymatic methods, as described previously (8,9), using a Hewlett Packard model 8450 spectrophotometer system set to read  $A_{335-345}$  minus  $A_{390-400}$ . A new specific method was used to assay ADP separate from GDP (36).

### Materials

RPMI 1640 media, penicillin, and streptomycin were from Gibco. Fetal calf serum was obtained from Hyclone Laboratories. Fura 2 (free acid) was obtained from Molecular Probes (Eugene, OR). Thapsigargin was from Calbiochem. All other biochemicals were from Sigma or Boehringer Mannheim.

## RESULTS

Figure 1 illustrates spontaneous oscillations in NADH fluorescence in islet extracts supplied with ATP, NAD, hexokinase, and glucose.

The three panels show data obtained from two different preparations of ~1,000 islets. Figures 1A and B were from the same preparation and Fig. 1C from a separate preparation. Figure 1A shows repetitive oscillations superimposed on an upward drift that may be due to mitochondrial contamination. To show the oscillations more clearly, the raw data in Fig. 1A were fitted with a polynomial function (narrow line), which was then subtracted out in Fig. 1B.

It should be noted that glycolytic oscillations in extracts show a considerable range of periods and amplitudes, depending on the particular extract and the experimental conditions, especially factors affecting phosphofructokinase activity (9,11,12,37). Furthermore, there may be a considerable lag after the reaction is started before the oscillations begin, most likely to allow suitable shifts in metabolite concentrations to bring them into the oscillatory region for phosphofructokinase. Thus, the data in Fig. 1 are consistent with our prior observations of oscillations in skeletal muscle extracts.

Oscillations in NADH fluorescence have been associated with every oscillating glycolytic system so far studied and have in fact been used for monitoring the yeast and heart extract systems. Oscillations in NADH fluorescence have also been reported in single  $\beta$ -cells on stimulation with glucose (5). Importantly, the initial rise in NADH preceded that of  $\text{Ca}^{2+}$ , consistent with  $\text{Ca}^{2+}$  changes being secondary to metabolic coupling factors. The data indicate that a cell-free islet extract, like the cell-free skeletal muscle extract, is capable of generating spontaneous oscillations of glycolysis. However, the large amount of material required made it impractical to use such islet extracts in the following studies of oxygen consumption by permeabilized clonal  $\beta$ -cells (HIT). To obviate this constraint, experiments were performed using skeletal muscle extract.

As illustrated in Figure 2, when the permeabilized HIT cells were incubated in a spontaneously oscillating glycolytic extract and oscillations in  $\text{Ca}^{2+}$  were observed (bottom trace), then  $\text{O}_2$  consumption also oscillated (top trace) and with the same period.

When the cells are permeabilized,  $\text{Ca}^{2+}$  is taken up by the endoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase and the free concentration in the medium declines and approaches a set point, which is in part determined by the ATP/ADP ratio (33,34); in this case, because of the oscillations in glycolysis and the ATP/ADP ratio,  $\text{Ca}^{2+}$  does not remain constant but oscillates. We previously showed a strong correlation between  $\text{Ca}^{2+}$  and the ATP/ADP ratio in permeabilized clonal  $\beta$ -cells, a high ATP/ADP ratio promoting  $\text{Ca}^{2+}$  sequestration (26,34). The inverse correlation of  $\text{Ca}^{2+}$  and oxygen changes during the oscillations in Fig. 2 suggests that oxygen consumption may be enhanced by low ATP/ADP ratios and relatively suppressed at high ATP/ADP ratios. This is consistent with our recent results of effects of different fixed ATP/ADP ratios on oxygen consumption in permeabilized HIT cells (38). The correlation of oxygen consumption and ATP/ADP ratio could not be determined directly in the oscillating system because sampling for measurement of the nucleotides would disrupt the oxygen measurements. As noted above, and illustrated in Fig. 2, there may be a considerable lag before oscillations begin, and furthermore, it takes some time after permeabilization for  $\text{Ca}^{2+}$  and oxygen to reach near plateaus around which they oscillate. Thus, the subsequent oxygen traces shown are portions from the plateau phases.

Removal of  $\text{Ca}^{2+}$  with EGTA (Fig. 3) had little effect on the oscillations in Oxygen use (amplitude,  $118 \pm 51\%$ ; period,  $85 \pm 23\%$  of same-day control; mean  $\pm$  SD,  $n = 5$ ), even

though EGTA lowers the free  $\text{Ca}^{2+}$  to  $<10$  nmol/l and eliminates  $\text{Ca}^{2+}$  oscillations. Under control conditions, free  $\text{Ca}^{2+}$  oscillates in the range of 100-200 nmol/l (26,33); however, it is likely that higher levels of  $\text{Ca}^{2+}$  may be reached in response to stimulatory concentrations of glucose in intact cells.

To assess the influence of higher  $\text{Ca}^{2+}$  concentrations on the oscillations of  $\text{O}_2$  consumption in our permeabilized cell system, we inhibited the endoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase with thapsigargin (39). Under these conditions the  $\text{Ca}^{2+}$  set point is presumably maintained by the mitochondria, rather than the endoplasmic reticulum, but at a much higher level of 500-1,000 nmol/l (40). We previously showed that thapsigargin greatly raises the  $\text{Ca}^{2+}$  concentration in permeabilized HIT cells under steady-state conditions (33). Like the removal of  $\text{Ca}^{2+}$ , this treatment had little effect on the oscillations in  $\text{O}_2$  consumption (Fig. 3) (amplitude,  $92 \pm 27\%$ ; period,  $93 \pm 17\%$  of control,  $n = 3$ ). These findings indicate that variations in free  $\text{Ca}^{2+}$  do not generally control the rate of respiration and are not responsible for the oscillations in  $\text{O}_2$  consumption. This conclusion is supported by the relatively small effects of  $\text{Ca}^{2+}$  on oxygen consumption by permeabilized HIT cells under steady-state conditions (38). Furthermore, when a suspension of intact HIT cells is stimulated with glucose, the increase in oxygen consumption precedes and is not further augmented by the increase in intracellular free  $\text{Ca}^{2+}$ (41).

In addition to oscillations in  $\text{Ca}^{2+}$  and the ATP/ADP ratio, the extract is characterized by pulsatile delivery of the mitochondrial substrates alpha-glycerophosphate and pyruvate. These can be metabolized by the permeabilized cells, and therefore their levels also oscillated (Fig. 4). alpha-glycerophosphate increased when its precursor, dihydroxyacetone-phosphate, was high. Pyruvate decreased as dihydroxyacetone-phosphate rose, perhaps in part because of the use of pyruvate to reoxidize the pulse of NADH that occurs when phosphofructokinase is activated and flux through glyceraldehyde-3-phosphate dehydrogenase suddenly increases (8,9). The repeated activation of phosphofructokinase that generates the metabolite oscillations is indicated by the decreases in glucose-6-phosphate (which is in equilibrium with fructose-6-phosphate) and corresponding increases in dihydroxyacetone-phosphate (which is in equilibrium with fructose-1,6- $\text{P}_2$ ).

To test the possibility that the fluctuations in mitochondrial fuels caused the observed oscillations in  $\text{O}_2$  consumption, we added high concentrations of pyruvate or alpha-glycerophosphate (10 times the levels in Fig. 4) to markedly dampen the influence of their pulsatile production from glucose. These additions did not inhibit the  $\text{O}_2$  oscillations (Fig. 5) (pyruvate: amplitude,  $103 \pm 30\%$ ; period,  $108 \pm 21\%$  of control,  $n = 5$ ; alpha-glycerophosphate: amplitude,  $101 \pm 21\%$ ; period,  $99 \pm 27\%$  of control,  $n = 4$ ). Thus, mitochondrial fuel supply does not appear to be the dominant factor regulating oscillations in our system under these conditions. Clearly although the rate of respiration can be affected by substrate supply, the oscillations here are not.

To demonstrate that the  $\text{O}_2$  oscillations could be inhibited in a rather simple manner that does not involve poisoning or greatly altering the metabolic state of the mitochondria, we added fructose-2,6- $\text{P}_2$ . High concentrations of fructose-2,6- $\text{P}_2$  prevent glycolytic oscillations by competing with fructose-1,6- $\text{P}_2$  for an activator site on phosphofructokinase, thereby preventing the autocatalytic activation of the enzyme; this leads to continuous flow through the glycolytic pathway rather than the normal pulsatile flux (10). Similar results are seen with another analog, glucose-1,6- $\text{P}_2$  (11). Importantly the associated oscillations in the ATP/ADP ratio and ADP concentration are eliminated, being replaced with steady-state values of intermediate magnitude (10,11). Addition of fructose-2,6- $\text{P}_2$  in the permeabilized cell system also blocked oscillations in  $\text{O}_2$  consumption (Fig. 6A). This experiment confirms

that the O<sub>2</sub> oscillations are metabolically generated and are not a consequence of artifacts such as pulsating air flow or electrical oscillations in the measuring equipment.

A high concentration of ADP (1 mmol/l) maximally stimulates mitochondrial respiration and should make it independent of the micromolar fluctuations in ADP caused by glycolytic oscillations. Indeed, addition of ADP increased respiration and blocked the oscillations (Fig. 6B). It should be noted that because these experiments were performed in an open system, the resupply of O<sub>2</sub> from the air prevented cells from becoming anoxic, even when respiring in the presence of large amounts of ADP. These data also indicate that ADP limits respiration under the conditions of these experiments and thus may be the dominant factor linking oscillatory glycolysis to oscillatory O<sub>2</sub> consumption.

## DISCUSSION

In the studies presented here, the oscillations in O<sub>2</sub> consumption by the permeabilized HIT cells were induced by the oscillatory glycolyzing extract in which they were placed. However, intact  $\beta$ -cells or islets also exhibit oscillations in O<sub>2</sub> consumption, as well as oscillations in membrane potential, intracellular free Ca<sup>2+</sup>, and insulin secretion. The pulsatile character of insulin secretion is seen in vivo in normal individuals and is lost in some type II diabetic patients and their near relatives (42), a fact that suggests its physiological importance.

We have proposed that the basis of all these oscillatory phenomena is the same as in the model system used here, namely oscillatory metabolism of glucose. We suggest that it is the changes in ADP associated with glycolytic oscillations that most likely cause repeated opening and closing of ATP-sensitive K channels (43) and thus the changes in membrane potential and Ca<sup>2+</sup> fluxes. The Ca<sup>2+</sup> rises in turn are important for the secretory process and also for activating mitochondrial dehydrogenases, shifting the metabolic stance to favor the production of additional signaling compounds, such as malonyl CoA and long-chain acyl CoA (44). However, it is also ADP that appears to be the major controller of the oscillations in O<sub>2</sub> consumption, rather than the mitochondrial substrates or Ca<sup>2+</sup> activation of the dehydrogenases. The overall increase in O<sub>2</sub> consumption in glucose-stimulated islets presumably reflects the increased ATP utilization by the exocytotic secretory process and perhaps ion pumping, as well as the increase in substrate supply. The apparent paradox is that enhanced O<sub>2</sub> consumption would require a rise in the ADP concentration, whereas closure of ATP-sensitive K channels would require a decrease in ADP. The resolution of this paradox may well lie in the oscillatory behavior of glycolysis and the energy state, whereby ADP concentrations alternate between high and low values. The observed oscillations in membrane potential and O<sub>2</sub> consumption in intact cells are consistent with such alternating control. Our recent demonstration of oscillations in the ATP/ADP ratio and glucose-6-phosphate in a suspension of  $\beta$ -cells, correlating with oscillations in intracellular free Ca<sup>2+</sup>, further supports this proposal (45). The present studies with permeabilized cells obviously do not test whether the glucose-induced oscillations in Ca<sup>2+</sup> and insulin release in intact islets are indeed due to glycolytic oscillations; they are useful for showing the responses of mitochondrial metabolism to putative coupling factors that may link O<sub>2</sub> consumption to these other oscillating parameters.

## Acknowledgments

This work was supported by a grant from the Juvenile Diabetes Foundation (to B.E.C.), U.S. Public Health Service Grants DK35914 (to B.E.C.) and DK31559 (to K.T.), and grants from the Upjohn Company (to B.E.C.).

## ABBREVIATIONS

<b>PFK</b>	phosphofructokinase
<b>ADP</b>	Adenosine Diphosphate
<b>Oxygen</b>	O <sub>2</sub>
<b>a-glycerophosphate</b>	alpha-glycerophosphate


## REFERENCES

1. Longo EA, Tornheim K, Deeney JT, Varnum BA, Tillotson D, Prentki M, Corkey BE. Oscillations in cytosolic free Ca<sup>2+</sup>, oxygen consumption, and insulin secretion in glucose-stimulated rat pancreatic islets. *J Biol Chem.* 1991; 266:9314–9319. [PubMed: 1902835]
2. Weigle DS. Pulsatile secretion of fuel-regulatory hormones. *Diabetes.* 1987; 36:764–775. [PubMed: 3552804]
3. Lefèbvre PJ, Paolisso G, Scheen AT, Henquin JC. Pulsatility of insulin and glucagon release: physiological significance and pharmacological implications. *Diabetologia.* 1987; 30:443–452. [PubMed: 3311858]
4. Valdeolmillos M, Santos RM, Contreras D, Soria B, Rosario LM. Glucose-induced oscillations of intracellular Ca<sup>2+</sup> concentration resembling bursting electrical activity in single mouse islets of Langerhans. *FEBS Lett.* 1989; 259:19–23. [PubMed: 2689228]
5. Pralong WF, Bartley C, Wollheim CB. Single islet β-cell stimulation by nutrients: relationship between pyridine nucleotides, cytosolic Ca<sup>2+</sup> and secretion. *EMBO J.* 1990; 9:53–60. [PubMed: 2403930]
6. Grapengiesser E, Gylfe E, Hellman B. Glucose-induced oscillations of cytoplasmic Ca<sup>2+</sup> in the pancreatic β-cell. *Biochem Biophys Res Commun.* 1988; 151:1299–1304. [PubMed: 3281672]
7. Grapengiesser E, Gylfe E, Hellman B. Three types of cytoplasmic Ca<sup>2+</sup> oscillations in stimulated pancreatic β-cells. *Arch Biochem Biophys.* 1989; 268:404–407. [PubMed: 2536264]
8. Tornheim K, Lowenstein JM. The purine nucleotide cycle IV: interaction with oscillations of the glycolytic pathway in muscle extracts. *J Biol Chem.* 1974; 249:3241–3247. [PubMed: 4275340]
9. Tornheim K, Lowenstein JM. The purine nucleotide cycle: control of phosphofructokinase and glycolytic oscillations in muscle extracts. *J Biol Chem.* 1975; 250:6304–6314. [PubMed: 169235]
10. Tornheim K. Fructose 2,6-bisphosphate and glycolytic oscillations in skeletal muscle extracts. *J Biol Chem.* 1988; 263:2619–2624. [PubMed: 2963814]
11. Andrés V, Schultz V, Tornheim K. Oscillatory synthesis of glucose 1,6-bisphosphate and frequency modulation of glycolytic oscillations in skeletal muscle extracts. *J Biol Chem.* 1990; 265:21441–21447. [PubMed: 2254306]
12. Tornheim K, Andrés V, Schultz V. Modulation by citrate of glycolytic oscillations in skeletal muscle extracts. *J Biol Chem.* 1991; 266:15675–15678. [PubMed: 1831450]
13. Frenkel R. Control of reduced diphosphopyridine nucleotide oscillations in beef heart, extracts. II. Oscillations of glycolytic intermediates and adenine nucleotides. *Arch Biochem, Biophys.* 1968; 125:157–165. [PubMed: 4296954]
14. Das J, Busse HG. Analysis of the dynamics of relaxation type oscillation in glycolysis of yeast extracts. *Biophys J.* 1991; 60:369–379. [PubMed: 1832975]
15. Chance B, Schoener B, Elsaesser S. Metabolic control phenomena involved in damped sinusoidal oscillations of reduced diphosphopyridine nucleotide in a cell-free extract of *Saccharomyces carlsbergensis*. *J Biol Chem.* 1965; 240:3170–3181. [PubMed: 14342349]
16. Hess B, Boiteux A. Oscillatory phenomena in biochemistry. *Anna Rev Biochem.* 1971; 40:237–258. [PubMed: 4330578]
17. Hess B, Boiteux A, Kruger J. Cooperation of glycolytic enzymes. *Adv Enz Regul.* 1969; 7:149–167.
18. Betz A, Chance B. Phase relationship of glycolytic intermediates in yeast cells with oscillatory metabolic control. *Arch Biochem Biophys.* 1965; 109:585–594. [PubMed: 14320501]

19. Ibsen KH, Schiller KW. Oscillations of nucleotides and glycolytic intermediates in aerobic suspensions of Erlich ascites tumor cells. *Biochim Biophys Acta*. 1967; 131:405–407. [PubMed: 4292845]
20. Foe LG, Kemp RG. Isolation and characterization of phosphofructokinase C from rabbit brain. *J Biol Chem*. 1985; 260:726–730. [PubMed: 3155729]
21. Dunaway GA, Kasten TP. Nature of the subunits of the 6-phosphofructo-1-kinase isoenzymes from rat tissues. *Biochem J*. 1987; 242:667–671. [PubMed: 2954542]
22. Tornheim K, Lowenstein JM. Control of phosphofructokinase from rat skeletal muscle: effects of fructose diphosphate, AMP, ATP and citrate. *J Biol Chem*. 1976; 251:7322–7328. [PubMed: 12161]
23. Tornheim K. Activation of muscle phosphofructokinase by fructose 2,6-bisphosphate and fructose 1,6-bisphosphate is differently affected by other regulatory metabolites. *J Biol Chem*. 1985; 260:7985–7989. [PubMed: 3159721]
24. Underwood AH, Newsholme EA. Properties of phosphofructokinase from rat liver and their relationship to the control of glycolysis and gluconeogenesis. *Biochem, J*. 1965; 95:868–875. [PubMed: 14342527]
25. Yaney GC, Schultz V, Cunningham BA, Dunaway GA, Corkey BE, Tornheim K. Phosphofructokinase isozymes in pancreatic islets and clonal  $\beta$ -cells (INS-1). *Diabetes*. 1995; 44:1285–1289. [PubMed: 7589825]
26. Corkey BE, Tornheim K, Deeney JT, Glennon MC, Parker JC, Matschinsky FM, Ruderman NB, Prentki M. Linked oscillations of free  $\text{Ca}^{2+}$  and the ATP/ADP ratio in permeabilized RINm5F insulinoma cells supplemented with a glycolyzing cell-free muscle extract. *J Biol Chem*. 1988; 263:4254–4258. [PubMed: 2831225]
27. McCormack JG, Halestrap AP, Denton RM. Role of calcium ions in regulation of mammalian intramitochondrial metabolism. *Physiol Rev*. 1990; 70:391–425. [PubMed: 2157230]
28. McCormack JG, Longo EA, Corkey BE. Glucose-induced activation of pyruvate dehydrogenase in isolated rat pancreatic islets. *Biochem J*. 1990; 267:527–530. [PubMed: 2185742]
29. Hansford RG. Dehydrogenase activation by  $\text{Ca}^{2+}$  in cells and tissues. *J Bioenerg Biomembr*. 1991; 23:823–854. [PubMed: 1778993]
30. MacDonald MJ. High content of mitochondrial glycerol-3-phosphate dehydrogenase in pancreatic islets and its inhibition by diazoxide. *J Biol Chem*. 1981; 256:8287–8290. [PubMed: 6790537]
31. MacDonald MJ. Calcium activation of pancreatic islet mitochondrial glycerol phosphate dehydrogenase. *Horm Metab Res*. 1982; 14:678–679. [PubMed: 6818120]
32. Rutter GA, Pralong WF, Wollheim CB. Regulation of mitochondrial glycerol phosphate dehydrogenase by  $\text{Ca}^{2+}$  within electroporated insulin-secreting cells (INS-1). *Biochim Biophys Acta*. 1992; 1175:107–113. [PubMed: 1482691]
33. Deeney JT, Tornheim K, Korchak HM, Prentki M, Corkey BE. Acyl-CoA esters modulate intracellular  $\text{Ca}^{2+}$  handling by permeabilized clonal pancreatic  $\beta$ -cells. *J Biol Chem*. 1992; 267:19840–19845. [PubMed: 1400300]
34. Corkey BE, Deeney JT, Glennon MC, Matschinsky FM, Prentki M. Regulation of steady-state free  $\text{Ca}^{2+}$  levels by the ATP/ADP ratio and orthophosphate in permeabilized RINm5F insulinoma cells. *J Biol Chem*. 1988; 263:4247–4253. [PubMed: 2831224]
35. Chance B, Legailais V, Sorge J, Graham N. A versatile time-sharing multichannel spectrophotometer. *Anal Biochem*. 1975; 66:498–504. [PubMed: 166573]
36. Tornheim K, Schultz V. Specific enzymatic spectrophotometry assay of adenosine 5'-diphosphate. *Anal Biochem*. 1993; 211:329–330. [PubMed: 8317712]
37. Tornheim K, Lowenstein JM. The purine nucleotide cycle III: oscillations in metabolite concentrations during the operation of the cycle in muscle extracts. *J Biol Chem*. 1973; 248:2670–2677. [PubMed: 4348916]
38. Civelek VN, Deeney JT, Shalosky NJ, Tornheim K, Hansford RG, Prentki M, Corkey BE. Regulation of pancreatic  $\beta$ -cell mitochondrial metabolism: influence of  $\text{Ca}^{2+}$ , substrate and ADP on oxygen consumption by permeabilized clonal  $\beta$ -cells (HIT). *Biochem J*. 1996; 318:615–621. [PubMed: 8809055]




39. Thastrup O, Cullen PJ, Drobak BK, Hanley MR, Dawson AP. Thapsigargin, a tumor promoter, discharges intracellular  $\text{Ca}^{2+}$  stores by specific inhibition of the endoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase. *Proc Natl Acad Sci USA*. 1990; 87:2466–2470. [PubMed: 2138778]
40. Prentki M, Matschinsky FM.  $\text{Ca}^{2+}$ , cAMP, and phospholipid-derived messengers in coupling mechanisms of insulin secretion. *Physiol Rev*. 1987; 67:1185–1248. [PubMed: 2825225]
41. Civelek VN, Deeney JT, Kubik K, Schultz V, Tornheim K, Corkey BE. Temporal sequence of metabolic and ionic events in glucose-stimulated clonal pancreatic  $\beta$ -cells (HIT). *Biochem J*. 1996; 315:1015–1019. [PubMed: 8645138]
42. O’Rahilly S, Turner RC, Matthews DR. Impaired pulsatile secretion of insulin in relatives of patients with non-insulin-dependent diabetes. *N Engl J Med*. 1988; 318:1225–1230. [PubMed: 3283553]
43. Hopkins WF, Fotherazi S, Peter-Riesch B, Corkey BE, Cook DL. Two sites for adenine-nucleotide regulation of ATP-sensitive potassium channels in mouse pancreatic  $\beta$ -cells and HIT cells. *J Membr Biol*. 1992; 129:287–295. [PubMed: 1433280]
44. Prentki M, Corkey BE. Are the  $\beta$ -cell signaling molecules malonyl-CoA and cytosolic long-chain acyl-CoA implicated in multiple tissue defects of obesity and NIDDM? *Diabetes*. 1996; 45:273–283. [PubMed: 8593930]
45. Nilsson T, Schultz V, Berggren P-O, Corkey BE, Tornheim K. Temporal patterns of changes in ATP/ADP ratio, glucose 6-phosphate and cytoplasmic free  $\text{Ca}^{2+}$  in glucose-stimulated pancreatic  $\beta$ -cells. *Biochem J*. 1996; 314:91–94. [PubMed: 8660314]




**Figure. 1. Oscillations in NADH fluorescence in a cell-free supernatant prepared from isolated rat islets**

A and B are traces obtained from the same extract, and C represents a separate extract. The raw data in A were fitted with a 6th order polynomial (narrow line), which was subtracted out in B to show the repetitive oscillations more clearly. Data are arbitrary fluorescence units. The reaction mixture contained 1 mmol/l ATP, 1 mmol/l  $MgCl_2$ , 20 mmol/l HEPES, pH 7.1, 100 mmol/l KCl, 7.5 mmol/l potassium phosphate, 10 mmol/l glucose, 0.5 mmol/l NAD, 0.06 U/ml crystalline yeast hexokinase (gel filtered in 20 mmol/l HEPES), 0.04 U/ml apyrase (an ATPase), and islet extract equivalent to 50% of the volume. In C, the HEPES buffer was pH 7.2. Reactions were started by adding hexokinase, apyrase, and islet extract in rapid succession.




**Figure 2. Oscillations in free  $\text{Ca}^{2+}$  and Oxygen consumption in permeabilized HIT cells induced by an oscillating glycolytic muscle extract**

The reaction mixture contained 2  $\mu\text{mol/l}$  fura 2 (free acid), 2 mmol/l sodium ATP, 3 mmol/l  $\text{MgCl}_2$ , 20 mmol/l HEPES adjusted to pH 7.1 with KOH, 97 mmol/l KCl, 5 mmol/l  $\text{KHCO}_3$ , 6.5 mmol/l potassium phosphate, 0.3 mmol/l GTP, 4 mmol/l sodium aspartate, 10 mmol/l glucose, 30  $\mu\text{mol/l}$  NAD, 0.2 U/ml crystalline yeast hexokinase (gel filtered in 20 mmol/l HEPES), rat muscle extract equivalent to 1 mg of protein per milliliter, ~80 mg/ml saponin, and HIT cells equivalent to ~1 mg protein per milliliter. The reaction was started by addition of muscle extract, saponin, and hexokinase in rapid succession. Free  $\text{Ca}^{2+}$  was measured using the fluorescent signals of fura 2 and  $\text{O}_2$  consumption using an  $\text{O}_2$  electrode. Note that air was blown over the surface of the solution to balance the average rate of  $\text{O}_2$  consumption; thus, a rise in the trace indicates a decreased rate of  $\text{O}_2$  consumption, and a fall in the trace indicates an increased rate of  $\text{O}_2$  consumption, compared with the average. These traces are representative of experiments that were repeated at least three times.




**Figure 3. Effect of different concentrations of  $\text{Ca}^{2+}$  on oscillations in  $\text{O}_2$  consumption**  
Conditions were similar to those described for Fig. 2, except that fura 2 was omitted and EGTA (2 mmol/l) or thapsigargin (3  $\mu\text{mol/l}$ ) was added as indicated. These traces are representative of experiments that were repeated at least three times.




**Figure 4. Oscillations in alpha-glycerophosphate (a-GP), pyruvate, dihydroxyacetone-phosphate (DHAP), glucose-6-phosphate (G6P), ATP, and ADP**

Conditions were similar to those described for Fig. 2. The development of oscillations was monitored by following changes in  $Ca^{2+}$  by fura 2 fluorescence. In one experiment, after several oscillations in  $Ca^{2+}$  were seen (A, trace a), samples were taken in interval S at the times indicated by the experimental points in B and C and deproteinized and assayed for the metabolites. The sampling disrupted the fluorescence trace. Trace b shows the  $Ca^{2+}$  oscillations in a similar reaction mixture run previously and not sampled for metabolites. The  $Ca^{2+}$  scale ranges from ~ 100-1,000 nmol/l, but the traces have been offset.



**Figure 5. Effect of high concentrations of alpha-glycerophosphate or pyruvate on oscillations in O<sub>2</sub> consumption**

Conditions were similar to those described for Fig. 2, except that fura 2 was omitted and 5 mmol/l alpha-glycerophosphate or 1 mmol/l pyruvate was added as indicated. These traces are representative of experiments that were repeated at least three times.



**Figure 6. Effect of fructose-2,6-P<sub>2</sub> (A) or ADP (B) on oscillations in O<sub>2</sub> consumption**  
 Conditions were similar to those described for Fig. 2, except that fura 2 was omitted and 100 μmol/l fructose-2,6-P<sub>2</sub> (A) or 1 mmol/l ADP (B) was added at the time indicated by the arrow. These traces are representative of experiments that were repeated at least three times.