

Haplotypes in candidate genes related to nitric oxide pathway and vascular permeability associated with migraine and aura

Flavia Magazoni Gonçalves · Marcelo Rizzatti Luizon ·
Jose G. Speciali

Received: 5 March 2012 / Accepted: 12 March 2012 / Published online: 25 March 2012
© The Author(s) 2012. This article is published with open access at Springerlink.com

Dear Editor,

In a recent article in *Journal of Headache and Pain*, Markus Schürks has reviewed current findings on migraine genetics identified by both candidate gene approaches and genome-wide association studies [1]. We would like to contribute to some candidate gene studies focused on endothelial function possibly involved in migraine pathophysiology, which reported associations of haplotypes with migraine with aura (MA) or with aura in migraine patients [2–4]. This common condition has gained much attention because it is now recognized as an established risk factor for cardiovascular disease and ischemic stroke [5].

We have studied three clinically relevant SNPs in the vascular endothelial growth factor (*VEGF*) promoter region; C⁻²⁵⁷⁸A (rs699947), G⁻¹¹⁵⁴A (rs1570360), and G⁻⁶³⁴C (rs2010963) and found that haplotype “AGC” was more frequent in MA than controls ($P = 0.0023$) [2]. We have also examined three clinically relevant endothelial nitric oxide synthase (*eNOS*) polymorphisms; T⁻⁷⁸⁶C (rs2070744), an Intron 4 VNTR and Glu298Asp (rs1799983), and two tag-SNPs rs3918226 and rs743506. Our findings suggest that

haplotypes “CCaGluG” and “CCbGluG” are associated with increased susceptibility to the presence of aura in patients with migraine (both $P < 0.001$) [3]. Finally, we have also examined two functionally relevant polymorphisms of inducible nitric oxide synthase (*iNOS*); C⁻¹⁰²⁶A (rs2779249) and G2087A (rs2297518). The haplotype “AA” was more commonly found in MA than in patients without aura ($P = 0.0027$) [4].

Gene–gene interaction studies among candidate genes are further required in order to better elucidate the genetic basis of migraine, as previously pointed out elsewhere [6]. Therefore, we are currently performing interaction analysis among the polymorphisms previously reported [2–4]. However, our findings on *eNOS*, *iNOS* and *VEGF* haplotypes must be replicated in populations with different genetic backgrounds, which will further validate the role of these candidate genes associated with migraine and aura and may provide clinically relevant information to migraine susceptibility.

Conflict of interest None.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

F. M. Gonçalves (✉)

Department of Pharmacology, Faculty of Medical Sciences,
State University of Campinas, Rua Tessália Vieira de Camargo,
126, Campinas, SP CEP: 13083-887, Brazil
e-mail: flaviamagazoni@yahoo.com.br; luizonmr@usp.br

M. R. Luizon

Department of Pharmacology, Faculty of Medicine of Ribeirão Preto, University of São Paulo, Ribeirão Preto, SP, Brazil

J. G. Speciali

Division of Neurology, Department of Neuroscience,
Faculty of Medicine of Ribeirão Preto,
University of São Paulo, Ribeirão Preto, SP, Brazil

References

- Schurks M (2012) Genetics of migraine in the age of genome-wide association studies. *J Headache Pain* 13(1):1–9
- Goncalves FM, Martins-Oliveira A, Speciali JG, Izidoro-Toledo TC, Luizon MR, Dach F, Tanus-Santos JE (2010) Vascular endothelial growth factor genetic polymorphisms and haplotypes in women with migraine. *DNA Cell Biol* 29(7):357–362

3. Goncalves FM, Martins-Oliveira A, Speciali JG, Luizon MR, Izidoro-Toledo TC, Silva PS, Dach F, Tanus-Santos JE (2011) Endothelial nitric oxide synthase haplotypes associated with aura in patients with migraine. *DNA Cell Biol* 30(6):363–369
4. de O S Mansur T, Goncalves FM, Martins-Oliveira A, Speciali JG, Dach F, Lacchini R, Tanus-Santos JE (2012) Inducible nitric oxide synthase haplotype associated with migraine and aura. *Mol Cell Biochem*. doi:[10.1007/s11010-012-1231-0](https://doi.org/10.1007/s11010-012-1231-0)
5. Bigal ME, Kurth T, Santanello N, Buse D, Golden W, Robbins M, Lipton RB (2010) Migraine and cardiovascular disease: a population-based study. *Neurology* 74(8):628–635
6. Maher BH, Griffiths LR (2011) Identification of molecular genetic factors that influence migraine. *Mol Genet Genomics* 285(6):433–446