



Published in final edited form as:

Tob Control. 2012 March ; 21(2): 162–170. doi:10.1136/tobaccocontrol-2011-050200.

Tobacco industry denormalisation as a tobacco control intervention: a review

Ruth E Malone¹, Quinn Grundy¹, and Lisa A Bero²

¹Department of Social and Behavioral Sciences, School of Nursing, University of California, San Francisco, CA, USA

²Department of Clinical Pharmacy, School of Pharmacy, and Institute for Health Policy Studies, School of Medicine, University of California, San Francisco, CA, USA

Abstract

Objective—To conduct a review of research examining the effects of tobacco industry denormalisation (TID) on smoking-related and attitude-related outcomes.

Methods—The authors searched Pubmed and Scopus databases for articles published through December 2010 (see figure 1). We included all peer-reviewed TID studies we could locate that measured smoking-related outcomes and attitudes toward the tobacco industry. Exclusion criteria included: non-English language, focus on tobacco use rather than TID, perceived ad efficacy as sole outcome, complex program interventions without a separately analysable TID component and non peer-reviewed literature. We analysed the literature qualitatively and summarised findings by outcome measured.

Results—After excluding articles not meeting the search criteria, the authors reviewed 60 studies examining TID and 9 smoking-related outcomes, including smoking prevalence, smoking initiation, intention to smoke and intention to quit. The authors also reviewed studies of attitudes towards the tobacco industry and its regulation. The majority of studies suggest that TID is effective in reducing smoking prevalence and initiation and increasing intentions to quit. Evidence is mixed for some other outcomes, but some of the divergent findings may be explained by study designs.

Conclusions—A robust body of evidence suggests that TID is an effective tobacco control intervention at the population level that has a clear exposure–response effect. TID may also contribute to other tobacco control outcomes not explored in this review (including efforts to ‘directly erode industry power’), and thus may enhance public support and political will for structural reforms to end the tobacco epidemic.

INTRODUCTION

Population level interventions have demonstrated over the past two decades that measures altering the social context of tobacco use can significantly reduce smoking prevalence.^{1–4}

Correspondence to Prof Ruth E Malone, Department of Social and Behavioral Sciences, School of Nursing, University of California, San Francisco, 3333 California St, Suite 455, San Francisco, California 94118, USA; ruth.malone@ucsf.edu.

Competing interests None.

Contributors RM originated the idea for the paper, reviewed studies, analysed findings, wrote sections of the paper and reviewed and edited all drafts. QG retrieved data, reviewed studies, analysed findings, developed tables and figures, wrote the first draft of the paper, and reviewed and edited all drafts. LB reviewed studies, analysed findings, wrote paper sections, and reviewed and edited all drafts.

Provenance and peer review Commissioned; externally peer reviewed.

Tobacco industry denormalisation (TID), a focus of several successful tobacco control initiatives in the US and elsewhere, is increasingly regarded as essential to effectively addressing tobacco at the population level. Beginning with the ‘vector analysis’⁵ which first emphasised tobacco industry activities, rather than smokers’ individual behaviours, as critical for tobacco control, a focus on the supply side in advocacy, research, policy and programme planning has appeared.^{2, 6–8} This emphasis, represented in specific provisions in the World Health Organisation’s Framework Convention on Tobacco Control (FCTC),⁹ has not been universally embraced. In some countries, efforts still focus primarily on health education and prohibitions on youth tobacco sales. Some TID efforts have met with aggressive tobacco industry responses, occasionally including lawsuits aimed at curtailing them.^{10, 11}

Mahood¹² distinguishes between the denormalisation of tobacco use (which focuses on the addicted individual) and the denormalisation of the industry, arguing that only the latter offers the prospect of addressing the chief structural cause of the tobacco disease epidemic: industry activity. In this paper we use ‘tobacco industry denormalisation’ to mean themes, campaigns and perspectives aiming towards ‘the reversal of the process of industry normalisation promoted by cigarette manufacturers for decades’.¹² The rationale for TID is captured by the first principle of the Guidelines for implementing Article 5.3 of the FCTC: ‘there is a fundamental and irreconcilable conflict between the tobacco industry’s interests and public health policy interests’.¹³

No previous reviews have specifically examined the effectiveness of TID as a tobacco control strategy. We review evidence on TID and smoking-related outcomes. We analyse why some findings appear to diverge from the bulk of published literature. The evidence suggests strongly that TID is an effective strategy that should be part of comprehensive tobacco control.

METHODS

Search

We searched the PubMed and Scopus databases for articles published through December 2010 using the following search terms: (‘Tobacco Industry’ (Mesh) OR ‘tobacco industry’) AND (delegitimization OR delegitimation OR denormalisation OR de-normalisation OR denormalization OR anti-industry OR counter-industry OR vilification OR industry manipulation); ‘tobacco industry’ AND (deception* OR mistrust* OR lie* OR lying OR false allegation); (‘tobacco industry’ (Mesh) OR ‘tobacco industry’ OR tobacco) AND (countermarketing OR counteradvertising); (‘tobacco industry’ (Mesh) OR ‘tobacco industry’ OR tobacco OR smoking) AND ‘truth campaign’; (‘tobacco industry’ (Mesh) OR ‘tobacco industry’ OR tobacco OR smoking) AND (truth OR ‘truth campaign’); and (‘tobacco industry’ (Mesh) OR ‘tobacco industry’) AND (opinion* OR perception* OR belie* OR ‘support for action’ OR attitude*) (see figure 1). Peer-reviewed research was included if it measured effects of TID on tobacco-related behaviours, attitudes towards industry and support for tobacco control policy (see figure 1). Studies were included from any country, involving any population that measured smoking or tobacco-industry related outcomes. Exclusion criteria included: non-English language, focus on tobacco use rather than TID, perceived ad efficacy as sole outcome, complex programme interventions without a separately analysable TID component and non-peer-reviewed literature.

The searches yielded 891 articles (figure 1). RM and QG independently identified 54 articles meeting inclusion criteria. LB reviewed discrepancies; final selection was achieved by consensus. Six additional articles were identified from reference lists. QG read the 60 articles and abstracted information using a data collection form.

Analysis

Studies were grouped for descriptive analysis by type of outcome measured. Six studies used structural equation modelling to theorise links between TID and smoking-related outcomes and were summarised separately.

RESULTS

Study characteristics

The 60 articles represented 56 unique studies (table 1). Most were conducted in the US (n = 46; 82%). Several regions and counter-industry campaigns were represented: the national 'truth' campaign (n = 15), the Florida 'truth' campaign (n = 9), the Minnesota Initiative, Target Market (n = 3), the Wisconsin Campaign (n = 2), the Mississippi campaign (n = 1) and a media literacy programme in Washington (n = 1).

Nine smoking-related outcomes were measured (see table 1). All were measured using self-report. Three studies employed qualitative methodologies. Cross-sectional designs dominated (n = 37; 66%); 23 of these used repeated measurements. Four studies were longitudinal. There were several quasi-experimental designs: controlled comparison (n = 1), controlled before and after (n = 3), historically controlled (n = 3) and controlled experiment (n = 1). Four studies were randomised controlled trials (RCTs).

In all, 38 studies employed random sampling (60%). The most commonly used sampling frame was a vendor-generated telephone list (n = 34; 54%). Middle and high schools were the next most common (n = 10; 16%). Several studies used the same data sets: the Legacy Media Tracking Surveys (LMTS), sponsored by the American Legacy Foundation (ALF) (n = 14); the Florida Youth Tobacco Survey (FYTS), sponsored by the Florida state Department of Health (n = 2), and the Florida Anti-tobacco Media Evaluation (FAME), through Florida State University (n = 5). These data sets were representative of their population; LMTS oversampled racial and ethnic minorities.

Youth aged 12–17 were the focus of most studies (n = 37; 59%). Three used an extended definition of youth: ages 12–25 years. Five examined young adults, defined as either 18–25 years (n = 4) or 18–29 years (n = 1). A total of 15 studies included adults, 2 requiring adults to have children aged 12–17. Three studies employed mixed youth/adult samples.

A conflict of interest exists when campaigns are evaluated by implementing agencies. In Florida, the Department of Health contracted for an independent evaluation.¹⁴ Twelve studies evaluating the national 'truth' campaign were funded by the ALF, 2 by other sources; 3 did not disclose funding. These studies' principal researchers are housed at ALF, RTI International, Centers for Disease Control, American Institute for Research, RAND Corporation and other academic research centres. It appeared that none of the researchers were responsible for data collection through FAME, FYTS or LMTS surveys and many acknowledged external survey management companies.

Smoking prevalence

TID's relation to smoking prevalence was examined in 13 studies. Measured by self-report, studies most commonly employed 30 day and 100-cigarettes-in-lifetime referents, placing respondents on a 3–5 classification continuum (table 1). This measure is widely used and appropriate for youth and young adults, capturing the construct of smoking initiation and progression to smoking dependence.

Smoking prevalence: youth—The first youth campaign to be highly funded and thoroughly evaluated was the ‘truth’ campaign piloted by the state of Florida, launched in April 1998, featuring a strong TID component. Surveys following Florida ‘truth’s’ implementation showed large decreases in smoking prevalence among youth ages 12–17^{14–16} and similar decreases for prevalence in all categories along the smoking behaviour continuum.¹⁵ Prevalence of never users and those defined as committed non-smokers rose significantly.¹⁵ Less than a year after campaign launch, non-smoking youth who remained non-smokers were 2.3 times more likely than those who started smoking to say they were influenced ‘a lot’ by the ‘truth’ campaign’s primary, industry manipulation message.¹⁴

At 2 years into the Florida campaign, national ‘truth’ was launched, permitting comparisons between Florida and national youth. Florida youth were significantly less likely than national youth to have smoked in the past month, or to have ever tried smoking.^{17, 18} Florida youth also held less favourable beliefs about the tobacco industry.¹⁷ Predictive of smoking behaviour in the past 30 days were two beliefs central to the campaign: ‘cigarette companies lie’ and ‘cigarette companies try to get young people to start smoking’.¹⁷ However, 4 years after campaign termination a reversal was seen: although most youth were still able to confirm ‘truth’ campaign awareness, smoking rates for youth had increased 6.8%.¹⁶

Three studies examined counter-industry media campaign effects on smoking prevalence among a national youth sample. One compared youth smoking among states with long-term funded, recently funded, or no counter-industry campaigns.¹⁹ The rate of decrease in youth smoking rates in states with established or new campaigns versus those without was nearly double; similarly, odds of current smoking were reduced significantly faster in states with counter-industry campaigns than in those without.¹⁹ Over time, negative beliefs and attitudes about the tobacco industry showed significantly stronger relationships with smoking status in campaign versus non-campaign states.¹⁹

In a large, random, national sample of youth ages 12–17, higher sensation seeking and weaker counter-industry attitudes were independent predictors of current smoking.²⁰ The relationship between counter-industry attitudes and smoking behaviour was consistent across risk groups, suggesting that counter-industry messages may be equally effective for high and low risk youth.²⁰ In Canada, a school-based survey of high school students showed that decreases in occasional and regular smoking behaviour were significantly related to TID beliefs.²¹

A historically controlled experiment examined the effects of increased exposure to the national ‘truth’ campaign.²² Researchers found a significant inverse relationship between ‘truth’ exposure and youth smoking prevalence; however, this effect diminished at higher exposure levels.²² Researchers concluded that roughly 22% of the 36% decline in youth smoking prevalence from 1997 to 2002 was attributable to the national ‘truth’ campaign.²²

ALF and the Washington State Department of Health partnered to implement and evaluate a tobacco media literacy curriculum about deceptive industry advertising tactics.²³ Although the programme had significant effects on other measures of tobacco attitudes, beliefs and behaviours, it had null effects on smoking prevalence.²³

Smoking prevalence: young adults—Three cross-sectional studies measured smoking prevalence in young adults. Counter-industry attitudes and beliefs, including support for action against the industry, were strongly, negatively associated with current smoking in all three.^{24–26} Among US Air Force recruits prior to a mandated cessation programme, the strongest predictor of smoking status was attitude towards the tobacco industry.²⁶ Along the smoking status continuum, former smokers relative to never smokers, experimental smokers

relative to never smokers and current smokers relative to former smokers were less likely to agree with the statement 'tobacco companies lied/misled the public'.²⁶

Smoking initiation

Five studies measured smoking initiation. Smoking initiation is generally only applicable in measuring youth smoking behaviours: four of the studies surveyed samples aged 12–20. However, one examined enforced cessation during Basic Military Training as an intervention, which allowed for measurement of relapse/initiation.²⁶ Relapse was less likely among baseline smokers who agreed with statements about the industry's deceptive practices.²⁶ However, attitude towards the industry was not a significant predictor of smoking initiation between baseline 'never' or 'experimental' smokers.²⁶

The relationship between smoking initiation and the national 'truth' campaign was examined in two longitudinal surveys. More frequent 'truth' recall was associated with both decreased likelihood of smoking initiation and tobacco dependence.²⁷ Increased cumulative campaign exposure was associated with a 20% decrease in initiation risk of over a period of 7 years.²⁸

Two repeated cross-sectional surveys examined progression of non-smokers to smokers in association with Florida's 'truth' campaign, finding evidence of an inverse relationship between increased exposure to truth and smoking initiation over a period of 22 months.²⁹ Rates of smoking initiation for baseline non-smokers varied inversely with the number of 'truth' ads recalled, the reported influence of the counter-industry theme and the strength of industry manipulation attitudes.²⁹ In another study, compared with those unaffected by the campaign, youth reporting low or high anti-industry ad effects were 1.3 and 1.7 times more likely to remain non-smokers by the second survey.³⁰

Intention to smoke

In all, 17 studies measured intention to smoke.

Intention to smoke: youth—Florida's 'truth' campaign was associated with increases in the proportion of youth ages 12–17 who identified as 'closed to smoking' and experimenters who stated they would not smoke again.^{15, 31} Florida teens were also less likely than their national counterparts to be open to future smoking.¹⁷ However, after the state cut funding for the programme in 1999, trends in non-smoking intentions were significantly reduced.³¹

Two components of the Minnesota counter-industry campaign Target Market (TM) were studied: TM/org (youth organising) and TM/ads (mass media). No significant relationship was found between exposure to TM/org and intention to smoke.³² However, youth with greater intentions to smoke scored significantly lower on certain attitudinal items such as 'teens have been influenced by the tobacco industry'.³² Termination of the TM/ads component had negative impacts on prevalence of respondents scoring highly on industry manipulation attitudes/beliefs, and prevalence of participants not intending to smoke in the next year.³³

A study of Wisconsin's counter-industry themed campaign found that positive attitudes towards the campaign and number of ads seen were significantly, negatively associated with smoking intentions.³⁴

In a repeated cross-sectional study during the first 3 years of the national 'truth' campaign, campaign recall was associated with greater odds of youth ages 12–17 ruling out future smoking.³⁵ Higher sensation-seeking traits and weaker counter-industry attitudes independently predicted intention to smoke.²⁰ Another repeated cross-sectional study of this same population found no statistically significant increase in the percentage of non-smokers

saying they probably/definitely would not smoke in the next year; however, six of nine counter-industry and empowerment attitudes were strongly associated with reduced smoking intentions.³⁶ Analysis of racial/ethnic data found a significant association between ‘truth’ exposure and belief and attitude indices, but the impact, though similar for white and African–American youth, was lower for Hispanic youth.³⁷ Though most youth across racial/ethnic subgroups do not intend to smoke at baseline (94%), exposure to ‘truth’ was associated with 2.0 greater odds of not intending to smoke among never smokers and 5.7 greater odds among ever smokers, across all racial/ethnic subgroups.³⁷

A longitudinal study of youth at low and high risk for smoking showed that those exposed to the national ‘truth’ campaign were more likely to hold anti-smoking beliefs at follow-up, and more frequent ‘truth’ recall was significantly associated with decreased likelihood of developing openness to smoking, intentions to smoke soon and in 5 years, but not for intentions to smoke within 1 year.²⁷

Three studies used RCTs to investigate the effectiveness of different message themes on adolescents’ intentions to smoke.^{38–40} These concluded that industry manipulation-themed ads were ineffective in decreasing intentions to smoke. One study randomised 7th and 10th grade students (44% Hispanic) to eight different ad conditions (two of which could be considered TID-themed) and a control group.³⁸ Two TID-themed ads enhanced health risk severity perceptions. One TID theme positively influenced 10th graders’ perceived vulnerability to social disapproval risks.³⁸

Ninth grade California students were randomised to nine ad conditions, three of which were ‘counter-industry’ type.³⁹ Among all participants, no ad type lowered intentions to smoke versus the control.

A third study randomised 16 groups of Virginia high school students to 1 of 3 message theme conditions (where 1 was industry manipulation), or a control.⁴⁰ Those viewing ads portraying negative life circumstances of smokers had lower intentions to smoke than either control or industry manipulation ad groups.⁴⁰

Intention to smoke: adults—Evidence for TID’s effects on intention to smoke among adults is less consistent, perhaps because most campaigns targeted youth. No significant association between campaign awareness and intention to not smoke was found for young adults aged 18–24 in relation to the national ‘truth’ campaign.⁴¹ However, several specific attitudes/beliefs promoted by ‘truth’ were associated with intention to not smoke such as, ‘I would like to see cigarette companies go out of business’.⁴¹ Another study surveying general counter-industry attitudes among adults found support for counter-industry action was negatively associated with intention to smoke within the year.²⁵

Two controlled before and after studies examined TID and intention to smoke in the context of movies. Surveyed Australian adults viewed *The Insider* (a film about tobacco industry duplicity) or the control *Erin Brokovich* (plot analogous but not about tobacco). The *The Insider* group showed a decline in intentions to smoke at the post-film survey and a divergent trend from the control group for current, former and non-smokers.⁴² However, when late responders were included, these effects were non-significant, suggesting the film’s impact may have been transitory.⁴² A historically controlled experiment with cinemagoers seeing an industry manipulation-themed advert prior to viewing their film found a greater proportion of non-smoking participants in the intervention group agreed that smoking in movies was ‘not OK’, but a significantly higher percentage of smokers in the intervention group said they would still be smoking in 12 months.⁴³

Intention to quit

Eight studies^{8, 24, 25, 34, 41, 44–49} examined TID's effects on intentions to quit smoking. All used a single-item measure with adults. Several cross-sectional studies found that beliefs about industry deceptiveness were positively related to consideration of quitting,^{34, 44, 45} as was support for counter-industry action.^{24, 25} Another study found that supporting action against the industry was positively associated with quit intentions, but not with a serious quit attempt.²⁵ In a longitudinal study in four Western nations, smokers reporting medium and high TID beliefs were more likely to intend to quit, and although TID beliefs at time 1 did not predict abstinence at time 2, smokers with stronger beliefs at time 2 were more likely to be abstinent than those whose beliefs did not increase.⁸

Although exposure to the national 'truth' campaign was not associated with intention to quit smoking among young adults, seven of nine counter-industry belief/attitude items targeted by the 'truth' campaign were associated with intention to quit among young adult smokers.⁴¹ Awareness of Florida's 'truth' campaign reached approximately 50% of adults, and the only variable significantly associated with quit intentions was awareness of the industry manipulation theme, independent of parental status.^{44, 45} However, in an evaluation of Wisconsin's campaign, beliefs about industry deceptiveness were not significantly related to quitting considerations.⁴⁹

One study examined effectiveness of ad themes on college students' tobacco use, including intention to quit. College students assigned to TID-themed ads were twice as likely to intend to quit as those who viewed social norms ads.⁴⁷

Other outcomes

Youth generally overestimate perceptions of peer smoking prevalence (PPSP); it may be a precursor to future smoking.⁵⁰ Two studies suggested that TID is associated with lower PPSP.^{23, 50} Evidence from two studies of smoking resistance self-efficacy (SRSE), which is predictive of youth smoking, was mixed: TID-themed ads were associated with lower SRSE scores than ads emphasising the effects of smoking.⁵¹ However, industry manipulation ads were associated with greater SRSE when they used less, versus more appealing actors.⁵¹ Explicit industry manipulation messages (vs implicit) were associated with stronger SRSE.⁵² One RCT found that no tested message themes affected SRSE or marketing resistance self-efficacy.³⁸ TID-related activities were associated with higher empowerment (the degree to which youth feel they would like to get involved in organising against the tobacco industry and feel youth can make a difference) scores.^{32, 53} However, youth with higher intentions to smoke were less likely to believe that youth could be effective.³²

Attitudes towards industry and its regulation

Views of industry—Across diverse samples, tobacco companies were regarded as dishonest, unethical and less trustworthy than other companies.^{42, 54–57} Negative views of industry appeared to be increasingly negative over time.^{55, 57} Australian smokers had stronger counter-industry beliefs than either Canadian or UK smokers; UK smokers reported weaker counter-industry beliefs than US or Canadian smokers.⁸ A minority of Russian adults felt tobacco companies behaved unethically, despite three-quarters agreeing that tobacco companies definitely or maybe bribed politicians.⁵⁸ Rather than being a TID marker, this behaviour was considered typical for a Russian corporation.

There is some evidence that support for the process of TID is weaker among smokers;^{42, 54–56} however,⁵⁹ in one study smokers were polarised on these beliefs.⁵⁵ Smokers from four countries reported relatively strong counter-industry beliefs, especially

among the older and more educated.⁸ Women smokers were less likely to report counter-industry beliefs than men.⁸

In a Canadian study, youth appeared less distrustful of tobacco companies than adults, a consistent finding across smoking categories and school grade.⁵⁷ African American youth focus group participants, however, expressed anger at tobacco industry targeting, considering it a form of racism.⁶⁰ A Florida survey found non-smoking youth more likely than smoking youth to espouse counter-industry attitudes.⁵⁹

One cross-sectional study surveyed adult African-Americans' views about black organisations accepting tobacco industry philanthropy.⁶¹ Despite most agreeing that philanthropy makes the industry money and encourages smoking, one-third said it also helps the community.⁶¹ Those with college educations and men were less likely to find this philanthropy acceptable; younger adults were less likely to agree that this philanthropy aimed to help communities.⁶¹

Views of regulation—There were varied opinions ranging from neutral to strong beliefs about industry responsibility for smoking's harms and its regulation. Descriptive research has assessed attitudes towards tobacco industry regulation in diverse samples including: adults in Ontario;⁵⁴ Mississippi parents before and after a youth-targeted, counter-industry state media campaign;⁶² Korean immigrant, male smokers;⁶³ school nurses⁶⁴ and New Zealand politicians.⁶⁵ In a sample of smokers in four countries, thinking about the conduct of tobacco companies and belief that the industry should take more responsibility for tobacco's harms were independently predictive of support for industry regulation,⁴⁶ and counter-industry beliefs were associated with noticing anti-smoking information, tobacco ads and secondhand smoke restrictions.⁸

Experimental studies: attitudes about industry

Two experimental studies examined how an intervention could shape perceptions of the industry. Public perceptions of the tobacco industry were assessed using a controlled before-and-after design, with the movie *The Insider* as the intervention.⁴² Post test, subjects viewing *The Insider* rated tobacco industry executives lower on ethics and honesty and higher on power, held more negative views of industry conduct and showed less acceptance of the industry than controls.⁴² An RCT investigated effects of perceptions of Philip Morris (PM) as a tobacco company on evaluations of the company's advertising among undergraduates.⁶⁶ There was no association between students' opinion of PM and awareness that it is a tobacco company; however, PM corporate advertisements were rated more favourably by students unaware of this fact.⁶⁶

Theorising TID

Six studies employed structural equation modelling to theorise TID's effects. One model suggested knowledge of industry deception leads to mistrust of the industry, which in turn is associated with support for action against the industry and reduced receptivity to advertising.²⁴ Models derived from counter-industry campaign data show that TID exposure leads to negative beliefs about the industry's conduct, predicting negative attitudes towards the industry; these are associated with lesser progression towards smoking and reduced receptivity to pro-tobacco influences.^{67, 68} Mistrust of the industry appears to strongly influence negative attitudes towards the industry, which in turn explains significant variability in smoking behaviour, suggesting these campaigns succeed because they resonate with trust-related values.⁶⁹ Social imagery, perceived tobacco independence and brand equity have also been shown to mediate the relationship between current smoking and exposure to the national 'truth' campaign.⁷⁰⁻⁷²

DISCUSSION

A robust body of evidence supports TID as an effective population-level tobacco control strategy that contributes to reduced smoking prevalence among youth and young adults, reduced smoking initiation among youth, increased intentions to quit and reduced perceived peer smoking prevalence. Evidence is mixed on TID's impact on intentions to smoke, youth empowerment and views of the industry and its regulation, but evidence from California suggests TID's importance as part of a comprehensive social norm change programme.²

Limitations

Because TID is not yet an established indexing term, we may have missed relevant studies. We reluctantly excluded literature on California's landmark programme, the first to feature a strong TID component, because published reports merged TID and other social norm change components into a single construct; these could not be separately analysed. However, California programme evaluations suggest that TID has been an important element in increasing quitting, reducing smoking prevalence and increasing support for tobacco control.^{2, 3, 73-76} Heterogeneity in TID interventions and outcome measures did not allow quantitative analyses. Most evidence is from cross-sectional studies, limiting the ability to draw causal conclusions. Most were US studies; national/cultural differences in attitudes towards industry and regulation could limit generalisability.

Interpreting contradictory trials findings

The RCTs' failure to find an association between TID and intent to smoke may be because the intensity and duration of the interventions were less than in observational studies. Processing of TID-related ads may require additional exposures.⁴⁰ Experimental studies may not capture TID's true effectiveness. In Sutfin's study,⁴⁰ the only ad type associated with decreased intention to smoke was also the only type that participants reported having previously seen, suggesting that repeated exposures might increase intervention effectiveness.

In addition, trials did not consistently control for understanding of the intervention. In one study, only 34% of participants correctly identified TID ads.⁴⁰ However, this was not included as a covariate for the outcome of intention to smoke. In larger studies,^{38, 39} most students recalling ads correctly identified themes.

None of these studies reported absolute numbers of participants expressing intention to smoke. Although Pechmann^{38, 39} reported that only one of nine ad types tested had a significant effect on smoking intent compared to control, differences between ad types appeared small, making it difficult to determine if there were meaningful differences in proportions of participants expressing intentions to smoke. It may also be that TID interventions are simply less effectively delivered at the individual level.

Research gaps

Many important questions remain unanswered by existing TID-related research. For example, the cost effectiveness of TID programmes relative to other tobacco control interventions remains largely unstudied. A cost-utility analysis of the national 'truth' campaign estimated that the campaign recovered the costs of development, delivery, evaluation and litigation, and averted nearly US\$1.9 billion in smoking-related medical costs.⁷⁷ However, the cost effectiveness of such mass media TID interventions compared with other tobacco control interventions is unknown. Similarly, comparative studies are needed to analyse potentially synergistic effects of TID and other tobacco control interventions; whether TID interventions are a necessary component of comprehensive

tobacco control; and whether they have an additional effect once other components are in place.

Three existing comparative studies tested the national ‘truth’ campaign against the Philip Morris-sponsored youth smoking prevention programme, ‘Think Don’t Smoke’.^{27, 35, 36, 50} Whereas the ‘truth’ campaign was associated with an increase in anti-tobacco beliefs and attitudes,^{27, 35, 36} ‘Think Don’t Smoke’ was associated with an increase in favourable attitudes towards the tobacco industry.^{35, 36} Similarly, the ‘truth’ campaign was associated with decreased perceived smoking prevalence,⁵⁰ decreased intentions to smoke and lower rates of smoking initiation,²⁷ while ‘Think Don’t Smoke’ was associated with increased intentions to smoke soon.²⁷ Arguably, an industry-sponsored programme does not offer a fair comparison, so studies aimed at teasing out certain effects of TID as compared with other programme components may be useful. However, TID is also likely to have broader, indirect effects on the policy climate, which constitutes another important area for research.

Most reviewed studies involved mass media interventions; however, TID does not only consist of or work through such relatively expensive interventions, but through wider tobacco control advocacy efforts, such as through earned or unpaid media.^{78–81} It is difficult, if not impossible, to fully and explicitly account for such efforts, which change the public discourse about tobacco use by reframing it away from individual behaviour change towards industry regulation.

Conclusion

Unpacking why TID is an effective tobacco control intervention is complex methodologically and theoretically.⁸² TID’s effectiveness is likely due to synergies between myriad political and cultural influences that cannot be isolated.⁸³ The evidence suggests that TID is most effectively delivered at the population level and that increased exposure is generally associated with increased effects. Regardless of how TID works, the industry’s aggressive responses suggest that TID passes the ‘scream test’, constituting a threat to the industry’s legitimacy and its continued success in normalising its business, its marketing, and its products.^{1, 11, 84–88} TID may contribute to efforts to ‘directly erode industry power’, making tobacco companies less able to thwart effective tobacco control.⁸⁹ TID could also enhance public support and political will to counter industry ‘makeover’ efforts and tackle fundamental structural reforms to end the tobacco epidemic.^{7, 90–93}

Acknowledgments

Funding National Cancer Institute, Grant Number CA120138 (REM); Flight Attendants’ Medical Research Institute (LAB).

References

1. Chapman, S. *Public Health Advocacy and Tobacco Control: Making Smoking History*. Oxford, UK: Blackwell Publishing; 2007.
2. Roeseler A, Burns D. The quarter that changed the world. *Tob Control*. 2010; 19 (Suppl 1):i3–15. [PubMed: 20382647]
3. Tang H, Abramsohn E, Park H-Y, et al. Using a cessation-related outcome index to assess California’s cessation progress at the population level. *Tob Control*. 2010; 19 (Suppl 1):i56–61. [PubMed: 20382652]
4. Pierce, JP.; Gilpin, EA.; Emery, SL., et al. *Tobacco Control in California: Who’s Winning the War? An Evaluation of the Tobacco Control Program, 1989–1996*. La Jolla, CA: University of California San Diego; 1998.
5. LeGresley E. A ‘vector analysis’ of the tobacco epidemic. *Medicus Mundi Schweiz*. 1999:72.

6. World Health Organization. Tobacco Industry Interference with Tobacco Control. Geneva, Switzerland: World Health Organization; 2009.
7. Callard C, Thompson D, Collishaw N. Transforming the tobacco market: why the supply of cigarettes should be transferred from for-profit corporations to non-profit enterprises with a public health mandate. *Tob Control*. 2005; 14:278–83. [PubMed: 16046692]
8. Hammond D, Fong GT, Zanna MP, et al. Among smokers from four countries. *Am J Prev Med*. 2006; 31:225–32. [PubMed: 16905033]
9. World Health Organization. Framework Convention on Tobacco Control. Geneva, Switzerland: World Health Organization; 2003.
10. Glantz, S.; Balbach, ED. Tobacco War: Inside the California Battles. Berkeley, CA: University of California Press; 2000.
11. Apollonio DE, Malone RE. Turning negative into positive: public health mass media campaigns and negative advertising. *Health Educ Res*. 2009; 24:483–95. [PubMed: 18948569]
12. Mahood, G. Tobacco Industry Denormalization: Telling the Truth About the Tobacco Industry's Role in the Tobacco Epidemic. Canada: Nonsmokers' Rights Association; 2004.
13. WHO. Guidelines for Implementation of Article 5.3 of the WHO Framework Convention on Tobacco Control. Geneva, Switzerland: World Health Organization (WHO); 2008. Framework Convention on Tobacco Control.
14. Zucker D, Hopkins RS, Sly DF, et al. Florida's "truth" campaign: a counter-marketing, anti-tobacco media campaign. *J Public Health Manag Pract*. 2000; 6:1–6. [PubMed: 10848476]
15. Bauer UE, Johnson TM, Hopkins RS, et al. Changes in youth cigarette use and intentions following implementation of a tobacco control program: findings from the Florida Youth Tobacco Survey, 1998–2000. *JAMA*. 2000; 284:723–8. [PubMed: 10927781]
16. Dietz NA, Westphal L, Arheart KL, et al. Changes in youth cigarette use following the dismantling of an antitobacco media campaign in Florida. *Prev Chronic Dis*. 2010; 7:A65. [PubMed: 20394704]
17. Niederdeppe J, Farrelly MC, Haviland ML. Confirming "truth": more evidence of a successful tobacco countermarketing campaign in Florida. *Am J Public Health*. 2004; 94:255–7. [PubMed: 14759936]
18. Sly DF, Heald GR, Ray S. The Florida "truth" anti-tobacco media evaluation: design, first year results, and implications for planning future state media evaluations. *Tob Control*. 2001; 10:9–15. [PubMed: 11226354]
19. Hersey JC, Niederdeppe J, Ng SW, et al. How state counter-industry campaigns help prime perceptions of tobacco industry practices to promote reductions in youth smoking. *Tob Control*. 2005; 14:377–83. [PubMed: 16319360]
20. Thrasher JF, Niederdeppe JD, Jackson C, et al. Using anti-tobacco industry messages to prevent smoking among high-risk adolescents. *Health Educ Res*. 2006; 21:325–37. [PubMed: 16492681]
21. Leatherdale ST, Sparks R, Kirsh VA. Beliefs about tobacco industry (mal)practices and youth smoking behaviour: insight for future tobacco control campaigns (Canada). *Cancer Causes Control*. 2006; 17:705–11. [PubMed: 16633918]
22. Farrelly MC, Davis KC, Haviland ML, et al. Evidence of a dose-response relationship between "truth" antismoking ads and youth smoking prevalence. *Am J Public Health*. 2005; 95:425–31. [PubMed: 15727971]
23. Austin EW, Pinkleton BE, Hust SJ, et al. Evaluation of an American Legacy Foundation/Washington State Department Of Health Media Literacy Pilot Study. *Health Commun*. 2005; 18:75–95. [PubMed: 15918791]
24. Ling PM, Neilands TB, Glantz SA. The effect of support for action against the tobacco industry on smoking among young adults. *Am J Public Health*. 2007; 97:1449–56. [PubMed: 17600255]
25. Ling PM, Neilands TB, Glantz SA. Young adult smoking behavior: a national survey. *Am J Prev Med*. 2009; 36:389–94. e2. [PubMed: 19269128]
26. Klesges RC, Sherrill-Mittleman DA, Debon M, et al. Do we believe the tobacco industry lied to us? Association with smoking behavior in a military population. *Health Educ Res*. 2009; 24:909–21. [PubMed: 19528314]

27. Davis KC, Farrelly MC, Messeri P, et al. The impact of national smoking prevention campaigns on tobacco-related beliefs, intentions to smoke and smoking initiation: results from a longitudinal survey of youth in the United States. *Int J Environ Res Public Health*. 2009; 6:722–40. [PubMed: 19440412]
28. Farrelly MC, Nonnemaker J, Davis KC, et al. The Influence of the National truth campaign on smoking initiation. *Am J Prev Med*. 2009; 36:379–84. [PubMed: 19211213]
29. Sly DF, Trapido E, Ray S. Evidence of the dose effects of an antitobacco counteradvertising campaign. *Prev Med*. 2002; 35:511–18. [PubMed: 12431900]
30. Sly DF, Hopkins RS, Trapido E, et al. Influence of a counteradvertising media campaign on initiation of smoking: the Florida “truth” campaign. *Am J Public Health*. 2001; 91:233–8. [PubMed: 11211631]
31. Niederdeppe J, Farrelly MC, Hersey JC, et al. Consequences of dramatic reductions in state tobacco control funds: Florida, 1998–2000. *Tob Control*. 2008; 17:205–10. [PubMed: 18390911]
32. Dunn CL, Pirie PL, Oakes JM. Outcomes of a statewide anti-tobacco industry youth organizing movement. *Am J Health Promot*. 2004; 19:3–11. [PubMed: 15460095]
33. Sly DF, Arheart K, Dietz N, et al. The outcome consequences of defunding the Minnesota youth tobacco-use prevention program. *Prev Med*. 2005; 41:503–10. [PubMed: 15917046]
34. Tangari AH, Burton S, Andrews JC, et al. How do antitobacco campaign advertising and smoking status affect beliefs and intentions? Some similarities and differences between adults and adolescents. *J Publ Pol Market*. 2007; 26:60–74.
35. Farrelly MC, Davis KC, Duke J, et al. Sustaining ‘truth’: changes in youth tobacco attitudes and smoking intentions after 3 years of a national antismoking campaign. *Health Educ Res*. 2009; 24:42–8. [PubMed: 18203679]
36. Farrelly MC, Heaton CG, Davis KC, et al. Getting to the truth: evaluating national tobacco countermarketing campaigns. *Am J Public Health*. 2002; 92:901–7. [PubMed: 12036775]
37. Cowell AJ, Farrelly MC, Chou R, et al. Assessing the impact of the national ‘truth’ antismoking campaign on beliefs, attitudes, and intent to smoke by race/ethnicity. *Ethn Health*. 2009; 14:75–91. [PubMed: 19152160]
38. Pechmann C, Zhao G, Goldberg ME, et al. What to convey in anti-smoking advertisements for adolescents: the use of protection motivation theory to identify effective message themes. *J Market*. 2003; 67:1–18.
39. Pechmann C, Reibling ET. Antismoking advertisements for youths: an independent evaluation of health, counter-industry, and industry approaches. *Am J Public Health*. 2006; 96:906–13. [PubMed: 16571709]
40. Sutfin EL, Szykman LR, Moore MC. Adolescents’ responses to anti-tobacco advertising: exploring the role of adolescents’ smoking status and advertisement theme. *J Health Commun*. 2008; 13:480–500. [PubMed: 18661389]
41. Richardson AK, Green M, Xiao H, et al. Evidence for truth® the young adult response to a youth-focused anti-smoking media campaign. *Am J Prev Med*. 2010; 39:500–6. [PubMed: 21084069]
42. Dixon HG, Hill DJ, Borland R, et al. Public reaction to the portrayal of the tobacco industry in the film *the Insider*. *Tob Control*. 2001; 10:285–91. [PubMed: 11544395]
43. Edwards C, Oakes W, Bull D. Out of the smokescreen II: will an advertisement targeting the tobacco industry affect young people’s perception or smoking in movies and their intention to smoke? *Tob Control*. 2007; 16:177–81. [PubMed: 17565137]
44. Delva J, Dietz NA, Perron B, et al. Adult awareness of a youth-focused anti-tobacco campaign: does having children matter? *Subst Use Misuse*. 2009; 44:763–74. [PubMed: 19444720]
45. Dietz NA, Delva J, Woolley ME, et al. The reach of a youth-oriented anti-tobacco media campaign on adult smokers. *Drug Alcohol Depend*. 2008; 93:180–4. [PubMed: 17964081]
46. Young D, Borland R, Siahpush M, et al. Australian smokers support stronger regulatory controls on tobacco: findings from the ITC Four-Country Survey. *Aust N Z J Public Health*. 2007; 31:164–9. [PubMed: 17461008]
47. Murphy-Hoefler R, Hyland A, Higbee C. Perceived effectiveness of tobacco countermarketing advertisements among young adults. *Am J Health Behav*. 2008; 32:725–34. [PubMed: 18442351]

48. Murphy-Hoefer R, Hyland A, Rivard C. The influence of tobacco countermarketing ads on college students' knowledge, attitudes, and beliefs. *J Am Coll Health*. 2010; 58:373–81. [PubMed: 20159761]
49. Netemeyer RG, Andrews JC, Burton S. Effects of antismoking advertising-based beliefs on adult smokers' consideration of quitting. *Am J Public Health*. 2005; 95:1062–66. [PubMed: 15914834]
50. Davis KC, Nonnemaker JM, Farrelly MC. Association between national smoking prevention campaigns and perceived smoking prevalence among youth in the United States. *J Adolesc Health*. 2007; 41:430–6. [PubMed: 17950162]
51. Shadel WG, Fryer CS, Tharp-Taylor S. Uncovering the most effective active ingredients of antismoking public service announcements: the role of actor and message characteristics. *Nicotine Tob Res*. 2009; 11:547–52. [PubMed: 19372574]
52. Shadel WG, Fryer CS, Tharp-Taylor S. Tobacco industry manipulation messages in anti-smoking public service announcements: the effect of explicitly versus implicitly delivering messages. *Addict Behav*. 2010; 35:526–9. [PubMed: 20071100]
53. Dunn CL, Pirie PL. Empowering youth for tobacco control. *Am J Health Promot*. 2005; 20:7–10. [PubMed: 16171155]
54. Ashley MJ, Cohen JE. What the public thinks about the tobacco industry and its products. *Tob Control*. 2003; 12:396–400. [PubMed: 14660775]
55. Durkin SJ, Germain D, Wakefield M. Adult's perceptions about whether tobacco companies tell the truth in relation to issues about smoking. *Tob Control*. 2005; 14:429–30. [PubMed: 16319369]
56. Wakefield M, Miller C, Woodward S. Community perceptions about the tobacco industry and tobacco control funding. *Aust N Z J Public Health*. 1999; 23:240–4. [PubMed: 10388165]
57. Waller BJ, Cohen JE, Ashley MJ. Youth attitudes towards tobacco control: a preliminary assessment. *Chronic Dis Can*. 2004; 25:97–100. [PubMed: 15841850]
58. Danishevski K, Gilmore A, McKee M. Public attitudes towards smoking and tobacco control policy in Russia. *Tob Control*. 2008; 17:276–83. [PubMed: 18653793]
59. Sly Heald G, Hopkins RS, DF, et al. The industry manipulation attitudes of smokers and nonsmokers. *J Public Health Manag Pract*. 2000; 6:49–56. [PubMed: 10848483]
60. Johnson DM, Wine LA, Zack S, et al. Designing a tobacco counter-marketing campaign for African American youth. *Tob Induc Dis*. 2008; 4:7. [PubMed: 18822164]
61. King G, Gebreselassie T, Mallett RK, et al. Opinions of African Americans about tobacco industry philanthropy. *Prev Med*. 2007; 45:464–70. [PubMed: 17889293]
62. Carver V, Reinert B, Range LM, et al. Media campaign influences parents' opinions about their children and tobacco. *J Public Health Manage Pract*. 2003; 9:72–8.
63. Kim SS, Nam KA. Korean male smokers' perceptions of tobacco control policies in the United States. *Public Health Nurs*. 2005; 22:221–9. [PubMed: 15982195]
64. Reinert B, Carver V, Range LM. School nurses' opinions about the prevention of tobacco use. *J Community Health Nurs*. 2005; 22:205–11. [PubMed: 16245972]
65. Hudson S, Thomson G, Wilson N. A pilot qualitative study of New Zealand policymakers' knowledge of, and attitudes to, the tobacco industry. *Aust New Zealand Health Policy*. 2007; 4:17. [PubMed: 17651500]
66. Henriksen L, Fortmann SP. Young adults' opinions of Philip Morris and its television advertising. *Tob Control*. 2002; 11:236–40. [PubMed: 12198275]
67. Hersey JC, Niederdeppe J, Evans WD, et al. The theory of "truth": how counterindustry media campaigns affect smoking behavior among teens. *Health Psychol*. 2005; 24:22–31. [PubMed: 15631559]
68. Hersey JC, Niederdeppe J, Evans WD, et al. The effects of state counterindustry media campaigns on beliefs, attitudes, and smoking status among teens and young adults. *Prev Med*. 2003; 37:544–52. [PubMed: 14636787]
69. Thrasher JF, Jackson C. Mistrusting companies, mistrusting the tobacco industry: clarifying the context of tobacco prevention efforts that focus on the tobacco industry. *J Health Soc Behav*. 2006; 47:406–22. [PubMed: 17240928]

70. Evans WD, Wasserman J, Bertolotti E, et al. Branding behavior: the strategy behind the truth campaign. *Soc Mark Q.* 2002; 8:17–29.
71. Evans WD, Price S, Blahut S. Evaluating the truth brand. *J Health Commun.* 2005; 10:181–92. [PubMed: 15804907]
72. Evans WD, Price S, Blahut S, et al. Social imagery, tobacco independence, and the truthsm campaign. *J Health Commun.* 2004; 9:425–41. [PubMed: 15513790]
73. Cowling DW, Modayil MV, Stevens C. Assessing the relationship between ad volume and awareness of a tobacco education media campaign. *Tob Control.* 2010; 19(Suppl 1):i37–42. [PubMed: 20382649]
74. Zhang X, Cowling DW, Tang H. The impact of social norm change strategies on smokers' quitting behaviours. *Tob control.* 2010; 19(Suppl 1):i51–5. [PubMed: 20382651]
75. Park HY, Dent C, Abramsohn E, et al. Evaluation of California's in-school tobacco use prevention education (TUPE) activities using a nested school-longitudinal design, 2003–2004 and 2005–2006. *Tob Control.* 2010; 19(Suppl 1):i43–50. [PubMed: 20382650]
76. Modayil MV, Cowling DW, Tang H, et al. An evaluation of the California community intervention. *Tob Control.* 2010; 19(Suppl 1):i30–6. [PubMed: 20382648]
77. Holtgrave DR, Wunderink KA, Vallone DM, et al. Cost-utility analysis of the National truth campaign to prevent youth smoking. *Am J Prev Med.* 2009; 36:385–8. [PubMed: 19211214]
78. Christofides N, Chapman S, Dominello A. The new pariahs: discourse on the tobacco industry in the Sydney press, 1993–97. *Aust N Z J Public Health.* 1999; 23:233–9. [PubMed: 10388164]
79. Chapman S. The news on tobacco control: time to bring the background into the foreground. *Tob Control.* 1999; 8:237–9. [PubMed: 10599562]
80. McLeod K, Wakefield M, Chapman S, et al. Changes in the news representation of smokers and tobacco-related media advocacy from 1995 to 2005 in Australia. *J Epidemiol Community Health.* 2009; 63:215–20. [PubMed: 19015222]
81. Miller CL, Hill DJ, Quester PG, et al. Response of mass media, tobacco industry and smokers to the introduction of graphic cigarette pack warnings in Australia. *Eur J Public Health.* 2009; 19:644–9. [PubMed: 19589850]
82. Agostinelli G, Grube JW. Tobacco counter-advertising: a review of the literature and a conceptual model for understanding effects. *J Health Commun.* 2003; 8:107–27. [PubMed: 12746036]
83. Chapman S, Freeman B. Markers of the denormalisation of smoking and the tobacco industry. *Tob Control.* 2008; 17:25–31. [PubMed: 18218803]
84. Balbach ED, Glantz SA. Tobacco control advocates must demand high-quality media campaigns: the California experience. *Tob Control.* 1998; 7:397–408. [PubMed: 10093175]
85. Ibrahim JK, Glantz SA. Tobacco industry litigation strategies to oppose tobacco control media campaigns. *Tob Control.* 2006; 15:50–8. [PubMed: 16436406]
86. Ibrahim JK, Glantz SA. The rise and fall of tobacco control media campaigns, 1967–2006. *Am J Public Health.* 2007; 97:1383–96. [PubMed: 17600257]
87. McDaniel PA, Malone RE. The role of corporate credibility in legitimizing disease promotion. *Am J Public Health.* 2009; 99:452–61. [PubMed: 19106419]
88. Mandel LL, Bialous SA, Glantz SA. Avoiding “truth”: tobacco industry promotion of life skills training. *J Adolesc Health.* 2006; 39:868–79. [PubMed: 17116518]
89. Thomson G, Wilson N. Directly eroding tobacco industry power as a tobacco control strategy: lessons for New Zealand? *N Z Med J.* 2005; 118:U1683. [PubMed: 16224507]
90. Smith EA, Malone RE. Altria means tobacco: Philip Morris's identity crisis. *Am J Public Health.* 2003; 93:553–6. [PubMed: 12660196]
91. Borland R. A strategy for controlling the marketing of tobacco products: a regulated market model. *Tob Control.* 2003; 12:374–82. [PubMed: 14660771]
92. Gilmore AB, Branston JR, Sweanor D. The case for OFSMOKE: how tobacco price regulation is needed to promote the health of markets, government revenue and the public. *Tob Control.* 2010; 19:423–30. [PubMed: 20876078]
93. Smith EA, Malone RE. Thinking the “unthinkable”: why Philip Morris's considered quitting. *Tob Control.* 2003; 12:208–13. [PubMed: 12773733]

What this paper adds

- Tobacco industry denormalisation (TID), a focus of successful tobacco control initiatives in the United States and elsewhere, is increasingly regarded as essential to effectively addressing tobacco at the population level. However, TID is not an established indexing term and its effects are challenging to measure. No previous reviews have examined the existing literature on the effectiveness of TID as a tobacco control strategy.
- This review analyzes the evidence on TID's effects on smoking prevalence, smoking initiation, intention to smoke, intention to quit, attitudes toward the tobacco industry and its regulation and other outcomes.
- Robust evidence, summarised here for the first time, shows that TID is an effective tobacco control intervention at the population level.




Figure 1. Derivation of 60 articles reviewed from PubMed and Scopus search. The following search strategies were employed: ('Tobacco Industry'[Mesh] OR 'tobacco industry') AND (delegitimization OR delegitimation OR denormalization OR de-normalization OR de-normalisation OR anti-industry OR counter-industry OR vilification OR industry manipulation); 'tobacco industry' AND (deception* OR mistrust* OR lie* OR lying OR false allegation); ('tobacco industry'[Mesh] OR 'tobacco industry' OR tobacco) AND (countermarketing OR counteradvertising); ('tobacco industry'[Mesh] OR 'tobacco industry' OR tobacco OR smoking) AND 'truth campaign'; ('tobacco industry'[Mesh] OR 'tobacco industry' OR tobacco OR smoking) AND (truth OR 'truth campaign'); and ('tobacco industry'[Mesh] OR 'tobacco industry') AND (opinion* OR perception* OR belie* OR 'support for action' OR attitude*).

Table 1

Characteristics of included studies

Author, year, country	n	Population	Design	Outcomes measured
Ashley and Cohen, 2003, Canada ⁵⁴	1607	Adult: 18+ years	Cross-sectional	Attitudes towards tobacco industry (TI)
Austin <i>et al</i> , 2005, USA ²³	119	Youth: 15–25 years	Controlled before and after	Current smoking prevalence [‡] ; intent to smoke [‡] ; perceived peer prevalence
Bauer <i>et al</i> , 2000, USA ¹⁵	20 978–23 745	Youth: 12–17 years	Repeated cross-sectional (three waves)	Current smoking prevalence [§] ; intent to smoke [¶]
Carver <i>et al</i> , 2003, USA ⁶²	800, 790	Adult: 18+ with children	Historically controlled	Attitudes towards TI
Cowell <i>et al</i> , 2009, USA ³⁷	31 785	Youth: 12–17 years	Repeated cross-sectional (seven waves)	Intent to smoke ^{**}
Danishevski <i>et al</i> , 2008, Russia ⁵⁸	1600	Adult: 18+ years	Cross-sectional	Attitudes towards TI
Davis <i>et al</i> , 2007, USA ⁵⁰ Farrelly <i>et al</i> , 2009 ³⁵	35 074	Youth: 12–17 years	Repeated cross-sectional (eight waves)	Intent to smoke; perceived peer smoking prevalence
Davis <i>et al</i> , 2009, USA ²⁷	16 327	Youth: 12–17 years	Longitudinal	Smoking initiation; intent to smoke [¶]
Dietz <i>et al</i> , 2008, USA ⁴⁵ Delva <i>et al</i> , 2009 ⁴⁴	2374	Adult: 18+ years	Cross-sectional	Intent to quit smoking
Dietz <i>et al</i> , 2010, USA ¹⁶	14 400	Youth: 12–17 years	Repeated cross-sectional (eight waves)	Current smoking prevalence [‡]
Dixon <i>et al</i> , 2001, Australia ⁴²	323 prior to study, 266 post study	Adult/youth: 15–60 years	Controlled before and after	Intent to smoke ^{‡‡} ; attitudes towards TI
Dunn <i>et al</i> , 2004, USA ³²	852	Youth: 15–17 years	Cross-sectional	Intent to smoke [*] ; empowerment
Dunn and Pirie, 2005, USA ⁵³	940	Youth: 12–25 years	Cross-sectional	Empowerment
Durkin <i>et al</i> , 2005, Australia ⁵⁵	1995–3001	Adult: 18+ years	Repeated cross-sectional (three waves)	Attitudes towards TI
Edwards <i>et al</i> , 2007, Australia ⁴³	3091	Youth: 12–25 years	Historically controlled	Intent to smoke ^{‡‡}
Evans <i>et al</i> , 2002, USA ⁷⁰ Evans <i>et al</i> , 2005 ⁷¹	2306	Youth: 12–17 years	Repeated cross-sectional (two waves)	Structural equation modelling
Evans <i>et al</i> , 2004, USA ⁷²	10 412	Youth: 12–17 years	Repeated cross-sectional (three waves)	Structural equation modelling
Farrelly <i>et al</i> , 2002, USA ²⁸	3439–6233	Youth: 12–17 years	Repeated cross-sectional (two waves)	Intent to smoke ^{**}
Farrelly <i>et al</i> , 2005, USA ²²	43 922	Youth: 12–17 years	Historically controlled	Current smoking prevalence [‡]
Farrelly <i>et al</i> , 2009, USA ²⁸	8904	Youth: 12–17 years	Longitudinal	Smoking initiation; intent to smoke ^{**}
Hammond <i>et al</i> , 2006, Canada/US/UK/ Australia ⁸ , Young <i>et al</i> , 2007 ⁴⁶	8222–9058	Adult: 18+ years, smoker [*]	Repeated cross-sectional (three waves)	Intent to quit; attitudes towards TI

Author, year, country	n	Population	Design	Outcomes measured
Henriksen and Fortmann, 2002, USA ⁶⁶	218	Young adult: 18–25 years	Randomised controlled trial	Attitudes towards TI
Hersey <i>et al</i> , 2003, USA ⁶⁸	6875	Youth: 12–25 years	Cross-sectional	Structural equation modelling
Hersey <i>et al</i> , 2005, USA ¹⁹	34 946	Youth: 12–17 years	Repeated cross-sectional (five waves)	Current smoking prevalence [‡]
Hersey <i>et al</i> , 2005, USA ⁶⁷	16 464	Youth: 12–17 years	Repeated cross-sectional (three waves)	Structural equation modelling
Hudson <i>et al</i> , 2007, New Zealand ⁶⁵	10	Adult: 18 years	Qualitative interviews	Attitudes towards TI
Johnson <i>et al</i> , 2008, USA ⁶⁰	28	Youth: 12–14 years	Qualitative focus groups	Attitudes towards TI
Kim and Nam, 2005, USA ⁶³	22	Adult: 18 years [*] , smoker	Qualitative focus groups	Attitudes towards TI
King <i>et al</i> , 2007, USA ⁶¹	410	Adult: 18+ years [*]	Cross-sectional	Attitudes towards TI
Klesges <i>et al</i> , 2009, USA ²⁶	36 013 prior to study, 20 672 post study	Adult: Air Force Recruits	Longitudinal	Current smoking prevalence [§] ; smoking initiation
Leatherdale <i>et al</i> , 2006, Canada ²¹	14 767	Youth: 14–18 years	Cross-sectional	Current smoking prevalence [§]
Ling <i>et al</i> , 2007, USA ²⁴	9455	Young adult: 18–29 years	Cross-sectional	Current smoking prevalence [‡] ; intent to quit; intent to smoke ^{**}
Ling <i>et al</i> , 2009, USA ²⁵	1731	Young adult: 18–29 years	Cross-sectional	Current smoking prevalence ^{‡‡‡} ; intent to quit
Murphy-Hoefer <i>et al</i> , 2008, USA ⁴⁷ Murphy-Hoefer <i>et al</i> , 2010, USA ⁴⁸	1011	Young adult: 18–24 years	Controlled experiment	Intent to quit
Netemeyer <i>et al</i> , 2005, USA ⁴⁹	1207	Adult: 18+ years, smoker	Cross-sectional	Intent to quit
Niederdeppe <i>et al</i> , 2004, USA ¹⁷	7478	Youth: 12–17 years	Repeated cross-sectional (three waves)	Current smoking prevalence [‡] ; intent to smoke ^{**}
Niederdeppe <i>et al</i> , 2008, USA ³¹	5010	Youth: 12–18 years	Repeated cross-sectional (five waves)	Intent to smoke [‡]
Pechmann <i>et al</i> , 2003, USA ³⁸	1667	Youth: 15–18 years	Randomised controlled trial	Intent to smoke [*] ; resistance self-efficacy
Pechmann <i>et al</i> , 2006, USA ³⁹	1725	Youth: 14–15 years	Randomised controlled trial	Intent to smoke [*] ; empowerment
Reinert <i>et al</i> , 2010, USA ⁶⁴	53	Adult: School nurses	Cross-sectional	Attitudes towards TI
Richardson <i>et al</i> , 2010, USA ⁴¹	19 701	Young adult: 18–24 years	Repeated cross-sectional (eight waves)	Intent to smoke ^{**} ; intent to quit
Shadel <i>et al</i> , 2009, USA ⁵¹ Shadel <i>et al</i> , 2010, USA ⁵²	110	Youth: 11–17 years	Controlled comparison	Resistance self-efficacy
Sly <i>et al</i> , 2000, USA ⁵⁹	1816	Youth: 12–17 years	Cross-sectional	Attitudes towards TI
Sly <i>et al</i> , 2001, USA ³⁰	1820	Youth: 12–17 years	Repeated cross-sectional (two waves)	Smoking initiation

Author, year, country	n	Population	Design	Outcomes measured
Sly <i>et al</i> , 2001, USA ¹⁸	Approximately 1800	Youth: 12–17 years	Repeated cross-sectional (four waves)	Current smoking prevalence [§]
Sly <i>et al</i> , 2002, USA ²⁹	1805	Youth: 12–17 years	Repeated cross-sectional (six waves)	Smoking initiation
Sly <i>et al</i> , 2005, USA ³³	1079–1150	Youth: 12–17 years	Repeated cross-sectional (four waves)	Intent to smoke ^{**}
Sutfin <i>et al</i> , 2008, USA ⁴⁰	488	Youth: 15–17 years	Randomised controlled trial	Intent to smoke [*]
Tangari <i>et al</i> , 2007, USA ³⁴	1208 adult, 900 youth	Adult/youth	Cross-sectional	Intent to smoke [*] ; intent to quit
Thrasher and Jackson, 2006, USA ⁶⁹	6670	Youth: 12–17 years	Repeated cross-sectional (two waves)	Structural equation modelling
Thrasher <i>et al</i> , 2006, USA ²⁰	10 035	Youth: 12–17 years	Repeated cross-sectional (two waves)	Current smoking prevalence [†] ; intent to smoke [*]
Wakefield <i>et al</i> , 1999, Australia ⁵⁶	808	Adult: 18+ years	Cross-sectional	Attitudes towards TI
Waller <i>et al</i> , 2004, Canada ⁵⁷	10 434, youth, 1607 adult	Adult/youth: 12–18 years	Repeated cross-sectional (three waves)	Attitudes towards TI
Zucker <i>et al</i> , 2000, USA ¹⁴	1247–1806	Youth: 12–17 years	Repeated cross-sectional	Current smoking prevalence

* Intent to smoke measured with three validated items on a five-point Likert scale: ‘in the future, you might smoke one puff or more of a cigarette’; ‘you might try out cigarette smoking for a while’; ‘if one of your best friends were to offer you a cigarette you would smoke it’.

† Current smoking prevalence measured using 30 day referent.

‡ Intent to smoke measured with one item on a four-point or seven-point scale: ‘if your best friend offered you a cigarette, would you smoke it?’.

§ Current smoking prevalence measured on 3–5 classification continuum: never smoker, experimenter, ever smoker, current smoker, former smoker.

¶ Intent to smoke measured with four items on a five-point scale: ‘do you think you will smoke a cigarette soon?’; ‘do you think you will smoke a cigarette anytime during the next year?’; ‘do you think you will be smoking cigarettes 5 years from now?’; ‘if one of your best friends offered you a cigarette would you smoke it?’.

** Intent to smoke measured with one item on a four-point Likert scale: ‘you will smoke a cigarette in the next year’.

†† Intent to smoke measured with one item on a seven-point scale: ‘do you think you will be smoking cigarettes this time next year?’.

††† Current smoking prevalence measured using 100-cigarettes referent.