
Isolation and mapping of a polymorphic DNA sequence (pTHI54) on chromosome 1p [D1S62]

T.Holm, Y.Nakamura, L.Ballard, M.Leppert, P.O'Connell, G.M.Lathrop, J.-M.Lalouel and R.White

The Howard Hughes Medical Institute, University of Utah Medical School, Salt Lake City, UT 84132, USA

SOURCE/DESCRIPTION: A 13.0 kb HindIII fragment from the phage THI54 isolated with the insulin oligonucleotide (1) was subcloned into the HindIII site of pBR322.

POLYMORPHISM: PvuII identifies a 2 allele polymorphism (P1 : 6.0 kb, P2 : 5.0 kb). BamHI, BglIII, EcoRI, HindIII, MspI, PstI, PvuII, RsaI, and TaqI also detect the same polymorphism.

FREQUENCY: Estimated from 92 unrelated Caucasians.
PvuII P1 : 0.51
P2 : 0.49

NOT POLYMORPHIC FOR: None detected.

CHROMOSOMAL LOCALIZATION: pTHI54 has been assigned to chromosome 1p by multipoint linkage analysis (2) with loci (FY, NRAS, NGFB, RH) known to span this region (3,4).

MENDELIAN INHERITANCE: Co-dominant segregation of this polymorphism was observed in 46 three generation families.

PROBE AVAILABILITY: Contact Y.N.

OTHER COMMENTS: RFLPs were observed under normal conditions of hybridization and washing.

REFERENCES:

1. Y. Nakamura et al., Science **235**: 1616-1622 (1987)
2. G.M. Lathrop et al., Am. J. Hum. Genet. **37**:482-498 (1985)
3. S. Povey et al., Cytogenet. Cell Genet. **40**:67-106(1985)
4. P. O'Connell et al., abstract submitted to the Human Gene Mapping Workshop 9