

Damaged Mitochondrial DNA Replication System and the Development of Diabetic Retinopathy

Shikha Tewari, Julia M. Santos, and Renu A. Kowluru

Abstract

Aim: In the pathogenesis of diabetic retinopathy, retinal mitochondria are damaged, superoxide levels are elevated, and mitochondrial DNA (mtDNA) biogenesis is impaired. mtDNA has a noncoding region, displacement loop (D-loop), which has essential transcription and replication elements, and this region is highly vulnerable to oxidative damage. The aim of this study is to investigate the effect of diabetes on the D-loop damage and the mtDNA replication machinery. **Results:** Using retina from wild-type (WT) and mitochondrial superoxide dismutase transgenic (Tg) mice, we have investigated the effect of diabetes on retinal D-loop damage and on the replication system. The results were confirmed in the isolated retinal endothelial cells in which the DNA polymerase gamma 1 (*POLG1*) function was genetically manipulated. Diabetes damaged retinal mtDNA, and the damage was more at the D-loop region compared with the cytochrome B region. Gene transcripts and mitochondrial accumulation of *POLG1*, *POLG2*, and mtDNA helicase, the enzymes that form replisome to bind/unwind and extend mtDNA, were also decreased in WT-diabetic mice compared with WT-normal mice. Tg-diabetic mice were protected from diabetes-induced damage to the D-loop region. Overexpression of *POLG1* prevented high glucose-induced D-loop damage. This was accompanied by a decrease in mitochondrial superoxide levels. **Innovation and Conclusions:** Integrity of the retinal D-loop region and the mtDNA replication play important roles in the mtDNA damage experienced by the retina in diabetes, and these are under the control of superoxide. Thus, the regulation of mtDNA replication/repair machinery has the potential to prevent mitochondrial dysfunction and the development of diabetic retinopathy. *Antioxid. Redox Signal.* 17, 492–504.

Introduction

HYPERGLYCEMIA INDUCES DAMAGE to the retinal microvasculature, resulting in the development of diabetic retinopathy, and oxidative stress is considered to play a major role (13, 16, 17, 26, 38, 40). Mitochondrial superoxide production acts as a single unifying mechanism for diabetic complications (3). We have shown that in diabetes, increased reactive oxygen species (ROS) impair mitochondria function, damage mitochondrial DNA (mtDNA), and accelerate the apoptosis of retinal capillary cells, and overexpression of mitochondrial superoxide dismutase (MnSOD) protects the damage of retinal mtDNA and the development of diabetic retinopathy (13, 17, 26, 25, 38).

MtDNA is a double-stranded circular molecule of 16,569 bp, and because of lack of protective histones, it is 10–20 times more vulnerable to oxidative damage than nuclear DNA (nDNA) (24, 33, 43). In the pathogenesis of diabetic retinopathy, retinal mtDNA damage is increased, mitochondria copy numbers are decreased, and the genes encoded by mtDNA become subnormal (25, 26, 38). MtDNA has a non-

coding region, the displacement loop (D-loop), which contains essential transcription and replication elements. The D-loop helps in the mtDNA replication by serving as a promoter for heavy strands (4). This, mainly unwound region, is highly vulnerable to ROS, and provides control sites for mtDNA

Innovation

In the pathogenesis of diabetic retinopathy, retinal mitochondria are dysfunctional, their copy numbers are decreased, and DNA is damaged. Here, we show that the damage to the mitochondrial DNA (mtDNA) is not uniform; the displacement loop region, the site for mtDNA replication and transcription, experiences more damage as compared with the cytochrome B, and the mtDNA replication/repair system is compromised. Targeting mtDNA replication/repair machinery has the potential to regulate mitochondria homeostasis, and to inhibit the development of diabetic retinopathy.

replication (28, 35, 41). Frequent mutations in the D-Loop regions are observed in various diseases including adenocarcinomas and other cancers (28, 32, 44). How diabetes affects the D-loop in the retina remains to be explored.

The maintenance/replication of the mtDNA is critical for the mitochondria to function properly, and mitochondria are equipped with excellent replication machinery (9, 12). The nuclear-encoded mtDNA replication enzymes, DNA polymerase gamma (POLG) (with two subunits-POLG1 and POLG2), and mtDNA helicase (Twinkle) form the minimal replisome required for replication (15, 39). POLG1 has two enzymatic activities: a C-terminal polymerase and an N-terminal "proofreading" exonuclease, and Twinkle serves as a DNA unwinding enzyme (42, 45). Mutations in POLG and Twinkle have been reported in patients with mitochondrial disorders; however, their role in the pathogenesis of diabetic retinopathy remains unclear.

The aim of this study is to investigate the effect of diabetes on the damage to the D-loop region, and on the replication/repair machinery in the retina. Our hypothesis is that in diabetes due to the damaged D-loop, the mtDNA replication system is impaired, and this plays a major role in retinal


mtDNA damage. Due to increased susceptibility of mtDNA to ROS, we have investigated D-loop damage in the diabetic mice in which superoxide levels are genetically manipulated by overexpressing MnSOD (Tg). To examine the effect of hyperglycemia specifically on retinal vascular cells, key parameters are investigated in the isolated retinal endothelial cells, the cells that demonstrate a histopathology characteristic of diabetic retinopathy, in which *POLG1* is manipulated.

Results

Diabetes damages retinal mitochondrial D-loop and impairs the replication machinery

Amplifications of D-loop, normalized to that of β -actin of nDNA, was reduced by $\sim 35\%$ in the retina of wild-type (WT) diabetic mice compared with WT-normal mice (Fig. 1a). Consistent with our previous report (38), the amplification of the cytochrome B (CytB), a marker of mitochondria copy number (27), normalized to that of β -actin, was also reduced by $\sim 50\%$ (Fig. 1b). Similarly, the ratio of semi-long fragment (~ 830 bp) to short fragment (~ 103 bp) of the D-loop region was decreased by 30% (Fig. 1c), but the ratio of 800 to 100 bp of

FIG. 1. Effect of diabetes on copy numbers and semi-long/short PCR fragments of D-loop and CytB regions. SYBR green PCR was performed in mouse retinal DNA (10 ng) using specific primers of ~ 100 bp for amplification of D-loop and of CytB regions of mtDNA, and copy numbers were calculated by ddCt method. Relative copy number of (a) D-loop region and (b) CytB region was calculated by normalizing their respective values with the nuclear gene, β -actin. For semi-long/short PCR fragments, semi-long fragment (800–900 bp) and small fragment (~ 100 bp) of D-loop or CytB regions were amplified using specific primers. The PCR products were analyzed on agarose gel, and the ratio of long/short PCR amplicons in (c) D-loop region and (d) CytB region were calculated. (e) 830 and 103 bp PCR amplicons at the D-loop region were analyzed in the retinal microvessels prepared by osmotic shock method, but without DNase treatment (18). (f) Linear amplification of 830 and 103 bp PCR amplicons at the D-loop region was determined using 4–12 ng DNA. Each measurement was made in duplicate in six or more mice in each group, and the values are represented as mean \pm SD. WT and WT-D = Wild-type normal and diabetic mice, respectively, Tg and Tg-D = MnSOD overexpressing normal and diabetic mice respectively; 4, 6, 8, 10, 12 = 4–12 ng DNA template. * $p < 0.05$ compared to WT and # $p < 0.05$ to WT-D. CytB, cytochrome B; D-loop, displacement loop; MnSOD, mitochondrial superoxide dismutase; mtDNA, mitochondrial DNA; PCR, polymerase chain reaction.



the CytB region remained unchanged (Fig. 1d), suggesting that the D-loop region experiences higher mutation/damage in diabetes compared with the CytB region. Consistent with the retinal D-loop damage, its microvasculature also showed increased D-loop damage in diabetes (Fig. 1e).

To explore the possible mechanism for increased damage at the D-loop region, gene and protein (total and mitochondrial) expressions of mtDNA replication enzymes were quantified. Diabetes decreased the gene and protein expressions of retinal

POLG1, POLG2, and Twinkle by 35%–45%, and decreased their mitochondrial accumulation, the site of their action (Fig. 2a–c).

MnSOD protects the retina from diabetes-induced damage to mtDNA and replication machinery

As shown in Figure 1a, b, MnSOD overexpression ameliorated diabetes-induced decrease in the amplification of both D-loop and CytB regions, and the values from




FIG. 2. Effect of diabetes on mtDNA replication machinery. Gene expressions of (a) *POLG1*, (b) *POLG2*, and (c) *Twinkle* were measured in the retina by qPCR. Ct values were normalized to β -actin for each sample. Western blot technique was used to quantify the expressions of *POLG1*, *POLG2*, and *Twinkle* in the (b) whole homogenate and (c) mitochondria using β -actin and CoxIV, respectively, as loading proteins. Cytosolic and nuclear contaminations in the mitochondrial fraction were detected by quantifying the expressions of lactate dehydrogenase (LDH) and H2B, respectively. (d) Retinal H₂O₂ levels were quantified fluorimetrically by the Amplex Red Hydrogen Peroxide/Peroxidase Assay kit using 120 μ g retinal protein. Values are represented as mean \pm SD from 5 to 6 mice in each group, each measurement made in duplicate. * p < 0.05 compared with WT and # p < 0.05 compared with WT-D. POLG, DNA polymerase gamma; Twinkle, mtDNA helicase; H2B, histone 2-B.

Tg-diabetic mice were significantly higher. Similarly, Tg-diabetic mice were also protected from D-loop damage (Fig. 1c, d), and the expression of *POLG1* and *POLG2* was significantly higher compared with the values obtained from WT-diabetic mice (Fig. 2a–c). Figure 2d shows that overexpression of MnSOD did not have any effect on retinal H₂O₂ levels.

Retinal endothelial cells

Abnormalities in retinal mtDNA replication machinery were confirmed in bovine retinal endothelial cells (BREC)s. As shown in Figure 3a–c, high glucose decreased transcripts of *POLG1*, *POLG2*, and *Twinkle* by 35%–75%, and their mitochondrial expressions were significantly decreased, compared with the values obtained from cells exposed to normal glucose ($p < 0.05$; Fig. 3). Scavenging of superoxide by MnSOD mimic manganese (III) tetrakis (4-benzoic acid) por-

phyrin chloride (MnTBAP), ameliorated glucose-induced decrease in the transcripts of *POLG1*, *POLG2*, and *Twinkle* and their mitochondrial accumulation compared with the cells incubated in high glucose alone (Fig. 3). Figure 4a shows 50% less *POLG1* immunostaining in the mitochondria region in cells incubated in 20 mM glucose compared with 5 mM glucose, confirming decreased *POLG1* mitochondrial localization.

To evaluate the effect of high glucose on functionally active *POLG*, newly synthesized mtDNA strands were quantified by incorporation of bromodeoxyuridine (5-bromo-2'-deoxyuridine) (BrDU). Addition of aphidicolin, a selective inhibitor of DNA polymerase α , resulted in BrDU-positive fluorescence in the mitochondria, suggesting the presence of newly synthesized mtDNA. However, exposure of cells to 20 mM glucose decreased BrDU-positive fluorescence by ~35%. As a control, cells in 5 mM glucose with an inhibitor of




FIG. 3. Effect of high glucose on the replication machinery and its amelioration by MnSOD mimic. Cells incubated in high glucose in the presence or absence of MnTBAP (200 μ M) were analyzed for gene expression (qPCR) and mitochondrial accumulation (by western blot technique) (a) *POLG1*, (b) *POLG2*, and (c) *Twinkle*. Each measurement was made in duplicate using cells from 3 to 5 different cell preparations, and the values are represented as mean \pm SD. 5 = 5 mM glucose, 20 = 20 mM glucose, MnT = 20 mM glucose + 200 μ M MnTBAP, and Mann = 20 mM mannitol. * $p < 0.05$ compared with 5 mM and # $p < 0.05$ compared with 20 mM glucose. MnTBAP, manganese (iii) tetrakis (4-benzoic acid) porphyrin chloride.




FIG. 4. Effect of high glucose on the POLG1 localization in the mitochondria and on its own function. **(a)** Localization of POLG1 in the mitochondria was evaluated using MitoTracker (green) for mitochondria and Texas Red-conjugated secondary antibody for POLG1. The cover slips were mounted using Vectashield containing DAPI (nucleus-blue), and examined under a Zeiss ApoTome using 40 \times magnification. Localization of POLG (green) in the mitochondria (red) was calculated using Axio vision software. **(b)** Functional assay for POLG1 was performed immunohistochemically by quantifying the incorporation of BrDU into mtDNA. The histograms represent mean \pm SD of 5–7 replicates. 5 = 5 mM glucose; 20 = 20 mM glucose, 5 mM + aphid or 20 mM + aphid = 5 or 20 mM glucose with 7 μ M aphidicolin; 5 mM + EtBR = 5 mM glucose with 1 μ g/ml EtBr. * p < 0.05 compared with 5 mM glucose. EtBR, ethidium bromide; BrDU, bromodeoxyuridine (5-bromo-2'-deoxyuridine).

POLG, ethidium bromide (EtBr), showed BrDU incorporation only in the nDNA (Fig. 4b).

To assess the binding of the D-loop to *POLG1*, chromatin immunoprecipitation (ChIP) of *POLG1* at the D-loop was performed, and Figure 5 shows that high glucose decreases the binding by 50%. Inclusion of MnTBAP prevented glucose-induced loss in the binding capacity of *POLG1*, the values obtained from the cells in high glucose supplemented with MnTBAP were similar to those from normal glucose.

Effect of genetic manipulation of *POLG1*

To further establish the role of *POLG1* in mtDNA damage, *POLG1* was silenced in BRECs using *POLG1*-siRNA. mRNA levels and mitochondrial expression of *POLG1* were 30%–35% lower in the cells transfected with *POLG1*-siRNA compared with the untransfected cells or cells transfected with scrambled RNA (Fig. 6a). *POLG1*-siRNA also increased D-loop damage, and this was similar to that observed in the untransfected cells exposed to high glucose (Fig. 6b). Similarly, the relative amplification of long (13.3 kb) to short (234 bp) fragments of mtDNA was also significantly decreased. The values obtained from the transfected cells in 5 mM glucose were similar to those obtained from untransfected cells in 20 mM glucose, but were significantly different from the untransfected cells in 5 mM glucose (Fig. 6c).

In contrast, when *POLG1* was overexpressed, glucose-induced D-loop damage, as detected by the ratio of semi-

long/short (891/91 bp) fragments, was prevented (p < 0.05; Fig. 7a), and *POLG1* levels in the mitochondria were increased (Fig. 7b). Damage of mtDNA was also prevented (Fig. 7c), suggesting a major role of *POLG1* in mtDNA replication and repair machinery. Glucose-induced increase in mitochondria superoxide levels, as evidenced by increased





FIG. 5. Effect of high glucose on the binding of *POLG1* to D-loop. Retinal cells were crosslinked, and DNA pull-down assay was performed using a *POLG1* antibody. PCR for the D-loop region in the Chip DNA was performed, and the Ct values were normalized to input DNA. Values represent 5 or more replicates in each group. * p < 0.05 compared with 5 mM and # p < 0.05 compared with 20 mM glucose.

FIG. 6. Regulation of mtDNA and D-loop damage by *POLG1* in endothelial cells. BRECs transfected with *POLG1*-siRNA or scrambled RNA were incubated in 5 or 20 mM glucose for 4 days. **(a)** *POLG1* mRNA levels were measured by qPCR, and its mitochondrial accumulation was quantified by western blot using Cox IV as loading control. **(b)** PCR for semi-long (800–900 bp) and short fragments (100 bp) of the D-loop was performed as described in Figure 1. **(c)** Extended length PCR for mtDNA and nDNA were performed by the semi-quantitative PCR method. Each measurement was made in duplicate in the cells obtained from 3 to 5 different cell preparations, and the values are represented as mean \pm SD. 5=5 mM glucose, 20=20 mM glucose, si5 and si20=*POLG1*-siRNA transfected cells incubated in 5 or 20 mM glucose, SC=Scrambled RNA transfected cells. * p <0.05 compared with 5 mM and # p <0.05 compared with 20 mM glucose. BREC, bovine retinal endothelial cell; Cox IV, cytochrome oxidase IV; nDNA, nuclear DNA.



immunostaining of MitoTracker Red, and by quantitative fluorometric method, was ameliorated by *POLG1* overexpression (Fig. 8a, b).

Discussion

In the development of diabetic retinopathy, mtDNA is damaged and mitochondria copy numbers and mass are decreased, and overexpression of MnSOD prevents these diabetes-induced retinal abnormalities (13, 17, 25, 26, 38). The maintenance/replication of the mtDNA has a critical role in mitochondria function, and the D-loop region contains the binding sites for major factors for mtDNA transcription. Here, we present exciting data to show that diabetes decreases the copy number of the D-loop region in the retina, and this region encounters more damage than other regions of the mtDNA. To make the bad situation worse, mtDNA replication enzymes become subnormal, hampering the replication/repair process. Regulation of mitochondrial superoxide levels by MnSOD ameliorates

diabetes-induced damage to the D-loop, and protects a decrease in mtDNA replication enzymes, suggesting that the D-loop damage and the replication of mtDNA are under the control of mitochondrial superoxide. Consistent with the results from the retina, high glucose also preferentially damages the D-loop region and the replication machinery in the retinal endothelial cells, the site of histopathology, and the binding of *POLG1* to the D-loop is also compromised. Thus, our results strongly support a crucial role of the D-loop integrity and the mtDNA replication in the development of diabetic retinopathy.

The D-loop region has a control site for mtDNA replication containing the leading strand origin of replication (23, 28, 35, 41), and it regulates transcription and replication of mtDNA. Variations in the D-loop region are associated with a number of diseases, including cancer, leukemia, diabetes, and diabetic neuropathy (28, 31, 32, 40, 44). We have shown that despite increased gene transcripts of DNA repair enzymes, retinal mtDNA is damaged in diabetes (20, 25, 26, 38), and here we present results showing increased D-loop damage compared




FIG. 7. Effect of overexpression of *POLG1* on regulation of D-loop and mtDNA damage. *POLG1*-transfected BRECs were incubated in high glucose for 4 days, and analyzed for (a) D-loop damage (PCR for semi-long and short fragments), (b) for mitochondrial accumulation by western blot technique, and for (c) mtDNA damage by extended-length PCR. Values are represented as mean \pm SD from cells obtained from 3 to 4 preparations, each measurement was made in duplicate. 5=5 mM glucose, 20=20 mM glucose, 5 and 20 *POLG*=cells overexpressing *POLG1* and incubated in 5 or 20 mM glucose, respectively. * $p < 0.05$ compared to 5 mM and # $p < 0.05$ to 20 mM glucose. R, reagent control cells in 5 mM glucose.

with the CytB region. These results suggest that the maintenance of the D-loop region is important in the pathogenesis of diabetic retinopathy.

POLG plays a crucial role in the maintenance of the mtDNA and function (9, 12). Overexpression of *POLG1* is associated

with increased expression of a mitochondrial transcriptional factor, TFAM, important in mitochondria biogenesis (5). In diabetes, the levels of TFAM are decreased in the retinal mitochondria, resulting in impaired mtDNA biogenesis and subnormal mtDNA copy number (36, 38). Furthermore, diabetes also decreases the number of mitochondria in the retinal vasculature (20, 36, 38, 48). Decreased expression of *POLG1* and *POLG2* in the retina in diabetes, thus, strongly suggests that impaired replication machinery further contributes to the decrease in the overall mitochondria number.


For replication of mtDNA, in addition to *POLG*, Twinkle also plays a significant role by serving as a replicative helicase (15). *POLG* and Twinkle together form replisome to bind, unwind, and extend mtDNA, and this process utilizes short stretches of RNA (RNA primers) synthesized by RNA polymerase to make a new complementary mtDNA strand (7, 21). Mutations in *Twinkle* and *POLG* genes have been reported to predispose mtDNA to multiple deletions, resulting in development of mitochondrial diseases, including progressive external ophthalmalgia, a condition with cytochrome C oxidase-deficient fibers (34, 46). Here, we show that diabetes significantly decreases the expression and mitochondrial accumulation of Twinkle, raising the possibility that the decreased mitochondrial levels could also be due to an impaired mitochondrial transport system, which the retina experiences in diabetes (48).

The development of diabetic retinopathy and abnormalities in mitochondria dysfunction are prevented in diabetic mice overexpressing MnSOD (13, 19, 30, 38). Our results clearly demonstrate that diabetes-induced damage to the D-loop and abnormalities in the replication machinery are also ameliorated in Tg-diabetic mice. In support, others have shown protection of UV-induced *POLG* damage by MnSOD (2). Furthermore, MnSOD overexpression has been shown by us to prevent decrease in other mtDNA repair glycosylases, TFAM, and mtDNA copy numbers (25, 26, 38). Thus, our results strongly suggest that the replication process is under the control of superoxide, and the prevention of the development of diabetic retinopathy in Tg mice could be due to protection of *POLG*. However, the role of NADPH oxidase-induced increase in retinal ROS in diabetes (1) cannot be ruled out.

The retina has multiple layers and cell types, and also a specialized vasculature that provides nutrients without interfering with light sensing (40). Retinal vasculature is the major sight of histopathology of diabetic retinopathy, and endothelial cells and pericytes are lost (8). To specifically understand the role of the mtDNA replication in the pathogenesis of diabetic retinopathy, our results from retinal microvessels and also isolated retinal endothelial cells confirm similar damage as observed in the retina from diabetic rodents. Subnormal mitochondria biogenesis and decreased copy number is also observed in retinal endothelial cells exposed to high glucose (36, 38). Furthermore, inhibition of mitochondrial superoxide accumulation by MnSOD mimic also prevents damage to the replication machinery, and this further confirms its importance in the development of diabetic retinopathy.

The balance between mtDNA repair and damage is important for mitochondria homeostasis, and oxidative stress-induced damage to *POLG* in human hepatoblast cells results in preferential D-loop damage (9, 11, 14, 47). Down-regulation of retinal *POLG* in diabetes suggests that the repair

FIG. 8. Effect of overexpression of *POLG* on mitochondrial superoxide. Mitochondrial superoxide levels were determined by MitoTracker Red (CM-H₂XROS) using both immunostaining and quantitative fluoremetric method. **(a)** For immunostaining, the cells were incubated with MitoTracker Red, fixed, mounted using Vectashield containing DAPI, and examined under a Zeiss ApoTome using 40 \times magnification (each picture is representative of 4–5 images). **(b)** For the quantitative method, the cell homogenate was incubated with MitoTracker Red, and the resultant fluorescence was quantified at 579 nm excitation and 599 nm emission wavelengths. Measurements in each group are represented as mean \pm SD obtained from 3 to 4 cell preparations, and each measurement is made at least in duplicate. * p < 0.05 compared with 5 mM glucose.



of oxidative lesions of mtDNA is compromised, resulting in increased accumulation of oxidatively modified DNA. In support, diabetes increases the levels of oxidatively modified DNA in the retinal mitochondria (25, 26). *POLG* also plays an important role in mtDNA stability by interacting with DNA ligase III (6), and others have shown that the expression of dominant negative DNA polymerase depletes mtDNA (11). Here, we show that the binding of *POLG1* to D-loop is compromised, further contributing to a decreased D-loop region. Consistent with this, *POLG* is the rate-limiting enzyme in mitochondrial base excision repair (BER), and its inactivation is likely to increase the buildup of cytotoxic BER pathway intermediates, including single-strand breaks (10, 24, 39). This raises the possibility that the decreased binding could be responsible for mtDNA synthesis, and could result in impaired retinal mitochondria biogenesis and subnormal copy number (36, 38, 48).

In conclusion, this is the first study demonstrating that the mtDNA presents differential damage in diabetes, the damage to the D-loop region is more extensive compared with the CytB region, and the replication/repair machinery is subnormal, further contributing to the dysfunctional mitochondria. Regulation of mitochondrial superoxide protects retinal mtDNA from such damage. Thus, regulation of mtDNA replication/repair machinery could present a plausible mechanism that prevents dysfunctional mitochondria, and the development of diabetic retinopathy, one of the most devastating complication of diabetes.

Methods

Animal models

Hemizygous MnSOD Tg mice with C57BL/6 background and their WT littermates were made diabetic by administering streptozotocin (55 mg/kg BW) for five consecutive days, a model of type I diabetes. Mice with blood glucose 250 mg/dl after 3 days of the last injection of streptozotocin were considered as diabetic. Age-matched normal WT mice and Tg mice served as controls. After \sim 10 months of diabetes, mice were euthanized by carbon dioxide narcosis, and the retina was isolated. Treatment of animals was carried out as per the guidelines of National Institute of Health Principles of Laboratory Animal Care, the Association for Research in Vision and Ophthalmology resolution on the use of animals in research, and the Institutional guidelines. Average blood glucose values for normal and diabetic mice (WT or Tg) were 125 ± 18 and 502 ± 106 mg/dl, respectively. These methods are in use in our laboratory (13, 19, 26), and as previously reported, overexpression of MnSOD prevented diabetes-induced increase in the formation of acellular capillaries in the retina (13).

Retinal endothelial cells

BRECs from 4th to 6th passage were incubated in 5 mM glucose or 20 mM glucose media for 4 days in the presence or absence of MnTBAP (200 μ M; Biomol, Plymouth Meeting, PA). The incubation media was changed every other day, and

supplemented with fresh MnTBAP (16, 26). Cells incubated in 20 mM mannitol, instead of 20 mM glucose, served as osmotic controls.

Manipulation of POLG1

To establish the role of POLG1 in regulation of mtDNA, BRECs from 3rd to 5th passage were transfected with *POLG1*-siRNA (Santa Cruz Biotechnology, Santa Cruz, CA) as previously described (20, 25). Briefly, cells were incubated with the transfection complex consisting of the siRNA and transfection reagent for 8 h at 37°C, washed with PBS, and incubated in 5 or 20 mM glucose media for 4 additional days. Parallel incubations were carried out using scrambled siRNA, and the transfection efficiency was evaluated by quantifying *POLG1* gene expression.

To investigate the effect of *POLG1* overexpression in improving glucose-induced mtDNA damage, the enzyme was overexpressed using 5 μ g *POLG1* expression plasmid (RC204456; OriGene, Rockville, MD) and TurboFectin 8.0 transfection reagent. After transfection, the cells were incubated in 5 or 20 mM glucose for 4 days. Parallel incubations were carried out using the transfection reagent alone.

Isolation of DNA and RNA

Total DNA was isolated from the retina or BRECs using Blood tissue DNA kit (Qiagen, Valencia, CA), and quantified by the Quant-iT dsDNA assay (Invitrogen, Carlsbad, CA). The primers were designed using primer express-3 software (Table 1) and synthesized by Integrated DNA Technologies (Coralville, IA).

RNA (1–2 μ g), isolated by TRIzol reagent (Invitrogen), was employed to make cDNA using High Capacity cDNA Reverse Transcription Kit (Applied Biosystem, Foster City, CA). Gene expressions for *POLG1*, *POLG2*, and *Twinkle* were measured by SYBR green-based real-time quantitative polymerase chain reaction (qPCR), with melting curve analysis on ABI 700 (Applied Biosystems) using their gene-specific primers (Table 1). *β -actin* was used as a housekeeping gene, and quantification of the transcripts was done by the ddCt method (38, 50).

Assessment of DNA damage

To evaluate mtDNA damage, three independent PCR-based methods were used. In the first method, mtDNA copy number of D-loop and that of CytB were measured by qPCR (SYBR green-based). Each reaction was performed in duplicate using 10 ng DNA and 0.5 μ M each of forward and reverse primers. The thermal conditions included 10 min at 95°C followed by 40 cycles of 15 s at 95°C and 60 s at 60°C. To assess the formation of specific PCR products, the amplicons were also analyzed on an agarose gel. The amplification of D-loop or CytB regions were normalized to that of *β -actin*, nDNA-encoded gene, and quantification was done by ddCT method.

The second method quantified the ratio of semi-long to short PCR amplicons at the D-loop and at the CytB regions of mtDNA using the primers spanning around 800–900 bp. Their expressions were normalized by their respective short fragments of ~100 bp, the fragments that are easy to amplify by DNA polymerase (35). This allowed us to analyze strand breaks in the long fragments, base modifications and also

identify bulky DNA adducts that cause an inhibition of the primer annealing and forward movement of POLG (22). Each PCR reaction was performed using GoTaq™ Reaction Buffer, 0.2 mM of dNTPs (Promega, Madison, WI), 0.2 μ M of each forward and reverse primers, GoTaq™ DNA polymerase, and 10 ng of template DNA. PCR conditions included 94°C for 5 min, 24 cycles of 94°C for 1 min, 60°C for 30 s, 72°C for 1 min, and a final extension at 72°C for 10 min. The linearity of the assay was performed using 1–12 ng DNA (Fig. 1f). The PCR products were separated on 2% agarose gel. The intensity of PCR products was measured using Un-Scan-It Gel digitizing software (26, 37, 38).

DNA damage was also evaluated in cells in which *POLG1* was either silenced or overexpressed. Briefly, 15 ng DNA was amplified using GeneAmp XL PCR kit (Applied Biosystems), 200 mM dNTP, 1.1 mM Mg(OAc)₂, 1 unit rTth DNA polymerase, and 0.1 mM primers. After 1 min at 75°C, polymerase was added followed by 1 min at 94°C, 24 cycles of 94°C for 15 s and 65°C for 12 min, and a final extension at 72°C for 12 min. PCR products were resolved on agarose gel.

Isolation of mitochondria and western blot

Mitochondria were isolated from the retina or BRECs using the Mitochondria Isolation kit from Invitrogen, as previously reported (29, 30, 38, 48). Protein content was determined by the Bicinchoninic Acid protein assay kit (Sigma-Aldrich, St. Louis, MO). The purity of the mitochondria was determined by measuring the expressions of nuclear protein, Histone 2-B, and of cytosolic protein, lactate dehydrogenase, using antibodies sc-31671 and sc-27235, respectively (Santa Cruz Biotechnology). As shown in Figure 2c, the method yielded largely purified mitochondria with little contamination of cytosolic or nuclear proteins.

Proteins (30–50 μ g) were separated on a 4%–16% SDS-PAGE, and identified using antibodies against *POLG1*, *POLG2*, and *Twinkle* (sc-48815, sc-160290, sc-134915 respectively; Santa Cruz Biotechnology). Cytochrome oxidase IV (Cox IV; Molecular Probe, Molecular Probes, Eugene, OR) was used as a loading control for mitochondria and *β -actin* for the homogenate. Routine dilutions for primary antibodies were 1:500 and for horseradish peroxidase-conjugated secondary antibodies, they were 1:2000. The band intensity of western blot was quantified using Un-Scan-It Gel digitizing software (Silk Scientific Inc., Orem, UT). The individual blots were normalized to Cox IV/ *β -actin*, and the values obtained from WT-normal mice or cells incubated in 5 mM glucose were considered as 100%.

Hydrogen peroxide

Retinal H₂O₂ levels were quantified using Amplex Red Hydrogen Peroxide/Peroxidase Assay kit (A-22188- Invitrogen). Homogenate (120 μ g) was incubated with 100 μ M Amplex red and 1 U/ml horse radish peroxidase for 30 min, and the resulting fluorescence was measured at 590/530 nm (emission and excitation). The assay was sensitive to as low as 0.01 pmol of H₂O₂.

Localization of POLG1 in the mitochondria

Localization of *POLG1* in the mitochondria was performed by fluorescence microscopy using MitoTracker green (Invitrogen) (20, 38). Briefly, the cells were incubated for 45 min

TABLE 1. LIST OF PRIMERS

Target mouse	Sequence	Target BRECs	Sequence
D-loop ^a		D-loop ^a	
Forward	5'-TGCCCCTCTTCTCGCTCCGG-3'	Forward	5'-TGCTTGACTCAGCTATGGCCG-3'
Reverse	5'-GGCGATAACGCATTGATGGCC-3'	Reverse	5'-TCATTATGCTGGTGCTCAAGATGCA-3'
CytB ^a		CytB ^a	
Forward	5'-AGACAAAGCCACCTTGACCCGAT-3'	Forward	5'-TGACCTTCCCCGCCCCATCCA-3'
Reverse	5'-ACGATTGCTAGGGCCGCGAT-3'	Reverse	5'-AGCCGTAGTTTACGTCTCGGCA-3'
β -Actin		β -Actin	
Forward	5'-GCGCAGTGTAGGCGGAGCTT-3'	Forward	5'-TGTTCCCTTCCACAGGGTGT-3'
Reverse	5'-AAAGGAAGCGCAGACCGGCC-3'	Reverse	5'-TCCCAGTTGGTAACAATGCCA-3'
D-loop (semi long)		D-loop (semi long)	
Forward	5'-TGGGGGCCAACCAGTAGAACA-3'	Forward	5'-AACAGACGCAATCCCAGGCCG-3'
Reverse	5'-TGCGTCTAGACTGTGTGCTGTCC-3'	Reverse	5'-AGCTGTTAACCCGACGGCGA-3'
CytB (semi long)		CytB (semi long)	
Forward	5'-TGACCTACCTGCCCCATCCAACA-3'	Forward	5'-CTGCCGAGACGTGAACTACGGC-3'
Reverse	5'-TGGGGATTGAGCGTAGAATGGCGT-3'	Reverse	5'-CGACTGTGCCGGCTGTGGTA-3'
POLG1		POLG1	
Forward	5'-AGGGGGAGGCCGAACCCGT-3'	Forward	5'-AGCCCCGCGAGCTGTTTGTG-3'
Reverse	5'-GGCGATAGCTGGCTGGTCAAG-3'	Reverse	5'-CACCGGGTGGGGACACCTCT-3'
POLG2		POLG2	
Forward	5'-TCTGGGAAACTACGGGCGACTCT-3'	Forward	5'-TGTTCCCGGTAGACGCCCTCC-3'
Reverse	5'-TGCTGGGTGTCTGATTGCTGTTT-3'	Reverse	5'-TCCAAGGCACCGTGAAGAAGG T-3'
TWINKLE		TWINKLE	
Forward	5'-CATCCCCCGCCTGG TGTCT-3'	Forward	5'-CAGAGGCGAGGGACCACGGA-3'
Reverse	5'-GCCCGG TCGACTGGCTCAAG-3'	Reverse	5'-GGCCGGTCCCTCCACAGCTA-3'
β -Actin (mRNA)		β -Actin (mRNA)	
Forward	5'-CCTCTATGCCAACACAGTGC-3'	Forward	5'-CGCCATGGATGATGATATTGC-3'
Reverse	5'-CATCGTACTCCTGCTGTGCTG-3'	Reverse	5'-AAGCCGGCCTTGCACAT-3'
mtDNA-long		mtDNA-long	
Forward	5'-AAAATCCCCGCAAACAATGACC ACCC-3'	Forward	5'-ATGAGTTGGTAGTTTCGGTT GGGGT-3'
Reverse	5'-GGCAATTAAGAGTGGGATGGA GCCAA-3'	Reverse	5'-ATTCTGGTCTGTGTATGG GCGTGT-3'
mtDNA-short		mtDNA-short	
Forward	5'-CCTCCCATTCATTATCGCCGCC CTTGC-3'	Forward	5'-CATACTCCTCTGTAAGCCACATAGC-3'
Reverse	5'-GTCTGGGTCTCCTAGTAGGTCT GGGAA-3'	Reverse	5'-AGACTTGCTAGTAGTCATCAGGTGCG-3'

^aThe primers used for real time PCR and also for short amplicons in semi long/short PCR. BRECs, bovine retinal endothelial cell; PCR, polymerase chain reaction.

at 37°C with 500 nM MitoTracker green, rinsed with PBS, fixed in 4% formaldehyde for 15 min, and incubated with 0.2% Triton-X 100 for 10 min. After blocking for 1 h with 1% BSA, the cells were incubated with anti-POLG1 for 2 h, washed, and blocked with 10% goat serum. They were then incubated with Texas Red-conjugated anti-rabbit IgG for 1 h, rinsed, mounted with DAPI-containing Vectashield mounting medium (Vector Laboratories, Burlingame, CA), and imaged using ZEISS ApoTome fluorescence microscope at 40 \times magnification (Carl Zeiss, Chicago, IL). The intensity of POLG1 in the mitochondria was calculated by the Axio-Vision software depending on the presence or absence of combined fluorescence of Texas red (POLG1) and MitoTracker green in a cell. The background for red and green

channels was normalized in each of the samples, and an intensity-scatter plot was generated displaying the Pearson's values and amount of green and red (yellow) co-localized channels.

Binding of POLG1 with D-loop

The binding affinity of POLG1 to its replication start site was assessed by ChIP of POLG1 at the D-loop of mtDNA. BRECs were cross linked with 1% formaldehyde for 10 min, and the DNA pull-down assay for the D-loop was performed as previously described (49). The protein was precleared with protein A agarose/salmon sperm DNA slurry, and 130 μ g of precleared protein was incubated with

2 μ g of POLG1 antibody. The antibody-protein-DNA complex was pulled down, washed, and reverse cross linked. The DNA was purified, and the D-loop region of mtDNA was quantified by qPCR.

Functional assay of POLG for mtDNA replication

After incubation of BRECs in 5 or 20 mM glucose for 4 days, they were incubated with aphidicolin (7 μ M) or EtBr (1 μ g/ml) for 4 h, rinsed and further incubated for 6 h with 15 μ M BrdU with aphidicolin or EtBr (23). The cells were washed with PBS, fixed with 2% paraformaldehyde, and permeabilized with 0.1% TritonX-100. DNA was denatured with 2N HCl at 37°C for 30 min to recover the BrdU epitope. Cells were blocked with 5% goat serum, and incubated with mouse anti-BrdU antibody (Sigma-Aldrich; 1:50 dilution). After washing the cells with PBS, they were incubated with Texas Red-conjugated anti-mouse IgG for 1 h (at 1:250 dilution). Coverslips were mounted with DAPI containing mounting solution, and the fluorescence was visualized using ZEISS ApoTome at 40 \times magnification.

Mitochondrial superoxide

Mitochondrial superoxide were detected using MitoTracker Red (CM-H2XRos; Molecular Probes), a cell permeable dye that sequesters in the mitochondria and emits fluorescence when oxidized. After the desired incubations, the cells were rinsed with DMEM to remove nonadherent cells, and incubated with 400 nM MitoTracker Red for 30 min at 37°C (25, 29). The excess MitoTracker was removed, and the cells were fixed in 1% formaldehyde, rinsed, and mounted using Vectashield mounting media. The fluorescence was visualized at 40 \times magnification.

Mitochondrial superoxide levels were also quantified fluorimetrically by incubating the cell homogenate (2–5 μ g protein) with 400 nM MitoTracker Red for 30 min at 37°C. The fluorescence was measured at 579 nm excitation and 599 nm emission wavelengths using an LS55 Fluorescence Spectrometer (25).

Statistical analysis

Data are expressed as mean \pm standard deviation. Statistical analysis was carried out using Sigma Stat software. The Shapiro–Wilk test was used to test for normal distribution of the data. For variables with normal distribution, ANOVA followed by Bonferroni test were applied, while Kruskal–Wallis test followed by Mann–Whitney U-test were used for data that did not present a normal distribution. A *p*-value < 0.05 was considered statistically significant.

Acknowledgments

The authors thank Doug Putt and Yakov Shamailov for their technical assistance. This study was supported in part by grants from the National Institutes of Health, Juvenile Diabetes Research Foundation, Thomas Foundation, and Research to Prevent Blindness.

Author Disclosure Statement

The authors declare that they do not have any competing financial interests.

References

- Al-Shabrawey M, Bartoli M, El-Remessy AB, Platt DH, Margetson S, Behzadian MA, Caldwell RW, and Caldwell RB. Inhibition of NAD(P)H oxidase activity blocks vascular endothelial growth factor overexpression and neovascularization during ischemic retinopathy. *Am J Pathol* 167: 599–607, 2005.
- Bakthavatchalu V, Dey S, Xu Y, Noel T, Jungsuwadee P, Holley AK, Dhar SK, Batinic-Haberle I, and St. Clair DK. Manganese superoxide dismutase is a mitochondrial fidelity protein that protects Poly γ against UV-induced inactivation. *Oncogene* 2011 [Epub ahead of print]; DOI: 10.1038/onc.2011.407.
- Brownlee M. The pathobiology of diabetic complications: a unifying mechanism. *Diabetes* 54: 1615–1625, 2005.
- Clayton DA. Replication and transcription of vertebrate mitochondrial DNA. *Annu Rev Cell Biol* 7: 453–478, 1991.
- Correia RL, Oba-Shinjo SM, Uno M, Huang N, and Marie SK. Mitochondrial DNA depletion and its correlation with TFAM, TFB1M, TFB2M and POLG in human diffusely infiltrating astrocytomas. *Mitochondrion* 1: 48–53, 2011.
- De A and Campbell C. A novel interaction between DNA ligase III and DNA polymerase gamma plays an essential role in mitochondrial DNA stability. *Biochem J* 402: 175–186, 2007.
- Falkenberg M, Larsson NG, and Gustafsson CM. DNA replication and transcription in mammalian mitochondria. *Annu Rev Biochem* 76: 679–699, 2007.
- Frank RN. Diabetic retinopathy. *N Engl J Med* 350: 48–58, 2004.
- Graziewicz MA, Longley MJ, and Copeland WC. DNA polymerase gamma in mitochondrial DNA replication and repair. *Chem Rev* 106: 383–405, 2006.
- Jarrett SG, Lin H, Godley BF, and Boulton ME. Mitochondrial DNA damage and its potential role in retinal degeneration. *Prog Retin Eye Res* 27: 596–607, 2008.
- Jeffrey DS and William CC. Mitochondrial DNA replication and disease: insights from DNA polymerase gamma mutations. *Cell Mol Life Sci* 68: 219–233, 2011.
- Kaguni LS. DNA polymerase gamma, the mitochondrial replicase. *Annu Rev Biochem* 73: 293–320, 2004.
- Kanwar M, Chan PS, Kern TS, and Kowluru RA. Oxidative damage in the retinal mitochondria of diabetic mice: possible protection by superoxide dismutase. *Invest Ophthalmol Vis Sci* 48: 3805–3811, 2007.
- Karunadharma PP, Nordgaard CL, Olsen TW, and Ferrington DA. Mitochondrial DNA damage as a potential mechanism for age-related macular degeneration. *Invest Ophthalmol Vis Sci* 51: 5470–5479, 2010.
- Korhonen JA, Pham XH, Pellegrini M, and Falkenberg M. Reconstitution of a minimal mtDNA replisome *in vitro*. *EMBO J* 23: 2423–2429, 2004.
- Kowluru RA and Abbas SN. Diabetes-induced mitochondrial dysfunction in the retina. *Invest Ophthalmol Vis Sci* 44: 5327–5334, 2003.
- Kowluru RA, Atasi L, and Ho YS. Role of mitochondrial superoxide dismutase in the development of diabetic retinopathy. *Invest Ophthalmol Vis Sci* 47: 1594–1599, 2006.
- Kowluru RA, Jirousek MR, Stramm L, Farid N, Engerman RL, and Kern TS. Abnormalities of retinal metabolism in diabetes or experimental galactosemia: V. Relationship between protein kinase C and ATPases. *Diabetes* 47: 464–469, 1998.
- Kowluru RA, Kowluru V, Ho YS, and Xiong Y. Overexpression of mitochondrial superoxide dismutase in mice protects the retina from diabetes-induced oxidative stress. *Free Radic Biol Med* 41: 1191–1196, 2006.

20. Kowluru RA, Mohammad G, Santos JM, and Zhong Q. Abrogation of MMP9 gene protects against the development of retinopathy in diabetic mice by preventing mitochondrial damage. *Diabetes* 60: 3023–3033, 2011.
21. Lee DY and Clayton DA. Properties of a primer RNA-DNA hybrid at the mouse mitochondrial DNA leading-strand origin of replication. *J Biol Chem* 271: 24261–2429, 1996.
22. Lee HC and Wei YH. Mitochondrial biogenesis and mitochondrial DNA maintenance of mammalian cells under oxidative stress. *Int J Biochem Cell Biol* 37: 822–834, 2005.
23. Lentz SI, Edwards JL, Backus C, McLean LL, Haines KM, and Feldman EL. Mitochondrial DNA (mtDNA) biogenesis: visualization and dual incorporation of BrdU and EdU into newly synthesized mtDNA *in vitro*. *J Histochem Cytochem* 58: 207–218, 2010.
24. Liu P and Demple B. DNA repair in mammalian mitochondria: much more than we thought? *Environ Mol Mutagen* 51: 417–426, 2010.
25. Madsen-Bouterse S, Zhong Q, Mohammad G, Ho YS, and Kowluru RA. Oxidative damage of mitochondrial DNA in diabetes, and its protection by manganese superoxide dismutase. *Free Radic Res* 44: 313–321, 2010.
26. Madsen-Bouterse SA, Mohammad G, Kanwar M, and Kowluru RA. Role of mitochondrial DNA damage in the development of diabetic retinopathy, and the metabolic memory phenomenon associated with its progression. *Antioxid Redox Signal* 13: 797–805, 2010.
27. Miller FJ, Rosenfeldt FL, Zhang C, Linnane AW, and Nagley P. Precise determination of mitochondrial DNA copy number in human skeletal and cardiac muscle by a PCR-based assay: lack of change of copy number with age. *Nucleic Acids Res* 31: e61, 2003.
28. Miyazono F, Schneider PM, Metzger R, Warnecke-Eberz U, Baldus SE, Dienes HP, Aikou T, and Hoelscher AH. Mutations in the mitochondrial DNA D-Loop region occur frequently in adenocarcinoma in Barrett's esophagus. *Oncogene* 21: 3780–3783, 2002.
29. Mohammad G and Kowluru RA. Matrix metalloproteinase-2 in the development of diabetic retinopathy and mitochondrial dysfunction. *Lab Invest* 90: 1365–1372, 2010.
30. Mohammad G and Kowluru RA. Novel role of mitochondrial matrix metalloproteinase-2 in the development of diabetic retinopathy. *Invest Ophthalmol Vis Sci* 52: 3832–3841, 2011.
31. Navaglia F, Basso D, Fogar P, Sperti C, Greco E, Zambon CF, Stranges A, Falda A, Pizzi S, Parenti A, Pedrazzoli S, and Plebani M. Mitochondrial DNA D-loop in pancreatic cancer: somatic mutations are epiphenomena while the germline 16519 T variant worsens metabolism and outcome. *Am J Clin Pathol* 126: 593–601, 2006.
32. Prior SL, Griffiths AP, and Lewis PD. A study of mitochondrial DNA D-loop mutations and p53 status in non-melanoma skin cancer. *Br J Dermatol* 161: 1067–1071, 2009.
33. Richter C. Oxidative damage to mitochondrial DNA and its relationship to ageing. *Int J Biochem Cell Biol* 27: 647–653, 1995.
34. Ronchi D, Fassone E, Bordoni A, Sciacco M, Lucchini V, Di Fonzo A, Rizzuti M, Colombo I, Napoli L, Ciscato P, Moggio M, Cosi A, Collotta M, Corti S, Bresolin N, and Comi GP. Two novel mutations in PEO1 (Twinkle) gene associated with chronic external ophthalmoplegia. *J Neurol Sci* 308: 173–176, 2011.
35. Rothfuss O, Gasser T, and Patenge N. Analysis of differential DNA damage in the mitochondrial genome employing a semi-long run real-time PCR approach. *Nucleic Acids Res* 38: e24, 2010.
36. Santos JM and Kowluru RA. Role of mitochondria biogenesis in the metabolic memory associated with the continued progression of diabetic retinopathy and its regulation by lipoic acid. *Invest Ophthalmol Vis Sci* 2: 8791–9798, 2011.
37. Santos JM, Mohammad G, Zhong Q, and Kowluru RA. Diabetic retinopathy, superoxide damage and antioxidant. *Curr Pharm Biotechnol* 12: 352–361, 2011.
38. Santos JM, Tewari S, Goldberg AFX, and Kowluru RA. Mitochondria biogenesis and the development of diabetic retinopathy. *Free Radic Biol Med* 51: 1849–1860, 2011.
39. Schultz RA, Swoap SJ, McDaniel LD, Zhang B, Koon EC, Garry DJ, Li K, and Williams RS. Differential expression of mitochondrial DNA replication factors in mammalian tissues. *J Biol Chem* 273: 3447–3451, 1998.
40. Shahidi AM, Sampson GP, Pritchard N, Edwards K, Russell A, Malik RA, and Efron N. Exploring retinal and functional markers of diabetic neuropathy. *Clin Exp Optom* 93: 309–323, 2010.
41. Sharma H, Singh A, Sharma C, Jain SK, and Singh N. Mutations in the mitochondrial DNA D-loop region are frequent in cervical cancer. *Cancer Cell Int* 5: 34, 2005.
42. Spelbrink JN, Toivonen JM, Hakkaart GA, Kurkela JM, Cooper HM, Lehtinen Sk, Lecrenier N, Back JW, Speijer D, Foury F, and Jacobs HT. *In vivo* functional analysis of the human mitochondrial DNA polymerase POLG expressed in cultured human cells. *J Biol Chem* 275: 24818–24828, 2000.
43. Suzuki S, Hinokio Y, Komatu K, Ohtomo M, Onoda M, Hirai S, Hirai M, Hirai A, Chiba M, Kasuga S, Akai H, and Toyota T. Oxidative damage to mitochondrial DNA and its relationship to diabetic complications. *Diabetes Res Clin Pract* 45: 161–168, 1999.
44. Tanwar M, Dada T, Sihota R, and Dada R. Mitochondrial DNA analysis in primary congenital glaucoma. *Mol Vis* 16: 518–533, 2010.
45. Tyynismaa H, Sembongi H, Bokori-Brown M, Granycome C, Ashmal N, Poulton J, Jalanko A, Spelbrink JN, Holt IJ, and Suomalainen A. Twinkle helicase is essential for mtDNA maintenance and regulates mtDNA copy number. *Hum Mol Genet* 13: 3219–3227, 2004.
46. Wanrooij S and Falkenberg M. The human mitochondrial replication fork in health and disease. *Biochim Biophys Acta* 1797: 1378–1388, 2010.
47. Wu Y, Li N, Zhang T, Wu H, Huang C, and Chen D. Mitochondrial DNA base excision repair and mitochondrial DNA mutation in human hepatic HuH-7 cells exposed to stavudine. *Mutat Res* 664: 28–38, 2009.
48. Zhong Q and Kowluru RA. Diabetic retinopathy and damage to mitochondrial structure and transport machinery. *Invest Ophthalmol Vis Sci* 52: 8739–8746, 2011.
49. Zhong Q and Kowluru RA. Epigenetic changes in mitochondrial superoxide dismutase in the retina and the development of diabetic retinopathy. *Diabetes* 60: 1304–1313, 2011.
50. Zhong Q and Kowluru RA. Role of histone acetylation in the development of diabetic retinopathy and the metabolic memory phenomenon. *J Cell Biochem* 110: 1306–1313, 2010.

Address correspondence to:

Dr. Renu A. Kowluru
Kresge Eye Institute
Wayne State University
4717 St. Antoine
Detroit, MI 48201

E-mail: rkowluru@med.wayne.edu

Date of first submission to ARS Central, October 13, 2011; date of final revised submission, December 8, 2011; date of acceptance, December 16, 2011.

Abbreviations Used

BER = base excision repair
BrDU = bromodeoxyuridine (5-bromo-2'-deoxyuridine)
BRECs = bovine retinal endothelial cells
ChIP = chromatin immunoprecipitation
CytB = cytochrome B
Cox IV = cytochrome oxidase IV
D-loop = displacement loop
EtBr = ethidium bromide

H2B = histone 2-B
LDH = lactate dehydrogenase
MnSOD = mitochondrial superoxide dismutase
MnTBAP = manganese (iii) tetrakis (4-benzoic acid) porphyrin chloride
MtDNA = mitochondrial DNA
nDNA = nuclear DNA
POLG = DNA polymerase gamma
qPCR = quantitative polymerase chain reaction
ROS = reactive oxygen species
SC = scrambled siRNA
Tg = mice overexpressing MnSOD
Twinkle = mtDNA helicase
WT = wild-type