

RESEARCH ARTICLE

Open Access

Genome-wide microarray analysis of tomato roots showed defined responses to iron deficiency

Anita Zamboni^{1†}, Laura Zanin^{2†}, Nicola Tomasi², Mario Pezzotti¹, Roberto Pinton², Zeno Varanini^{1*} and Stefano Cesco³

Abstract

Background: Plants react to iron deficiency stress adopting different kind of adaptive responses. Tomato, a *Strategy I* plant, improves iron uptake through acidification of rhizosphere, reduction of Fe³⁺ to Fe²⁺ and transport of Fe²⁺ into the cells. Large-scale transcriptional analyses of roots under iron deficiency are only available for a very limited number of plant species with particular emphasis for *Arabidopsis thaliana*. Regarding tomato, an interesting model species for *Strategy I* plants and an economically important crop, physiological responses to Fe-deficiency have been thoroughly described and molecular analyses have provided evidence for genes involved in iron uptake mechanisms and their regulation. However, no detailed transcriptome analysis has been described so far.

Results: A genome-wide transcriptional analysis, performed with a chip that allows to monitor the expression of more than 25,000 tomato transcripts, identified 97 differentially expressed transcripts by comparing roots of Fe-deficient and Fe-sufficient tomato plants. These transcripts are related to the physiological responses of tomato roots to the nutrient stress resulting in an improved iron uptake, including regulatory aspects, translocation, root morphological modification and adaptation in primary metabolic pathways, such as glycolysis and TCA cycle. Other genes play a role in flavonoid biosynthesis and hormonal metabolism.

Conclusions: The transcriptional characterization confirmed the presence of the previously described mechanisms to adapt to iron starvation in tomato, but also allowed to identify other genes potentially playing a role in this process, thus opening new research perspectives to improve the knowledge on the tomato root response to the nutrient deficiency.

Background

Iron (Fe) deficiency is a yield-limiting factor for a variety of field crops all around the world and generally results from the interaction of limited soil Fe bioavailability and susceptible genotype cultivation [1]. Iron is an important microelement for plant life due to its involvement as redox-active metal in photosynthesis, mitochondrial respiration, nitrogen assimilation, hormone biosynthesis, production and scavenging of reactive oxygen species, osmoprotection and pathogen defence [2].

Under aerated conditions at neutral alkaline pH, the soluble Fe concentration in soil solution is very low. To cope with Fe shortage plants have developed two

strategies for its acquisition. The *Strategy I* (all higher plants except grasses) relies on improvement of Fe uptake through acidification of soil solution by excretion of protons via a plasmalemma P-type ATPase resulting in an increased Fe solubility, reduction of Fe³⁺ to the more soluble Fe²⁺ by a Fe^{III}-chelate reductase and plasmalemma transport of Fe²⁺ by the activity of a Fe transporter [3]. Some model plants used to study *Strategy I* are dicots such as *Arabidopsis thaliana*, *Solanum lycopersicum* and *Pisum sativum* [4].

Plant responses to Fe deficiency have been recently analyzed on the basis of large-scale changes not only in transcriptome [5-14], but also in proteome [15-20] and metabolome [17]. Results of transcriptome analysis are influenced by differences in experimental plans, plant species and microarray platforms, and thus difficult to compare and be generalized. Notwithstanding this drawback, recently, a set of 92 transcripts that robustly

* Correspondence: zeno.varanini@uniud.it

† Contributed equally

¹Department of Biotechnology, University of Verona, via delle Grazie 15, 37134 Verona, Italy

Full list of author information is available at the end of the article

reflect the transcriptional response of *Arabidopsis* to Fe deficiency [21], has been described as the “ferrome” by Schmidt and Buckhout [21]. The “ferrome” consists of a list of transcripts considered to be involved in the basic response to iron deficiency. The ferrome is particularly enriched in genes related to heavy metal cation transport and metal homeostasis. Focusing on tomato, a plant often used as a model to study Fe deficiency (*Strategy I*) and a crop of economic importance, no information is available at genome-wide transcriptional level. Two proteomic characterizations of tomato roots in response to 1-week of Fe deprivation showed 23 [15] and 15 [16] differentially expressed protein spots respectively. Modifications in proteome suggest changes in energy metabolism, sulfur metabolism, response to oxidative stress and signal transduction.

In the present work a genome-wide transcriptional characterization of tomato roots in response to Fe deficiency is presented. This approach allowed indentifying 97 differentially expressed transcripts involved in the responses to the nutritional stress. Transcriptional changes, mainly related to positive modulation of glycolysis, TCA and methionine cycle, suggest that tomato roots behave similarly to *Arabidopsis* under Fe deficiency. Furthermore, flavonoid biosynthesis and root morphological changes are revealed as specific tomato responses to Fe shortage.

Results and discussion

Responses to Fe-deficiency

Typical responses of Fe-deficiency [22] were observed in tomato plants grown for 14 d in the presence of a low amount of Fe and thereafter subjected to 7 d of Fe deprivation. The chlorophyll content (SPAD index value) was reduced in Fe-deficient plants (Table 1). A concomitant increase in root Fe^{III}-chelate reductase activity (Table 1) was also observed with values similar to those commonly found in roots of Fe-deficient tomato plants [23]. Furthermore, Fe-deprived tomato plants developed more lateral roots and showed an abundant production of root-hairs (Figures 1 and 2).

Table 1 Leaf SPAD index values and root Fe^{III}-chelate reductase activity

Sample	SPAD index ^a	Fe ^{III} -chelate reductase (mol g ⁻¹ root FW h ⁻¹) ^b
Fe-sufficient	29.5 ± 0.3	0.37 ± 0.04
Fe-deficient	16.8 ± 0.6	1.41 ± 0.06

^aSPAD index value of fully expanded young leaves was determined using a SPAD-502 meter (Minolta, Osaka, Japan); mean and SD using data of the three biological replicates.

^bMean and SD of three biological replicates.




Figure 1 Shoot and root apparatus of tomato plants grown under different Fe-supply conditions. Leaf detail of Fe-deficient (A) and Fe-sufficient (B) plants. Shoot (C) and roots (E) of Fe-deficient plants and shoot (D) and roots (F) of Fe-sufficient plants.

Comparison of root gene expression profiles in Fe-deficient and Fe-sufficient tomato plants

Differences in root gene expression between Fe-sufficient and Fe-deficient tomato plants were obtained by a genome-wide gene expression analysis using a tomato chip developed through Combimatrix technology [24]. This chip allows monitoring simultaneously the expression of more than 25,000 tomato transcripts. Ninety-




Figure 2 Root apparatus of tomato plants grown under different Fe-supply condition. Detail of root apparatus of A) Fe-deficient and B) Fe-sufficient plants.

seven differentially expressed transcripts between Fe-deficient and Fe-sufficient tomato roots (75 up-regulated and 22 down-regulated) were identified by Linear Models for MicroArray (LIMMA) [25] (adjusted p-value ≤ 0.05 ; $|FC| \geq 2$). This result obtained using a large-scale chip reinforce the idea that plant transcriptional response to Fe shortage is based on the modulation of a relative small set of genes as previously observed for the *Arabidopsis* “ferrome” [21].

Manually curated annotation of the 97 differentially expressed transcripts was based on results of BlastP analysis against UniProt [26] database (Figure 3; Additional file 1) using terms of biological process of Gene Ontology (GO) [27]. Sequence grouping in functional categories according to the GO terms revealed that the most abundant functional category was “metabolic process” both for up-regulated and down-regulated transcripts (35% and 45% respectively; Figure 3). Other up-regulated transcripts belonged to “establishment of localization” (12%) and “cell wall organization and biogenesis” (8%), while for the down-regulated transcripts “response to stimulus” (9%) was one of the most representative main functional categories (Figure 3). Only up-regulated transcripts are present in the “secondary metabolic process” category (Table 2). Transcripts encoding proteins with no sequence homology to known proteins were defined as “no hits found” (12% and 5% for up-regulated and down-regulated transcripts respectively), while a similar percentage of transcripts showed homology to proteins involved in “unknown” biological process (24% and 23% respectively). A selection of differentially expressed and discussed in relation to Fe deficiency is reported in Table 2. The up-regulation and

down-regulation of six differentially expressed transcripts in response to Fe deficiency were confirmed through Real-time RT-PCR experiments (Table 3).

TC ID, description, Real-time *RT-PCR* relative expression value (Fe-deficient vs. Fe-sufficient) and microarray fold change value (Fe-deficient vs. Fe-sufficient). Real time data were normalized on the *EF1a* gene and were performed 4 times on 3 independent experiments. Real-time RT-PCR data are expressed as mean \pm SD.

Sixty-one of 97 transcripts are ascribable to adaptive responses to Fe deficiency involving Fe homeostasis, metabolic process, oxidative stress responses, root morphological modification, transport processes, hormone metabolism and signaling. The others are hardly related to specific role showing homology to protein without a specific biological process or lacking homology to known protein (“no hits found”).

Fe homeostasis

Our transcriptional analysis confirmed that roots of Fe-deficient tomato plants overexpressed genes involved in Fe uptake and reduction, including the transcripts encoding IRON-REGULATED TRANSPORTER (IRT) [28-30] and Fe^{III}-chelate reductase (FRO) [31]. The tomato bHLH protein (encoded by *LeFER*) plays a role in Fe-deficiency responses through the expression of these two tomato Fe mobilization genes belonging to the Fe uptake systems of the *Strategy I* plants [29,32,33]. Our data show the up-regulation of the *FER* transcript (#24) in Fe-deficient roots, which is in agreement with the positive modulation of Fe-uptake-related transcripts such as *LeFRO1* (#68), *LeIRT1* (#39) and a transcript encoding a NATURAL RESISTANCE-ASSOCIATED




Table 2 List of transcripts modulated in response to Fe-deficiency and reported in the Discussion

#	ProbeID ^a	Description ^b	UniProtID ^c	TC-ID ^d	FC ^e	p, value, adj ^f
<i>Biological process GO:0008150</i>						
1	TC192724_853_37_S	SRC2	B6SND5	TC192724	3.02	0.014
2	TC192763_241_40_S	Putative uncharacterized protein	C6T3H9	TC192763	2.11	0.030
3	TC193319_801_34_X2	Putative uncharacterized protein	Q40127	TC193319	-3.35	0.016
4	TC195341_998_35_S	Putative uncharacterized protein	Q40127	TC195341	-3.33	0.011
5	TC196753_1279_37_S	D-protein	Q8VWY8	TC196753	6.11	0.028
6	TC197095_638_34_X2	NtEIG-E80 protein (elicitor inducible gene product)	Q9FXS6	TC197095	4.69	0.024
7	TC198323_947_40_S	Predicted protein	B9MWQ1	TC198323	2.51	0.032
8	TC199253_1439_39_S	Zinc finger protein. putative	B9SLY6	TC199253	2.82	0.029
9	TC202360_487_40_S	EF-1 alpha-like protein	O49604	TC202360	2.15	0.024
10	TC204571_463_40_S	ATBET12, putative	B9STJ3	TC204571	-2.62	0.023
11	TC205207_890_35_S	Putative uncharacterized protein	Q40127	TC205207	-3.09	0.048
12	TC207200_893_35_S	VIT_00038707001	E0CVH7	TC207200	2.44	0.042
13	TC207407_739_37_S	Predicted protein	B9HQW6	TC207407	2.36	0.042
14	TC207665_362_36_S	Putative uncharacterized protein	A9PCS8	TC207665	-2.39	0.048
15	TC208712_583_40_S	Germin-like protein	Q5DT23	TC208712	2.35	0.046
16	TC208745_692_37_S	Putative uncharacterized protein	A5C0F7	TC208745	4.84	0.016
17	TC209321_482_35_S	VITISV_041870	A5C9V2	TC209321	6.94	0.011
18	TC209504_302_40_S	Hydrolase	Q4PSL3	TC209504	2.78	0.013
19	TC211515_728_35_S	Amino acid binding protein, putative	B9RBU8	TC211515	3.66	0.014
20	TC212954_1137_35_S	Predicted protein	B9HZ36	TC212954	2.02	0.046
21	TC213456_100_34_S	Putative D-protein	Q6K482	TC213456	4.97	0.035
22	TC214599_1004_37_S	Nodulin-like proteinAt2g16660/T24121.7	Q9SLF1	TC214599	6.87	0.042
23	TC215994_489_36_S	Putative aminotransferase, class V family protein	Q1H8R9	TC215994	2.08	0.036
<i>Biological regulation GO: 0065007</i>						
24	TC191806_907_40_S	BHLH transcriptional	Q5GA67	TC191806	6.56	0.024
25	TC191963_1058_37_S	Ferritin	Q308A9	TC191963	-4.71	0.008
26	TC194645_664_37_S	DNA binding protein	B9SZX2	TC194645	12.34	0.015
27	TC198138_1325_39_S	Thioredoxin peroxidase 1	Q7Y240	TC198138	7.06	0.046
28	TC203853_451_35_S	Heterogeneous nuclear ribonucleoprotein 27C	B9SSS5	TC203853	2.90	0.028
29	TC206202_500_40_S	Thioredoxin II	B9RLX0	TC206202	-2.47	0.048
30	TC215976_330_38_X2	Ferritin	Q308A9	TC215976	-3.64	0.005
<i>Cellular component organization or biogenesis GO:0071840</i>						
31	NP000231_1302_40_S	Extensin-like protein Dif54	Q43505	NP000231	8.22	0.011
32	TC191669_1238_40_S	Extensin-like protein Ext1	Q8VWM5	TC191669	5.53	0.003
33	TC204863_245_40_S	Extensin-like protein Ext1	Q8VWM5	TC204863	8.98	0.013
34	TC212258_415_40_S	Extensin-like protein Dif54	Q43505	TC212258	9.13	0.011
35	TC212487_279_40_S	Extensin-like protein Ext1	Q8VWM5	TC212487	6.44	0.0004
36	TC214133_1133_40_S	Extensin-like protein Dif10	Q43504	TC214133	5.42	0.024
<i>Cellular process GO:0009987</i>						
37	TC207486_546_40_S	Pollen specific actin-depolymerizing factor 2	Q8H2B6	TC207486	4.81	0.005
<i>Establishment of localization GO: 0051234</i>						
38	NP796451_1558_36_S	Root-specific metal transporter	Q84LR1	NP796451	12.00	0.024
39	TC191581_1150_36_S	Iron-regulated transporter 1	Q9XFB2	TC191581	9.42	0.013
40	TC192292_1560_39_S	Hippocampus abundant transcript 1 protein	B9SG70	TC192292	14.79	0.0004
41	TC200857_1001_40_S	Ammonium transporter 1 member 1	P58905	TC200857	2.05	0.043
42	TC205660_580_35_S	Metal tolerance protein	B9GLJ8	TC205660	3.77	0.049
43	TC206149_966_36_S	Aluminum-activated malate transporter 9	Q9LS46	TC206149	2.41	0.048
44	TC208376_922_36_S	Oligopeptide transporter, putative	B9SA63	TC208376	17.67	0.004
45	TC215768_1111_35_X2	Aquaporin	Q8W506	TC215768	-2.03	0.023
46	TC215874_553_40_S	Sec14 cytosolic factor, putative	B9S6A7	TC215874	3.5	0.009

Table 2 List of transcripts modulated in response to Fe-deficiency and reported in the Discussion (Continued)

47	TC216882_1121_38_X2	Hippocampus abundant transcript 1 protein <i>Metabolic process GO:0008152; Carbon utilization GO:0015976</i>	B9SG70	TC216882	9.29	0.003
48	TC204225_1412_35_S	Phosphoenolpyruvate carboxylase	P27154	TC204225	3.31	0.039
49	TC214978_700_36_S	Phosphoenolpyruvate carboxylase <i>Metabolic process GO:0008152; Catabolic process GO:0009056</i>	P27154	TC214978	3.74	0.036
50	TC194584_1854_37_S	Cysteine-type peptidase, putative	B9STX0	TC194584	3.61	0.011
51	TC203350_1573_40_S	Vacuolar processing enzyme 1	B2M1T0	TC203350	2.18	0.026
52	TC208154_2059_38_S	Beta-amylase PCT-BMYI	Q94EU9	TC208154	-3.06	0.042
53	TC215970_3405_40_S	Protease Do-like 7 <i>Metabolic process GO:0008152; Cellular metabolic process GO:0044237</i>	Q8RY22	TC215970	4.01	0.032
54	TC192049_1645_40_S	Sulfate adenyltransferase	Q43183	TC192049	-2.48	0.029
55	TC193934_828_38_X2	Methylthioribose kinase, putative	B9RY82	TC193934	2.12	0.023
56	TC194380_182_36_S	<i>S-adenosylmethionine-dependent methyltransferase, putative</i>	B9SZS6	TC194380	2.03	0.045
57	TC195032_699_35_S	CBL-interacting serine/threonine-protein kinase 11	O22932	TC195032	-2.62	0.030
58	TC198109_708_34_X2	Phosphofructokinase, putative	B9RRX6	TC198109	2.00	0.036
59	TC199972_1164_40_S	Fructose-bisphosphate aldolase	Q2PYX3	TC199972	2.53	0.022
60	TC201350_645_36_S	Protein phosphatase-2C	O82469	TC201350	2.10	0.028
61	TC201692_547_40_S	Xyloglucan endotransglucosylase/hydrolase 14	B9RKL5	TC201692	7.29	0.013
62	TC206357_361_38_S	Catalase isozyme 1	P30264	TC206357	-2.79	0.046
63	TC212978_951_34_S	Phosphoenolpyruvate carboxylase	Q8S915	TC212978	3.62	0.030
64	TC214826_1684_35_S	<i>ATP binding protein, putative</i>	B9RII2	TC214826	12.39	0.011
65	TC214837_697_40_S	Cytokinin oxidase/dehydrogenase	C3VPM8	TC214837	4.64	0.015
66	TC216529_896_39_S	Peroxidase 7	Q9SY33	TC216529	9.25	0.011
67	TC216572_542_40_S	Avr9/Cf-9 rapidly elicited protein 216 <i>Metabolic process GO:0008152; Oxidation reduction GO:0055114</i>	Q84QE0	TC216572	-2.61	0.042
68	TC191412_2284_38_S	Ferric-chelate reductase	Q6EMC0	TC191412	15.21	0.008
69	TC191893_9_37_S	Superoxide dismutase	Q7YK44	TC191893	-2.29	0.030
70	TC194139_2083_38_S	Ferric-chelate reductase	B9RIU2	TC194139	-10.76	0.023
71	TC194227_76_41_X2	Superoxide dismutase	Q7YK44	TC194227	-2.20	0.038
72	TC196465_645_40_X2	Gibberellin 20 oxidase	B9RUX2	TC196465	3.46	0.009
73	TC199400_1132_37_S	Peroxidase 2, putative	B9SZA0	TC199400	-2.05	0.022
74	TC201832_1136_40_S	Gibberellin 20 oxidase	B9RUX2	TC201832	3.00	0.013
75	TC205699_961_40_S	<i>Chlorophyll synthase, putative</i>	B9RJ38	TC205699	7.12	0.019
76	TC207549_14_41_S	Cytochrome C oxidase polypeptide vib	B9RJN9	TC207549	3.05	0.016
77	TC208767_662_38_S	Cationic peroxidase 1	B9SWU3	TC208767	4.75	0.022
78	TC213071_429_40_S	Peroxidase <i>Metabolic process GO:0008152; Primary metabolic process GO:0044238</i>	Q07446	TC213071	-3.29	0.032
79	TC197609_880_35_S	<i>Zinc finger protein, putative</i>	B9T6Q0	TC197609	3.54	0.014
80	TC192838_1967_36_S	Endo-1,4-beta-glucanase <i>Metabolic process GO:0008152; Secondary metabolic process GO:0019748</i>	B9RLZ9	TC192838	2.45	0.042
81	TC197109_626_37_S	Flavonoid 3-hydroxylase, putative	B9T1C6	TC197109	2.56	0.024
82	TC198786_1057_37_S	UDP-glucose:glucosyltransferase	B6EWY6	TC198786	2.19	0.022
83	TC203267_704_38_S	UDP-glucose:flavonoid glucoside 1,6-glucosyltransferase	C5NN14	TC203267	2.76	0.013
84	TC212095_566_35_S	UDP-glucose:flavonoid glucoside 1,6-glucosyltransferase <i>Response to stimulus GO:0050896</i>	C5NN14	TC212095	3.69	0.018
85	TC193192_66_41_X2	Peroxidase 4	B7UCP4	TC193192	-3.86	0.029
86	TC208216_282_40_S	<i>Pit1 protein</i>	Q40539	TC208216	4.71	0.019
87	TC195700_1019_40_S	Peroxidase <i>No hits found</i>	B9VRK9	TC195700	-2.67	0.013
88	TC203837_663_35_S	<i>No hits found</i>		TC203837	2.06	0.046
89	TC204355_604_36_S	<i>No hits found</i>		TC204355	-3.09	0.038
90	TC207055_310_37_S	<i>No hits found</i>		TC207055	3.05	0.030

Table 2 List of transcripts modulated in response to Fe-deficiency and reported in the Discussion (Continued)

91	TC209134_260_40_S	<i>No hits found</i>	TC209134	30.06	0.007
92	TC209988_335_40_S	<i>No hits found</i>	TC209988	6.82	0.013
93	TC211287_216_38_S	<i>No hits found</i>	TC211287	7.63	0.011
94	TC212074_241_40_S	<i>No hits found</i>	TC212074	6.44	0.023
95	TC212933_306_37_S	<i>No hits found</i>	TC212933	9.59	0.011
96	TC214074_254_40_S	<i>No hits found</i>	TC214074	2.22	0.041
97	TC215128_252_35_S	<i>No hits found</i>	TC215128	3.86	0.029

^aID of TomatoArray2.0 probes

^bDescription of each transcript. Bold discussed; italics undiscussed.

^cUniProtID [26] of the first hit obtained by BlastP analysis

^dID of the TC of DFCI Tomato Gene Index (Release 12.0) [92]

^eFold change value

^fadjusted *p*-value

MACROPHAGE PROTEIN1 (LeNRAMP1; #38) [29,32]. A positive modulation of FER and Fe mobilization proteins (IRT1, NRAMP1 and FRO1) was not found in two proteomics studies performed in tomato roots grown in conditions similar to those used in the present work [15,16]. Authors justified these results as related to the features of proteomic approach, which was not sensitive enough to detect FER and not well suited for membrane-bound proteins. However, functional characterizations of *fer* mutant proved that FER controls the expression of the iron-uptake genes [29,32,33]. A transcriptional behaviour similar to that described in the present work was observed for *Arabidopsis* orthologous genes in Fe-starved roots [5,8,11,34].

Our data show a strong down-regulation of another ferric chelate reductase (#70). Previous results indicated that the same transcript (TC194139 of the Release 12.0, corresponding to TC124302 of the Release 9.0) specific to the *Solanum lycopersicum* genome, is only slightly regulated by Fe and that its function is not essential for Fe uptake [34]. Our results showing a negative regulation of this FRO transcript in response to Fe-deficiency in roots, also quantified by Real-time RT-PCR experiment, confirm that this gene does not play a crucial role in deficiency-induced Fe uptake and suggest the involvement in other biological process.

Together with the positive modulation of *LeFER*, *LeFRO1* and *LeIRT1*, we observed for the first time a

high up-regulation of another bHLH transcript (#26); this result suggests that like in *Arabidopsis* [35] also in tomato plants the response to Fe deficiency through FER activity may need the interaction with another bHLH protein. BlastP analysis against TAIR database [36] using protein sequence obtained from the predicted coding sequence of the TC194645 showed the highest sequence homology with the protein encoded by *AtbHLH38* (score: 120; Evalue: 9 E-28; identity: 35%; positives: 57%) known to interact with FIT, the *Arabidopsis* orthologous of tomato FER [35].

The down-regulation of two ferritin transcripts (#25 and 30) is in line with the negative regulation of ferritin genes observed in roots of Fe-depleted *Arabidopsis* plants [5,8,11]. It has been suggested that ferritins can be involved in Fe homeostasis [37] with a main role of plastidial Fe [11]. *Arabidopsis* nodulin-like genes have been recently described to be putatively involved in Fe transport and storage under metal cation sufficiency [38]. Transcriptional levels of *Arabidopsis* nodulin-like genes were down-regulated at least until 72-h of Fe-deficiency, while two other nodulin-like genes were not modulated in response to different Fe conditions [38]. On the other hand in tomato roots we recorded a positive modulation of two transcripts encoding protein with a nodulin-like domain (#9 and 22). BlastP analysis against TAIR database [36] using protein sequence obtained from the predicted coding sequence of the

Table 3 Real-time RT-PCR validation of a set of genes differentially expressed in microarray analysis

TC ID	Description	Real-time RT-PCR (ratio)	Microarray
TC208376	Oligopeptide transporter, putative	17.87 ± 4.35	17.67
NP796451	Root-specific metal transporter	15.11 ± 3.97	12.00
TC191581	Iron-regulated transporter 1	3.07 ± 0.59	9.42
TC216882	Hippocampus abundant transcript 1 protein, putative	8.69 ± 1.53	9.29
TC205660	Metal tolerance protein	1.55 ± 0.44	3.77
TC194139	Ferric-chelate reductase, putative	-12.14 ± 3.15	-10.76

TC214599 (#22) showed the highest sequence homology with the protein encoded by At3g43660 (score: 325; Evalue: 2 E-29; identity: 54%; positives: 68%) which is one of the two *Arabidopsis* genes not modulated by iron [38].

Metabolic processes

Carbohydrate metabolism

As observed in *Arabidopsis* root microarray analyses [5,8,11] glycolysis-related genes are positively modulated in Fe-deficient roots. These transcriptional data fit well with the increased activity of glyceraldehyde-3-phosphate dehydrogenase (GADPH), pyruvate kinase (PK), and phosphofructokinase (PFK) recorded in response to Fe starvation in cucumber roots [39]. Furthermore, increased levels of protein related to glycolysis under Fe shortage were recently reported in sugar beat [17] and *Medicago truncatula* [20] roots. All together these evidences are consistent with the idea of a shift from anabolic to catabolic metabolism. In the present work, a fructose-bisphosphate aldolase (FBP aldolase) (#59) and a PFK (#58) transcripts were up-regulated under Fe deficiency, further confirming changes in primary metabolism in response to Fe starvation. The positive modulation of transcript encoding a PFK, an enzyme catalysing a protogenic reaction, supports the role of glycolysis in different process such as production of ATP and H⁺ for H⁺-ATPase, reducing equivalents for ferric chelate reductase and of phosphoenolpyruvate (PEP) [40]. Up-regulation of three phosphoenolpyruvate carboxylase (PEPC) transcripts (#48, 49 and 63) agrees with results of several proteomic and physiological studies on the response to Fe deficiency in tomato [15,16,41], sugar beat [17] and *Medicago truncatula* roots [20], showing a positive modulation of proteins involved both in glycolysis and TCA cycle. PEPC activity, through pyruvate consumption, can keep active the glycolytic pathway and give a contribution to the control of cytosolic pH [40,41]. In Fe deficiency, starch catabolism was reported to be enhanced both at transcriptional [5] and protein level [16]. Our analysis revealed a down-regulation of a transcript encoding a protein showing homology to a potato chloroplastic β -amylase (#52) involved into starch degradation in plastids [42]. Since starch catabolism mediated by this enzyme occurs in plastids, the negative modulation of β -amylase could be ascribed to other causes than an accelerated glycolysis.

The positive modulation of a cytochrome C oxidase (#76) is in line with the transient induction of electron-transport-chain genes observed in *Arabidopsis* [5] and consistent with the enhancement of respiration rate observed in cucumber [39] and sugar beat [43] Fe-deficient roots. This behaviour was interpreted as an attempt to

increase energy production through oxidative phosphorylation. However, more recently it has been suggested that the increased respiration rate in root segments of Fe-starved cucumber plants should not be interpreted as an increase in mitochondrial activity but rather as the result of an increase in the number of less efficient mitochondria and of the induction of different O₂-consuming reaction [44].

Methionine cycle

Nicotianamine (NA) is considered to be a key molecule for long-distance transport of Fe in plants [45]. Proteomic analysis of Fe-starved tomato roots [16] showed a positive modulation of proteins related to metabolism of methionine (e.g. methionine synthase), a precursor of nicotianamine. An up-regulation of a transcript encoding a methylthioribose kinase (MTK, #55), another enzyme of the methionine cycle, was observed in our transcriptional analysis. A positive modulation of MTK transcripts was also recorded in roots of rice, a *Strategy II* plant species, under Fe deficiency [46,47], although in this case, related to the necessity to increase the production of mugineic acid.

A down-regulation of a sulfate adenylyltransferase (ATPS; ATP sulfurylase) gene (#54) was observed in this work. However, the opposite was found in *Arabidopsis* [11]. The connection between the sulphur nutritional status and capability to respond to Fe shortage has been demonstrated in both *Strategy I* and *Strategy II* plants [30,48]. Recently an increase in methionine content related to phytosiderophore synthesis without significant changes in ATPS activity was described in Fe-deficient barley roots [48]. Interestingly, in a recent study of *Medicago truncatula* root proteome [20] two enzymes related to biosynthesis of cysteine and methionine were negatively affected by Fe deficiency.

Protein turnover

Response to Fe starvation induced the accumulation of gene transcripts related to protein turn-over, including protease (#53) and peptidase (#50 and 51) involved in protein degradation, and a gene encoding a heterogeneous nuclear ribonucleoprotein (#28) that can act in the pre-mRNA metabolism preceding protein synthesis.

The activity of these genes can be related to molecular events controlling plant responses to abiotic stresses such as nutritional deficiencies [49]. A general increase in protein synthesis was reported as a response to Fe-deficiency in cucumber roots [50]; furthermore, protein recycling in response to Fe starvation was suggested by analysis of expression profiles of soybean [13] and proteomic changes in cucumber [18] and *Medicago truncatula* [20] roots. Transcriptomic data presented here are in line with the idea that, under Fe deficiency, N recycling reactions take place, possibly related to the necessity of additional anaplerotic source of C and N [18,20].

Secondary metabolism

Phenolic compounds are reported as components of root exudates in Fe-deficient *Strategy I* plants. These molecules are involved in chelation and/or reduction of rhizospheric insoluble Fe [51,52]. Recently, phenolics have been proposed to selectively influence rhizospheric microorganisms and be involved in the reutilization of apoplastic Fe [53]. Our results showed an up-regulation of a flavonoid-3-hydroxylase (#81) gene and of three genes putatively involved into flavonoid glycosylation (#82, 83 and 84). Two out of the last three genes showed sequence homology to a *Catharanthus roseus* flavonoid glucoside 1,6-glucosyltransferase catalysing 1,6-glucosylation of flavonol and flavone glucosides [54]. A positive modulation of genes related to general phenylpropanoid pathway (e.g. genes encoding PAL and 4CL) was reported in Fe-deficient *Arabidopsis* roots [11]. Data of the present work underline the up-regulation of transcripts involved in a more specific branch of phenolic pathway supporting the idea that Fe-starved roots might operate flavonoid secretion into the rhizosphere in order to promote Fe acquisition [55].

Oxidative stress responses

Many proteins involved in antioxidative defence response contain Fe in heme group or coordinated to the thiol group of cysteine. The modulation of transcripts and proteins related to oxidative stress response in roots of Fe-starved plants seems to depend on the species and the experimental conditions [5,13,17,19]. However, catalase (CAT) and peroxidase (POX) activities are known to be depressed under Fe deficiency conditions [56] in tomato leaves. Our data showed a main down-regulation of transcripts encoding thioredoxin (TRX) (#29) and detoxifying enzymes catalase (CAT; #62), superoxide dismutase (#69 and 71), peroxidase (POX; #73, 78, 85 and 87). Two other POX transcripts (#66 and 77) and a thioredoxin peroxidase gene (#27) were, conversely, up-regulated. At protein level, a decrease of a CAT was reported while some POXs showed higher levels in response to Fe deficiency in tomato roots [15]; a different response of peroxidase isoforms has also been reported in sunflower [57]. Taken together, data of the present work suggest a different role in response to nutrient stress condition between the thioredoxin (ABB) and POX isogenes and are in agreement with previous results obtained in *Medicago truncatula* [20] and sugar beat [17] Fe-deficient roots.

A germin-like transcript (#15) was up-regulated under Fe deficiency, similarly to what has been observed in the tomato root proteome analysis [15]. The positive modulation of a germin protein reported in this proteomic study was justified hypothesizing its

role in producing hydrogen peroxide for apoplastic Fe reduction or in other stress response on the basis of sequence similarity to a *Nicotiana attenuata* germin protein [15,58].

Root morphological adaptation

Morphological modifications in roots of Fe-deficient plants are well documented [59,60]. Root hairs proliferation and development of transfer cells were described in Fe-starved tomato plants [29,61-63]. Enhanced formation of lateral roots and root hairs was also recorded in our experiment (Figure 1). Extensin proteins seem to be involved in this latter phenomenon; in fact, we observed a strong positive modulation of LeExt1 (#32, 33 and 35), LeDif10 (#36) and LeDif54 (#31 and 34). It was reported that these three genes encode extensin, a structural protein putatively conferring physical characteristics of the cell wall [64], and act during root hair formation in tomato due their predominant expression in root hair cells [64,65].

The overexpression of an endo-1,4- β -glucanase (#80) and a xyloglucan endotransglucosylase/hydrolase protein (#61) transcripts could be involved into the cell wall loosening [66] associated to root morphological adaptation to Fe deficiency, as previously observed in a tomato root proteomic analysis [16]. These changes in root morphology are also supported by the up-regulation of a transcript showing homology to a tobacco actin-depolymerizing factor (#37) related to the pollen tube elongation [67]. Here, we report a negative modulation of a transcript (#45) showing homology to the tobacco aquaporin PIP2 [68] suggesting a role of the PIP2 tomato protein in Fe-related morphological root changes. It has been suggested that PIP aquaporins can play a role not only in root water uptake but also in root development [68]. Transgenic plants exhibiting RNAi of PIP2 aquaporins showed a significant increase in the length of primary roots [68].

Transport processes

Among the positively modulated transcripts belonging to "transport" functional category, a stronger modulation (more than 17 times) of an oligopeptide transporter (OTP) gene (#44) was observed under Fe shortage. BlastP analysis against TAIR database [36] using protein sequence obtained from the predicted coding sequence of the TC208376 (#44) showed the highest sequence homology with the protein encoded by *AtOPT3* (score: 541; Evaluate: E-154; identity: 83%; positives: 90%). Positive modulation of *AtOPT2* and *AtOPT3* has been recorded in Fe-deficient *Arabidopsis* roots [8,11,21,69]. The plant members of OTP family have been described to have different functions in transport physiology such as long-distance sulphur distribution, nitrogen

mobilization, metal homeostasis, and heavy metal sequestration. OPTs can transport glutathione, metal-chelates and peptides [70]. It was hypothesized that some plants OPTs are able to transport Fe-chelates and Fe-NA suggesting a role of these proteins in long-distance metal transport *in planta* [71,72]. A similar function can be hypothesized for the tomato up-regulated OPT transcript, due to the previously described positive modulation of genes related to NA synthesis (see methionine cycle paragraph).

Another up-regulated gene involved in transport phenomena in response to Fe-deficiency is *LeAMT1* (#41). This tomato gene, firstly isolated from a root hair cDNA library [73] was root-specifically expressed and positively regulated by ammonium (NH_4^+) in root hairs [73,74]. This result suggests the presence of a linkage between NH_4^+ uptake and Fe shortage. It has been demonstrated that NH_4^+ -dependent rhizosphere acidification can improve Fe availability in the rhizosphere [75]. Interestingly, nitrate acquisition is limited under Fe deficiency [20,76].

Furthermore, favouring ammonium uptake with respect to nitrate could reduce competition for reducing equivalent between nitrogen and Fe acquisition. This is also in agreement with the hypothesized N-recycling in Fe-deficient plant roots [20].

Fe-deficient tomato roots strongly overexpressed two transcripts (#40 and 47) encoding a protein sharing features of major facilitator superfamily (MSF) with putative transport activities. However, on the basis of their sequence homology it was not possible to hypothesize an involvement in transport of a specific metabolite or mineral nutrient.

A positive modulation of a gene encoding a metal tolerance protein (MTP; #42) was also recorded. A similar behaviour was described in Fe-deficient *Arabidopsis* roots [8,11] and interpreted in the light of low specificity of IRT1 transporters, which can transport different metals into the Fe-deficient plants. MTP genes might therefore play a role in the detoxification of zinc ions taken up absorbed under Fe deficiency conditions [11].

Fe deficiency in tomato induced the expression of a transcript (#43) encoding a protein showing sequence homology to *Arabidopsis* vacuolar aluminium-activated malate transporter (ALMT) 9 [77]. An increase in organic acids concentration mainly citric and malic acids in plant roots under Fe starvation has been reported for many plant species [78]. However, a decrease of malate levels was observed in tomato root tips after 15-d of Fe deficiency while higher contents were recorded in leaves and xylem sap [41]. The tomato malate transporter gene might be involved in malate fluxes at intracellular level and/or in long-distance transport *in planta*.

Hormone metabolism and signaling

The role of plant hormones in the regulation of Fe deficiency responses has been extensively studied [79,80]. As for *Arabidopsis* Col-0 accession, we identified the positive modulation of a methionine cycle gene, MTK gene, that could be related not only to NA synthesis (see above) but also to the recycling of methylthioadenosine during ethylene production [11]. This is in agreement with the hypothesized involvement of ethylene and/or auxin in control of hair root production under Fe deficiency [79]. Furthermore, data of the present work suggest the involvement of other hormones (such as gibberellin and cytokinin) in Fe deficiency response in tomato roots. Concerning gibberellin, we recorded the up-regulation of two gibberellin oxidase 20 (GA20OX) transcripts (#72 and 74). They can regulate root morphological changes through the synthesis of active GA1. Indeed, it has been reported that GAs are related to tomato root growth [81]. In addition, it was shown that the expression of a tomato *SIGA20ox1:GUS* construct in *Arabidopsis* localized also in columella of secondary roots [82], thus suggesting a role in secondary root formation. The same authors also reported that the expression of the construct was positively affected in cotyledons, hypocotyls and roots by benziladenine, a synthetic cytokinin [82]. However cytokinins were described as negative regulators of root Fe uptake mechanism in *Arabidopsis* through a root-growth dependent pathway [83]. Cytokinins could play a similar role also in tomato roots. This hypothesis is supported by the up-regulation of a cytokinin oxidase/dehydrogenase (CKX) gene (#65) under Fe starvation. CKX genes are involved in regulation of plant growth and development through the control of the cytokinin concentration [84].

Focusing on signal transduction, we observed the up-regulation of a transcript (#60) encoding a protein with sequence homology to the MCP5, a protein phosphatase-2C (PP2C) of *Mesembryanthemum crystallinum* [85]. McMCP5 is expressed in roots and is induced in response to salt and drought stresses [85]. The tomato PP2C gene could play a role in signal transduction of stress nutrient condition as Fe starvation. The positive modulation of a gene encoding a SRC2 protein containing a C2 domain (#1) suggests a putative involvement of Ca as secondary messenger. Focusing on Ca, in our experiment we also identified a down-regulated gene (#57) in Fe-starved roots showing homology to an *Arabidopsis* gene (*AtSOS2*) encoding a CBL-interacting serine/threonine-protein kinase 11. *Arabidopsis* SOS2 interacts with the Ca binding protein SOS3 (SALT OVERLY SENSITIVE 3), thus controlling K and Na homeostasis and the response to salt stress [86,87]. In addition a negative modulation of a gene (#67) showing

homology to a tobacco Avr/Cf-9 rapidly elicited (ACRE) transcript encoding a PP2C was observed [88]. Our data confirm the results obtained with proteome analysis of tomato roots in response to Fe-deficiency, where changes in the levels of proteins involved in signal transduction were reported [16]. The observed transcriptional changes in tomato roots can be the result of the perception of nutrient stress condition and the following signal transduction.

Taken together, these results suggest that Fe deficiency responses are, at least in part, dependent on hormonal balance modifications possibly resulting from signal perception and transduction.

Conclusion

Ninety-seven differentially expressed transcripts were identified comparing root transcriptional profiles of Fe-deficient and Fe-sufficient tomato plants. Tomato roots respond to Fe deficiency by modulating the expression of a number of transcripts similar to the model plant *Arabidopsis*. The comparison of tomato Fe-responsive transcript set with the *Arabidopsis* “ferrome” [21], encompassing 92 transcripts that robustly represent the response to Fe shortage, confirms the involvement of the well know homologous key regulatory elements (e.g. bHLHs) controlling the expression of transcripts related to Fe uptake and translocation (e.g. IRT and FRO). As showed by *Arabidopsis* “ferrome” [21], tomato roots modulate transcripts involved in homeostasis of Fe and heavy metal cations (e.g. IRT, NRAMP, MTP, ferritin) and others cation (e.g. AMT). Both plant species require the up-regulation of transcripts related to glycolysis (e.g. PFK) and methionine cycle (e.g. MTK), the latter pathway being putatively linked to NA biosynthesis in response to Fe deficiency. Fe-NA complexes could be transported both in tomato and *Arabidopsis* plants through OPTs during the response to Fe shortage. Here we describe, for the first time, the modulation of a specific branch of phenolic (flavonoids) biosynthesis in response to Fe deficiency. In addition, tomato roots seem to be more characterized by root morphological adaptation, mainly linked to hair root production, as suggested by the strong up-regulation of extensin transcripts.

Therefore, this transcriptional study, while confirming evidence coming from proteomic studies, allowed identifying new putative targets for further functional investigations on the response to Fe deficiency in tomato roots.

Methods

Plant material, growth conditions and RNA extractions

Tomato seedling (*Solanum lycopersicum* L., cv. ‘Marmande superprecoce’ from DOTTO Spa, Italy),

germinated for 6 days on filter paper moistened with 1 mM CaSO₄, were grown for 14 days in a continuously aerated nutrient solution (pH adjusted at 6.0 with 1 M KOH) as reported by Tomasi et al. [22] with 5 μM Fe (Fe-EDTA); thereafter, most of the plants were transferred for a further week to a Fe-free nutrient solution (Fe-deficient) and some tomato plants were transferred for a week to a nutrient solution containing 100 μM Fe-EDTA (Fe-sufficient plants) as control. Nutrient solutions were renewed every three days. The controlled climatic conditions were the following: day/night photoperiod, 16/8 h; light intensity, 220 μE m⁻²s⁻¹; temperature (day/night) 25/20°C; RH 70 to 80%.

At the end of the growing period (27 days), Fe-deficient tomato plants clearly showed visible symptoms of Fe deficiency yellowing of the fully expanded apical leaves and proliferation of lateral roots and root hairs and increase in the diameter of the sub-apical root zone. 24 hours before harvesting, all nutrient solutions (both for Fe-deficient and Fe-sufficient plants) were renewed and the pH was adjusted to 7.5 with 10 mM Hepes-KOH. The pH of the growing medium was adjusted to this value to mimic as close as possible the conditions that are occurring in Fe-deficiency-inducing soil. However, in order to favour an equilibrate development of tomato plants growing in the Fe-free nutrient solution, the exposure to the pH of 7.5 was limited to the last two days.

Roots of Fe-deficient and Fe-sufficient tomato plants (27 d-old) were harvested five hours after the beginning of light phase. The collected roots were immediately frozen in liquid nitrogen and stored until further processing at -80°C. The collection was repeated in three independent cultivations and the roots from six plants were pooled for each treatment.

Ferric-chelate reduction

To determine the root capacity to reduce Fe(III)-EDTA, accordingly to Pinton et al. [89] roots of a single intact (Fe-sufficient or Fe-deficient) tomato plants were incubated in the dark at 25°C for 60 min in 50 mL of an aerated solution containing CaSO₄ 0.5 mM, BPDS 0.5 mM, Hepes-KOH 10 mM (pH 5.5) and 0.25 mM of Fe (III)-EDTA. Thereafter, the absorbance of the solutions at 535 nm was measured at intervals of 15 min and the amount of Fe(III) reduced calculated by the concentration of the Fe(II)-BPDS₃ complex formed, using an extinction coefficient of 22.1 mM⁻¹ cm⁻¹.

Microarray analysis

Transcriptional analysis was carried out using a Combi-matrix [24], produced by the Plant Functional Genomics Center, University of Verona [90]. The chip (TomatoArray2.0) carries 25,789 nonredundant probes (23,282

unique probes and 2,507 probes with more than one target) randomly distributed in triplicate across the array, each comprising a 35-40-mer oligonucleotide designed using the program oligoarray 2.1 [91]. The source of sequence information included tentative consensus sequences (TCs) derived from the DFCI Tomato Gene Index [92] Release 12.0 and expressed sequence tags. Eight bacterial oligonucleotide sequences provided by CombiMatrix, 8 probes designed on 8 Ambion spikes and 40 probes based on *Bacillus anthracis*, *Haemophilus ducreyi* and *Alteromonas phage* sequences were used as negative controls. Complete description of chip is available at the Gene Expression Omnibus [93] under the series entry (GPL13934).

Total RNA was isolated using the Spectrum™ Plant Total RNA kit (Sigma-Aldrich) and quantified by spectrophotometry using NanoDrop™ 1000 (Thermo Scientific). RNA quality was evaluated using Agilent 2100 Bioanalyzer (Agilent). Total RNA (1 µg) was amplified and labelled using the RNA ampULSe kit (Kreatech). After checking the quantity and quality of aRNA by spectrophotometry using NanoDrop™ 1000 (Thermo Scientific) and the quality subsequent labelling, 4 µg of labelled aRNA was hybridized to the array according to the manufacturer's recommendations [24]. Pre-hybridization, hybridization, washing and imaging were performed according to the manufacturer's protocols. The array was scanned with an Axon GenePix® 4400A scanner (MDS Analytical Technologies).

Analysis of raw data was performed using the open source software of the Bioconductor project [94,95] with the statistical R programming language [96,97]. Background adjustment, summarization and quantile normalization were performed using limma package [25]. Differentially expressed probes were identified by linear models analysis [25] using limma package and applying Bayesian correction, adjusted p-value of 0.05 and a |FC| ≥ 2. All microarray expression data are available at the Gene Expression Omnibus [93] under the series entry (GSE31112). Genes were grouped in main functional categories according to the "biological" terms of the Gene Ontology [27] assigned to each tomato TC or EST (Release 12.0) on the basis of the results of BlastP analysis [98] against the UniProt database [26] (Additional file 1). Genes without significant BlastP results were classified as "no hits found" (Evalue < 1e-8; identity > 40%).

Real-time RT-PCR experiments

0.5 µg of total RNA (checked for quality and quantity using a spectrophotometer NanoDrop™ 1000 (Thermo Scientific), followed by a migration in an agarose gel) of each sample was retrotranscribed using 1 pmol of Oligo d(T)23VN (New England Biolabs, Beverly, USA) and 10 U M-MulV RNase H for 1 h at 42°C (Finnzymes,

Helsinki, Finland) following the application protocol of the manufacturers. After RNA digestion with 1 U RNase A (USB, Cleveland, USA) for 1 h at 37°C, gene expression analyses were performed by adding 0.16 µl of the cDNA to the real-time PCR complete mix, FluoCycle™ sybr green (20 µl final volume; Euroclone, Pero, Italy), in a DNA Engine Opticon Real-Time PCR Detection (Biorad, Hercules, USA). Specific primers (Tm = 58°C) were designed to generate 80-140 bp PCR products (Additional file 2). Three genes were used as housekeeping to normalized the data: *elongation factor 1-alpha EF1a* (X14449; TC203463; forward: 5'- TGGATATGCTCCAGTGCTTG-3'; reverse: 5'- TTCCTTACCTGAACGCCTGT-3'), *histone H1* (AJ224933; TC192148; forward: 5'- CAAAGGC-CAAACTGCTACC-3'; reverse: 5'-AGGCTTTA-CAGCTGCTTTTCG-3') and *ubiquitin Ubi3* (X58253; TC196208; forward: 5'-AGCCAAAGAAGATCAAG-CACA-3'; reverse: 5'-GCCTCTGAACCTTTCCAGTG-3'). Each Real-Time RT-PCR was performed 4 times on 3 independent experiments; analyses of real-time result were performed using Opticon Monitor 2 software (Biorad, Hercules, USA) and R [93] with the qPCR package [99]. Efficiencies of amplification were calculated following the authors' indications [100]: PCR efficiencies were 99.15%, 89.16% and 87.25%, for *EF1a*, *H1* and *Ubi3* genes, respectively. The efficiencies for TC191581, TC192292, TC194139, TC216882, TC205660, TC208376 and NP796451 were respectively 94.75, 85.03, 98.92, 91.55, 92.70, 96.61 and 92.43%. The reported Real time data were normalized on the *EF1a* gene. Gene expression data were illustrated considering the differences in the amplification efficiency of PCR and using the gene expression levels in roots of Fe-sufficient plants as reference; applying the following formula:

$$\text{gene exp resion } x_{-Fe} = \frac{(2 \times E_x)^{[C_t(x+Fe) - C_t(x-Fe)]}}{(2 \times E_y)^{[C_t(y+Fe) - C_t(y-Fe)]}}$$

Where: E_x or E_y is the percentage value of PCR efficiency for the amplification of the gene x or y, respectively; $C_t(x_{+Fe})$ C_t for the control treatment (+Fe) and the considered gene (x); $C_t(x_{-Fe})$ C_t for the treated roots (-Fe) and the considered gene (x); $C_t(y_{+Fe})$ C_t for the control treatment (+Fe) and the housekeeping gene (y); $C_t(y_{-Fe})$ C_t for the treated roots (-Fe) and the housekeeping gene (y).

Additional material

Additional file 1: Functional annotation of 97 differentially expressed transcripts. ProbelD, Fold change, adjusted p-value, reference Tentative Consensus sequence in DFCI Tomato Gene Index (Release 12.0) <http://compbio.dfci.harvard.edu/cgi-bin/tgi/gimain.pl?gudb=tomato>.

description. UniProtID <http://www.uniprot.org/>. Biological process GO term and E-value are reported for each probe.

Additional file 2: Primer sequences of Real-time RT-PCR experiment.
TC ID, description and sequences of forward and reverse primers are reported.

Acknowledgements

This work was supported by grants: MIUR (FIRB-Futuro in Ricerca and PRIN), Cariverona and Internal Projects UniBZ.

Author details

¹Department of Biotechnology, University of Verona, via delle Grazie 15, 37134 Verona, Italy. ²Department of Agriculture and Environmental Sciences, University of Udine, via delle Scienze 208, 33100 Udine, Italy. ³Faculty of Science and Technology, Free University of Bolzano, piazza Università 5, 39100 Bolzano, Italy.

Authors' contributions

AZ and LZ made a substantial contribution to data collection and interpretation and manuscript drafting. NT participated in the project's design, data analysis and manuscript revision. MP critically revised the manuscript. RP contributed to data interpretation and critically revised the manuscript. ZV contributed to data interpretation and manuscript writing. SC participated in the project's design and coordination and critically revised the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 3 August 2011 Accepted: 20 March 2012

Published: 20 March 2012

References

- Hansen NC, Hopkins BG, Ellsworth JW, Jolley VD: **Iron nutrition in field crops.** In *Iron nutrition in plants and rhizospheric microorganisms*. 1 edition. Edited by: Barton LL, Abadia J. Dordrecht: Springer; 2007:23-59.
- Hansch R, Mendel RR: **Physiological functions of mineral micronutrients (Cu, Zn, Mn, Fe, Ni, Mo, B, Cl).** *Curr Opin Plant Biol* 2009, **12**:259-266.
- Marschner H, Romheld V, Kissel M: **Different strategies in higher plants in mobilization and uptake of iron.** *J Plant Nutr* 1986, **9**:695-713.
- Ivanov R, Brumbarova T, Bauer P: **Fitting into the Harsh Reality: Regulation of Iron-deficiency Responses in Dicotyledonous Plants.** *Mol Plant* 2012, **5**:27-42.
- Thimm O, Essigmann B, Kloska S, Altmann T, Buckhout TJ: **Response of Arabidopsis to iron deficiency stress as revealed by microarray analysis.** *Plant Physiol* 2001, **127**:1030-1043.
- Colangelo EP, Guerinot ML: **The essential basic helix-loop-helix protein FIT1 is required for the iron deficiency response.** *Plant Cell* 2004, **16**:3400-3412.
- Dinneny JR, Long TA, Wang JY, Jung JW, Mace D, Pointer S, Barron C, Brady SM, Schiefelbein J, Benfey PN: **Cell identity mediates the response of Arabidopsis roots to abiotic stress.** *Science* 2008, **320**:942-945.
- Buckhout TJ, Yang TJ, Schmidt W: **Early iron-deficiency-induced transcriptional changes in Arabidopsis roots as revealed by microarray analyses.** *BMC Genomics* 2009, **10**:147.
- Forner-Giner MA, Llosá MJ, Carrasco JL, Pérez-Amador MA, Navarro L, Ancillo G: **Differential gene expression analysis provides new insights into the molecular basis of iron deficiency stress response in the citrus rootstock Poncirus trifoliata (L.) Raf.** *J Exp Bot* 2010, **61**:483-490.
- García MJ, Lucena C, Romera FJ, Alcántara E, Pérez-Vicente R: **Ethylene and nitric oxide involvement in the up-regulation of key genes related to iron acquisition and homeostasis in Arabidopsis.** *J Exp Bot* 2010, **61**:3885-3899.
- Yang TJ, Lin WD, Schmidt W: **Transcriptional profiling of the Arabidopsis iron deficiency response reveals conserved transition metal homeostasis networks.** *Plant Physiol* 2010, **152**:2130-2141.
- O'Rourke JA, Graham MA, Vodkin L, Gonzalez DO, Cianzio SR, Shoemaker RC: **Recovering from iron deficiency chlorosis in near-isogenic soybeans: a microarray study.** *Plant Physiol Biochem* 2007, **45**:287-292.
- O'Rourke JA, Charlson DV, Gonzalez DO, Vodkin LO, Graham MA, Cianzio SR, Grusak MA, Shoemaker RC: **Microarray analysis of iron deficiency chlorosis in near-isogenic soybean lines.** *BMC Genomics* 2007, **8**:476.
- O'Rourke JA, Nelson RT, Grant D, Schmutz J, Grimwood J, Cannon S, Vance CP, Graham MA, Shoemaker RC: **Integrating microarray analysis and the soybean genome to understand the soybeans iron deficiency response.** *BMC Genomics* 2009, **10**:376.
- Brumbarova T, Matros A, Mock HP, Bauer P: **A proteomic study showing differential regulation of stress, redox regulation and peroxidase proteins by iron supply and the transcription factor FER.** *Plant J* 2008, **54**:321-334.
- Li J, Wu XD, Hao ST, Wang XJ, Ling HQ: **Proteomic response to iron deficiency in tomato root.** *Proteomics* 2008, **8**:2299-2311.
- Rellán-Alvarez R, Andaluz S, Rodríguez-Celma J, Wohlgemuth G, Zocchi G, Alvarez-Fernández A, Fiehn O, López-Millán AF, Abadía J: **Changes in the proteomic and metabolic profiles of Beta vulgaris root tips in response to iron deficiency and resupply.** *BMC Plant Biol* 2010, **10**:120.
- Donnini S, Prinsi B, Negri AS, Viganì G, Espen L, Zocchi G: **Proteomic characterization of iron deficiency responses in Cucumis sativus L. roots.** *BMC Plant Biol* 2010, **10**:268.
- Lan P, Li W, Wen TN, Shiao JY, Wu YC, Lin W, Schmidt W: **iTRAQ protein profile analysis of Arabidopsis roots reveals new aspects critical for iron homeostasis.** *Plant Physiol* 2011, **155**:821-834.
- Rodríguez-Celma J, Lattanzio G, Grusak MA, Abadía A, Abadía J, López-Millán AF: **Root responses of Medicago truncatula plants grown in two different iron deficiency conditions: changes in root protein profile and riboflavin biosynthesis.** *J Proteome Res* 2011, **10**:2590-2601.
- Schmidt W, Buckhout TJ: **A hitchhiker's guide to the Arabidopsis ferrome.** *Plant Physiol Biochem* 2011, **49**:462-470.
- Tomasi N, Rizzardo C, Monte R, Gottardi S, Jelali N, Terzano R, Vekemans B, De Nobili M, Varanini Z, Pinton R, Cesco S: **Micro-analytical, physiological and molecular aspects of Fe acquisition in leaves of Fe-deficient tomato plants re-supplied with natural Fe-complexes in nutrient solution.** *Plant Soil* 2009, **325**:25-38.
- Zouari M, Abadía A, Abadía J: **Iron is required for the induction of root ferric chelate reductase activity in iron-deficient tomato.** *J Plant Nutr* 2001, **24**:383-396.
- CombiMatrix Diagnostics. [<http://www.combimatrix.com>].
- Smyth GK: **Linear models and empirical Bayes methods for assessing differential expression in microarray experiments.** *Stat Appl Genet Mol Biol* 2004, **3**, Article 3.
- The UniProt Database. [<http://www.uniprot.org/>].
- The Gene Ontology. [<http://www.geneontology.org>].
- Eckhardt U, Mas Marques A, Buckhout TJ: **Two iron-regulated cation transporters from tomato complement metal uptake-deficient yeast mutants.** *Plant Mol Biol* 2001, **45**:437-448.
- Bereczky Z, Wang HY, Schubert V, Ganai M, Bauer P: **Differential regulation of nramp and irt metal transporter genes in wild type and iron uptake mutants of tomato.** *J Biol Chem* 2003, **278**:24697-24704.
- Zuchi S, Cesco S, Varanini Z, Pinton R, Astolfi S: **Sulphur deprivation limits Fe-deficiency responses in tomato plants.** *Planta* 2009, **230**:85-94.
- Li L, Cheng X, Ling HQ: **Isolation and characterization of Fe(III)-chelate reductase gene LeFRO1 in tomato.** *Plant Mol Biol* 2004, **54**:125-136.
- Ling HQ, Bauer P, Bereczky Z, Keller B, Ganai M: **The tomato fer gene encoding a bHLH protein controls iron-uptake responses in roots.** *Proc Natl Acad Sci USA* 2002, **99**:13938-13943.
- Brumbarova T, Bauer P: **Iron-mediated control of the basic helix-loop-helix protein FER, a regulator of iron uptake in tomato.** *Plant Physiol* 2005, **137**:1018-1026.
- Bauer P, Thiel T, Klatter M, Bereczky Z, Brumbarova T, Hell R, Grosse I: **Analysis of sequence, map position, and gene expression reveals conserved essential genes for iron uptake in Arabidopsis and tomato.** *Plant Physiol* 2004, **136**:4169-4183.
- Yuan Y, Wu H, Wang N, Li J, Zhao W, Du J, Wang D, Ling HQ: **FIT interacts with AtbHLH38 and AtbHLH39 in regulating iron uptake gene expression for iron homeostasis in Arabidopsis.** *Cell Res* 2008, **18**:385-397.
- The Arabidopsis Information Resource. [<http://www.arabidopsis.org>].

37. Briat JF, Duc C, Ravet K, Gaymard F: **Ferritins and iron storage in plants.** *Biochim Biophys Acta* 2010, **1800**:806-814.
38. Gollhofer J, Schläwicke C, Jungnick N, Schmidt W, Buckhout TJ: **Members of a small family of nodulin-like genes are regulated under iron deficiency in roots of *Arabidopsis thaliana*.** *Plant Physiol Biochem* 2011, **49**:557-564.
39. Espen L, Dell'Orto M, De Nisi P, Zocchi G: **Metabolic responses in cucumber (*Cucumis sativus* L.) roots under Fe-deficiency: a ³¹P-nuclear magnetic resonance in vivo-study.** *Planta* 2000, **210**:985-992.
40. Zocchi G: **Metabolic changes in iron-stressed dicotyledonous plants.** In *Iron nutrition in plants and rhizospheric microorganisms*. Edited by: Barton LL, Abadía J. Dordrecht, Springer; 2007:359-370.
41. López-Millán AF, Morales F, Gogorcena Y, Abadía A, Abadía J: **Metabolic responses in iron deficient tomato plants.** *J Plant Physiol* 2009, **166**:375-384.
42. Scheidig A, Fröhlich A, Schulze S, Lloyd JR, Kossmann J: **Down-regulation of a chloroplast-targeted beta-amylase leads to a starch-excess phenotype in leaves.** *Plant J* 2002, **30**:581-591.
43. López-Millán AF, Morales F, Andaluz S, Gogorcena Y, Abadía A, De Las Rivas J, Abadía J: **Responses of sugar beet roots to iron deficiency. Changes in carbon assimilation and oxygen use.** *Plant Physiol* 2000, **124**:885-898.
44. Viganì G, Maffi D, Zocchi G: **Iron availability affects the function of mitochondria in cucumber roots.** *New Phytol* 2009, **182**:127-136.
45. Kobayashi T, Nishizawa NK, Mori S: **Iron nutrition in field crops.** In *Iron nutrition in plants and rhizospheric microorganisms*. 1 edition. Edited by: Barton LL, Abadía J. Dordrecht: Springer; 2007:395-435.
46. Kobayashi T, Suzuki M, Inoue H, Itai RN, Takahashi M, Nakanishi H, Mori S, Nishizawa NK: **Expression of iron-acquisition-related genes in iron-deficient rice is co-ordinately induced by partially conserved iron-deficiency-responsive elements.** *J Exp Bot* 2005, **56**:1305-1316.
47. Zheng L, Huang F, Narsai R, Wu J, Giraud E, He F, Cheng L, Wang F, Wu P, Whelan J, Shou H: **Physiological and transcriptome analysis of iron and phosphorus interaction in rice seedlings.** *Plant Physiol* 2009, **151**:262-274.
48. Astolfi S, Zuchi S, Hubberten HM, Pinton R, Hoefgen R: **Supply of sulphur to S-deficient young barley seedlings restores their capability to cope with iron shortage.** *J Exp Bot* 2010, **61**:799-806.
49. Mazzucotelli E, Mastrangelo AM, Crosatti C, Guerra D, Stanca AM, Cattivelli L: **Abiotic stress response in plants: When post-transcriptional and post-translational regulations control transcription.** *Plant Sci* 2008, **174**:420-431.
50. Pontiggia A, De Nisi P, Zocchi G: **Effect of iron deficiency on RNA and protein synthesis in cucumber roots.** *J Plant Nutr* 2003, **10-11**:2177-2186.
51. Römheld V, Marschner H: **Mobilization of iron in the rhizosphere of different plant species.** *Adv Plant Nutr* 1986, **2**:155-204.
52. Tomasi N, Weisskopf L, Renella G, Landi L, Pinton R, Varanini Z, Nannipieri P, Torrent J, Martinoia E, Cesco S: **Flavonoids of white lupin roots participate in phosphorus mobilization from soil.** *Soil Biol Biochem* 2008, **40**:1971-1974.
53. Jin CW, You GY, He YF, Tang CX, Wu P, Zheng SJ: **Iron-deficiency-induced secretion of phenolics facilitates the reutilization of root apoplastic iron in red clover (*Trifolium pratense* L.).** *Plant Physiol* 2007, **144**:278-285.
54. Masada S, Terasaka K, Oguchi Y, Okazaki S, Mizushima T, Mizukami H: **Functional and structural characterization of a flavonoid glucoside 1,6-glucosyltransferase from *Catharanthus roseus*.** *Plant Cell Physiol* 2009, **50**:1401-1415.
55. Cesco S, Neumann G, Tomasi N, Pinton R, Weisskopf L: **Release of plant-borne flavonoids into the rhizosphere and their role in plant nutrition.** *Plant Soil* 2010, **329**:1-25.
56. Machold O: **Einfluss der Ernährungsbedingungen auf den Zustand des Eisens in den Blättern, den Chlorophyllgehalt und die Katalase- sowie Peroxydaseaktivität.** *Flora (Jena), Abt A* 1968, **159**:1-25.
57. Ranieri A, Castagna A, Baldan B, Soldatini GF: **Iron deficiency differently affects peroxidase isoforms in sunflower.** *J Exp Bot* 2001, **52**:25-35.
58. Lou Y, Baldwin IT: **Silencing of a germin-like gene in *Nicotiana attenuata* improves performance of native herbivores.** *Plant Physiol* 2006, **140**:1126-1136.
59. Römheld V, Marschner H: **Iron deficiency stress induced morphological and physiological changes in root tips of sunflower.** *Physiol Plant* 1981, **53**:354-360.
60. Römheld V, Kramer D: **Relationship between proton efflux and rhizodermal transfer cells induced by iron deficiency.** *Z Pflanzenphysiol* 1983, **113**:73-83.
61. Schmidt W: **Mechanisms and regulation of reduction-based iron uptake in plants.** *New Phytol* 1999, **141**:1-26.
62. Schikora A, Schmidt W: **Iron stress-induced changes in root epidermal cell fate are regulated independently from physiological responses to low iron availability.** *Plant Physiol* 2001, **125**:1679-1687.
63. Schikora A, Schmidt W: **Formation of transfer cells and (H⁺)-ATPase expression in tomato roots under P and Fe deficiency.** *Planta* 2002, **215**:304-311.
64. Bucher M, Brunner S, Zimmermann P, Zardi GI, Amrhein N, Willmitzer L, Riesmeier JW: **The expression of an extensin-like protein correlates with cellular tip growth in tomato.** *Plant Physiol* 2002, **128**:911-923.
65. Bucher M, Schroeder B, Willmitzer L, Riesmeier JW: **Two genes encoding extensin-like proteins are predominantly expressed in tomato root hair cells.** *Plant Mol Biol* 1997, **35**:497-508.
66. Cosgrove DJ: **Enzymes and other agents that enhance cell wall extensibility.** *Annu Rev Plant Physiol Plant Mol Biol* 1999, **50**:391-417.
67. Chen CY, Wong EI, Vidali L, Estavillo A, Hepler PK, Wu HM, Cheung AY: **The regulation of actin organization by actin-depolymerizing factor in elongating pollen tubes.** *Plant Cell* 2002, **14**:2175-2190.
68. Bots M, Vergeldt F, Wolters-Arts M, Weterings K, van As H, Mariani C: **Aquaporins of the PIP2 class are required for efficient anther dehiscence in tobacco.** *Plant Physiol* 2005, **137**:1049-1056.
69. Wintz H, Fox T, Wu YY, Feng V, Chen W, Chang HS, Zhu T, Vulpe : **Expression Profiles of *Arabidopsis thaliana* in Mineral Deficiencies Reveal Novel Transporters Involved in Metal Homeostasis.** *J Biol Chem* 2003, **278**:47644-47653.
70. Lubkowitz M: **The oligopeptide transporters: a small gene family with a diverse group of substrate and functions?** *Mol Plant* 2011, **4**:407-415.
71. Stacey MG, Osawa H, Patel A, Gassmann W, Stacey G: **Expression analyses of *Arabidopsis* oligopeptide transporters during seed germination, vegetative growth and reproduction.** *Planta* 2006, **223**:291-305.
72. Vasconcelos MW, Li GW, Lubkowitz MA, Grusak MA: **Characterization of the PT clade of oligopeptide transporters in rice.** *Plant Genome* 2008, **1**:77-88.
73. Lauter FR, Ninnemann O, Bucher M, Riesmeier JW, Frommer WB: **Preferential expression of an ammonium transporter and of two putative nitrate transporters in root hairs of tomato.** *Proc Natl Acad Sci USA* 1996, **93**:8139-8144.
74. von Würlén N, Lauter FR, Ninnemann O, Gillissen B, Walch-Liu P, Engels C, Jost W, Frommer WB: **Differential regulation of three functional ammonium transporter genes by nitrogen in root hairs and by light in leaves of tomato.** *Plant J* 2000, **21**:167-175.
75. Tomson CJ, Marschner H, Römheld V: **Effect of nitrogen fertilizer form on pH of the bulk soil and rhizosphere, and on the growth, phosphorus, and micronutrient uptake of bean.** *J Plant Nutr* 1993, **16**:493-506.
76. Nikolic M, Cesco S, Römheld V, Varanini Z, Pinton R: **Short-term interactions between nitrate and iron nutrition in cucumber.** *Funct Plant Biol* 2007, **34**:402-408.
77. Kovermann P, Meyer S, Hörtensteiner S, Picco C, Scholz-Starke J, Ravera S, Lee Y, Martinoia E: **The *Arabidopsis* vacuolar malate channel is a member of the ALMT family.** *Plant J* 2007, **52**:1169-1180.
78. Abadía J, López-Millán AF, Rombolà A, Abadía A: **Organic acid and Fe deficiency: a review.** *Plant Soil* 2002, **241**:75-86.
79. Romera FJ, Lucena C, Alcántara E: **Plant hormones influencing iron uptake in plants.** In *Iron nutrition in plants and rhizospheric microorganisms*. 1 edition. Edited by: Barton LL, Abadía J. Dordrecht: Springer; 2007:251-278.
80. Romera FJ, García MJ, Alcántara E, Pérez-Vicente R: **Latest findings about the interplay of auxin, ethylene and nitric oxide in the regulation of Fe deficiency responses by Strategy I plants.** *Plant Signal Behav* 2011, **6**:167-170.
81. van der Merwe MJ, Osorio S, Moritz T, Nunes-Nesi A, Fernie AR: **Decreased mitochondrial activities of malate dehydrogenase and fumarase in tomato lead to altered root growth and architecture via diverse mechanisms.** *Plant Physiol* 2009, **149**:653-669.
82. Martí E, Carrera E, Ruiz-Rivero O, García-Expósito JL: **Hormonal regulation of tomato gibberellin 20-oxidase1 expressed in *Arabidopsis*.** *J Plant Physiol* 2010, **167**:1188-1196.

83. Séguéla M, Briat JF, Vert G, Curie C: Cytokinins negatively regulate the root iron uptake machinery in *Arabidopsis* through a growth-dependent pathway. *Plant J* 2008, **55**:289-300.
84. Werner T, Köllmer I, Bartrina I, Holst K, Schmölling T: New insights into the biology of cytokinin degradation. *Plant Biol* 2006, **8**:371-381.
85. Miyazaki S, Koga R, Bohnert HJ, Fukuhara T: Tissue- and environmental response-specific expression of 10 PP2C transcripts in *Mesembryanthemum crystallinum*. *Mol Gen Genet* 1999, **261**:307-316.
86. Guo Y, Halfter U, Ishitani M, Zhu JK: Molecular characterization of functional domains in the protein kinase SOS2 that is required for plant salt tolerance. *Plant Cell* 2001, **13**:1383-1400.
87. Gong D, Guo Y, Schumaker KS, Zhu JK: The SOS3 family of calcium sensors and SOS2 family of protein kinases in *Arabidopsis*. *Plant Physiol* 2004, **134**:919-926.
88. Navarro L, Zipfel C, Rowland O, Keller I, Robatzek S, Boller T, Jones JD: The transcriptional innate immune response to flg22. Interplay and overlap with Avr gene-dependent defense responses and bacterial pathogenesis. *Plant Physiol* 2004, **135**:1113-1128.
89. Pinton R, Cesco S, Santi S, Agnolon F, Varanini Z: Water-extractable humic substances enhance iron deficiency responses by Fe-deficient cucumber plants. *Plant Soil* 1999, **210**:145-157.
90. Functional Genomic Center. [<http://dclab.sci.univr.it/FunctionalGenomics>].
91. Rouillard JM, Zuker M, Gulari E: OligoArray 2.0: design of oligonucleotide probes for DNA microarrays using a thermodynamic approach. *Nucleic Acid Res* 2003, **31**:3057-3062.
92. The DFCI Tomato Gene Index. [<http://compbio.dfci.harvard.edu/tgi/cgi-bin/tgi/gimain.pl?gudb=tomato>].
93. Gene Expression Omnibus. [<http://www.ncbi.nlm.nih.gov/geo>].
94. Gentleman RC, Carey VJ, Bates DM, Bolstad B, Dettling M, Dudoit S, Ellis B, Gautier L, Ge Y, Gentry J, Hornik K, Hothorn T, Huber W, Iacus S, Irizarry R, Leisch F, Li C, Maechler M, Rossini AJ, Sawitzki G, Smith C, Smyth G, Tierney L, Yang JY, Zhang J: Bioconductor: open software development for computational biology and bioinformatics. *Genome Biol* 2004, **5**:R80.
95. The Bioconductor project. [<http://www.bioconductor.org/>].
96. Ihaka R, Gentleman R: R: A language for data analysis and graphics. *J Comput Graph Stat* 1996, **5**:299-314.
97. The R Project for Statistical Computing. [<http://www.r-project.org/>].
98. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ: Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acid Res* 1997, **25**:3389-3402.
99. The qpcR library. [<http://www.dr-spiess.de/qpcR.html>].
100. Ritz C, Spiess AN: qpcR: an R package for sigmoidal model selection in quantitative real-time polymerase chain reaction analysis. *Bioinformatics* 2008, **24**:1549-155.

doi:10.1186/1471-2164-13-101

Cite this article as: Zamboni *et al.*: Genome-wide microarray analysis of tomato roots showed defined responses to iron deficiency. *BMC Genomics* 2012 **13**:101.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

