

Published in final edited form as:

Neuro Endocrinol Lett. 2011 ; 32(6): 763–768.

Dehydration increases sodium-dependent glutamate uptake by hypothalamic paraventricular nucleus synaptosomes

Thomas S. King, PhD¹, Glenn M. Toney², Pei-Yu Tian⁴, and Martin A. Javors^{3,4}

¹ Department of Cellular and Structural Biology, The University of Texas Health Science Center, San Antonio, Texas, USA

² Department of Physiology, The University of Texas Health Science Center, San Antonio, Texas, USA

³ Department of Pharmacology, The University of Texas Health Science Center, San Antonio, Texas, USA

⁴ Department of Psychiatry, The University of Texas Health Science Center, San Antonio, Texas, USA

Abstract

OBJECTIVES—The purpose of this study was to identify and characterize Na⁺-dependent, high affinity glutamate transporter (GLUT) activity in the hypothalamic paraventricular nucleus (PVN) and to compare GLUT activity in PVN of euhydrated versus water-deprived rats.

METHODS—Sprague-Dawley rats were deprived of water for two days before sacrifice. Control rats received water ad libitum. After sacrifice, PVN and cerebrum were removed and synaptosomes were prepared using standard techniques. Glutamate uptake was measured using [³H]-glutamate as substrate, physiological buffer, approximately 100 mg of synaptosomal tissue per assay and a Brandel cell harvester.

RESULTS—Glutamate uptake was saturable in PVN synaptosomes from euhydrated, control rats with a V_{max} of 541 ± 22 pmol/min-mg protein (SEM) and K_m of 17.6 ± 3.8 nM (SEM). In contrast, V_{max} of glutamate uptake was 808 ± 58 pmol/min-mg protein in PVN of rats deprived of water for 2 days. This was significantly higher than controls (*p*<0.001). K_m was 21.2 ± 7.3 nM and not significantly different from controls (NS).

CONCLUSIONS—Our results suggest that water deprivation of rats results in significantly higher synaptosomal glutamate uptake in PVN. Although the exact mechanism is unknown, increased transcription of the GLUT gene and/or increased cell surface expression of GLUT may contribute to the observed increase of glutamate uptake in dehydrated rats. Increased glutamate uptake may serve to restrict dehydration-induced activation of PVN efferent pathways specifically involved in release of neurohypophysial hormones and activation of sympathetic outflow that operate to maintain body fluid balance and cardiovascular function.

Keywords

Glutamate; glutamate uptake; paraventricular nucleus; dehydration

INTRODUCTION

As one of the primary brain areas responsive to changes in osmolality, the hypothalamic paraventricular nucleus (PVN) plays a central role in regulating fluid homeostasis. This is accomplished through progressive recruitment of neuroendocrine and sympathetic nervous system responses. Discharge of PVN neurons is increased by acute elevations of plasma osmolality and plasma angiotensin II and is decreased by blood volume expansion (Toney *et al.* 2003; Bains & Ferguson 1995; Chen & Toney 2001; Chen & Toney 2003; Ferguson 1988; Ferguson & Washburn 1998; Lovick & Coote 1988). Water deprivation increases plasma osmolality and circulating angiotensin II and these effects are accompanied by activation of PVN neurons (Stocker *et al.* 2004). Activation of PVN neurons by water deprivation may also be facilitated by loss of inhibitory input due to the decrease of intravascular volume. Among numerous and varied output projections of the PVN are two that play particularly important roles in maintaining body fluid homeostasis. These are vasopressinergic and oxytocinergic neurons that project to the posterior pituitary and a neurochemically diverse group of neurons with axonal projections to the intermediolateral cell column (IML) of the spinal cord (Pyner & Coote 2000; Saper *et al.* 1976; Shafton *et al.* 1998; Swanson & Kuypers 1980) as well as to the rostral ventrolateral medulla (RVLM) (Pyner & Coote 1999; 2000; Shafton *et al.* 1998). Activity of the premotor neurons of the RVLM maintains sympathetic vasomotor tone and regulates various vasomotor reflexes including activation of the sympathoadrenal system, increased arterial blood pressure, elevated heart rate (Dampney 1994a,b; Guyenet *et al.* 1996).

Dehydration-induced changes in the PVN includes enhanced synaptic glutamate release, increased synaptic glutamate receptor density and activation and enhanced glutamate receptor-effector coupling (Li & Tasker 2004). Glutamate activity within individual synapses is terminated by uptake of the neurotransmitter via selective glutamate transporters located in neuronal and astrocyte cell membranes (Nicholls & Attwell 1990). An inability to effectively clear glutamate could result in its diffusion to heterosynaptic pathways. Such glutamate “spillover” could result in abnormal activation of functionally unrelated pathways. Pathway specificity is maintained through activation of glutamate uptake via selective glutamate transporters (Tsvetkov *et al.* 2004; Nie & Weng 2009). Limiting glutaminergic activity to specific synaptic pathways for defined time periods is accomplished both by reducing glutamate release and by reducing the amount of glutamate in the synaptic cleft. The latter is typically accomplished via glutamate diffusion and/or by enhanced transporter-mediated uptake (Nicholls & Attwell 1990).

Increased glutaminergic drive associated with dehydration leads to enhanced excitability of magnocellular neurosecretory neurons of the PVN and supraoptic nuclei (Hatton 1997; Miyata *et al.* 1994; Theodosis *et al.* 1993; Li & Tasker 2004) as well as to increased angiotensin II and glutaminergic drive to pre-autonomic neurons of the PVN (Chen *et al.* 2001; Freeman & Brook 2007). In this study, we sought to identify and partially characterize Na⁺-dependent, high affinity glutamate transporter (GLUT) activity in the PVN and to compare GLUT activity in PVN of euhydrated versus water-deprived rats. We also examined effects of angiotensin II on glutamate uptake in the rat PVN.

MATERIALS AND METHODS

Animals and Water Deprivation Study

Adult male Sprague–Dawley rats (Charles River Laboratories) weighing 250–375 g were housed in a temperature-controlled room (22–23 °C) with an automatically controlled cycle of 14 h:10 h light:dark (lights on at 0700 h daily). Control (euhydrated) rats were provided tap water and laboratory chow (Harlan Teklad LM-485, 0.3% NaCl) ad libitum.

Experimental (dehydrated) rats were deprived of water but not food for two days before sacrifice in accordance with IACUC approval. All experimental procedures conformed to National Institutes of Health Guidelines and were approved by the Institutional Animal Care and Use Committee (IACUC) of the University of Texas Health Science Center at San Antonio.

Collection of Periventricular Nucleus Tissue

Rats were decapitated and the brain rapidly removed. The hypothalamus was dissected from the surrounding brain. The paraventricular nucleus (PVN) was identified visually and removed by microdissection. For comparison with PVN glutamate uptake measurements, cerebellum and cerebrum tissues were also collected from some rats.

Preparation of Synaptosomes

Synaptosomes are prepared as previously described (Whittaker & Barker 1972; Robinson *et al.* 1991) using pooled rat PVN. Synaptosomes were also prepared from rat cerebrum and cerebellum for comparison to the rat PVN. Briefly, fresh 80–90 mg dissections of the PVN from two rats were homogenized by 12 strokes of a fitted Teflon homogenizer in 10 ml of 0.32 M sucrose buffer (pH 7.4) in a glass homogenizer at 4 °C. The supernatant was next centrifuged at 1,000 × g for 10 minutes. The supernatant from this step was further centrifuged at 11,000 × g for 20 minutes at 4 °C. The resultant pellet was gently resuspended in assay buffer to a concentration of 20 mg original tissue weight/mL and stored briefly on ice before use.

Determination of Synaptosomal Glutamate Uptake

Glutamate uptake was measured as previously described (Robinson *et al.* 1991) with modifications. Briefly, duplicate assays were done in a final volume of 2 mL assay buffer containing (in mM) NaCl (126), KCl (4.8), CaCl₂ (1.25), dextrose (11), NaH₂PO₄ (3.7), Na₂HPO₄ (12.7), MgSO₄ (1.4) at pH 7.4. Nonspecific uptake was determined in a second set of samples to which a 1,000-fold increase in cold glutamate over the expected K_m (3–5 μM) was added. Glutamate uptake was initiated by addition of 1 μCi of [³H]L-glutamate and various amounts of carrier L-glutamate (2.5–80 μM). Glutamate uptake was stopped by addition of ice-cold phosphate buffer and placing the tubes in an ice-cold water bath, followed immediately by rapid filtration of the suspension through Whatman GFB glass fiber filters using a Brandel vacuum apparatus. The filters were then washed three times with 0.32 sucrose buffer (pH 7.4) at 4 °C to remove extracellular radioactivity from the filters.

Importantly, we avoided the commonly used polyethylenimine (PEI) during harvest of synaptosomes on the glass fiber filters. PEI is a strongly cationic polymer. Because of its ability to neutralize excess anionic colloidal charge on membranes, PEI is often used to increase synapsome attachment to filters (Cohen & Nadler 1997; Vancha *et al.* 2004; Whittaker & Barker 1972). However, we found that use of PEI resulted in greater background filter counts in comparison to water washes. Instead, we included five 32 M sucrose washes to remove non-specific ³H-glutamate binding to assay filters.

Filters were placed in 5 mL of cytosint ES (ICN Biochemicals Inc., Irvine, CA) and radioactivity determined by scintillation spectrometry (Beckman Instruments, LS 6500) at an approximate efficiency of 45%. Protein was determined by the BioRad version of the Bradford protein assay (Bradford 1976). Concentration-dependence of glutamate uptake was determined at six concentrations (2.5, 5, 10, 20, 40 and 80 μM) of glutamate. All values are presented as means ± SEM. Eadie-Hofstee transformations of concentration-dependence

data and Hill plots were fitted by linear regression analysis (Prism, Graphpad Software Inc., La Jolla, CA).

Na⁺-dependent uptake was determined by measuring the difference between radiolabeled glutamate accumulated in Na⁺ and in choline (absent of any Na⁺) medium.

Statistics

Displacement curves were analyzed by ANOVA using the Graphpad PRISM 4.0 (Graphpad USA). For comparison of Km and Vmax between groups, two-tailed statistics were used throughout, with preset alpha level of significance of $p < 0.05$. Data in the text are expressed as mean \pm standard error of the mean (SEM). Each experiment was repeated six times (n=6).

RESULTS

Glutamate uptake was saturable (Figure 1) and Na⁺-dependent in synaptosomes from the rat PVN. Elimination of Na⁺ (by substitution with choline) from the uptake assay medium inhibited glutamate uptake (Figure 2). Glutamate uptake was also linear with time through 15 minutes. Metabolism of glutamate was considered negligible. Similar values for Km and Vmax of glutamate uptake were determined for synaptosomes prepared from rat cerebrum and cerebellum (data not shown).

Two-days of water deprivation increased Vmax of glutamate uptake but had no effect on Km of glutamate uptake in the male rat PVN. Glutamate uptake exhibited a Km of 17.6 ± 3.8 nM (SEM) and a Vmax of 541 ± 22 pmol/min-mg protein (SEM) in euhydrated (control) rats. In rats deprived of water for 3 days, Vmax of glutamate uptake was 808 ± 58 pmol/min-mg protein, which was significantly higher than controls ($p < 0.001$) whereas Km was 21.2 ± 7.3 nM, not significantly different from controls (NS). Angiotensin II ($2 \mu\text{M}$) increased Vmax (314.8 ± 10.6 pmol/min-mg protein vs. 254.7 ± 6.6 pmol/min-mg protein for controls/no angiotensin II), but not Km (6.5 ± 0.9 nM vs 4.1 ± 0.5 nM for controls/no angiotensin II), for glutamate uptake in the PVN of control rats.

DISCUSSION

The primary findings of this study are that two-day water deprivation results in increased Vmax, but no difference in Km, for glutamate uptake in the male rat PVN. The Vmax of glutamate uptake was higher in dehydrated rats (808 ± 58 pmol/min-mg protein) than in euhydrated rats (Vmax of 541 ± 22 pmol/min-mg protein). This suggests an increase in the number of available glutamate transporters in the dehydrated rats. In contrast, the Km for glutamate uptake did not differ between dehydrated rats (21.2 ± 7.3 nM) and euhydrated rats (Km of 17.6 ± 3.8 nM), suggesting no change in the affinity of the transporters for glutamate. Glutamate uptake into synaptosomes in the rat PVN was found to be both saturable and Na⁺ dependent.

PVN neurons play pivotal roles in maintenance of homeostasis by regulating pituitary release of hormones as well as CNS sympathetic outflow during periods of stress and fluid imbalance with associated hypertension (Allen 2002; de Wardener 2001, Miyakubo *et al.* 2002; Ranson *et al.* 1998). Glutamatergic, GABAergic, and noradrenergic afferents to the PVN finely tune PVN output (Boudaba *et al.* 1997; Li *et al.* 2003) and define pathway specificity during conditions of heightened glutamatergic activity. Dehydration is associated with increased glutamatergic tone and enhanced excitability of magnocellular neurons (Hatton 1997; Miyata *et al.* 1994; Theodosis *et al.* 1993; Li & Tasker 2004). Li and Tasker (2004) showed dehydration-induced increases in postsynaptic excitatory potentials interpreted as increased glutamate release sites based on paired-pulse facilitation analysis

indicating a lack of change in the probability of glutamate release. Dehydration induced swelling of magnocellular neurons, glial cell retraction away from these neurons and the formation of new glutaminergic and GABAergic synapses within the SON and PVN (Miyata *et al.* 1994; Tweedle & Hatton 1977). Increased release of glutamate normally induces increased glutamate uptake, either by increases in transporter production and membrane insertion or by translocation of extrasynaptic transporters to synaptic membrane sites (Danbolt 2001; Nicholls & Attwell 1990; Otis *et al.* 1996b; Takagaki 1976). During dehydration states selectively activated PVN pathways include those projecting to brain stem autonomic centers and to sympathetic preganglionic neurons located within the intermediolateral cell column of the spinal cord (Ranson *et al.* 1998; Shafton *et al.* 1998; Toth *et al.* 1999) and thus sympathetic outflow regulating cardiovascular as well as end organ functions; *e.g.*, renal sodium excretion.

Given the complexity of PVN outflow regulating various autonomic and neuroendocrine functions as well as the ubiquitous distribution of glutamate within and outside the PVN, activation of specific glutaminergic pathways is critical to body fluid regulation. Pathway specificity during conditions of heightened glutaminergic activity may depend on tight control over both duration and synapse-specific glutamate release. An inability to effectively clear such glutamate could result in diffusion of transmitter from the original synapse to heterosynaptic pathways. Such glutamate “spillover” might produce undesirable activation or even eventual synaptic plasticity in those alternate pathways. Because pathway specificity is maintained through activation of glutamate uptake via selective glutamate transporters located in glial and neuronal cell membranes (Tsvetkov *et al.* 2004; Nie & Weng 2009), an increase in glutamate uptake is predicted to avoid extrasynaptic diffusion of glutamate from the active synapse (Asztely *et al.* 1997) and thus possible activation of neighboring but otherwise functionally unrelated glutamate synapses and pathways (Asztely *et al.* 1997). Increases in glutamate uptake may also be important for regulating the duration of postsynaptic pathway activation (Mennerick & Zorumski 1994) and preventing local depletion of the neurotransmitter and/or desensitization of postsynaptic glutamate receptors (Dudel *et al.* 1988; Otis *et al.* 1996a; Otis *et al.* 1996b; Tang *et al.* 1989; Turecek *et al.* 2000; Trussell *et al.* 1989; Trussell & Fischbach 1989). Through such mechanisms, glutamate uptake in dehydration may play a key role in maintaining efficacy of transmission within neuroendocrine and autonomic pathways that preserve cardiovascular function and the ability of animals to forage for sources of water. Our observation of increased V_{max} , but not K_m , in glutamate uptake within the PVN in dehydrated rats is in keeping with increased glutamate release and maintenance of pathway specificity. Increased glutamate uptake thus serves to define pathway specificity between PVN and downstream neurons related to neuroendocrine and sympathetic outflow.

We also found that angiotensin II (2 μM) minimally increases V_{max} , but not K_m , for glutamate uptake in the rat PVN. Angiotensin II (AT1) receptors are densely distributed in the PVN (Gehlert *et al.* 1991) with numerous direct synaptic contacts between AT1 receptor expressing fibers and spinally projecting neurons (Li *et al.* 2003). Sympathetic output induced either by cardiac stimulation or by hyperosmolality can be attenuated by PVN infusion of the selective angiotensin II receptor antagonist losartan (Chen & Toney 2001). The relatively small increases observed in V_{max} in response to application of angiotensin II to synaptosomal tissue may be a reflection of suboptimal distribution of angiotensin II in the synaptosomal preparation or because glutamate transporter numbers were already near-maximally increased in the dehydrated animals. Alternatively, G-protein signaling pathways downstream of AT1R receptors that are dominantly expressed in the PVN may be optimal in synaptosomal preparations. Dehydration decreases intravascular volume and increases plasma osmolality and circulating Ang II levels (Stocker *et al.* 2002; Brooks *et al.* 2004). Angiotensin II (AT1) receptors are densely distributed in the PVN (Gehlert *et al.* 1991) with

numerous direct synaptic contacts between AT1 receptor expressing fibers and spinally projecting neurons (Li *et al.* 2003). Angiotensin II has been shown to increase the discharge of PVN neurons *in vivo* (Ferguson 1988) and *in vitro* (Cato & Toney 2005; Li *et al.* 2003). Excitation by angiotensin II involves both presynaptic inhibition of GABAergic synaptic activity (Li *et al.* 2003) as well as postsynaptically to activate a non-selective cation current (Cato & Toney 2005). Thus, angiotensin II has the potential to act in the PVN to increase sympathetic outflow during dehydration-induced homeostatic stress, as well as during cardiovascular diseases such as heart failure and hyper-tension in which angiotensin II actions are increased in the periphery and CNS. What role angiotensin II might play on glutaminergic tone within the PVN in dehydrated rats remains to be determined.

In summary, we found that two-day water deprivation increased glutamate uptake by PVN synaptosomes reflected as an increased V_{max} with no difference in K_m . Increased glutamate uptake may be a response to increased glutamatergic activity within the PVN following short-term dehydration. Increased glutamate uptake may serve to maintain pathway specificity within the PVN, thereby preserving relatively selective activation of sympathetic outflow and pituitary neuro-peptide (*i.e.*, vasopressin and oxytocin) release.

Acknowledgments

The authors want to acknowledge the technical assistance of Mr. Alfredo S. Calderon, Department of Physiology, UTHSCSA. These studies were funded in part by HL102310 (GMT).

REFERENCES

1. Allen AM. Inhibition of the hypothalamic paraventricular nucleus in spontaneously hypertensive rats dramatically reduces sympathetic vasomotor tone. *Hypertension*. 2002; 39:275–280. [PubMed: 11847197]
2. Anderson CM, Swanson RA. Astrocyte glutamate transport: review of properties, regulation, and physiological functions. *Glia*. 2000; 32:1–14. [PubMed: 10975906]
3. Asztely F, Erdemli G, Kullmann DM. Extrasynaptic glutamate spillover in the hippocampus: dependence on temperature and the role of active glutamate uptake. *Neuron*. 1997; 18:281–293. [PubMed: 9052798]
4. Bains JS, Ferguson AV. Paraventricular nucleus neurons projecting to the spinal cord receive excitatory input from the subfornical organ. *Am J Physiol Regul Integr Comp Physiol*. 1995; 268:R625–R633.
5. Boudaba C, Schrader LA, Tasker JG. Physiological evidence for local excitatory synaptic circuits in the rat hypothalamus. *J Neurophysiol*. 1997; 77:3396–3400. [PubMed: 9212285]
6. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem*. 1976; 72:248–254. [PubMed: 942051]
7. Brooks VL, Freeman KL, O'Donoghue TL. Acute and chronic increases in osmolality increase excitatory amino acid drive of the rostral ventrolateral medulla in rats. *Am J Physiol*. 2004; 287:R1359–R1368.
8. Cato JM, Toney GM. Angiotensin II excites paraventricular nucleus neurons that innervate the rostral ventrolateral medulla: an *in vitro* patch-clamp study in brain slices. *J Neurophysiol*. 2005; 93:403–413. [PubMed: 15356186]
9. Chen QH, Toney GM. AT(1)-receptor blockade in the hypothalamic PVN reduces central hyperosmolality-induced renal sympathoexcitation. *Am J Physiol*. 2001; 281:R1844–R1853.
10. Chen QH, Toney GM. Identification and characterization of two functionally distinct groups of spinal cord-projecting paraventricular nucleus neurons with sympathetic-related activity. *Neurosci*. 2003; 118:797–807.
11. Cohen SM, Nadler JV. Sodium-dependent proline and glutamate uptake by hippocampal synaptosomes during post-natal development. *Dev Brain Res*. 1997; 100:230–233. [PubMed: 9205813]

12. Danbolt NC. Glutamate uptake. *Prog Neurobiol.* 2001; 65:1–105. [PubMed: 11369436]
13. Dampney RA. The subretrofacial vasomotor nucleus: anatomical, chemical and pharmacological properties and role in cardiovascular regulation. *Prog Neurobiol.* 1994; 42:197–227. [PubMed: 8008825]
14. Dampney RA. Functional organization of central pathways regulating the cardiovascular system. *Physiol Rev.* 1994; 74:323–364. [PubMed: 8171117]
15. de Wardener HE. The hypothalamus and hypertension. *Physiol Rev.* 2001; 81:1599–1658. [PubMed: 11581498]
16. Ferguson AV. Paraventricular nucleus neurons projecting to the dorsomedial medulla are influenced by systemic angiotensin. *Brain Res Bull.* 1988; 20:197–201. [PubMed: 2836037]
17. Ferguson AV, Washburn DL. Angiotensin II: a peptidergic neurotransmitter in central autonomic pathways. *Prog Neurobiol.* 1998; 54:169–192. [PubMed: 9481797]
18. Freeman KL, Brooks VL. AT(1) and glutamatergic receptors in paraventricular nucleus support blood pressure during water deprivation. *Am J Physiol Integr Comp Physiol.* 2007; 292:R1675–R1682.
19. Gehlert DR, Gackenhaimer SL, Schober DA. Autoradio-graphic localization of subtypes of angiotensin II antagonist binding in the rat brain. *Neuroscience.* 1991; 44:501–514. [PubMed: 1944896]
20. Guyenet, PG.; Koshiya, N.; Huangfu, D.; Baraban, SC.; Stornetta, RL.; Li, YW. Role of medulla oblongata in generation of sympathetic and vagal outflows.. In: Holstege, G.; Bandler, R.; Saper, CB., editors. *The Emotional Motor System.* Prog Brain Res. Vol. 107. Elsevier; Amsterdam: 1996. p. 127-144.
21. Hatton GI. Function-related plasticity in hypothalamus. *Annu Rev Neurosci.* 1997; 20:375–397. [PubMed: 9056719]
22. Li DP, Chen SR, Pan HL. Angiotensin II stimulates spinally projecting paraventricular neurons through presynaptic disinhibition. *J Neurosci.* 2003; 23:5041–5049. [PubMed: 12832527]
23. Li S, Tasker JG. Dehydration-induced synaptic plasticity in magnocellular neurons of the hypothalamic supraoptic nucleus. *Endocrinology.* 2004; 145:5141–5149. [PubMed: 15297447]
24. Lovick TA, Coote JH. Effects of volume loading on paraventriculo-spinal neurones in the rat. *J Auton Nerv Syst.* 1988; 25:135–140. [PubMed: 3235776]
25. Mennerick S, Zorumski CF. Glial contributions to excitatory neurotransmission in cultured hippocampal cells. *Nature (London).* 1994; 368:59–62. [PubMed: 7906399]
26. Miyakubo H, Hayashi Y, Tanaka J. Enhanced response of subfornical organ neurons projecting to the hypothalamic paraventricular nucleus to angiotensin II in spontaneously hypertensive rats. *Auton Neurosci.* 2002; 95:131–136. [PubMed: 11871778]
27. Miyata S, Nakashima T, Kiyohara T. Structural dynamics of neural plasticity in the supraoptic nucleus of the rat hypothalamus during dehydration and rehydration. *Brain Res Bull.* 1994; 34:169–175. [PubMed: 8055345]
28. Nicholls D, Attwell D. The release and uptake of excitatory amino acids. *Trends Pharmacol Sci.* 1990; 11:462–468. [PubMed: 1980041]
29. Nie H, Weng H-R. Glutamate transporters prevent excessive activation of NMDA receptors and extrasynaptic glutamate spillover in the spinal dorsal horn. *J Neurophysiol.* 2009; 101:2041–2051. [PubMed: 19211657]
30. Otis T, Zhang S, Trussell LO. Direct Measurement of AMPA Receptor Desensitization Induced by glutamatergic synaptic transmission. *J Neurosci.* 1996a; 16:7496–7504. [PubMed: 8922405]
31. Otis TS, Wu YC, Trussell LO. Delayed clearance of transmitter and the role of glutamate transporters at synapses with multiple release sites. *J Neurosci.* 1996b; 16:1634–1644. [PubMed: 8774432]
32. Pyner S, Coote JH. Identification of an efferent projection from the paraventricular nucleus of the hypothalamus terminating close to spinally projecting rostral ventrolateral medullary neurons. *Neuroscience.* 1999; 88:949–957. [PubMed: 10363830]
33. Pyner S, Coote JH. Identification of branching paraventricular neurons of the hypothalamus that project to the rostroventrolateral medulla and spinal cord. *Neuroscience.* 2000; 100:549–556. [PubMed: 11098118]

34. Ranson RN, Motawei K, Pyner S, Coote JH. The paraventricular nucleus of the hypothalamus sends efferents to the spinal cord of the rat that closely appose sympathetic preganglionic neurons projecting to the stellate ganglion. *Exp Brain Res.* 1998; 120:164–172. [PubMed: 9629958]
35. Robinson MB, Hunte-Ensor M, Sinor J. Pharmacologically distinct sodium-dependent L-[³H] glutamate transport processes in rat brain. *Brain Research.* 1991; 44:196–202. [PubMed: 2039937]
36. Saper CB, Loewy AD, Swanson LW, Cowan WM. Direct hypothalamo-autonomic connections. *Brain Res.* 1976; 117:305–312. [PubMed: 62600]
37. Shafton AD, Ryan A, Badoer E. Neurons in the hypothalamic paraventricular nucleus send collaterals to the spinal cord and to the rostral ventrolateral medulla in the rat. *Brain Res.* 1998; 801:239–243. [PubMed: 9729407]
38. Stocker SD, Stricker EM, Sved AF. Arterial baroreceptors mediate the inhibitory effect of acute increases in arterial blood pressure on thirst. *Am J Physiol.* 2002; 282:R1718–R1729.
39. Stocker SD, Cunningham JT, Toney GM. Water deprivation increases Fos immunoreactivity in OVN autonomic neurons with projections to the spinal cord and rostral ventrolateral medulla. *Am J Physiol Regul Integr Comp Physiol.* 2004; 287:R1172–R1183. [PubMed: 15271657]
40. Swanson LW, Kuypers HG. The paraventricular nucleus of the hypothalamus: cytoarchitectonic subdivisions and organization of projections to the pituitary, dorsal vagal complex, and spinal cord as demonstrated by retrograde fluorescence double-labeling methods. *J Comp Neurol.* 1980; 194:555–570. [PubMed: 7451682]
41. Takagaki G. Properties of the uptake and release of glutamic acid by synaptosomes from rat cerebral cortex. *J Neurochem.* 1976; 27:1417–1425. [PubMed: 1003217]
42. Theodosis DT, Poulain DA. Activity-dependent neuronal glial and synaptic plasticity in the adult mammalian hypothalamus. *Neuroscience.* 1993; 57:501–535. [PubMed: 8309521]
43. Toney GM, Chen QH, Cato MJ, Stocker SD. Central osmotic regulation of sympathetic nerve activity. *Acta Physiol Scand.* 2003; 177:43–55. [PubMed: 12492778]
44. Toth ZE, Gallatz K, Fodor M, Palkovits M. Decussations of the descending paraventricular pathways to the brainstem and spinal cord autonomic centers. *J Comp Neurol.* 1999; 414:255–266. [PubMed: 10516595]
45. Trussell LO, Fischbach GD. Glutamate receptor desensitization and its role in synaptic transmission. *Neuron.* 1989; 3:209–218. [PubMed: 2576213]
46. Trussell LO, Thio LL, Zorumski CF, Fischbach GD. Rapid desensitization of glutamate receptors in vertebrate central neurons. *Proc Natl Acad Sci USA.* 1988; 85:4562–4566. [PubMed: 2898144]
47. Tsvetkov E, Ryong-Moon Shin R, Bolshakov VY. Glutamate uptake determines pathway specificity of long-term potentiation in the neural circuitry of fear conditioning. *Neuron.* 2004; 41:139–151. [PubMed: 14715141]
48. Tweedle CD, Hatton GI. Ultrastructural changes in the rat hypothalamic neurosecretory cells and their associated glia during minimal dehydration and rehydration. *Cell Tissue Res.* 1977; 181:59–72. [PubMed: 880623]
49. Vancha AR, Govindaraju S, Parsa VL, Jasti M, Gonzalez-Garcia M, Ballesteros RP. Use of polyethyleneimine polymer in cell culture as attachment factor and lipofection enhancer. *BMC Biotechnology.* 2004; 4:23–35. [PubMed: 15485583]
50. Whittaker, VP.; Barker, LA. The subcellular fractionation of brain tissue with special reference to the preparation of synaptosomes and their component organelles.. In: Fried, R., editor. *Methods of Neurochemistry.* Vol. 2. Marcel Dekker; New York: 1972. p. 1-52.




Fig. 1. Glutamate uptake in rat PVN synaptosomes. Data expressed as mean \pm SEM (n=6).




Fig. 2. Glutamate uptake in rat PVN synaptosomes: Na⁺ dependence. Data expressed as mean ± SEM (n=6).




Fig. 3.

Glutamate uptake was saturable in PVN synaptosomes from euhydrated, control rats with a K_m of 17.6 ± 3.8 nM (SEM) and a V_{max} of 541 ± 22 pmol/min-mg protein (SEM). In rats deprived of water for 3 days, V_{max} of glutamate uptake was 808 ± 58 pmol/min-mg protein, which was significantly higher than controls ($p < 0.001$). K_m was 21.2 ± 7.3 nM which was not significantly different from controls (NS). Data expressed as mean \pm SEM ($n=6$)