

Purification and Characterization of the Acetone Carboxylase of *Cupriavidus metallidurans* Strain CH34

Caroline Rosier,^a Natalie Leys,^b Céline Henoumont,^c Max Mergeay,^b and Ruddy Wattiez^a

Department of Proteomics and Microbiology, Interdisciplinary Center of Mass Spectrometry (CISMa), University of Mons, Mons, Belgium^a; Expert Group for Molecular and Cellular Biology, SCK•CEN, Mol, Belgium^b; and Department of General, Organic and Biomedical Chemistry, NMR and Molecular Imaging Laboratory, University of Mons, Mons, Belgium^c

Acetone carboxylase (Acx) is a key enzyme involved in the biodegradation of acetone by bacteria. Except for the *Helicobacteraceae* family, genome analyses revealed that bacteria that possess an Acx, such as *Cupriavidus metallidurans* strain CH34, are associated with soil. The Acx of CH34 forms the heterohexameric complex $\alpha_2\beta_2\gamma_2$ and can carboxylate only acetone and 2-butanone in an ATP-dependent reaction to acetoacetate and 3-keto-2-methylbutyrate, respectively.

Acetone is a toxic compound found in air, water, and soil, both naturally as a pollutant (5, 17) and also as a result of being produced by mammals and bacteria (8, 11, 12, 23). Acetone can also be degraded by various bacteria using a CO₂-dependent pathway including acetone carboxylase as a key enzyme, a property of increasing interest for bioremediation purposes (1–3, 6, 7, 9, 10, 15, 16, 18–22). Acetone carboxylase (Acx) is a member of a protein family that also contains acetophenone carboxylase and ATP-dependent hydantoinases/oxoprolinases. While the members of this family share several similar characteristics, they differ with respect to the substrates, the products of ATP hydrolysis, and structural properties (16).

Genome analyses revealed a lot of bacterial species that possess the acetone carboxylase and thus are potentially able to detoxify acetone (see Fig. S1 in the supplemental material). Most of these bacteria, such as *Cupriavidus metallidurans*, were found in soil or in contact with soil (e.g., by plant symbiosis) and belong to *Proteobacteria* and especially *Betaproteobacteria*. In the *Alphaproteobacteria* class, the *Rhizobiales* and the *Rhodobacterales* orders were found to contain an Acx. In the *Deltaproteobacteria* class, only one species (*Geobacter uraniireducens* Rf4), up to now, was discovered to contain an Acx, which had around 30% amino acid (aa) sequence identity with the CH34 Acx depending on the subunit. The only pathogenic species that possess the enzyme are those belonging to the *Helicobacteraceae* family (*Epsilonproteobacteria*) which are found in the mammalian stomach (4).

In general, similar gene organizations for the Acx operon were found, with three genes, *acxA*, *acxB*, and *acxC*, encoding the three acetone carboxylase subunits (β , α , and γ subunits, respectively) and one regulator, *acxR*, which was identified as a σ^{54} - or σ^{70} -specific transcriptional regulator and can be divergently transcribed (19). The only known paralogous enzyme with similar biochemical function is acetophenone carboxylase from *Aromatoleum aromaticum* (16). This enzyme consists of a hetero-octamer of four subunits whose corresponding genes *apcABCDE* are clustered as an operon. Acetone carboxylase does not contain a paralogue of *ApcE*.

Acetone carboxylase induction in *Cupriavidus metallidurans*. A recent study focused on acetone metabolism in *C. metallidurans* CH34, a betaproteobacterium found in industrial biotopes highly contaminated with metals (14), showing an overexpression of the acetone carboxylase when grown in spaceflight

conditions (13). As observed in *Rhodobacter capsulatus* and *Xanthobacter autotrophicus*, the *C. metallidurans* Acx subunits were induced at a high level (19% \pm 4% of the total proteins) when acetone was present in the culture (Fig. 1) (19, 20). An *acxR* knockout mutant was constructed in this study. This mutant, in which no acetone carboxylase was produced (Fig. 1), was unable to grow with acetone or isopropanol. High expression of this enzyme may compensate for a low turnover number for catalysis, allowing a reasonable rate of acetone carboxylation to support growth with a relatively low doubling time (4 to 20 h for *X. autotrophicus*, *R. capsulatus*, and *C. metallidurans* CH34) (19).

Acetone carboxylase purification and characterization. The partial characterization of acetone carboxylase was conducted in *X. autotrophicus* strain Py2, *Rhodococcus rhodochrous* strain B276, *R. capsulatus* strain B10, *Alicyclophilus denitrificans* strain K601, two species of *Paracoccus*, and very recently *Aromatoleum aromaticum*, showing high structural similarities (1, 6, 7, 15, 16, 18–20).

In this study, the Acx of CH34 was purified according to a two-step procedure consisting of anion DEAE-Sepharose chromatography followed by Sephacryl S300 molecular filtration (18, 19). The native molecular mass of the acetone carboxylase complex was determined by gel filtration and was estimated to be 388 \pm 15 kDa, corresponding to an $\alpha_2\beta_2\gamma_2$ configuration (86, 76, and 19 kDa for the α , β , and γ subunits, respectively), as described previously in other bacteria (6, 7, 16, 18, 19). The absorption spectrum of acetone carboxylase between 250 and 350 nm exhibited a maximal peak at 287.2 nm, which is close to the value obtained in *X. autotrophicus* (281 nm) (18).

Enzymatic activity. Depending on the species, the properties of Acx enzymes differ with regard to the substrates and cofactors required to support the carboxylation reaction (1, 6, 16, 18–20).

Received 23 December 2011 Accepted 22 March 2012

Published ahead of print 6 April 2012

Address correspondence to Ruddy Wattiez, ruddy.wattiez@umons.ac.be.

Supplemental material for this article may be found at <http://aem.asm.org/>.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/AEM.07974-11




FIG 1 Acetone carboxylase expression. SDS-PAGE of protein extracts (10 μ g) from CH34 grown in the presence of 9 mM gluconate (1), 25 mM acetone (2), 25 mM isopropanol (3), and 25 mM *n*-propanol (4). (5) Protein extract (10 μ g) from the *acxR* knockout mutant cultivated in the presence of acetone 25 mM. (6) Purified acetone carboxylase (5 μ g). Molecular mass is indicated on the left.

The *C. metallidurans* enzyme showed poor stability and maximum activity in a pH range of 6.5 to 8.0.

Of all the tested high-energy compounds (ATP, ITP, UTP, or GTP), only Mg-ATP supported acetone carboxylation in CH34, resulting in acetoacetate formation (Fig. 2). Similar results were obtained in *X. autotrophicus*, *A. aromaticum*, and *R. capsulatus*, while in *R. rhodochrous*, no activity was observed with ATP (6, 16, 19). As observed in these bacteria, the acetone carboxylase reaction in CH34 showed that ATP is hydrolyzed into AMP and 2 inorganic phosphates.

NH_4^+ ions have also been shown to increase acetone carboxylase activity (19). Yet tests performed in the presence of NH_4Cl (100 mM) showed no significant increase in acetoacetate production by the Acx of CH34. In contrast, as observed in *X. autotroph-*

cus, potassium (40 mM) and CO_2 (KHCO_3) sources stimulated the CH34 acetone carboxylase activity.

The specific activities obtained with the purified enzyme of CH34 were 0.4 to 0.6 U/mg, compared to 0.08 to 0.240 U/mg for the other Acx enzymes (6, 16, 18, 19). Nevertheless, the comparison in terms of activity has to be taken cautiously due to the differences observed with the stability of the purified enzymes.

Among all the tested substrates, only acetone and 2-butanone were identified as substrates of the *C. metallidurans* Acx (Fig. 2). Interestingly, we showed that *C. metallidurans* CH34 was also able to grow in the presence of 2-butanone as the sole carbon source. Studies in *X. autotrophicus* and *A. aromaticum* also revealed that 2-butanone was the only alternative substrate of acetone carboxylase (16, 18). In *R. rhodochrous*, the acetone carboxylase was found to utilize a wider range of substrates, including 2-butanone, which was consumed at a rate identical to that of acetone, and also 2-pentanone, 3-pentanone, and 2-hexanone, which were degraded at rates that were 70, 40, and 42% of the rate of acetone, respectively (6). Nuclear magnetic resonance (NMR) analyses revealed that carboxylation of 2-butanone by the Acx of CH34 produces 3-keto-2-methylbutyrate (Fig. 3). Recently, Schühle and Heider suggested that the Acx of *A. aromaticum* carboxylates only methyl groups adjacent to carbonyl and proposed that butanone was transformed to 3-oxopentanoic acid (16). Nevertheless, no experimental characterization of the carboxylated product was realized from butanone by the Acx of *A. aromaticum* (16).

We propose for *C. metallidurans* that 3-keto-2-methylbutyrate obtained by carboxylation of butanone was then activated to coenzyme A (CoA) thioester and thiolitically cleaved to propionyl-CoA and acetyl-CoA, as observed in the leucine catabolism pathway.

	Acetone carboxylase			Acetophenone carboxylase
	<i>A. aromaticum</i> ¹ <i>X. autotrophicus</i> ² <i>R. capsulatus</i> ³	<i>C. metallidurans</i> CH34	<i>R. rhodochrous</i> ⁴	<i>A. aromaticum</i> ⁵
Subunit composition:	$\alpha_2\beta_2\gamma_2$ (85/75/20 kDa in <i>A. aromaticum</i>) (85/78/20 kDa in <i>X. autotrophicus</i>) (85/78/20 kDa in <i>R. capsulatus</i>)	$\alpha_2\beta_2\gamma_2$ (86/76/19 kDa)	$\alpha_2\beta_2\gamma_2$ (85/74/16 kDa)	($\alpha\beta\beta'\gamma$) ₂ + ϵ_2 (85/75/70/15 + 34 kDa)
Cofactors: ATP	++++	++++	-	++++
GTP	-	-	++++	-
ITP	-	-	++++	-
UTP	-	-	+++	-
CTP	-	ND	++	-
XTP	ND	ND	+	ND
Substrates:	acetone \rightarrow acetoacetate 2-butanone \rightarrow 2-ketovalerate ^a	acetone \rightarrow acetoacetate 2-butanone \rightarrow 3-keto-2-methylbutyrate	acetone \rightarrow acetoacetate 2-butanone ^b 2-pentanone ^b 3-pentanone ^b 2-hexanone ^b	acetophenone \rightarrow benzoylacetate propiophenone ^b 4-acetyl pyridine ^b
No-substrates:	2-pentanone 3-pentanone 2-hexanone Chloroacetone	2-pentanone 3-pentanone 2-hexanone <i>n</i> -propanol isopropanol acetaldehyde dimethylsulfoxide		ring substituted acetophenone derivatives aliphatic ketones

FIG 2 Comparison of the acetone and acetophenone carboxylases from various species. 1, information obtained from reference 16; 2, information obtained from references 18 and 19; 3, information obtained from reference 19; 4, information obtained from reference 6; 5, information obtained from reference 10. A superscript "a" indicates that the products of the enzymatic reaction were not identified in *X. autotrophicus* and *R. capsulatus*; for *A. aromaticum*, the nature of the product was suggested by the authors but not experimentally identified. A superscript "b" indicates that the product of the enzymatic reaction was not identified. ND, not determined. +, supports the Acx reaction at different levels; -, does not support the Acx reaction.




FIG 3 Determination by NMR analyses of acetone carboxylase reaction products with 2-butanone as the substrate. (A and B) Enzymatic reactions realized without substrates. (C, D, E, and F) Enzymatic reactions realized in the presence of 81.5 μg of pure Acx with 8 mM, 12 mM, 16 mM, and 24 mM ATP, respectively. (G) Enzymatic reaction realized with 16 mM ATP and 163 μg of pure Acx. ppm, parts per million.

In conclusion, *C. metallidurans* CH34 is able to degrade acetone and, besides acetone, only 2-butanone using an ATP-dependent pathway including the Acx enzyme. The corresponding *acx* genes are located on the second chromosome or chromid.

ACKNOWLEDGMENTS

We thank C. s'Heeren from the University of Mons for her technical assistance and W. Heylen and R. Van Houdt from SCK•CEN for the *acxR* knockout mutant construction.

This work was supported by the European Space Agency ESA/ESTEC through the PRODEX program in collaboration with the Belgian Science Policy through the MESSAGE-1 project agreement.

REFERENCES

- Birks SJ, Kelly DJ. 1997. Assay and properties of acetone carboxylase, a novel enzyme involved in acetone-dependent growth and CO_2 fixation in *Rhodobacter capsulatus* and other photosynthetic and denitrifying bacteria. *Microbiology* 143:755–766.
- Bonnet-Smits EM, Robertson LA, Van Dijken JP, Senior E, Kuenen JG. 1988. Carbon dioxide fixation as the initial step in the metabolism of acetone by *Thiosphaera pantotropha*. *J. Gen. Microbiol.* 134:2281–2289.
- Boyd JM, Ellsworth H, Ensign SA. 2004. Bacterial acetone carboxylase is a manganese-dependent metalloenzyme. *J. Biol. Chem.* 279:46644–46651.
- Brahmachary P, et al. 2008. The human gastric pathogen *Helicobacter pylori* has a potential acetone carboxylase that enhances its ability to colonize mice. *BMC Microbiol.* 8:14.
- Chowdhury BR, Das SK. 2005. Acetone—a toxic chemical. *Indian Chem. Eng. B* 47:277–279.
- Clark DD, Ensign SA. 1999. Evidence for an inducible nucleotide-dependent acetone carboxylase in *Rhodococcus rhodochrous* B276. *J. Bacteriol.* 181:2752–2758.
- Dullius CH, Chen CY, Schink B. 2011. Nitrate-dependent degradation of acetone by *Alicyclophillus* and *Paracoccus* strains and comparison of acetone carboxylase enzymes. *Appl. Environ. Microbiol.* 77:6821–6825.
- Inui M, et al. 2008. Expression of *Clostridium acetobutylicum* butanol synthetic genes in *Escherichia coli*. *Appl. Microbiol. Biotechnol.* 77:1305–1316.
- Janssen PH, Schink B. 1995. Catabolic and anabolic enzyme activities and energetics of acetone metabolism of the sulfate-reducing bacterium *Desulfococcus biacutus*. *J. Bacteriol.* 177:277–282.
- Jobst B, Schühle K, Linne U, Heider J. 2010. ATP-dependent carboxylation of acetophenone by a novel type of carboxylase. *J. Bacteriol.* 192:1387–1394.
- Kalapos MP. 2003. On the mammalian acetone metabolism: from chemistry to clinical implications. *Biochim. Biophys. Acta* 1621:122–139.
- Kotani T, Yamamoto T, Yurimoto H, Sakai Y, Kato N. 2003. Propane monooxygenase and NAD^+ -dependent secondary alcohol dehydrogenase in propane metabolism by *Gordonia* sp. strain TY-5. *J. Bacteriol.* 185:7120–7128.
- Leys N, et al. 2009. The response of *Cupriavidus metallidurans* CH34 to spaceflight in the international space station. *Antonie Van Leeuwenhoek* 96:227–245.
- Mergeay M, et al. 1985. *Alcaligenes eutrophus* CH34 is a facultative chemolithotroph with plasmid-bound resistance to heavy metals. *J. Bacteriol.* 162:328–334.
- Nocek B, Boyd J, Ensign SA, Peters JW. 2004. Crystallization and preliminary X-ray analysis of an acetone carboxylase from *Xanthobacter autotrophicus* strain Py2. *Acta Crystallogr. D Biol. Crystallogr.* 60:385–387.
- Schühle K, Heider J. 2012. Acetone and butanone metabolism of the denitrifying bacterium “*Aromatoleum aromaticum*” demonstrates novel biochemical properties of an ATP-dependent aliphatic ketone carboxylase. *J. Bacteriol.* 194:131–141.
- Singh HB, et al. 1994. Acetone in the atmosphere: distribution, sources, and sinks. *J. Geophys. Res.* 99:1805–1819.
- Sluis MK, Ensign SA. 1997. Purification and characterization of acetone carboxylase from *Xanthobacter* strain Py2. *Proc. Natl. Acad. Sci. U. S. A.* 94:8456–8461.
- Sluis MK, et al. 2002. Biochemical, molecular, and genetic analyses of the acetone carboxylases from *Xanthobacter autotrophicus* strain Py2 and *Rhodobacter capsulatus* strain B10. *J. Bacteriol.* 184:2969–2977.
- Sluis MK, Small FJ, Allen JR, Ensign SA. 1996. Involvement of an ATP-dependent carboxylase in a CO_2 -dependent pathway of acetone metabolism by *Xanthobacter* strain Py2. *J. Bacteriol.* 178:4020–4026.
- Sydatk C, May O, Altenbuchner J, Mattes R, Siemann M. 1999. Microbial hydantoinases—industrial enzymes from the origin of life? *Appl. Microbiol. Biotechnol.* 51:293–309.
- Taylor DG, Trudgill PW, Cripps RE, Harris PR. 1980. The microbial metabolism of acetone. *J. Gen. Microbiol.* 118:159–170.
- Zheng YN, et al. 2009. Problems with the microbial production of butanol. *J. Ind. Microbiol. Biotechnol.* 36:1127–1138.