

Effect of Metformin on Serum Ferritin Level in Women with Polycystic Ovary Syndrome

S Behradmanesh¹, GhH Ranjbar Omrani², F Ghazanfarpour³, A Baradaran^{4*}

¹Department of Internal Medicine and Endocrinology, Shahr-e-Kord University of Medical Sciences, Shahr-e-Kord, Iran ²Department of Internal Medicine and Endocrinology, Endocrine and Metabolic Research Center, Nemazee Hospital, Shiraz University of Medical Sciences, Shiraz, Iran ³Skin Diseases and Leishmaniasis Research Center, Isfahan University of Medical Sciences, Isfahan, Iran ⁴Department of Pathology, Isfahan University of Medical Sciences, Isfahan, Iran

Abstract

Background: Polycystic ovary syndrome (PCOS) is one of the most common diseases among women associated with various inflammatory reactants such as C-reactive protein (CRP) and ferritin. This study aimed to investigate the effect of metformin on probable reduction of serum ferritin in patients with PCOS.

Methods: This study was conducted on 45 patients with PCOS who had not other systemic diseases and did not take any medications. Weight, waist and hip circumferences (WHR), body mass index (BMI), metabolic indexes, CRP, ferritin and "Homeostasis Model Assessment of Insulin Resistance (HOMA-IR)" were measured before the study. Metformin (500 mg/tid) tablets were prescribed for three months and then same above parameters were re-measured.

Results: Of 45 patients, 19 (42.2%) were overweight and 14 (31.1%) were obese. After drug therapy, there was a significant reduction in waist circumference and serum ferritin. This reduction was significant only in the lean and overweight groups but not in the obese group. There was not significant association between serum ferritin and CRP, HOMA-IR, BMI and WHR. There was not significant correlation between CRP and HOMA-IR and also BMI.

Conclusion: The effect of metformin on reduction of serum ferritin was not significant just in obese group and was not associated with metabolic and anthropometric indexes.

Keywords: Ferritin; Polycystic Ovary Syndrome; Metformin

Introduction

Polycystic ovary syndrome (PCOS) is one of the most common diseases with prevalence of 6-7% among women in their reproductive age.¹⁻⁴ Obesity, hyperandrogenism, ovarian dysfunction and insulin resistance are features of this disease.⁵ In addition, other diseases such as type 2 diabetes mellitus (T2DM), metabolic syndrome (MetS) and cardiovascular diseases (CVD) are related to this syndrome.⁶⁻⁹ It has been recently found that acute phase reactants of in-

flammation including C-reactive protein (CRP) are higher than normal in T2DM, hyperinsulinemia¹⁰ and PCOS.¹¹ Serum level of CRP has a higher value in obese PCOS subjects than those non-obese with PCOS. A treatment with metformin for six months lowers CRP serum level in all subjects as well as obese and non-obese.¹² Another acute phase reactants of inflammation is ferritin which is higher than normal in patients with metabolic syndrome, T2DM patients and PCOS.¹³⁻¹⁵ Serum ferritin levels in overweight and obese women with PCOS are significantly higher compared to overweight and obese women without this disorder.¹⁵⁻¹⁸

Metformin causes improvement of insulin resistance in patients with PCOS.^{19,20} Regarding the prevalence of PCOS among women at reproductive age⁴ and accompany of acute phase reactants of in-

*Correspondence: Azar Baradaran, MD, Associate Professor of Pathology, Al-Zahra Medical Center, Isfahan University of Medical Sciences, Isfahan, Iran. Tel: +98-913-3278027, e-mail: azarbaradaran@yahoo.com

Received: October 15, 2010;

Accepted: February 12, 2011

flammation with the risk of cardiovascular diseases,²¹ we sought to investigate the effect of metformin on lowering serum ferritin level in PCOS, obese and non-obese patients.

Materials and Methods

This is a before-after quasi experimental study on 45 PCOS women. The PCOS subjects were selected based on Homburg criteria and others.^{12,22} The study was carried out in clinics related to Shiraz University of Medical Sciences, Shiraz Iran, from January 2006 and lasted for six months.

Exclusion criteria were pregnancy, diabetes mellitus, psychotic disorders, hypertension, cardiovascular diseases, smoking, alcohol drinking and oral contraceptive pill taking, anti-inflammatory drugs and any other medication for hyperlipidemia, hypertension, and glucose intolerance in the past six months were also the exclusion criteria. Patients who were also on special diet or on standard sports to lose weight, all were excluded too. The secondary PCOS was also ruled out.

All subjects signed the consent forms and the Ethics Committee of the university approved the study. Basic measurements on anthropometric indexes included waist and hip (cm) (waist was calculated by the greatest circumference between the last rib and iliac crest and the hip was calculated by the greatest circumference from both greater femoral trochanters),²³ height (m), weight (kg), BMI (kg/m^2) and WHR.

BMI ≤ 25 was considered as normal, $25 < \text{BMI} < 30$ overweight and $\text{BMI} \geq 30$ was considered as obese. $\text{WHR} > 0.8$ was considered as a sign of central obesity.²⁴ In addition, fasting insulin was measured with immunotech kit (made in Check by IRMA method), while fasting plasma glucose (FPG), triglyceride (Tg), total cholesterol (chol), HDL and LDL [through enzymatic calorimetric method], serum ferritin and high sensitive C reactive protein (hsCRP) [by IRMA method] levels that were measured using standard kits.

Homeostasis Model Assessment of Insulin Resistance (HOMA-IR) was calculated by fasting serum insulin (pmol/L) multiplied by FBS (mmol/L) divided to 22.5.²⁵ Blood pressure was measured by standard method too. In the first week, apo-metformin (Apotex Inc., Toronto, Canada) was prescribed (500 mg/bid) to lower gastrointestinal complications and then reached 500 mg/tid and continued for 12 weeks. The subjects were asked to have a normal diet (30-35 kcal

/kg/day) together with ordinary physical activity. At the end of three months (12 weeks), anthropometric indexes were re-measured and recorded. The above-mentioned lab tests were repeated. Insulin resistance was calculated and systolic diastolic BP was re-measured by the same method too.

Data were expressed as mean \pm SD and frequency distribution. Normal distribution of the data was checked with Kolmogorov Smirnov test. Paired t-test and ANOVA by Tukey posthoc were used to investigate the changes in variables before and after treatment as well as in subgroups. Correlation among some variables was done by Pearson's bivariate correlation coefficient test. Data were analyzed by SPSS software (version 13.0, Chicago, IL, USA). A P -value < 0.05 was considered significant.

Results

The age of the subjects was 22.5 ± 4.5 years. Of 45 subjects, 12 (26.7%) had normal BMI, 19 (42.2%) were overweight and 14 (31.1%) were obese. Forty four (97.8%) had central obesity based on WHR. Comparison of the changes in anthropometric indexes before and after three months with metformin showed that hip circumference had a significant decrease while other parameters showed no significant difference (Table 1).

Table 2 shows the comparison of metabolic parameters. Serum ferritin level had a significant decrease ($p < 0.001$, $t = 4.42$). Comparison of basic and after three months treatment of hsCRP, ferritin and HOMA-IR with metformin showed that basic serum ferritin was not different between groups (normal, overweight and obese) (Not Significant). Comparison of serum ferritin, hsCRP and HOMA-IR before and after three months treatment with metformin among mentioned above sub-groups of BMI showed a significant difference of HOMA-IR ($p = 0.04$), serum ferritin ($p = 0.02$) in normal BMI group, while there was not significant difference of hsCRP ($p = 0.4$). In overweight group, there was only significant difference of serum ferritin ($p = 0.001$). There was not significant difference of parameters before and after treatment in obese sub-group.

Data analysis by ANOVA showed no significant difference of CRP ($p = 0.8$), ferritin ($p = 0.1$) and HOMA-IR ($p = 0.07$) among three groups of obese, non-obese and overweight subjects.

Table 1: Comparison of the changes in anthropometric indexes before and after three months treatment (n=45).

	Before	After 3-months	P value
Weight (kg)	70.9±12.2	70.2±11.9	0.08
BMI (kg/m ²)	27.7±4.1	27.4±4.2	0.1
Waist (cm)	96.9±11	95.8±11.4	0.2
Hip (cm)	105±7.3	104±7.3	0.03
WHR	0.92±0.06	0.91±0.07	0.7
Systolic BP (mmHg)	112.3±8.5	113.3±8.1	0.4
Diastolic BP (mmHg)	72.1±8.2	74.4±7.5	0.09

Table 2: comparison of the changes in metabolic parameters before and three months after treatment (n=45)^a.

	Before	After 3-months	P value
Insulin (μU/mL)	11.8±9.6	11.6±8.3	0.9
FPG (mg/dL)	77.3±8.4	79.6±15.1	0.3
TG (mg/dL)	156.3±76.5	157.5±78.3	0.4
Total cholesterol (mg/dL)	191.8±38.5	193.8±40	0.5
LDL	90.3±31.4	90.3±28.3	0.3
HDL	69.8±19.5	71.6±15	0.5
Ferritin (μg/L)	68.5±51.3	35.1±33.9	0.00
hs CRP (mg/L)	3.2±3.7	2.3±2.2	0.05

^aFPG, TG and hsCRP denotes fasting plasma glucose, triglyceride and highly sensitive CRP respectively.

Pearson's correlation test showed an association between CRP and insulin resistance indexes before intervention ($p=0.02$, $r=0.34$) (Table 3). There was not association between serum ferritin with CRP and insulin resistance either before or after three months of treat-

ment ($p>0.05$). Associations between CRP, HOMA-IR with BMI and WHR before and after the treatment has been presented in Table 4. There was not association between ferritin and these anthropometric indexes either before or three months after treatment ($p>0.05$).

Table 3: Mean±SD of CRP, ferritin and insulin resistance indexes in three groups of obese, overweight and lean before and after treatment.

	Obese (n=14)		Overweight (n=19)		Lean (n=12)	
	Before	After 3 months	Before	After 3 months	Before	After 3 months
hs CRP	5.07±5.7	2.9±1.9	2.8±2.4	2.5±2.8	1.4±0.73	1.3±0.38
Ferritin	77.6±43	59.3±48.7	52.3±24.8	23.7±13.5	83.5±80.9	24.8±20.7
HOMA-IR	2.9±1.3	3.5±2.4	1.8±0.9	2.5±1.2	1.3±0.87	0.8±0.25

Table 4: Association of CRP, HOMA-IR with WHR and BMI before and after treatment.

	CRP		HOMA-IR	
	Before	After 3 months	Before	After 3 months
BMI				
r	0.33	0.39	0.43	0.54
p	0.02	0.008	0.003	0.00
WHR				
r	0.23	0.2	0.14	0.41
p	0.12	0.09	0.35	0.005

Discussion

In this study, we found that three months intake of metformin in PCOS patients, irrespective to BMI groups, caused a reduction in serum ferritin. In a comparison between two groups of obese PCOS, reduction of serum ferritin during treatment with metformin and diene was investigated by Luque-Ramírez *et al.*¹⁸ They showed that taking metformin for three months and also six months resulted in a reduction in serum ferritin among these patients. After data analysis based on BMI, they found that intake of metformin for three months reduced ferritin just in overweight and lean groups but this reduction was not significant in the obese group. They concluded that the reason for this difference in obese group can possibly be due to the comparison of metformin with diene.¹⁸ To investigate the reduction of CRP in PCOS during metformin treatment, Morin-Papunen *et al.* showed that CRP significantly decreased after three months and 6 months post treatment respectively compared to the beginning of the study. These finding was not in accordance with our findings. In the present study, there was not significant reduction in any of the subgroups possibly due to the sample size compared to our findings.¹²

Previous studies showed that obese PCOS had higher insulin resistance compared to normal weight PCOS.^{26,27} Insulin resistance and hyperinsulinemia resulted into an increase in iron storage through bowel absorption, as indicated in metabolic syndrome and T2D.^{17,28,29} It was thought that metformin with respect to its effect on central obesity and facilitating insulin function³⁰ reduces insulin sensitivity index together with a decrease in serum ferritin that does not affect CRP in comparison to the obese PCOS taking diene.¹⁷ In the present study, there was not any association between BMI, WHR, HOMA-IR and serum ferritin not only before but also after treatment possibly due to lack of control group or sample size difference in these two studies (19 vs 45). While in the present study, there was not any association between CRP and ferritin either before or after the treatment, former studies showed that the increase in ferritin is independent to the increase in CRP.^{15,18} Previous studies showed an increase in CRP serum level in obese people³¹ and an association between abdominal obesity, insulin resistance and CRP.³² It is was found that anthropometric parameters especially waist and WHR are modified with metformin in PCOS resulting in a decrease in CRP.¹² Although, there was not a signifi-

cant decrease in CRP with metformin among all subjects and in subgroups in the present study, there was a direct significant association between CRP with BMI (before and after treatment). There was not a significant decrease in HOMA-IR in all subjects while a significant decrease was noticed among subgroups.

Among other anthropometric indexes and metabolic parameters of our study, hip circumference showed a significant reduction, although among others a decrease was seen but was not significant. In an another study in obese PCOS patients conducted by Morin-Papunen *et al.*, intake of metformin for three straight months showed no significant decrease for BMI, WHR and insulin sensitivity index whereas a significant decrease was observed for WHR after six months. In addition, FPG and fasting insulin showed a significant reduction after three months.³³ The effect of metformin on non-obese PCOS was also investigated in an another study by Morin-Papunen *et al.* They found a significant reduction in BMI, WHR, FPG and fasting insulin after three months of therapy with metformin, while the reduction in insulin sensitivity index was significant. The different results of two above studies compared to our findings was the existence of a control group and low sample size.³³ There was not a significant difference between our findings and those two above regarding the HOMA-IR and the reduction of anthropometric indexes especially in non-obese subjects. To investigate the basic CRP, ferritin and HOMA-IR in the related subgroups, there was a significant difference in CRP and HOMA-IR, in obese patients compared to normal weights ($p=0.03$), something consistent with former studies.¹² The level of ferritin was not different in three groups. In a comparative research serum ferritin in PCOS group (compared to healthy group according to identical weight) serum ferritin was found not to be different in those with normal weight, while, mean serum ferritin level in overweight and obese PCOS subjects was significantly higher compared to the control group with identical weight, apparently not consistent with the present study possibly due to lack of a control group.¹² In this study, we can concluded that intake of metformin for three months causes a reduction in serum ferritin in all PCOS subjects. This reduction was independent of subjects' weight group, CRP and HOAM-IR. Although hyperinsulinemia was investigated as an iron loading agent in PCOS obese subjects in former studies,¹⁸ regarding the results of the present study, the reason for iron overload still

needs further consideration by other investigators.

Conflict of interest: None declared.

References

- Azziz R. Controversy in clinical endocrinology: diagnosis of polycystic ovary syndrome: the Rotterdam criteria are premature. *J Clin Endocrinol Metab* 2006;**91**:781-5. [1641821] [doi:10.1210/jc.2005-2153]
- Diamanti-Kandarakis E, Kouli CR, Bergiele AT, Filandra FA, Tsianateli TC, Spina GG, Zapanti ED, Bartzis MI. A survey of the polycystic ovary syndrome in the Greek island of Lesbos: hormonal and metabolic profile. *J Clin Endocrinol Metab* 1999;**84**:4006-11. [10566641] [doi:10.1210/jc.84.11.4006]
- Asunción M, Calvo RM, San Millán JL, Sancho J, Avila S, Escobar-Morreale HF. A prospective study of the prevalence of the polycystic ovary syndrome in unselected Caucasian women from Spain. *J Clin Endocrinol Metab* 2000;**85**:2434-8. [10902790] [doi:10.1210/jc.85.7.2434]
- Azziz R, Woods KS, Reyna R, Key TJ, Knochenhauer ES, Yildiz BO. The prevalence and features of the polycystic ovary syndrome in an unselected population. *J Clin Endocrinol Metab* 2004;**89**:2745-9. [15181 052] [doi:10.1210/jc.2003-032046]
- Azziz R, Carmina E, Dewailly D, Diamanti-Kandarakis E, Escobar-Morreale HF, Futterweit W, Janssen OE, Legro RS, Norman RJ, Taylor AE, Witchel SF; Androgen Excess Society. Positions statement: criteria for defining polycystic ovary syndrome as a predominantly hyperandrogenic syndrome: an Androgen Excess Society guideline. *J Clin Endocrinol Metab* 2006;**91**:4237-45. [16940456] [doi:10.1210/jc.2006-0178]
- Dunaif A. Insulin resistance and the polycystic ovary syndrome: mechanism and implications for pathogenesis. *Endocr Rev* 1997;**18**:774-800. [9408743]
- Barber TM, McCarthy MI, Wass JA, Franks S. Obesity and polycystic ovary syndrome. *Clin Endocrinol (Oxf)* 2006;**65**:137-45. [16886951] [doi:10.1111/j.1365-2265.2006.02587.x]
- Wild S, Pierpoint T, McKeigue P, Jacobs H. Cardiovascular disease in women with polycystic ovary syndrome at long-term follow-up: a retrospective cohort study. *Clin Endocrinol (Oxf)* 2000;**52**:595-600. [10792339] [doi:10.1046/j.1365-2265.2000.01000.x]
- Mehrabian F, Khani B, Kelishadi R, Kermani N. The prevalence of metabolic syndrome and insulin resistance according to the phenotypic subgroups of polycystic ovary syndrome in a representative sample of Iranian females. *J Res Med Sci* 2011;**16**:1-7.
- Festa A, D'Agostino R Jr., Tracy RP, Haffner SM; Insulin Resistance Atherosclerosis Study. Elevated levels of acute-phase proteins and plasminogen activator inhibitor-1 predict the development of type 2 diabetes: the insulin resistance atherosclerosis study. *Diabetes* 2002;**51**:1131-7. [11916936] [doi:10.2337/diabetes.51.4.1131]
- Kelly CC, Lyall H, Petrie JR, Gould GW, Connell JM, Sattar N. Low grade chronic inflammation in women with polycystic ovarian syndrome. *J Clin Endocrinol Metab* 2001;**86**:2453-5. [11397838] [doi:10.1210/jc.86.6.2453]
- Morin-Papunen L, Rautio K, Ruokonen A, Hedberg P, Puukka M, Tapanainen JS. Metformin reduces serum C-reactive protein levels in women with polycystic ovary syndrome. *J Clin Endocrinol Metab* 2003;**88**:4649-54. [14557435] [doi:10.1210/jc.2002-021688]
- Bozzini C, Girelli D, Olivieri O, Martinelli N, Bassi A, De Matteis G, Tenuti I, Lotto V, Friso S, Pizzolo F, Corrocher R. Prevalence of body iron excess in the metabolic syndrome. *Diabetes Care* 2005;**28**:2061-3. [16043762] [doi:10.2337/diacare.28.8.2061]
- Jiang R, Manson JE, Meigs JB, Ma J, Rifai N, Hu FB. Body iron stores in relation to risk of type 2 diabetes in apparently healthy women. *JAMA* 2004;**291**:711-7. [14871914] [doi:10.1001/jama.291.6.711]
- Escobar-Morreale HF, Luque-Ramírez M, Alvarez-Blasco F, Botella-Carretero JI, Sancho J, San Millán JL. Body iron stores are increased in overweight and obese women with polycystic ovary syndrome. *Diabetes Care* 2005;**28**:2042-4. [16043756] [doi:10.2337/diacare.28.8.2042]
- Botella-Carretero JI, Luque-Ramírez M, Alvarez-Blasco F, San Millán JL, Escobar-Morreale HF. Mutations in the hereditary hemochromatosis gene are not associated with the increased body iron stores observed in overweight and obese women with polycystic ovary syndrome. *Diabetes Care* 2006;**29**:2556. [17065702] [doi:10.2337/dc06-1655]
- McCarty MF. Hyperinsulinemia may boost both hematocrit and iron absorption by up-regulating activity of hypoxia-inducible factor-1alpha. *Med Hypotheses* 2003;**61**:567-73. [14592787]
- Luque-Ramírez M, Alvarez-Blasco F, Botella-Carretero JI, Sanchón R, San Millán JL, Escobar-Morreale HF. Increased body iron stores of obese women with polycystic ovary syndrome are a consequence of insulin resistance and hyperinsulinism and are not a result of reduced menstrual losses. *Diabetes Care* 2007;**30**:2309-13. [17536071] [doi:10.1016/S0306-9877(03)00231-7]
- Nestler JE, Jakubowicz DJ. Lean women with polycystic ovary syndrome respond to insulin reduction with decreases in ovarian P450c17 alpha activity and serum androgens. *J Clin Endocrinol Metab* 1997;**82**:4075-9. [9398716] [doi:10.1210/jc.82.12.4075]
- Velazquez EM, Mendoza S, Hamer T, Sosa F, Glueck CJ. Metformin therapy in polycystic ovary syndrome reduces hyperinsulinemia, insulin resistance, hyperandrogenemia, and systolic blood pressure, while facilitating normal menses and pregnancy. *Metabolism* 1994;**43**:647-54. [8177055] [doi:10.1016/0026-0495(94)90209-7]
- Ridker PM, Hennekens CH, Buring JE, Rifai N. C-reactive protein and other markers of inflammation in the prediction of cardiovascular disease in women. *N Engl J Med* 2000;**342**:836-43. [10733371] [doi:10.1056/NEJM200003233421202]
- Homburg R. Polycystic ovary syndrome-from gynaecological curiosity to multisystem endocrinopathy. *Hum Reprod* 1996;**11**:29-39. [8671153]
- Bues JB, Polonsky KS and Burant CF. Type 2 diabetes mellitus. In: Larsen, Kronenberg, Melmed, Polonsky (editors): *Williams Textbook of Endocrinology*. Saunders, Philadelphia, 2003; p.1441.
- Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults--The Evidence Report. National Institutes of Health. *Obes Res* 1998;**6**:51S-209S. [9813653]
- Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner

- RC. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 1985;**28**:412-9. [3899825] [doi:10.1007/BF00280883]
- 26 Dunaif A, Graf M, Mandeli J, Laumas V, Dobrjansky A. Characterization of groups of hyperandrogenic women with acanthosis nigricans, impaired glucose tolerance, and/or hyperinsulinemia. *J Clin Endocrinol Metab* 1987;**65**:499-507. [3305551] [doi:10.1210/jcem-65-3-499]
- 27 Dunaif A, Segal KR, Futterweit W, Dobrjansky A. Profound peripheral insulin resistance, independent of obesity, in polycystic ovary syndrome. *Diabetes* 1989;**38**:1165-74. [2670645] [doi:10.2337/diabetes.38.9.1165]
- 28 Jehn M, Clark JM, Guallar E. Serum ferritin and risk of the metabolic syndrome in U.S. adults. *Diabetes Care* 2004;**27**:2422-8. [1545191] [doi:10.2337/diacare.27.10.2422]
- 29 Ford ES, Cogswell ME. Diabetes and serum ferritin concentration among U.S. adults. *Diabetes Care* 1999;**22**:1978-83. [10587829] [doi:10.2337/diacare.22.12.1978]
- 30 Pasquali R, Gambineri A, Biscotti D, Vicennati V, Gagliardi L, Colitta D, Fiorini S, Cognigni GE, Filicori M, Morselli-Labate AM. Effect of long-term treatment with metformin added to hypocaloric diet on body composition, fat distribution, and androgen and insulin levels in abdominally obese women with and without the polycystic ovary syndrome. *J Clin Endocrinol Metab* 2000;**85**:2767-74. [10946879] [doi:10.1210/jc.85.8.2767]
- 31 Festa A, D'Agostino R Jr, Williams K, Karter AJ, Mayer-Davis EJ, Tracy RP, Haffner SM. The relation of body fat mass and distribution to markers of chronic inflammation. *Int J Obes Relat Metab Disord* 2001;**25**:1407-15. [11673759] [doi:10.1038/sj.ijo.0801792]
- 32 Pannacciulli N, Cantatore FP, Minenna A, Bellacicco M, Giorgino R, De Pergola G. C-reactive protein is independently associated with total body fat, central fat, and insulin resistance in adult women. *Int J Obes Relat Metab Disord* 2001;**25**:1416-20. [11673760] [doi:10.1038/sj.ijo.0801719]
- 33 Morin-Papunen LC, Vauhkonen I, Koivunen RM, Ruokonen A, Martikainen HK, Tapanainen JS. Endocrine and metabolic effects of metformin versus ethinyl estradiol-cyproterone acetate in obese women with polycystic ovary syndrome: a randomized study. *J Clin Endocrinol Metab* 2000;**85**:3161-8. [10999803] [doi:10.1210/jc.85.9.3161]
- 34 Morin-Papunen L, Vauhkonen I, Koivunen R, Ruokonen A, Martikainen H, Tapanainen JS. Metformin versus ethinyl estradiol-cyproterone acetate in the treatment of nonobese women with polycystic ovary syndrome: a randomized study. *J Clin Endocrinol Metab* 2003;**88**:148-56. [12519844] [doi:10.1210/jc.2002-020997]