

Complete Genome Sequence of the European Sheatfish Virus

Carla Mavian,^a Alberto López-Bueno,^a María Pilar Fernández Somalo,^b Antonio Alcamí,^a and Ali Alejo^c

Centro de Biología Molecular Severo Ochoa (Consejo Superior de Investigaciones Científicas-Universidad Autónoma de Madrid), Madrid, Spain^a; Ministerio de Agricultura, Alimentación y Medio Ambiente, Algete, Spain^b; and Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria, Valdeolmos, Spain^c

Viral diseases are an increasing threat to the thriving aquaculture industry worldwide. An emerging group of fish pathogens is formed by several ranaviruses, which have been isolated at different locations from freshwater and seawater fish species since 1985. We report the complete genome sequence of European sheatfish ranavirus (ESV), the first ranavirus isolated in Europe, which causes high mortality rates in infected sheatfish (*Silurus glanis*) and in other species. Analysis of the genome sequence shows that ESV belongs to the amphibian-like ranaviruses and is closely related to the epizootic hematopoietic necrosis virus (EHNV), a disease agent geographically confined to the Australian continent and notifiable to the World Organization for Animal Health.

Ranaviruses are emerging pathogens causing serious disease in wild and farmed cold-blooded animals (6). They are icosahedral viruses of the *Iridoviridae* family with circular double-stranded DNA (dsDNA) genomes ranging from 105 to 140 kbp and a complex nucleocytoplasmic replication cycle. One of their most intriguing biological properties is their wide host tropism, which includes amphibians, fish, and reptiles.

Complete ranavirus genomes from different hosts and locations show the existence of two groups, one corresponding to the fish viruses grouper iridovirus (GIV) and Singapore grouper iridovirus (SGIV), isolated in Asia, and the second to the amphibian-like ranaviruses (ALRVs), which include viruses isolated from amphibians or reptiles in North America, Asia, and Europe as well as one species, the epizootic hematopoietic necrosis virus (EHNV), isolated from diseased fish in Australia.

The first fish ranavirus to be isolated in Europe was European sheatfish ranavirus (ESV), which was obtained from moribund sheatfish (*Silurus glanis*) fry in Germany in 1989 (2, 8). A second ranavirus isolated from catfish (*Ameiurus melas*) in France (12) and Italy (4) was found to be very closely related to the former (3, 9, 10) and may represent an isolate of the same virus. Experimental ESV infection in sheatfish results in 100% mortality by 8 days after challenge (1) and causes high mortality in pike (*Esox lucius*) (11), too. Other species, such as black bullhead (*Ameiurus melas*) (7) and rainbow trout (3), are apparently not affected, although viral replication has been detected.

To obtain the complete genome sequence of ESV, the reference virus stock from the Spanish Central Veterinary Laboratory was plaque purified on ZF4 cells to ensure clonality and DNA from viral particles was extracted. Sequencing was performed using Genome Sequencer FLX Titanium equipment (Roche 454 Life Sciences) at the Parque Científico de Madrid. A total of 71,527 viral reads with an average length of 355 bp was assembled with Newbler 2.5.3 in 7 contigs flanked by repeated regions that were joined into a single contig by PCR and Sanger sequencing.

The final genome sequence of ESV is 127,732 bp long, which is slightly longer than that of the EHNV genome, and it has a GC content of 54.23%. Annotation of the ESV genome was performed with genome annotation transfer utility (GATU) software (13) using the genome of EHNV as a template. A total of 136 putative open reading frames, including all of the iridovirus- and ranavirus-specific genes as well as 12 out of 13 ALRV-specific genes, were identified. Global multiple alignments with other ranaviruses us-

ing LAGAN software (5) show that ESV has the highest degree of conservation with EHNV (88%) and *Ambystoma tigrinum* virus (ATV) (79%). Dot plot analyses show large colinearity stretches with both the EHNV and ATV genomes. Compared to ATV, ESV shows 9 major sequence insertions, while in comparison with EHNV, ESV shows three insertions, one deletion, and one inversion. Altogether, these results place ESV as a novel ALRV belonging to the ATV/EHNV group.

Nucleotide sequence accession number. The complete genome sequence of European sheatfish virus was deposited in GenBank under accession number [JQ724856](https://www.ncbi.nlm.nih.gov/nuccore/JQ724856).

ACKNOWLEDGMENTS

The ESV reference virus strain was provided by the European Union Reference Laboratory for Fish Diseases, which, in turn, received it from W. Ahne, Munich, Germany.

This work was supported by grant AGL 2009-08711 from the Spanish Ministerio de Ciencia e Innovación. Alberto López-Bueno and Carla Mavian are recipients of Ramón y Cajal and Formación de Personal Investigador fellowships, respectively, from the same institution.

REFERENCES

1. Ahne W, Ogawa M, Schlotfeldt HJ. 1990. Fish viruses: transmission and pathogenicity of an icosahedral cytoplasmic deoxyribovirus isolated from sheatfish (*Silurus glanis*). *Zentralbl. Veterinarmed.* B 37:187–190.
2. Ahne W, Schlotfeldt HJ, Thomsen I. 1989. Fish viruses: isolation of an icosahedral cytoplasmic deoxyribovirus from sheatfish (*Silurus glanis*). *Zentralbl. Veterinarmed.* B 36:333–336.
3. Ariel E, Holopainen R, Olesen NJ, Tapiovaara H. 2010. Comparative study of ranavirus isolates from cod (*Gadus morhua*) and turbot (*Psetta maxima*) with reference to other ranaviruses. *Arch. Virol.* 155:1261–1271.
4. Bovo G, et al. 1993. Isolamento di un agente virale irido-like da pesce gatto (*Ictalurus melas*) dall'allevamento. *Boll. Soc. Ital. Patol. Ittica* 11:3–10. (In Italian.)
5. Brudno M, et al. 2003. LAGAN and Multi-LAGAN: efficient tools for large-scale multiple alignment of genomic DNA. *Genome Res.* 13:721–731.
6. Chinchar VG. 2002. Ranaviruses (family Iridoviridae): emerging cold-blooded killers. *Arch. Virol.* 147:447–470.

Received 10 March 2012 Accepted 13 March 2012

Address correspondence to Ali Alejo, alejo@inia.es.

C.M. and A.L.-B. contributed equally to this article.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JVI.00618-12

7. Gobbo F, Cappellozza E, Pastore MR, Bovo G. 2010. Susceptibility of black bullhead *Ameiurus melas* to a panel of ranavirus isolates. *Dis. Aquat. Organ.* 90:167–174.
8. Hedrick R, McDowell TS, Ahne W, Torhy C, de Kinkelin P. 1992. Properties of three iridovirus-like agents associated with systemic infections of fish. *Dis. Aquat. Organ.* 13:203–209.
9. Holopainen R, Ohlemeyer S, Schutze H, Bergmann SM, Tapiovaara H. 2009. Ranavirus phylogeny and differentiation based on major capsid protein, DNA polymerase and neurofilament triplet H1-like protein genes. *Dis. Aquat. Organ.* 85:81–91.
10. Hyatt AD, et al. 2000. Comparative studies of piscine and amphibian iridoviruses. *Arch. Virol.* 145:301–331.
11. Jensen BB, Ersboll AK, Ariel E. 2009. Susceptibility of pike *Esox lucius* to a panel of Ranavirus isolates. *Dis. Aquat. Organ.* 83:169–179.
12. Pozet F, Morand M, Moussa A, Torhy C, de Kinkelin P. 1992. Isolation and preliminary characterisation of a pathogenic icosahedral deoxyribovirus from the catfish *Ictalurus melas*. *Dis. Aquat. Organ.* 14:35–42.
13. Tcherepanov V, Ehlers A, Upton C. 2006. Genome Annotation Transfer Utility (GATU): rapid annotation of viral genomes using a closely related reference genome. *BMC Genomics* 7:150.