



Published in final edited form as:

Tetrahedron. 2012 June 17; 68(24): 4797–4804. doi:10.1016/j.tet.2012.03.121.

A new protecting group and linker for uridine ureido nitrogen

Yong Wang and Michio Kurosu

Pharmaceutical Sciences, College of Pharmacy, University of Tennessee Health Science Center, 881 Madison, Memphis, TN 38163, USA

Abstract


(2,6-Dichloro-4-methoxyphenyl) (2,4,6-trichlorophenyl) methoxymethyl chloride [**1**, monomethoxydiphenylmethoxymethyl chloride (MDPM-Cl)] shows a significant relative stability and **1** reacts with uridine ureido nitrogen in the presence of DBU to form the corresponding protected uridine **8** in 95% yield. The MDPM-protected uridines are stable to a wide variety of conditions utilized for the synthesis of analogs of capuramycin and muraymycins. Significantly, the MDPM protecting group can conveniently be deprotected by using 30% TFA in CH₂Cl₂. In addition, polymer-bound MDPM-Cl **23** is useful for immobilization of uridine derivatives.

Keywords


Ureido nitrogen; Uridine; Monomethoxydiphenylmethoxymethylgroup; Polymer-bound; linker

1. Introduction

Uridine is an essential biological compound for multiple biosynthetic processes and found in all cells.¹ To date, a large number of uridine-containing natural products that show significant biological activities have been isolated.² Uridine-containing antibiotics such as liposidomycin, caprazamycin, muraymycins, and capuramycin have been of increasing interest to the development of new antibacterial agents for MDR-bacterial infections.³



In our ongoing program of development of drug leads for MDR *Mycobacterium tuberculosis*, we have synthesized analogs of muraymycins A₁ and capuramycin.⁴ Protection of the uridine ureido nitrogen was indispensable to achieve the synthesis of wide range of uridine-containing molecules. Benzyloxymethyl (BOM) group has been utilized as a convenient protecting group for the uridine ureido nitrogen (P₁).⁵ However, we and other groups observed that BOM deprotection of uridine derivatives under hydrogenation conditions often resulted in poor yield with over-reduction product(s).⁶ In general, hydrogenolytic deprotection is the only method to remove the BOM group of the uridine ureido nitrogen.



We have sought an alternative methoxymethyl-type protecting group for the uridine ureido nitrogen that 1) is stable to reaction conditions for the syntheses of a wide range of muraymycin and capuramycin analogs, but 2) can be cleaved with mild and volatile acids. In this paper we report our studies of (2,6-dichloro-4-methoxyphenyl) (2,4,6-trichlorophenyl) methoxymethyl chloride [**1**, monomethoxydiphenyl methoxymethyl chloride (MDPM-Cl)] for the protection of the uridine ureido nitrogen and application of **1** to the polymer-bound MDPM-Cl for synthesis of analogs on polymer-support.

2. Results and discussion

We have previously reported utilities of *rac*-**5a** and optically pure **5a** for protections of alcohols, amines, and carboxylic acids, and as a chiral derivatizing agent.⁷ (2,6-Dichloro-4-methoxyphenyl)(2,4-dichlorophenyl)methanol (**5a**) could efficiently be synthesized via a

Friedel-Crafts reaction followed by LiBH₄ reduction. Similarly, a large quantity of (2,6-dichloro-4-methoxyphenyl) (2,4,6-trichlorophenyl) methanol (**5b**) could be synthesized efficiently. Stability tests of **5a** and **5b** against a wide variety of Lewis and Brønsted acids revealed that **5b** exhibited longer half-life in the acidic conditions summarized in Table 1; **5a** was converted to the corresponding TFA ester in 30 min. when exposed to 30% TFA in CH₂Cl₂ at room temperature, on the other hand, **5b** required more than 1h to form its TFA ester under the same reaction conditions. In addition, we have recognized that *rac*-**5a** forms diastereomers in NMR spectra when reacted with chiral substrates. However, **5b** has not formed noticeable diastereomers in NMR spectra. In addition, generated diastereomers have not been separated via HPLC.⁸ Because of the reasons stated above, we decided to utilize **5b** for further investigation. As illustrated in Scheme 1 (2,6-dichloro-4-methoxyphenyl) (2,4,6-trichlorophenyl) methanol (**5b**) could be converted to the corresponding MDPM-Cl **1**. The alcohol **5b** was first transformed to the (alkoxymethyl)methyl sulfide **6** in 98% yield, which was then subjected to a Cl-displacement reaction with SO₂Cl₂ to yield **1** in greater than 95% yield.⁹ MDPM-Cl **1** was stable at room temperature and can be stored for several months without loss of purity. The uridine ureido nitrogen was efficiently protected with **1** in the presence of DBU in DMF to afford **8** in 95% yield (Table 1). The MDPM group of **8** exhibited excellent stability against a variety of Brønsted and Lewis acids such as 20% TFA, 10% TMSOTf, 10% HCl, 30% HF, 80% AcOH, 30% TsOH, La(OTf)₃, BF₃•OEt₂, and TiCl₄ at room temperature. The MDPM-protected uridine **8** also showed stability under basic conditions; **8** was intact under NH₄OH (40% in aq MeOH), LiOH (10% in aq THF-MeOH), and DBU (10% in toluene) at room temperature for over 24 h. Selected examples are summarized in Table 1. Moreover, the MDPM protecting group of **8** was stable to the reduction conditions such as Al(Hg), ⁿBu₃SnH/AIBN, and Raney Ni. The MDPM-protected uridine **8** was stable to NBS, and was photolytically stable (200~350 nm for over 6 h). The MDPM protecting group of **8** was not cleaved by hydrogenation conditions using Pd-C or Pd black even at 100 psi. The MDPM group could not be cleaved by using the standard conditions for the deprotection of PMB (*p*-methoxybenzyl) ether groups, however, could be deprotected with 30% TFA in CH₂Cl₂ to afford uridine (**7**). The (2,4-dichloro-4-methoxyphenyl)(2,4,6-trichlorophenyl)methyl cation generated by the treatment of **8** with 30% TFA is a sterically hindered species and stabilized by the electron-withdrawing Cl atoms, being efficiently reacted with the trifluoroacetate ion to afford (2,4-trichloro-4-methoxyphenyl)(2,4,6-trichlorophenyl)methyl 2,2,2-trifluoroacetate (**9**) in quantitative yield. The treatment of **9** with NH₃/MeOH of **7** gave rise to the parent alcohol **5b** in quantitative yield (Scheme 2). Thus, the MDPM-Cl **1** can be regenerated through the chemical steps illustrated in Scheme 1.

In order to demonstrate robustness of MDPM group as a protecting group for the uridine ureido nitrogen, the MDPM-protected uridine **8** was transformed to a wide range of uridine derivatives that can be utilized for the syntheses of analogs of muraymycins and capuramycin.^{4d} Selected examples are summarized in Table 2. Ketal formations of **10** or **8** under acidic conditions provided the corresponding 2,3-protected derivatives in good to high yields (entries 1 and 5). The primary TBS and TIPS group of **11a** or **13** could be deprotected selectively with 50% HF in CH₃CN to furnish **12** and **8**, respectively without cleavage of the MDPM group (entries 2 and 4). The trityl group of **11b** was selectively removed by using BF₃•OEt₂ in the presence of TolSH without affecting the MDPM group (entry 3). Selective hydrogenolytic deprotection of the benzyl group of **15** was carried out via 10% Pd-C in ⁱPrOH-water to furnish **13** without over-reduction of the uracil double bond (entry 6). Olefination of the carbonodithioate **16** was achieved by using ⁿBu₃SnH and AIBN in toluene at refluxing temperature to provide 2',3'-didehydro-2',3'-dideoxy derivative **17** in a reasonable yield (entry 7). Hydrogenation of the double bond of **17** was also achieved under a standard hydrogenation condition with 10% Pd-C within 1 h to provide the MDPM-

protected uridine-2',3'-dideoxy derivative **18** in high yield (entry 8). Thus, it was experimentally proved that MDPM group is a robust protecting group for the uridine ureido nitrogen to synthesize a wide range of uridine derivatives.

We have previously developed a novel ester linker **27**, whose esters are stable against Brønsted and Lewis acids, Brønsted bases and a wide variety of nucleophiles.^{7c} If the uridine ureido nitrogen can be immobilized onto polymer-resin, systematic syntheses of capuramycin and muraymycin analogs would be dramatically enhanced. As summarized in Scheme 3, the (2,6-dichloro-4-hydroxyphenyl) (2,4,6-trichlorophenyl) methanone (**20**)¹⁰ could efficiently be linked with hydroxymethylpolystyrene (PS) (~2 mmol/g) via a Mitsunobu reaction.¹¹ The carbonyl group of **21** was reduced by LiBH₄ in THF to afford the PS-alcohol **22**. Available alcohol-linkers on the polymer surface were determined to be 1.8~2.0 mmol/g by coupling of the linkers with Fmoc-β-Ala-OH and subsequent release of Fmoc chromophore and elemental analyses of the Cl atoms for **22**. According to the procedure summarized in Scheme 1, the PS-alcohol was transformed to the PS-MDPM-Cl **23**, whose available chloromethoxy group was determined to be 1.5~1.8 mmol/g by its elemental analysis. Uridine (**7**) and 2,3-isopropylidene uridine **24** could be loaded onto the linker resin in 6 h via a 2 fold excess of **7** or **24** and DBU in DMF. The loaded uridine or 2,3-isopropylidene uridine were cleaved with 30% TFA in CH₂Cl₂ in 3 h to afford uridine (**7**) in greater than 85% yields.

3. Conclusion

In conclusion, we have developed a new protecting group, (2,6-dichloro-4-methoxyphenyl) (2,4,6-dichlorophenyl) methoxymethyl chloride (**1**) for the uridine ureido nitrogen. MDMP protecting group has significant advantages over BOM protecting group for the syntheses of muraymycin and capuramycin analogs systematically in that MDMP group 1) is stable to a wide variety of acids, 2) is also stable to hydrogenation conditions, and 3) can efficiently be deprotected by solvolytic cleavage with TFA (at 30% concentrations) at room temperature within 2 h without the addition of a cation scavenger.¹¹ Similarly, the MDMP resin **23** has been developed to immobilize uridine and a uridine derivative. In this article we have demonstrated robustness of the MDMP group in uridine derivatives and utility of the linker resin **23** with a limited number of molecules. Moreover, as BOM group has been widely utilized in organic syntheses, a new protecting group **1** and linker resin **23** described here will be valuable assets to protect not only for ureido nitrogens, but also for *primary*, *secondary*, and phenolic alcohols, and carboxylic acids.¹³ It is worth mentioning that immobilization of the uridine ureido nitrogen on polymer-support is not possible with previously reported linker resins. Utility of **1** and **23** in generation of optimized libraries of uridine-containing antibiotics in solution or on polymer-support will be reported elsewhere.

4. Experimental section

4.1. General

All glassware were oven dried, assembled hot and cooled under a stream of nitrogen before use. Reactions with air sensitive materials were carried out by standard syringe techniques. Commercially available reagents were used as received without further purification. Thin layer chromatography was performed using 0.25 mm silica gel 60 plates visualizing at 254 nm, or stained with anisaldehyde solution by heating with a hot-air gun. Specified products were purified by flash column chromatography using silica gel 60. IR absorptions were performed on NaCl plates. ¹H NMR spectral data were recorded on 500 or 400 MHz NMR spectrometer. The residual solvent signal was utilized as an internal reference CDCl₃ (7.26). ¹³C NMR spectral data were recorded at 125, 100 MHz instruments. The residual

solvent signal was utilized as an internal reference CDCl_3 (77.23). For all NMR spectra, δ values are given in ppm and J values in Hz.

4.2. (2,6-Dichloro-4-methoxyphenyl)(2,4,6-trichlorophenyl) methanone (4b)

Anhydrous AlCl_3 (450 mg, 3.4 mmol) was added to PhNO_2 (10 mL). The reaction mixture was cooled to 0 °C, and 2, 4, 6-trichlorobenzoyl chloride (0.53 mL, 3.4 mmol) and 3,5-dichloroanisole (500 mg, 2.82 mmol) were added. The reaction mixture was stirred at rt for 24 h, then it was diluted with Et_2O (10 mL) at 0 °C and quenched by 1 N NaOH (~3 mL). The mixture was stirred vigorously until precipitate was formed. Filter and wash the precipitate with DCM. The organic layer was dried over Na_2SO_4 and concentrated to give the crude product. Purification by silica gel chromatography (hexanes/ EtOAc = 20/1) provided **4b** (867 mg, 80%). ^1H NMR (CDCl_3 , 500 MHz) 7.36 (s, 2H), 6.88 (s, 2H), 3.85 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 194.25, 163.33, 141.89, 138.12, 136.31, 129.29, 127.78, 117.65, 57.03; IR (film): 1633, 1557, 1388, 1341 cm^{-1} ; HRMS (EI) m/z = 382.8967 calcd for $\text{C}_{14}\text{H}_8\text{Cl}_5\text{O}_2$ (MH^+); found: 382.8975.

4.3. (2,6-Dichloro-4-methoxyphenyl)-(2,4,6-trichloro-phenyl) methanol (rac-5b)

A stirred solution of **4b** (385 mg, 1 mmol) in THF (2 mL) was added LiBH_4 (2 mL, 2.0 M in THF) dropwise at 0 °C. After 12 h at rt, the reaction mixture was quenched by sat. aq. NH_4Cl (4 mL) at 0 °C. The water phase was extracted with Et_2O . The combined extract was washed with brine, dried over Na_2SO_4 and evaporated *in vacuo*. Purification by silica gel chromatography (hexanes/ EtOAc = 5/1) gave *rac*-**5b** (368 mg, 95%). ^1H NMR (CDCl_3 , 400 MHz) 7.31 (s, 2H), 6.85 (s, 2H), 6.69 (d, J = 10 Hz, 1H), 3.79 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 159.23, 135.91, 135.04, 133.92, 129.57, 127.85, 115.55, 72.82, 55.88; IR (film): 3476, 1457, 1431, 1309 cm^{-1} ; HRMS (EI) m/z = 384.9123 calcd for $\text{C}_{14}\text{H}_{12}\text{Cl}_5\text{O}_2$ [$\text{M}+\text{H}$]; found: 384.9116.

4.4. (2,6-Dichloro-4-methoxyphenyl)-(2,4,6-trichlorophenyl) methoxymethyl methyl sulfide (6)

To a stirred suspension of NaH (31 mg, 60% in oil, 0.47 mmol) in THF (0.4 mL) *rac*-**5b** (100 mg, 0.26 mmol) in THF (0.3 mL) was added at 0 °C. After 30 min, chloromethyl methyl sulfide (0.044 mL, 0.52 mmol) was added. Reaction mixture was stirred at 0 °C for 3 h, and quenched by sat. aq. NH_4Cl . The aqueous layer was extracted with Et_2O , and the combined extract was washed with brine, dried over Na_2SO_4 , and evaporated *in vacuo*. Purification by silica gel chromatography (hexanes/ EtOAc = 10/1) gave **6** (114 mg, 98%). ^1H NMR (CDCl_3 , 400 MHz) 7.31 (s, 2H), 6.86 (s, 2H), 6.68 (s, 1H), 4.72 (q, J = 11.6 Hz, 2H), 3.79 (s, 3H), 2.19 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 159.26, 136.77, 133.95, 132.79, 129.57, 125.47, 115.53, 75.67, 74.29, 55.70, 14.88; IR (film): 3469, 1544, 1368, 1351 cm^{-1} ; HRMS (EI) m/z = 446.8977 calcd for $\text{C}_{16}\text{H}_{13}\text{Cl}_5\text{O}_2\text{SNa}$ ($[\text{M}+\text{Na}]^+$); found: 446.8985.

4.5. (2,6-Dichloro-4-methoxyphenyl)(2,4,6-trichlorophenyl)-methoxy methyl chloride (1)

To a stirred solution of **6** (447 mg, 1.0 mmol) in CH_2Cl_2 (2.5 mL) was added sulfuryl chloride (0.08 mL, 1.0 mmol) at rt. The reaction mixture was stirred for 1 h and all volatiles were evaporated to provide the crude product which was solidified by addition of hexanes (2 mL). The white solid was washed with hexanes twice to afford **1** (418 mg, 96%). ^1H NMR (CDCl_3 , 400 MHz) 7.33 (s, 2H), 6.88 (s, 2H), 6.77 (s, 1H), 5.57 (q, J = 6.4 Hz, 2H), 3.80 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 159.56, 136.83, 136.63, 134.40, 131.73, 129.68, 124.31, 115.62, 80.11, 55.75; IR (film): 3473, 1445, 1309 cm^{-1} ; Anal. Calcd. $\text{C}_{15}\text{H}_{10}\text{Cl}_6\text{O}_2$, C 41.42; H 2.32; Cl 48.91; found C 41.81; H 2.41; Cl 48.97.

4.6. 3-[(2,6-Dichloro-4-methoxyphenyl)(2,4,6-trichlorophenyl) methoxymethyl]-1-(3,4-dihydroxy-5-hydroxymethyl tetrahydrofuran-2-yl)-1H-pyrimidine-2,4-dione (8)

To a stirred solution of uridine (**7**, 1.83 g, 7.5 mmol) in DMF (15 mL) at 0 °C DBU (1.5 mL, 10.0 mmol) and **1** (2.18 g, 5.0 mmol) were added. After 1 h, MeOH (2 ml) was added, and the reaction mixture was partitioned between EtOAc and water. The aqueous layer was extracted with EtOAc, and the combined extract was washed with brine, dried over Na₂SO₄, and evaporated *in vacuo*. Purification by silica gel chromatography (DCM/MeOH = 15/1) afforded **8** (2.70 g, 84%). ¹H NMR (CDCl₃, 400 MHz) 7.67 (d, *J* = 6.8 Hz, 1H), 7.30 (d, *J* = 3.6 Hz, 2H), 6.83 (d, *J* = 4.8 Hz, 2H), 6.57 (s, 1H), 5.77 (d, *J* = 8.4 Hz, 1H), 5.59 (m, 3H), 4.32 (m, 2H), 4.24 (s, 1H), 3.97 (d, *J* = 12.0 Hz, 1H), 3.90 (s, 1H), 3.83 (m, 1H), 3.78 (s, 3H), 3.05 (s, 1H), 2.20 (s, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 162.94, 159.36, 151.86, 140.07, 136.64, 134.06, 132.46, 129.49, 125.15, 115.49, 101.61, 93.18, 85.71, 77.85, 74.79, 70.39, 69.14, 61.69, 55.73, 36.61, 31.51; IR (film): 3435, 1719, 1665, 1440, 1081 cm⁻¹; HRMS (EI) *m/z* = 640.9819 calcd for C₂₄H₂₂Cl₅N₂O₈ [M+H]; found: 640.9825.

4.7. 1-[5-(*tert*-Butyldimethylsilyloxymethyl)-3,4-dihydroxy-tetrahydrofuran-2-yl]-3-[(2,6-dichloro-4-methoxyphenyl)(2,4,6-trichlorophenyl)methoxymethyl]-1H-pyrimidine-2,4-dione (10a)

To a stirred solution of **8** (64 mg, 0.1 mmol) in CH₂Cl₂ (0.5 mL) was added imidazole (14 mg, 0.2 mmol) and TBSCl (18 mg, 0.12 mmol). After 4 h, the reaction mixture was quenched with water and extracted with EtOAc. The combined extract was dried over Na₂SO₄, evaporated *in vacuo*. Purified by silica gel chromatography (hexanes/EtOAc = 5/1) provide **10a** (62 mg, 82 %). ¹H NMR (CDCl₃, 400 MHz) δ 7.89 (d, *J* = 8.4 Hz, 1H), 7.28 (s, 2H), 6.81 (s, 2H), 6.56 (s, 1H), 5.76 (t, *J* = 4.0 Hz, 1H), 5.72 (d, *J* = 8.4 Hz, 1H), 5.55 (s, 1H), 4.18 (bs, 3H), 4.12 (m, 1H), 3.96 (d, *J* = 11.6 Hz, 1H), 3.79 (d, *J* = 11.6 Hz, 1H), 3.75 (s, 3H), 3.13 (bs, 1H), 0.90 (s, 9H), 0.13 (s, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 162.80, 159.33, 152.24, 138.47, 136.65, 134.03, 132.48, 129.47, 125.22, 115.47, 101.46, 91.72, 85.99, 77.93, 76.44, 70.51, 69.17, 62.29, 55.69, 36.57, 31.6, -5.48, -5.56; [α]_D²⁰ = +28 (*c* 2.5 in CHCl₃); IR (film): 3420, 1701, 1659, 1439, 1255, 1088 cm⁻¹; HRMS (EI) *m/z* = 755.0684 calcd for C₃₀H₃₆Cl₅N₂O₈Si [M+H]; found: 755.0681.

4.8. 3-[(2,6-Dichloro-4-methoxy-phenyl)(2,4,6-trichlorophenyl) methoxymethyl]-1-(3,4-dihydroxy-5-trityloxymethyl tetrahydrofuran-2-yl)-1H-pyrimidine-2,4-dione (10b)

To a stirred solution of **8** (128 mg, 0.2 mmol) in pyridine (0.7 mL) was added trityl chloride (67 mg, 0.24 mmol) and DMAP (2 mg). The reaction mixture was stirred at 60 °C for 4 h and cooled to rt. All volatiles were removed. The partition between EtOAc and water was conducted. EtOAc phase was washed with 1 NHCl and brine. The organic phase was dried over Na₂SO₄ and concentrated *in vacuo*. Purification by silica gel chromatography (hexanes/EtOAc = 6/1) gave the **10b** (155 mg, 88%). ¹H NMR (500 MHz, CDCl₃) δ 7.71 (t, *J* = 8.5 Hz, 1H), 7.31 (d, *J* = 7.5 Hz, 6H), 7.25 (m, 6H), 7.22 (m, 5H), 6.74 (s, 2H), 6.47 (s, 1H), 5.64 (s, 1H), 5.51 (m, 2H), 5.39 (d, *J* = 7.0 Hz, 1H), 4.27 (bs, 1H), 4.14 (bs, 2H), 4.10 (bs, 1H), 3.65 (d, *J* = 4.5 Hz, 3H), 3.42 (d, *J* = 11.0 Hz, 1H), 3.33 (m, 1H), 2.94 (bs, 1H); ¹³C NMR (125 MHz, CDCl₃) δ 162.70, 159.35, 152.09, 143.18, 138.42, 136.68, 134.05, 132.56, 129.51, 128.64, 128.09, 127.48, 125.25, 115.50, 101.65, 91.90, 87.56, 84.51, 77.86, 76.19, 70.61, 69.17, 62.40, 55.71, 25.64; [α]_D²⁰ = +23 (*c* 1.5 in CHCl₃); IR (film): 3425, 1718, 1669, 1446, 1096 cm⁻¹; HRMS (EI) *m/z* = 883.0914 calcd for C₄₃H₃₆Cl₅N₂O₈ [M+H]; found: 883.0916.

4.9. 3-[(2,6-Dichloro-4-methoxy-phenyl)(2,4,6-trichloro-phenyl)methoxymethyl]-1-(3,4-dihydroxy-5-triisopropylsilanyloxymethyltetrahydrofuran-2-yl)-1H-pyrimidine-2,4-dione (**13**)

The same procedure for the synthesis of **10a** was applied, but TIPSCl was used. Yield: 90%. ¹H NMR (500 MHz, CDCl₃) δ 7.90 (q, *J* = 8.0 Hz, 1H), 7.30 (s, 2H), 6.83 (d, *J* = 1.5 Hz, 2H), 6.57 (s, 1H), 5.74 (m, 2H), 5.57 (m, 2H), 4.32 (bs, 1H), 4.19 (m, 2H), 4.06 (d, *J* = 12.5 Hz, 1H), 3.77 (s, 3H), 2.92 (bs, 1H), 1.16 (m, 3H), 1.08 (d, *J* = 6.5 Hz, 9H); ¹³C NMR (125 MHz, CDCl₃) δ 162.68, 159.35, 152.37, 138.26, 136.69, 134.05, 132.58, 129.50, 125.25, 115.50, 101.48, 92.01, 86.31, 77.95, 76.64, 70.59, 69.17, 62.55, 55.72, 25.65, 18.02, 11.85, -3.58; [α]_D²⁰ = +27 (*c* 5.5 in CHCl₃); IR (film): 3422, 1700, 1657, 1441, 1243, 1093 cm⁻¹; HRMS (EI) *m/z* = 797.1153 calcd for C₃₃H₄₂Cl₅N₂O₈Si [M+H]; found: 797.1159.

4.10. General procedure of desilylations

To a stirred solution of **13** (26 mg, 0.03 mmol) in MeCN (0.5 mL) at rt was added 50% HF (100 μL). After 5 h, the reaction mixture was quenched by aq. NaHCO₃. The organic layer was washed with brine, and then dried over Na₂SO₄. Purification by silica gel chromatography (hexanes/EtOAc = 4/1) gave the product **8** (19 mg, 88%).

4.11. Cyclohexylidenation of **10a**

To a stirred solution of **10a** (75.6 mg, 0.1 mmol) in MeCN (0.5 mL) were added dipent-4-enyl acetal (30 mg, 0.12 mmol), NBS (47 mg, 0.26 mmol), and BF₃-OEt₂ (0.01 mmol). The reaction mixture was stirred at rt and protected from light with an aluminum foil for 15 min. The reaction mixture was quenched with Et₃N (50 μL), and extracted with CH₂Cl₂. The organic extract was washed with 10% Na₂S₂O₃, and sat. aq. NaHCO₃, and dried over Na₂SO₄. Purified by silica gel chromatography (hexanes/EtOAc = 10/1) afforded **11a** (63 mg, 75%). ¹H NMR (500 MHz, CDCl₃) δ 7.69 (d, *J* = 8.5 Hz, 1H), 7.29 (d, *J* = 3.5 Hz, 2H), 6.82 (d, *J* = 3.5 Hz, 2H), 6.59 (d, *J* = 8.5 Hz, 1H), 5.94 (d, *J* = 4.0 Hz, 1H), 5.72 (d, *J* = 8.0 Hz, 1H), 5.59 (m, 2H), 4.73 (m, 1H), 4.65 (m, 1H), 4.29 (d, *J* = 2.5 Hz, 1H), 3.93 (d, *J* = 11.5 Hz, 1H), 3.79 (d, *J* = 11.5 Hz, 1H), 3.77 (s, 3H), 1.77 (m, 2H), 1.67 (m, 2H), 1.57 (m, 4H), 1.40 (m, 2H); ¹³C NMR (125 MHz, CDCl₃) δ 162.53, 159.24, 151.44, 141.79, 136.70, 133.94, 132.58, 129.47, 125.18, 115.50, 114.94, 102.13, 97.64, 87.33, 83.16, 79.85, 78.03, 69.25, 63.28, 55.68, 37.17, 34.89, 25.89, 24.91, 23.95, 23.63, 18.33, -5.40, -5.45; Yield: 75%. [α]_D²⁰ = +19 (*c* 2.0 in CHCl₃); IR (film): 1726, 1703, 1656, 1442, 1261, 1088 cm⁻¹; HRMS (EI) *m/z* = 835.1310 calcd for C₃₆H₄₄Cl₅N₂O₈Si [M+H]; found: 835.1316.

4.12. 3-[(2,6-Dichloro-4-methoxyphenyl)-(2,4,6-trichloro phenyl)methoxymethyl]-1-(2-cyclohexyl-6-trityloxymethyltetrahydrofuro[3,4-d][1,3]dioxol-4-yl)-1H-pyrimidine-2,4-dione (**11b**)

Procedure, see **4.11**. Yield: 90%. ¹H NMR (400 MHz, CDCl₃) δ 7.50 (dd, *J* = 4.0 Hz, 1H), 7.39 (d, *J* = 7.6 Hz, 7H), 7.29 (m, 9H), 6.82 (d, *J* = 3.2 Hz, 2H), 6.54 (d, *J* = 8.4 Hz, 1H), 5.85 (d, *J* = 3.2 Hz, 1H), 5.55 (t, *J* = 11.6 Hz, 1H), 5.46 (t, *J* = 10.0 Hz, 1H), 5.36 (bs, 1H), 4.78 (d, *J* = 6.4, 2H), 4.35 (s, 1H), 3.75 (s, 3H), 3.40 (bs, 2H), 1.74 (m, 2H), 1.59 (m, 2H), 1.25 (m, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 162.60, 159.33, 152.07, 143.19, 138.02, 136.48, 134.15, 132.76, 129.50, 128.69, 128.09, 127.48, 125.28, 115.51, 101.66, 97.63, 91.91, 87.56, 84.51, 77.80, 76.19, 70.51, 69.19, 62.42, 55.71, 37.15, 34.60, 25.04, 23.55; [α]_D²⁰ = +17 (*c* 2.3 in CHCl₃); IR (film): 1736, 1719, 1674, 1444, 1231 1075 cm⁻¹; HRMS (EI) *m/z* = 963.1540 calcd for C₄₉H₄₄Cl₅N₂O₈ [M+H]; found: 963.1544.

4.13. 3-[(2,6-Dichloro-4-methoxy-phenyl)-(2,4,6-trichloro-phenyl)-methoxymethyl]-1-(2-cyclohexyl-6-hydroxymethyl-tetrahydrofuro[3,4-d][1,3]dioxol-4-yl)-1H-pyrimidine-2,4-dione (12)

To a stirred solution of **11b** (35 mg, 0.04 mmol) in CH₂Cl₂ (0.4 mL) at 0 °C was added TolSH (14 mg, 0.12 mmol) and BF₃•Et₂O (100 μL). After 1 h at rt., the reaction was quenched with sat. aq. NaHCO₃. The aqueous layer was extracted with CH₂Cl₂ three times. The combined organic extract was dried over Na₂SO₄, and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (hexanes/EtOAc = 4/1) to provide **12** (26 mg, 90%). ¹H NMR (400 MHz, CDCl₃) δ 7.32 (s, 1H), 7.30 (d, *J* = 5.2 Hz, 2H), 6.83 (d, *J* = 6.4 Hz, 2H), 6.58 (s, 1H), 5.77 (dd, *J* = 8.0 Hz, 1H), 5.57 (s, 2H), 5.47 (dd, *J* = 5.6 Hz, 1H), 5.00 (d, *J* = 2.8 Hz, 1H), 4.93 (dd, *J* = 3.6 Hz, 1H), 4.28 (d, *J* = 2.8 Hz, 1H), 3.88 (d, *J* = 8.0 Hz, 1H), 3.79 (d, *J* = 8.0 Hz, 1H), 3.77 (s, 3H), 2.50 (bs, 1H), 1.67 (m, 2H), 1.64 (m, 2H), 1.59 (m, 4H), 1.41 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 162.56, 159.27, 151.34, 141.88, 136.70, 133.96, 132.55, 129.46, 125.17, 115.52, 115.04, 102.16, 97.66, 87.43, 83.15, 79.88, 78.00, 69.26, 62.86, 55.70, 37.12, 34.66, 24.92, 23.98, 23.54; [α]_D²⁰ = +17 (c 1.0 in CHCl₃); IR (film): 3464, 1743, 1710, 1665, 1456, 1222 cm⁻¹; HRMS (EI) *m/z* = 721.0445 calcd for C₃₀H₃₀Cl₅N₂O₈ [M+H]; found: 721.0450.

4.14. Acetonization of 8

To a stirred solution of **8** (64 mg, 0.1 mmol) in acetone (0.6 mL) at 0 °C were added PTSA (2 mg) and 2,2-dimethoxypropane (15 μL, 0.12 mmol). The mixture was stirred for 4 h at the same temperature and concentrated *in vacuo*. Purification by silica gel chromatography (hexanes/EtOAc = 4/1) afford **14** (59 mg, 87%). ¹H NMR (400 MHz, CDCl₃) δ 7.33 (s, 1H), 7.30 (d, *J* = 5.6 Hz, 2H), 6.83 (d, *J* = 6.0 Hz, 2H), 6.58 (s, 1H), 5.76 (d, *J* = 8.0 Hz, 1H), 5.55 (bs, 2H), 5.47 (dd, *J* = 5.6 Hz, 1H), 5.02 (d, *J* = 2.0 Hz, 1H), 4.94 (d, *J* = 5.6 Hz, 1H), 4.28 (d, *J* = 2.8 Hz, 1H), 3.90 (dd, *J* = 4.8 Hz, 1H), 3.89 (s, 1H), 3.88 (s, 3H), 2.52 (bs, 1H), 1.57 (s, 3H), 1.35 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 162.55, 159.28, 151.35, 141.74, 136.69, 133.97, 132.53, 129.46, 125.18, 115.46, 114.24, 102.13, 97.58, 87.30, 83.67, 80.34, 78.03, 69.24, 62.84, 55.70, 27.25, 25.25; [α]_D²⁰ = +17 (c 1.5 in CHCl₃); IR (film): 3462, 1740, 1713, 1666, 1448, 1221 cm⁻¹; HRMS (EI) *m/z* = 681.0132 calcd for C₂₇H₂₆Cl₅N₂O₈ [M+H]; found: 681.0138.

4.15. 1-(6-Benzyloxymethyl-2,2-dimethyl-tetrahydro-furo[3,4-d][1,3]dioxol-4-yl)-3-[(2,6-dichloro-4-methoxy-phenyl)-(2,4,6-trichloro-phenyl)methoxymethyl]-1H-pyrimidine-2,4-dione (15)

To a stirred suspension of NaH (5 mg, 60% in oil, 0.125 mmol) in DMF (0.5 mL) at 0 °C was added **14** (78 mg, 0.1 mmol) in DMF (0.5 mL). After 5 min. BnBr (36 μL, 0.3 mmol) was added. After 3 h, the reaction mixture was quenched with H₂O, and extracted with EtOAc. The organic phase was washed with brine, dried over Na₂SO₄, and evaporated *in vacuo* to give the crude product. This was used for the study of debenzoylation.

4.16. General procedure of hydrogenation

To a solution of **15** (76 mg, 0.1 mmol) in ⁱPrOH/H₂O (2:1, 3 mL) was added 10% Pd/C (10 mg) under N₂. H₂ was intruded by using a balloon. After 2 h, Pd/C was filtered with a celite pad. The filtrate was concentrated *in vacuo*. Purification by silica gel chromatography (hexanes/EtOAc = 4/1) afforded **14** (65 mg, 95%). Physical data for **14**, see **4.14**.

4.17 O,O'-((2R,3S,4S,5R)-2-(((tert-butyl)dimethylsilyloxy)methyl)-5-(3-(((2,6-dichloro-4-methoxyphenyl)(2,4,6-trichlorophenyl)methoxy)methyl)-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)tetrahydrofuran-3,4-diyl) S,S'-dimethyl dicarbonodithioate (16)

To a stirred solution of **10a** (160 mg, 0.21 mmol) in DMSO (0.3 mL) was added 5 N NaOH (0.2 mL). After 20 min., CS₂ (0.2 mL) was added. After an additional 30 min., MeI (0.3 mL) was added. Then the reaction mixture was stirred for 2 h, and quenched by sat. aq. NH₄Cl, and extracted with EtOAc. The combined organic extract was washed with brine, dried over Na₂SO₄, concentrated *in vacuo*. Purification by silica gel chromatography (hexanes/EtOAc = 10/1) gave **15** (135 mg, 69%). ¹H NMR (400 MHz, CDCl₃) δ 7.88 (d, *J* = 8.0 Hz, 1H), 7.28 (d, *J* = 11.6 Hz, 2H), 6.82 (d, *J* = 9.2 Hz, 2H), 6.60 (t, *J* = 7.6 Hz, 1H), 6.55 (d, *J* = 9.2 Hz, 1H), 6.25 (t, *J* = 5.2 Hz, 1H), 6.05 (m, 1H), 5.77 (t, *J* = 6.8 Hz, 1H), 5.60 (m, 2H), 4.45 (s, 1H), 4.03 (dd, *J* = 27.2 Hz, 2H), 3.76 (s, 3H), 2.59 (s, 3H), 2.52 (d, *J* = 15.2 Hz, 3H); 0.95 (s, 9H), 0.19 (d, *J* = 12.0 Hz, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 214.70, 162.61, 159.23, 151.29, 138.36, 136.69, 134.58, 133.93, 132.88, 129.38, 125.23, 115.53, 102.83, 85.97, 84.37, 80.01, 78.77, 77.81, 69.47, 63.35, 55.69, 26.01, 19.11, 18.40, -5.25, -5.59; [α]_D²⁰ = +23 (*c* 2.5 in CHCl₃); IR (film): 1711, 1648, 1446, 1259, 1079 cm⁻¹; HRMS (EI) *m/z* = 956.9699 calcd for C₃₄H₃₉Cl₅N₂O₈S₄SiNa ([M+Na]⁺); found: 956.9702.

4.18. 1-[5-(tert-Butyl-dimethylsilyloxymethyl)-2,5-dihydrofuran-2-yl]-3-[(2,6-dichloro-4-methoxyphenyl)(2,4,6-trichloro-phenyl)methoxymethyl]-1H-pyrimidine-2,4-dione (17)

To a stirred solution of **15** (47 mg, 0.05 mmol) in toluene (0.5 mL) were added AIBN (4 mg) and ⁿBu₃SnH (0.07 mL, 0.25 mmol) at 100 °C. After 30 min, the reaction mixture was cooled to rt and all volatiles were evaporated *in vacuo*. Purification by silica gel chromatography (hexanes/EtOAc = 10/1) gave **16** (26 mg, 71%). ¹H NMR (400 MHz, CDCl₃) δ 7.78 (m, 1H), 7.31 (d, *J* = 2.0 Hz, 2H), 6.88 (bs, 1H), 6.75 (d, *J* = 4.8 Hz, 2H), 6.23 (d, *J* = 9.2 Hz, 1H), 5.81 (m, 1H), 5.69 (m, 1H), 5.61 (m, 2H), 4.88 (bs, 1H), 3.87 (m, 1H), 3.78 (m, 1H), 3.69 (s, 1H), 3.68 (s, 3H), 0.90 (s, 9H), 0.17 (d, *J* = 12.0 Hz, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 163.10, 159.26, 139.28, 136.78, 133.94, 132.83, 129.42, 126.83, 125.24, 115.43, 101.80, 90.59, 87.41, 77.80, 69.21, 64.14, 55.68, 25.92, 18.52, -5.36, -5.50; [α]_D²⁰ = +20 (*c* 1.5 in CHCl₃); IR (film): 1705, 1658, 1443, 1255, 1079 cm⁻¹; HRMS (EI) *m/z* = 721.0629 calcd for C₃₀H₃₄Cl₅N₂O₆Si [M+H]; found: 721.0623.

4.19. 1-[5-(tert-Butyl-dimethylsilyloxymethyl)-tetrahydro furan-2-yl]-3-[(2,6-dichloro-4-methoxyphenyl)(2,4,6-trichlorophenyl)methoxymethyl]-1H-pyrimidine-2,4-dione (18)

The procedure described for **15** was applied. Yield: 95%. ¹H NMR (400 MHz, CDCl₃) δ 7.70 (m, 1H), 7.29 (d, *J* = 2.0 Hz, 2H), 6.77 (d, *J* = 4.8 Hz, 2H), 6.21 (d, *J* = 9.2 Hz, 1H), 5.85 (m, 1H), 5.78 (m, 1H), 5.63 (m, 1H), 4.80 (bs, 1H), 3.82 (m, 1H), 3.71 (m, 1H), 3.68 (s, 1H), 3.66 (s, 3H), 2.15 (m, 2H), 1.85 (m, 2H), 0.90 (s, 9H), 0.17 (d, *J* = 12.0 Hz, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 163.11, 159.22, 139.25, 136.79, 133.96, 132.85, 125.21, 115.40, 101.87, 90.64, 87.39, 77.82, 69.19, 64.11, 55.66, 28.01, 25.95, 21.33, 18.55, -5.35, -5.51; [α]_D²⁰ = +21 (*c* 1.3 in CHCl₃); IR (film): 1707, 1650, 1440, 1245, 1083 cm⁻¹; HRMS (EI) *m/z* = 723.0785 calcd for C₃₀H₃₆Cl₅N₂O₆Si [M+H]; found: 723.0789.

4.20. Polymer-supported (2,6-dichloro-4-methoxyphenyl) (2,4,6-trichlorophenyl)methanone 21

Hydromethylstyrene resin **19** (purchased from Aldrich, 1.0 g, ~2 mmol) was washed with THF. **19** was suspended in THF (20 mL) and TPP (5.24 g, 20 mmol), DIAD (3.93 mL, 20 mmol) and **20** (1.11 g, 3 mmol) were added. The reaction mixture was gently stirred for 12 h. The polymer-resins were filtered, and thoroughly washed with THF/H₂O (4:1), THF, EtOAc, and hexanes, and dried under high vacuum to give **21** (1.75 g).

4.21. Polymer-supported (2,6-dichloro-4-methoxyphenyl) (2,4,6-trichlorophenyl)methanol **22**

To a gently stirred polymer-resin **21** (1.75 g) in THF (8 mL) was added LiBH₄ (10 ml, 2.0 M in THF) at 0 °C. After 12 h at rt, the reaction mixture was quenched with H₂O (5 mL). The polymer-resins were thoroughly washed with THF/1% HCl (4:1), THF, EtOAc and hexanes, and dried under high vacuum to afford the polymer-resin **22** (1.66 g).

4.22. Polymer-supported MDPM-CI **23**

To a suspension of **22** (0.44 g) in THF (10 mL) was added NaHMDS (0.6 mL, 1.0 M in THF, 0.6 mmol). After 30 min, chloromethyl methyl sulfide (0.09 ml, 1.3 mmol) was added. After 8 h, the polymer was thoroughly washed with THF/H₂O (4:1), THF, EtOAc, and hexanes, and dried *in vacuo* to give the polymer-supported methyl sulfide (464 mg). This was suspended in CH₂Cl₂ (2 mL) and sulfonyl chloride (0.17 mL, 4.1 mmol) was added. After 5 h, the polymer resins were washed thoroughly with CH₂Cl₂, and dried under high vacuum to afford **23**. Anal. found. Cl 232mg/g.

4.23. Loading uridine on PS-MDPM-CI resin **23**

To a suspension of PS-MDPM-CI **23** (800 mg) in DMF (1 mL) were added DBU (30μL, 0.2 mmol) and uridine (49 mg, 0.2 mmol). After 3 h, the polymer resins were thoroughly washed with THF/H₂O (4:1), THF, EtOAc and hexanes, and then dried under high vacuum to afford **25** (55 mg).

4.24. Loading **24** on PS-MDPM-CI resin **23**

The procedure was same as described in **4.23**.

4.25. Cleavage of linker

The resin **25** was washed with CH₂Cl₂, and added 30 % TFA in CH₂Cl₂ (2 mL). After 4 h at rt., the polymer resins were washed thoroughly with CH₂Cl₂. The filtrate was concentrated *in vacuo*. Purification by silica gel chromatography (CHCl₃:MeOH = 10:1) furnished uridine (**7**) (19 mg, 90%). The resin **26** was cleaved via the same procedure to afford **7** (21 mg, 85%)


Acknowledgments

The authors thank the National Institutes of Health (NIAID grant AI084411-02) and The University of Tennessee for generous financial supports. NMR data were obtained on instruments supported by the NIH Shared Instrumentation Grant.


References and notes

- (a) Yamamoto T, Koyama H, Kurajoh M, Shoji T, Tsutsumi Z, Moriwaki Y. *Clinica Chimica Acta*. 2011; 412:1712–1724.(b) Guillemette C, Levesque E, Harvey M, Bellemare J, Menard V. *Drug Metabo Rev*. 2010; 42:24–44.
- (a) Bryskier A, Dini AC. *Agents: Antibacter Antifung*. 2005:377–398.(b) Timothy DH, Lloyd AJ, Roper DI. *Infect Disorders: Drug Targets*. 2006; 6:85–106.
- (a) Reddy VM, Einck L, Nacy CA. *Antimicro Agents Chemother*. 2008; 52:719–721.(b) Ichikawa S. *Chem Pharm Bull*. 2008; 56:1059–1072. [PubMed: 18670104] (c) Dini, Christophe. *Curr Top Med Chem*. 2005; 5:1221–1236. [PubMed: 16305528] (d) Dini C, Drochon N, Feteanu S, Guillot JC, Peixoto C, Aszodi J. *Bioorg Med Chem Lett*. 2001; 11:529–531. [PubMed: 11229763]
- (a) Kurosu M, Li K, Crick DC. *Org Lett*. 2009; 11:2393–2397. [PubMed: 19405507] (b) Kurosu M, Li K. *J Org Chem*. 2008; 73:9767–9771. [PubMed: 18783274] (c) Kurosu M, Biswas K, Crick DC. *Org Lett*. 2007; 9:1141–1145. [PubMed: 17311394] (d) Kurosu M, Narayanasamy P, Crick DC.


- Heterocycles. 2007; 72:339–352.(e) Kurosu M, Mahapatra S, Narayanasamy P, Crick DC. *Tetrahedron Lett.* 2007; 48:799–803.
5. (a) Hanessian S, Kloss J, Sugawara T. *J Am Chem Soc.* 1986; 108:2758–2759.(b) Goff DA, Harris RN, Bottaro JC, Bedford CD. *J Org Chem.* 1986; 51:4711–4714.(c) Connor DS, Klein GW, Taylor GN, Boeckman RK, Medwid JB. *Org Syn.* 1972; 52:C10–19.
 6. Spork AP, Wiegmann D, Granitzka M, Stalke D, Ducho C. *J Org Chem.* 2011; 76:10083–10098. [PubMed: 22059552]
 7. (a) Kurosu M, Li K. *Org Lett.* 2009; 11:911–914. [PubMed: 19170578] (b) Kurosu M, Li K. *Synthesis.* 2009; 21:3633–3641.(c) Kurosu M, Biswas K, Crick DC. *Org Lett.* 2007; 9:1141–1144. [PubMed: 17311394] (d) Kurosu M, Biswas K, Narayanasamy P, Crick DC. *Synthesis.* 2007; 16:2513–2516.
 8. The esters and carbamates synthesized with 5b showed single isomers in their ¹H-NMRs and single peaks in their HPLC chromatography analyses.
 9. Goómez C, Macià B, Lillo VJ, Yus M. *Tetrahedron.* 2006; 62:9832–9839.
 10. (2,6-Dichloro-4-hydroxyphenyl) (2,4,6-trichlorophenyl) methanone (20) was synthesized by demethylation of 4b with HBr in AcOH.
 11. Mitsunobu O. *Synthesis.* 1981; 29:1–28.
 12. Deprotections of alkoxybenzyl ethers have resulted in good yields when a cation scavenger was added into reaction mixture.
 13. The BOM group has been utilized to protect alcohols and carboxylic acids. We realized that 1 and 23 could protect or immobilize primary alcohols, phenolic alcohols, and carboxylic acids.



Scheme 1.
Syntheses of monomethoxydiphenylmethoxymethyl chlorides **1**.




Scheme 2.
Deprotection of MDPM group.

**Scheme 3.**

The linker **23** for immobilizations of uridine derivatives.

Table 1

Protection of the uridine ureido nitrogen with **1** and its relativities against representative reagents.

















Conditions ^a	8 (t _{1/2})	Conditions ^a	8 (t _{1/2})
20% TFA, CH ₂ Cl ₂	>6h	DBU, Toluene	>24h
10% TMSOTf, CH ₂ Cl ₂	>2h	Al(Hg), 1,4-Dioxane	>12h
10% HCl, MeCN	>6h	Raney Ni, 1,4-Dioxane	>12h
10% HCl, 1,4-dioxane	>6h	^t Bu ₃ SnH, AIBN, Toluene, reflux	>6h
30% HF, MeCN	>12h	NBS, THF	>6h
80% AcOH	>12h		
30% TsOH, 1,4-dioxane	>12h	hv, MeCN	>6h
La(OTf) ₃ , THF	>12h	H ₂ /Pd-C, 1 atm MeOH	>12h ^b
10% BF ₃ -OEt ₂ , CH ₂ Cl ₂	>12h	H ₂ /Pd-C, 50 psi MeOH	>12h ^b
10% TiCl ₄ , CH ₂ Cl ₂	>4h	H ₂ /Pd-C, 100 psi MeOH	>12h ^b
DDQ, CH ₂ Cl ₂ -water	>12h	H ₂ /Pd black MeOH	>12h ^b





^aReaction was carried out at room temperature.

^bthe uracil double bond of **8** was reduced.

Table 2

Functionalizations of MDPM-protected uridines.

Entry	Starting material	Conditions	Product	Yield (%) ^d
1	 <p>MDPM: </p> <p>10a : R = TBS 10b : R = Tr</p>	NBS, TMSOTf/ CH ₃ CN, rt, 1h	 <p>11a : R = TBS 11b : R = Tr</p>	75 (95) ^b for 10a 90 (98) ^b for 10b
2	 <p>11a</p>	50% HF/ CH ₃ CN, rt, 1h	 <p>12</p>	80
3	 <p>11b</p>	BF ₃ -OEt ₂ (3 equiv), TolSH/ CH ₂ Cl ₂ , rt, 1h	 <p>12</p>	90
4	 <p>13</p>	50% HF/ CH ₃ CN, rt, 12h	 <p>8</p>	88
5	 <p>8</p>	 TsOH-H ₂ O/ acetone, rt, 4h	 <p>14</p>	90
6	 <p>15</p>	H ₂ (1 atm) / 10% Pd-C iPrOH-watert, 2h	 <p>14</p>	95 ^c

Entry	Starting material	Conditions	Product	Yield (%) ^a
7	 16	ⁿ BuSnH, AIBN / toluene, reflux, 2h	 17	75
8	 17	H ₂ (1 atom) / 10% Pd-C MeOH, 2h,	 18	95 ^c

^aProduct was isolated by SiO₂ chromatography or PTLC.,

^byield based on recovering starting material.,

^cno over-reduction was observed.