

Published in final edited form as:

*Oncol Rep.* 2010 July ; 24(1): 171–176.

## A variant of estrogen receptor- $\alpha$ , ER- $\alpha$ 36 is expressed in human gastric cancer and is highly correlated with lymph node metastasis

Hao Deng<sup>1</sup>, Xuan Huang<sup>1</sup>, Jun Fan<sup>1</sup>, Liubai Wang<sup>1</sup>, Qin Xia<sup>1</sup>, Xiuping Yang<sup>2</sup>, Zhaoyi Wang<sup>3</sup>, and Lijiang Liu<sup>1,4</sup>

<sup>1</sup>Department of Pathology and Pathophysiology, School of Medicine, Jiangnan University, Wuhan, Hubei Province 430056

<sup>2</sup>Department of Pathology, Xiehe Hospital, Tongji Medical College, HuaZhong University of Science and Technology, Wuhan, Hubei Province 430030, P.R. China

<sup>3</sup>Departments of Medical Microbiology and Immunology and Pathology, Creighton University Medical School, 2500 California Plaza, Omaha, NE 68178, USA

<sup>4</sup>Jiangda Pathology Center, Jiangnan University, Wuhan, Hubei Province 430056, P.R. China

### Abstract

Gastric cancer is one of the most common cancers in the world; almost two-thirds of gastric cancer cases and deaths occur in less developed regions. The molecular and cellular events during development of gastric cancer remain unclear. Herein, we examined the expression of ER- $\alpha$ 36, an ER- $\alpha$  variant, in established gastric cancer cell lines and specimens from 22 gastric cancer patients. RT-PCR, Western blot and immunohistochemistry methods were used to assess expression levels of ER- $\alpha$ 36. ER- $\alpha$ 36 localization in gastric cancer cells was determined with an immunofluorescence assay. Both mRNA and protein of ER- $\alpha$ 36 were detected in all established gastric cancer cell lines examined. Higher levels of ER- $\alpha$ 36 mRNA were expressed in 17 of 22 (77.27%) tumor specimens examined compared to the paired normal tissues ( $p < 0.05$ ). ER- $\alpha$ 36 protein was expressed mainly on the plasma membrane and in the cytoplasm of the established gastric cancer cells. ER- $\alpha$ 36 expression is highly correlated with lymph node metastasis in human gastric cancer ( $P < 0.05$ ). The estrogen receptor variant ER- $\alpha$ 36 is highly expressed in human gastric cancer. ER- $\alpha$ 36 expression may be used as a predictive marker for lymph node metastasis of gastric cancer.

### Keywords

gastric cancer; ER- $\alpha$ 36; lymph node metastasis

### Introduction

Estrogen signaling is pivotal in the establishment and maintenance of reproductive function in men and women. It is also involved in normal development and physiology of bone, cardiovascular and neuronal systems. It is believed that dysregulated estrogen signaling is involved in the initiation and development of neoplasia in at least breast and endometrial

cancers (1). However, the involvement of estrogen signaling in other types of human cancer has not been well established.

The diverse physiological functions of estrogens are mediated by specific nuclear receptors designated as estrogen receptors (ERs). In 1986, the cloning of the estrogen receptor (ER) was first reported (2,3). Until 1996, it was assumed that there was only one ER responsible for all of the physiological and pharmacological effects of natural and synthetic estrogens and antiestrogens. However, in 1996, a second ER was cloned (4). Currently, the first ER discovered is referred to as ER- $\alpha$ , while the second is called ER- $\beta$ .

ER- $\alpha$  and ER- $\beta$  share a common structural architecture (5,6). Both are composed of three independent but interacting functional domains: the N-terminal A/B domain, the C or DNA-binding domain, and the D/E/F or ligand-binding domain. The N-terminal domain of ER- $\alpha$  encodes a ligand-independent activation function (AF-1), a region involved in interaction with co-activators, and transcriptional activation of target-genes. The DNA-binding domain or C domain contains a two zinc-finger structure, which plays an important role in receptor dimerization and binding to specific DNA sequences. The C-terminal E/F domain is a ligand-binding domain that mediates ligand binding, receptor dimerization, nuclear translocation, and a ligand-dependent transactivation function (AF-2).

It has been well documented that the ERs are ligand-activated transcription factors that stimulate target gene transcription (1). Stimulation of target gene expression by ERs in response to estrogens is prevalently thought to be responsible for cell proliferation. Despite the clarity with which ERs have been shown to act as transcription factors, it became apparent that not all of the physiological effects mediated by estrogens are accomplished through a direct effect on gene transcription. Another signaling pathway (also known as a 'non-classic', 'non-genomic' or 'membrane signaling' pathway) exists that involves cytoplasmic proteins, growth factors and other membrane-initiated signaling pathways (7).

Previously, we identified and cloned a novel variant of ER- $\alpha$ , ER- $\alpha$ 36, with a molecular weight of 36 kDa (8,9). ER- $\alpha$ 36 lacks both transactivation domains AF-1 and AF-2 of the 66-kDa full-length ER- $\alpha$  (ER- $\alpha$ 66) (8,9). The transcripts of ER- $\alpha$ 36 are initiated from a previously unidentified promoter located in the first intron of the ER- $\alpha$ 66 gene (8). Further research found that ER- $\alpha$ 36 is subjected to a transcriptional regulation totally different from the ER- $\alpha$ 66 (10).

Estrogens have been proposed to provide protection against gastric cancer of women before menopause (11); the incidence of gastric cancer is higher in men than in women before menopause (12). The incidence of gastric cancer in women after menopause reaches a level similar to men (13). These findings suggested an involvement of estrogen signaling in protection against human gastric cancer. Tokunaga *et al* (14) first reported ER- $\alpha$ 66 expression in human gastric cancer. Since then, a number of laboratories have reported expression of ER- $\alpha$  in gastric cancer and correlated ER status to tumor grade (15,16). In addition, the effects of estrogens on the growth of human gastric cancer xenografts in nude mice were studied (17). It was found that ER- $\alpha$ 66 was expressed mainly in the cytoplasm of gastric carcinoma cells while it is predominantly expressed in the cell nuclear of breast cancer cells (18).

In the present study, we examined the expression and localization of ER- $\alpha$ 36 in established gastric cancer cells and specimens from gastric cancer patients. We also examined correlations between ER- $\alpha$ 36 expression and lymph node metastasis of human gastric adenocarcinoma.

## Materials and methods

### Cell culture

Human gastric adenocarcinoma cell lines BGC-823 and SGC7901 were obtained from Cell Center of Basic Medicine, Chinese Academy of Medical Sciences (Beijing, China). Human gastric cancer cell line AGS and breast cancer cell line MCF-7 were obtained from American Type Culture Collection (ATCC, Manassas, VA, USA). Human gastric cancer cell line MKN-45 was kindly provided by the laboratory of Molecular and Immunopharmacology of Tongji Medical College. All cells were maintained in RPMI-1640 medium plus 10% FCS at 37°C in a 5% CO<sub>2</sub> atmosphere.

### Gastric tumor samples

Tumor tissues, corresponding normal gastric tissues and tumor-adjacent tissues of 22 gastric cancer patients (between January and July, 2007) were obtained from the Xiehe Hospital of Tongji Medical College, after approval of the Institutional Review Board. Tumor tissues were divided into two groups. One group was fixed in 10% formalin, embedded in paraffin and stained with hematoxylin and eosin and immunohistochemistry. The other group was snap-frozen in liquid nitrogen and stored at -80°C for RNA extraction. The patients comprised 13 men and eight women aged 36–79 (mean: 56.9 years). Prior to surgery, none of the patients received any anti-cancer treatment. The tumor size, histological differentiation, T stage and N stage were evaluated according to the clinicopathological classification of the WHO (2003).

### RT-PCR amplification

For RT-PCR analysis, gastric cancer cells and frozen tissues of gastric adenocarcinoma were disrupted in TRIzol reagent (Invitrogen, San Diego, CA, USA) and total RNA was isolated according to the manufacturer's instructions. Total RNA was reversely transcribed into cDNA by RT-PCR kit (Takara, Dalian, China). Following reverse transcription, PCR reaction was carried out with 35 cycles using the following conditions: 94°C for 40 sec, 60°C for 1 min and 72°C for 40 sec. ER- $\alpha$ 36 primers and GAPDH Primers were designed by Primer 5.0 (Primer Biosoft International, Palo Alto, CA, USA) and were used simultaneously in the same reaction. The following primers were used: ER- $\alpha$ 36 forward primer 5'-CAAGTGGTTTCCTCGTGTCTAAAG-3'; ER- $\alpha$ 36 reverse primer 5'-TGTTGAGTGTTGGTTGCC AGG-3'; GAPDH forward primer 5'-ACCACAGTCCATGCCATCAC-3'; GAPDH reverse primer 5'-TCCACCACCC TGTTGCTGTA-3'. The PCR products (219 bp for ER- $\alpha$ 36 and 452 bp for GAPDH) were separated on 1.5% agarose gel and stained with GoldView (SBS Genetech Co., Beijing, China). The densities of DNA bands were assessed with the analysis software (Biostep Photoimpact, Beijing, China) and the relative values were determined.

### Western blot assay

Cells were lysed with the RIPA buffer (Beyotime, Shanghai, China). Proteins concentration was determined with an Enhanced BCA Protein assay kit (Beyotime). Cell lysates were mixed with gel-loading buffer, separated by 12% SDS-PAGE gels and transferred to a PVDF membrane (Millipore, Billerica, MA, USA). The membrane was probed with the indicated antibodies and visualized with the appropriate secondary antibodies (Beyotime) and enhanced chemiluminescence (ECL) reagents (Beyotime). The densities of protein bands were determined with the Totallab analysis software (Nonlinear Dynamics Technical, NC, USA). The rabbit polyclonal ER- $\alpha$ 36 antibody was development as described before (9). The anti- $\beta$ -actin (C4) antibody was purchased from Santa Cruz Biotechnology (CA, USA).

## Immunofluorescence and immunohistochemistry assays

Gastric cancer cells cultured on cover glass were fixed with acetone for 10 min. After washing with PBS, cells were permeabilized with 0.5% (V/V) Triton X-100 for 10 min, washed with PBS and blocked with 3% serum in PBS at room temperature for 1 h. The cover glasses were incubated with the rabbit polyclonal ER- $\alpha$ 36 antibody at room temperature for 1 h and washed 3 times with PBS containing 0.5% Triton X-100 (PBST), then incubated with a Cy3-labeled secondary antibody (Beyotime). The specificity of staining was verified by omitting the primary antibody. Finally, the cover glasses were washed 3 times with PBST, one time with PBS, then mounted with anti-fade medium (Molecular Probes, Eugene, OR, USA) and examined under a Olympus microscope at  $\times 400$  magnification. Hoechst 33258 was used for nuclear staining.

Immunohistochemical staining was performed on 4- $\mu$ m thick tumor sections via a 'two-step' assay. Briefly, tissue slides were deparaffinized with xylene and rehydrated through a gradual alcohol series. The endogenous peroxidase activity was blocked by incubation in a 3% hydrogen peroxide/methanol buffer for 10 min. Antigen retrieval was carried out by immersing the slides in EDTA buffer (pH 8.0) and boiling in a water-bath at 95°C for 25 min. The slides were rinsed in phosphate-buffered saline (PBS) and incubated with normal goat serum to block non-specific staining. The slides were then incubated with the polyclonal anti-ER- $\alpha$ 36 antibody at a dilution of 1:50 overnight at 4°C in a humidified chamber. The slides were incubated with the second antibody (horseradish peroxidase-conjugated goat anti-rabbit immunoglobulin; 1:100, Dako, CA, USA) for 45 min. Diaminobenzidine was used as a chromagen and slides were counterstained with hematoxylin. Scoring for ER- $\alpha$ 36 staining was graded as follows: no staining or staining observed in < 10% of tumor cells was given a score 0; faint/barely perceptible staining detected in 10% of tumor cells was scored as 1+; a moderate or strong staining observed in 10% of tumor cells was scored as 2+ or 3+, respectively. A score of 0 and 1+ was considered negative whereas 2+ and 3+ were considered positive. Immunostained slides were evaluated by two pathologists independently in a blinded manner. In most cases, the evaluations of the two pathologists were identical; any discrepancies were resolved by re-examination and consensus.

## Statistical analysis

The correlation between ER- $\alpha$ 36 expression and clinic pathologic characteristics was determined using Pearson's  $\chi^2$  test.  $P < 0.05$  was considered statistical significance. The statistical analyses were performed using SPSS 12.0 software.

## Results

### ER- $\alpha$ 36 is expressed in established gastric cancer cell lines

To determine whether the estrogen receptor variant, ER- $\alpha$ 36 is expressed in gastric cancer, we examined ER- $\alpha$ 36 expression in four gastric cancer cell lines including SGC-7901, AGS, BGC-923 and MKN-45 with Western blot analysis. Fig. 1 shows that higher levels of ER- $\alpha$ 36 protein were detected in all four gastric cancer cell lines compared to ER-positive breast cancer MCF-7 cells. In addition, RT-PCR analysis confirmed ER- $\alpha$ 36 mRNA was also highly expressed in all cell lines (Fig. 2). These results demonstrated that the novel variant of ER- $\alpha$ 36 is highly expressed in established gastric cancer cells.

Previously, we demonstrated that ER- $\alpha$ 36 is mainly expressed on the plasma membrane and in the cytoplasm of breast cancer cells (19). We decided to determine the localization of ER- $\alpha$ 36 in gastric cancer cells. Indirect immunofluorescence assay with a specific anti-ER- $\alpha$ 36

antibody revealed that ER- $\alpha$ 36 is highly expressed outside of the cell nuclei; mainly on the plasma membrane and in the cytoplasm of the gastric cancer cells (Fig. 3).

### ER- $\alpha$ 36 expression is detected in human gastric adenocarcinoma tissues

To determine the expression patterns of ER- $\alpha$ 36 in gastric adenocarcinoma tissues, the immunohistochemistry (IHC) assay was performed in specimens from 22 gastric cancer patients. The results showed that ER- $\alpha$ 36 is expressed in 18 of 22 (81.8%) specimens of primary gastric adenocarcinoma examined consistent with the results with established gastric cancer cell lines. We observed that ER- $\alpha$ 36 exhibited cytoplasmic and membranous patterns in tumor specimens (Fig. 4).

We also performed RT-PCR analysis to assess expression levels of ER- $\alpha$ 36 transcripts. ER- $\alpha$ 36 transcripts were detected in tumor tissues (T), adjacent non-malignant tissues (A) and corresponding normal tissues (N) of all 22 gastric cancer patients (data not shown). Higher-levels of ER- $\alpha$ 36 mRNA expression were detected in 17 of 22 (77.27%) tumor specimens compared with the paired normal tissues ( $p < 0.01$ ). Fig. 5 shows RT-PCR results from four representative cases. In comparison with corresponding normal tissues, levels of ER- $\alpha$ 36 mRNA expression were increased in tumor tissues of cases 1, 3 and 4 human gastric cancers (Fig. 5).

### ER- $\alpha$ 36 expression is correlated with lymph node metastasis

We also assessed whether ER- $\alpha$ 36 expression correlated with the clinicopathological properties of gastric adenocarcinoma. Table I shows that ER- $\alpha$ 36 expression was not associated with gender, age, tumor size, histological differentiation and tumor stage. However, ER- $\alpha$ 36 expression was significantly associated with lymph node metastasis ( $P = 0.0372$ ).

## Discussion

Previous studies have reported conflicting expression results for the classical ER- $\alpha$  (ER- $\alpha$ 66) in gastric cancer (20–25). Using the immunohistochemical method, gastric tumor tissues were often considered as ER-negative or low-expression tissues (2–28% of the tumors) (20,21). It was reported that in gastric cancer, ER- $\alpha$ 66 expression showed no difference between cancerous and normal tissues and no correlation with tumor grade (22). However, in other studies, it was found that ER- $\alpha$ 66 expression in gastric cancer correlated with a worse prognosis (23) and more widespread lymph node metastasis (24,25). Thus, the expression and possible function of ER- $\alpha$ 66 in development of human gastric cancer is controversial.

In this study, we found that ER- $\alpha$ 36 is highly expressed in established gastric cancer cell lines including SGC-7901, AGS, BGC-923, and MKN-45. Using indirect immunofluorescence staining, we demonstrated that ER- $\alpha$ 36 was mainly expressed on the plasma membrane and in the cytoplasm of gastric cancer cells. Both ER- $\alpha$ 36 mRNA and protein were also detected in most specimens from human gastric cancer patients; exhibiting higher levels of expression compared to their matched normal tissues. Our results thus for the first time indicated that the ER- $\alpha$  variant ER- $\alpha$ 36 is highly expressed in human gastric adenocarcinoma tissues and suggested that the non-genomic estrogen signaling mediated by ER- $\alpha$ 36 may be involved in gastric carcinogenesis.

We also found that ER- $\alpha$ 36 protein was mainly expressed on the plasma membrane and in the cytoplasm of gastric cancer specimens. It is worth noting that a weak cytoplasmic pattern of ER- $\alpha$ 66 expression was frequently observed in ER- $\alpha$ 36-expressing breast cancer tissues while mainly in the cell nuclei in ER- $\alpha$ 36-negative breast cancer cells (19), suggesting that

ER- $\alpha$ 36 expression may influence ER- $\alpha$ 66 cellular localization. Previously, it was reported that ER- $\alpha$ 66 expression was observed predominantly in the cytoplasm of gastric cancer cells (18). Thus, it is possible that these cells might also express ER- $\alpha$ 36.

We further found that high levels of ER- $\alpha$ 36 expression significantly correlated with lymph node metastasis in gastric cancer. Lymph node metastasis is one of the most important prognostic factors in gastric cancer. Thus, enhanced ER- $\alpha$ 36 expression in gastric adenocarcinoma may be involved in increased metastasis, suggesting that ER- $\alpha$ 36 expression can be used as a predictive marker for metastasis.

## Acknowledgments


This work was supported by National Natural Science Foundation of China grant 30870981 (L.J. Liu); Huibei Health Foundation grant NX200727 (L.J. Liu); Wuhan Young Chenguang Foundation grant 20045006071-7 (H. Deng); and Hubei Natural Science Foundation grant 2006AB191 (H. Deng). This work was also supported by NIH grant DK070016 (Z.Y. Wang).

## References

1. Nilsson S, Makela S, Treuter E, Tujague M, Thomsen J, Andersson G, Enmark E, Pettersson K, Warner M, Gustafsson JA. Mechanisms of estrogen action. *Physiol Rev.* 2001; 81:1535–1565. [PubMed: 11581496]
2. Greene GL, Gilna P, Waterfield M, Baker A, Hort Y, Shine J. Sequence and expression of human estrogen receptor complementary DNA. *Science.* 1986; 231:1150–1154. [PubMed: 3753802]
3. Greene GL, Press MF. Structure and dynamics of the estrogen receptor. *J Steroid Biochem.* 1986; 24:1–7. [PubMed: 3517497]
4. Kuiper GG, Enmark E, Peltö-Huikko M, Nilsson S, Gustafsson JA. Cloning of a novel receptor expressed in rat prostate and ovary. *Proc Natl Acad Sci USA.* 1996; 93:5925–5930. [PubMed: 8650195]
5. Zhang WH, Andersson S, Cheng G, Simpson ER, Warner M, Gustafsson JA. Update on estrogen signaling. *FEBS Lett.* 2003; 546:17–24. [PubMed: 12829231]
6. Kong EH, Pike AC, Hubbard RE. Structure and mechanism of the oestrogen receptor. *Biochem Soc Trans.* 2003; 31:56–59. [PubMed: 12546653]
7. Segars JH, Driggers PH. Estrogen action and cytoplasmic signaling cascades. Part: I membrane-associated signaling complexes. *Trends Endocrinol Metab.* 2002; 13:349–354. [PubMed: 12217492]
8. Wang ZY, Zhang XT, Shen P, Loggie BW, Chang YC, Deuel TF. Identification, cloning, and expression of human estrogenreceptor- $\alpha$ 36, a novel variant of human estrogen receptor- $\alpha$ 66. *Biochem Biophys Res Commun.* 2005; 336:1023–1027. [PubMed: 16165085]
9. Wang ZY, Zhang XT, Shen P, Loggie BW, Chang YC, Deuel TF. A variant of estrogen receptor- $\alpha$ , hER- $\alpha$ 36: transduction of estrogen- $\alpha$  and antiestrogen-dependent membrane-initiated mitogenic signaling. *Proc Natl Acad Sci USA.* 2006; 103:9063–9068. [PubMed: 16754886]
10. Zou Y, Ding L, Coleman M, Wang Z. Estrogen receptor-alpha (ER-alpha) suppresses expression of its variant ER-alpha36. *FEBS Lett.* 2009; 583:1368–1374. [PubMed: 19328202]
11. La Vecchia C, D'Avanzo B, Franceschi S, Negri E, Parazzini F, Decarli A. Menstrual and reproductive factors and gastric-cancer risk in women. *Int J Cancer.* 1994; 59:761–764. [PubMed: 7989115]
12. Brenner H, Rothenbacher D, Arndt V. Epidemiology of stomach cancer. *Methods Mol Biol.* 2009; 472:467–477. [PubMed: 19107449]
13. Palli D, Cipriani F, Decarli A, Galli M, Saieva C, Fraumeni JF Jr, Blot WJ, Buiatti E. Reproductive history and gastric cancer among post-menopausal women. *Int J Cancer.* 1994; 56:812–815. [PubMed: 8119771]
14. Tokunaga A, Nishi K, Matsukura N, Tanaka N, Onda M, Shiota A, Asano G, Hayashi K. Estrogen and progesterone receptors in gastric cancer. *Cancer.* 1986; 57:1376–1379. [PubMed: 3948119]



15. Matsui M, Kojima O, Uehara Y, Takahashi T. Characterization of estrogen receptor in human gastric cancer. *Cancer*. 1991; 68:305–308. [PubMed: 2070329]
16. Singh S, Poulson R, Wright NA, Sheppard MC, Langman MJ. Differential expression of oestrogen receptor and oestrogen inducible genes in gastric mucosa and cancer. *Gut*. 1997; 40:516–520. [PubMed: 9176081]
17. Furukawa H, Iwanaga T, Koyama H, Taniguchi H. Effect of sex hormones on carcinogenesis in the stomachs of rats. *Cancer Res*. 1982; 42:5181–5182. [PubMed: 7139622]
18. Nishi K, Tokunaga A, Shimizu Y, Yoshiyuki T, Wada M, Matsukura N, Tanaka N, Onda M, Asano G. Immunohistochemical study of intracellular estradiol in human gastric cancer. *Cancer*. 1987; 59:1328–1332. [PubMed: 3545434]
19. Lee LM-J, Cao J, Chen P, Gatalica Z, Wang Z-Y. ER- $\alpha$ 36, a novel variant of ER- $\alpha$ , is expressed in ER- $\alpha$  positive and negative human breast carcinomas. *Anticancer Res*. 2008; 28:479–484. [PubMed: 18383888]
20. Takano N, Iizuka N, Hazama S, Yoshino S, Tangoku A, Oka M. Expression of estrogen receptor-alpha and -beta mRNAs in human gastric cancer. *Cancer Lett*. 2002; 176:129–135. [PubMed: 11804739]
21. Wang M, Pan JY, Song GR, Chen HB, An LJ, Qu SX. Altered expression of estrogen receptor alpha and beta in advanced gastric adenocarcinoma: Correlation with prothymosin alpha and clinicopathological parameters. *Eur J Surg Oncol*. 2007; 133:195–201. [PubMed: 17046193]
22. Cui G, Yuan A, Qvigstad G, Waldum HL. Estrogens receptors are not expressed in esophagogastric carcinomas that come from a high incidence area of China. *Hepatogastroenterology*. 2001; 48:594–595. [PubMed: 11794139]
23. Harrison JD, Jones JA, Ellis IO, Morris DL. Oestrogen receptor D5 antibody is an independent negative prognostic factor in gastric cancer. *Br J Surg*. 1991; 78:334–336. [PubMed: 2021850]
24. Zhao XH, Gu SZ, Liu SX, Pan BR. Expression of estrogen receptor and estrogen receptor messenger RNA in gastric carcinoma tissues. *World J Gastroenterol*. 2003; 9:665–669. [PubMed: 12679906]
25. Matsui M, Kojima O, Kawakami S, Uehara Y, Takahashi T. The prognosis of patients with gastric cancer possessing sex hormone receptors. *Surg Today*. 1992; 22:421–425. [PubMed: 1421863]




**Figure 1.**


Western blot analysis of ER- $\alpha$ 36 protein expression in human gastric cancer cell lines.

Expression levels of ER- $\alpha$ 36 protein were assessed in four human gastric cancer cell lines, SGC7901, BGC823, MNN45 and AGS. ER-positive breast cancer MCF-7 cell line was used as a positive control. The levels of ER- $\alpha$ 36 and  $\beta$ -actin protein expression were determined with a video-densitometry system and the band densities relative to the density of  $\beta$ -actin band are shown.







**Figure 2.** RT-PCR analysis of ER- $\alpha$ 36 mRNA expression in human gastric cancer cell lines. Expression levels of ER- $\alpha$ 36 mRNA were examined in four human gastric cancer cell lines, SGC7901, BGC823, MKN45 and AGS. ER-positive breast cancer MCF-7 cell line was used as a positive control. The levels of mRNA expression of ER- $\alpha$ 36 and  $\beta$ -actin were determined with a video-densitometry system and the band densities relative to the density of  $\beta$ -actin band are shown.



**Figure 3.** Indirect immunofluorescence analysis of ER- $\alpha$ 36 expression in human gastric cancer cell lines. Merged images (original magnification,  $\times 400$ ) of SGC-7901 (A), MKN-45 (B), AGS (C) and BGC-823 (D) stained with an anti-ER- $\alpha$ 36 antibody (red) and with Hoechst 33258 (blue). Scale bar, 10  $\mu$ m.



**Figure 4.** Immunohistochemical staining of ER- $\alpha$ 36 in primary gastric adenocarcinoma, ER- $\alpha$ 36 expression in high-differentiation (A) and low-differentiation (B) gastric adenocarcinoma (magnification,  $\times 200$ ). Scale bar, 20  $\mu$ m.



**Figure 5.** RT-PCR analysis of ER-α36 mRNA level in specimens of gastric adenocarcinoma. ER-α36 expression in tumor tissues (T), adjacent non-malignant tissues (A) and corresponding normal tissues (N) was assessed with RT-PCR method.

**Table I**Relationship between expression level of ER- $\alpha$ 36 mRNA and clinicopathological characteristics.

<b>Factors</b>	<b>Low expression</b>	<b>High expression</b>	<b>P-value</b>
Age			0.3649
60 years	2	9	
>60 years	3	8	
Gender			0.391
Men	3	10	
Female	2	7	
Tumor size			0.3628
5 cm	2	5	
>5 cm	3	12	
Histological differentiation			0.4416
High differentiation	1	2	
High differentiation	4	15	
T stage			0.4519
T2-3	4	13	
T4	1	4	
N stage			0.0372
N0	4	4	
N1-3	1	13	