

# In silico prediction of exposure amino acid sequences of outer inflammatory protein A of *Helicobacter pylori* for surface display on *Escherichia coli*

Omid Teymounejad, Ashraf M. Mobarez, Zuhair M. Hassan<sup>1</sup>, Seyed M. Moazzeni<sup>1</sup>, Bagher Yakhchali<sup>2</sup>, Vajihe Eskandari<sup>2</sup>

Departments of Bacteriology and <sup>1</sup>Immunology, Tarbiat Modares University, Tehran, <sup>2</sup>Department of Industrial and Environmental Biotechnology, National Institute of Genetic Engineering and Biotechnology, Tehran, Iran

**BACKGROUND:** Outer inflammatory protein A (OipA) is an outer membrane protein of *Helicobacter pylori* that is involved in inducing IL-8 and intracellular signaling. In this study, we have predicted exposure amino acid sequences of OipA for insertion in permissive sites of CstH subunit of *Escherichia coli* CS3 pilli for bacterial surface display.

**MATERIALS AND METHODS:** Databases: National Center for Biotechnology Institute and Protein Data Bank. Servers: PHD, SABLE, GOR 4, SignalP3.0, TBBpred, PRODIV-TMHMM, TMRPres2D, CPH Models, PHYRE, GETAREA, VADAR, Pep state and pep window. Software: Swiss PDB viewer and Discovery studio.

**RESULTS:** In silico prediction of exposure amino acid sequences of OipA led to detection of six sequences of amino acid, 76-87, 106-112, 170-182, 222-230, 242-258, and 278-290. These sequences inserted between amino acid sequences 66-67, 100-101, and 109-110 of CstH that were predicted by Eskandari et al. as permissive sites of CstH.

**CONCLUSION:** OipA has the ability to induce IL-8 from gastric epithelial cells and some papers are mentioned that this outer membrane protein involve to attachment and intracellular signaling. Receptor of OipA and adhesion motifs on this protein is unknown. Detection of exposure motifs aids to recognition of adhesion motifs and receptor of OipA on gastric epithelial cells. In this study, we have predicted exposure amino acid sequences for insert to subunit CstH of CS3 pilli *E. coli* for surface display.

**Key words:** CstH, in silico prediction, OipA, surface sequences

## Introduction

*Helicobacter pylori* is a gram negative spiral bacterium that colonizes stomach for long time, and this colonization can lead to gastritis, peptic ulcer, and gastric cancer.<sup>[1]</sup> World Health Organization and International Agency for Research on Cancer have classified this bacterium as type one carcinogen for gastric cancer.<sup>[2,3]</sup> *H. pylori* colonizes half of the world population and this colonization reaches up to 90% in developing country.<sup>[4]</sup> Outer membrane proteins have pivotal role in pathogenesis of this bacterium.<sup>[5]</sup> Most of these outer membrane proteins increase attachment to gastric epithelial cell and some of them induce host inflammatory responses.<sup>[5]</sup> Outer inflammatory protein A (OipA) is an outer membrane protein that is involved to increase pathogenesis of *H. pylori*.<sup>[6]</sup> Presence of this protein is linked to proinflammatory signaling of gastric epithelial cell, duodenal ulceration, gastric cancer, increase of *H. pylori* density, and neutrophil infiltration.<sup>[6,7]</sup> *OipA* mutant of *H. pylori* revealed that this outer membrane protein associated with bacterial adhesion to gastric epithelial cell. Inducing IL-8 secretion by this protein is controversial; some reports showed that *oipA* mutants had not reduced IL-8 secretion but the others have reported about main role of OipA in induction of IL-8.<sup>[5,6,8]</sup>

The aim of this research is computational studies for prediction of OipA exposure sequences of amino acid. Results of this study apply for insert in permissive sites

### Access this article online

#### Quick Response Code:



#### Website:

www.ijhg.com

#### DOI:

10.4103/0971-6866.96659

**Address for correspondence:** Dr. Ashraf M. Mobarez, Department of Bacteriology, Faculty of medicine, Tarbiat Modares University. E-mail: mmmobarez@modares.ac.ir.

Dr. Zuhair M. Hassan, Department of Immunology, Faculty of medicine, Tarbiat Modares University. E-mail: hasan\_zm@modares.ac.ir.

of CstH subunit of *Escherichia coli* CS3 pilli for bacterial surface display of outer inflammatory protein A on *E. coli* for detection adhesion motif and role of OipA in induction of IL-8 and proinflammatory signaling.

**Materials and Methods**

Databases: Amino acid sequences of OipA and CstH were retrieved from National Center for Biotechnology Institute and Uni-prot KB/Swiss-Prot. Accession number of CstH in NCBI and Uni-prot KB/Swiss-Prot were X16944 and P15488.<sup>[9,10]</sup> Tertiary structure of CstH presents in Protein Data Bank.<sup>[11]</sup>

Servers: Secondary structure of proteins predicted by PHD, SABLE, and GOR 4 servers.<sup>[12,13]</sup> SignalP3.0 server detected signal sequence of OipA and CstH.<sup>[14]</sup> TBBpred, PRODIV-TMHMM, and TMRPres2D servers used for transmembrane topology prediction in bilayer lipid and external loops of OipA. CPH Models and PHYRE servers applied for finding of homology modeling and tertiary structure prediction.<sup>[15-18]</sup> GETAREA and VADAR servers applied for prediction of accessible surface area amino acids among sequences of external loops. Pep state and pep window used for probability of expression in inclusion bodies and hydropathy plot of two proteins.

Software: Swiss PDB viewer and Discovery studio applied for data analysis.<sup>[19,20]</sup>

**Results**

In silico prediction of exposure amino acid sequences of OipA led to detection of six sequences of amino acid; 76-87, 106-112, 170-182, 222-230, 242-258, and 278-290. These sequences inserted between amino acid sequences 66-67, 100-101 and 109-110 of CstH were predicted by Eskandari *et al.* as permissive sites of CstH.

**Discussion**

Protein engineering is a branch of bioinformatics that is useful for progress of biological research.<sup>[21]</sup> In this study, we used from some tools of protein engineering for process of outer inflammatory protein A and prediction of exposure amino acid sequences for detection of adhesion motifs and sequences involve to

proinflammatory signaling after bacterial surface display on *E. coli*.

Carrier protein for OipA on surface of *E. coli* is CstH subunit of CS3 pili that previously was analyzed by Eskandari *et al.* for permissive sites. Prediction for secondary, tertiary structure, and hydropathy plot showed three sites for insertion of our sequences.<sup>[21]</sup>

We applied three servers for prediction of secondary structure and compared these results with together [Figures 1- 3].

SignalP3.0 server predicted seventeen N-terminal amino acids of OipA which do not present in mature OipA on the cell surface and this sequence was deleted before further analysis by other tools [Figure 4].

Secondary structure of transmembrane proteins in prokaryotes has a little difference from eukaryotes. In outer membrane of prokaryotes, hydrophobic beta barrel integrate in bilayer lipid and famous tool for prediction of transmembrane topology is TBBpred server [Figure 5].<sup>[22]</sup>

Tertiary structure of protein causes that some

<pre> MKKALLTLTSLFSLWHAERNGFYLGFLNPLEGSIYQGGSIGKKALENTLNEAINNAKNSLFPEQNTKAI CchHHHHHHHHHHHeccccCceeeeeCCCCCCCCcchHHHHHHHHHHHHHHHHcCCCCHHHHHH RDAQNALNAVKDKSTKIANRFAGNGGSGGIFNELSLGYKFLGKGIIGFRHSLFFGYQLGGVSVPGSGL HHHHHHcCchHHHHHHHHcCCCCCCCCCceeeceeeEceeeEceEeeEeeccccccccCceE IAFLPYGFNTDLLINWNTDKRASQEVYVRRVKGSLIFYKDMTGRGLDANTLKKASRHIIRKSSGLVIGMD EEEECCCCceEeCCCCcHHHHHHHHcCCeEccccccccchHHHHcCCceeecccccehhhe IGASTWFASNNLTPFNQVKSHTIFQLQKGFVWRWSDGYDIDRYGDEIYLGSSVELGVKVPFAKVNYYG cchhhhhcCCCCCCCCCceEeecccccccccCchhhcCCCCceeeccCCEEEEceeeeeece DDYGDKLDYKRVVSVLYNLYNFKNKH CCCCCceEEEEEEEEHHcCC                     </pre>
--

**Figure 1: Secondary structure prediction of OipA by PHD server E: beta sheet, H: alpha helices, C: random coil**

<pre> MKKALLTLTSLFSLWHAERNGFYLGFLNPLEGSIYQGGSIGKKALENTLNEAINNAKNSLFPEQNTKAI ccceeehhhhhhhhhhcceecc RDAQNALNAVKDKSTKIANRFAGNGGSGGIFNELSLGYKFLGKGIIGFRHSLFFGYQLGGVSVPGSGL hhhhhhhhhhcchhhheeeccccccccceehhhcccccccccccccccccccccccccccccccccccc IAFLPYGFNTDLLINWNTDKRASQEVYVRRVKGSLIFYKDMTGRGLDANTLKKASRHIIRKSSGLVIGMD eeeecc IGASTWFASNNLTPFNQVKSHTIFQLQKGFVWRWSDGYDIDRYGDEIYLGSSVELGVKVPFAKVNYYG ccceeecc DDYGDKLDYKRVVSVLYNLYNFKNKH cc                     </pre>
--

**Figure 2: Secondary structure prediction of OipA by GOR4 server E: beta sheet, H: alpha helices, C: random coil**

<pre> MKKALLTLTSLFSLWHAERNGFYLGFLNPLEGSIYQGGSIGKKALENTLNEAINNAKNSLFPEQNTKAI CCHHHHHHHHHHHHHCCCCCECCCHH RDAQNALNAVKDKSTKIANRFAGNGGSGGIFNELSLGYKFLGKGIIGFRHSLFFGYQLGGVSVPGSGL HHHHHHCC IAFLPYGFNTDLLINWNTDKRASQEVYVRRVKGSLIFYKDMTGRGLDANTLKKASRHIIRKSSGLVIGMD ECCCCCCHHHHHHHCCCCCHHHCCCCCCCCCCCCCCCCCCCCCCCCCHHHCCCCCCCCCCCCCEEE IGASTWFASNNLTPFNQVKSHTIFQLQKGFVWRWSDGYDIDRYGDEIYLGSSVELGVKVPFAKVNYYG ECCCCCCCCCCCCCCCCCCCCCEEEEEECEEECCCCCCCCCCCCCCCCCCCCCEEEEEECCCCCCC DDYGDKLDYKRVVSVLYNLYNFKNKH CCCCCEEEEEEEEEEEEEEECCCC                     </pre>
---

**Figure 3: Secondary structure prediction of OipA by SABLE server E: beta sheet, H: alpha helices, C: random coil**

sequences of external loops get surface and the others get inside of protein.<sup>[22]</sup> Next step was tertiary structure

prediction of OipA and detection of exposure amino acid sequences in external loops [Figures 6-8].




Figure 4: SignalP3.0 server for prediction of signal sequence of OipA




Figure 5: Prediction of beta barrel of OipA in *H. pylori* outer membrane by TBBpred server "n" refers to residues in non barrel regions and "b" refers to residues in barrel region




Figure 6: Hydropathy plot of OipA by pepwindow server




Figure 7: Prediction of accessible surface area amino acids by vadar server




Figure 8: Tertiary structure prediction of OipA by CPH model server and shown with Discovery Studio Visualizer





Figure 9: Prediction of external and internal loops sequence of OipA by PRODIV-TMHMM server



**Figure 10: Prediction of external and internal loops sequence of OipA by TMRPres2D server**

The CPH model server revealed that tertiary structure of OipA, same as VacA, was most likely an auto display protein and its insertion in the outer membrane was carried out by type V secretion system (T5SS). N-terminal long hydrophilic region confirms this finding [Figures 6-10].

## Conclusion

Exposure amino acid sequences of OipA and other outer membrane proteins have important role in interaction with stomach epithelial cell, and detection of these sequences is very useful for finding of receptors on surface of Gastric epithelial cell.<sup>[5]</sup>

Receptor of OipA and adhesion motifs on this protein is unknown. Detection of exposure motifs aids to recognition of adhesion motifs and receptor of OipA on gastric epithelial cells.<sup>[23]</sup> In this study, we have predicted exposure amino acid sequences for insert to subunit CstH of CS3 pilli *E. coli* for surface display.

## References

1. Wei J, O'Brien D, Vilgelm A, Piazuolo MB, Correa P, Washington MK, et al. Interaction of *Helicobacter pylori* with gastric epithelial cells is mediated by the p53 protein family. *Gastroenterology* 2008;134:1412-23.
2. Zhang H, Fang DC, Wang RQ, Yang SM, Liu HF, Luo YH. Effect of *Helicobacter pylori* infection on expression of Bcl-2 family members in gastric adenocarcinoma. *World J Gastroenterol* 2004;10:227-30.
3. Franco AT, Johnston E, Krishna U, Yamaoka Y, Israel DA, Nagy TA, et al. Regulation of gastric carcinogenesis by *Helicobacter pylori* virulence factors. *Cancer Res* 2008;68:379-87.
4. Zhang ZW, Farthing MJ. Molecular mechanisms of *H. pylori* associated gastric carcinogenesis. *World J*

5. Gastroenterol 1999;5:369-74.
6. Dossumbekova A, Prinz C, Mages J, Lang R, Kusters JG, Van Vliet AH, et al. *Helicobacter pylori* HopH (OipA) and bacterial pathogenicity: Genetic and functional genomic analysis of hopH gene polymorphisms. *J Infect Dis* 2006;194:1346-55.
7. Clyne M, Dolan B, Reeves EP. Bacterial factors that mediate colonization of the stomach and virulence of *Helicobacter pylori*. *FEMS Microbiol Lett* 2007;268:135-43.
8. Yamaoka Y, Kwon DH, Graham DY. A M(r) 34,000 proinflammatory outer membrane protein (oipA) of *Helicobacter pylori*. *Proc Natl Acad Sci U S A* 2000;97:7533-8.
9. Shao SH, Wang H, Chai SG, Liu LM. Research progress on *Helicobacter pylori* outer membrane protein. *World J Gastroenterol* 2005;11:3011-3.
10. Bairoch A. The Universal Protein Resource (UniProt) 2009. *Nucleic Acids Res* 2009;37:D169-74.
11. Jalajakumari MB, Thomas CJ, Halter R, Manning PA. Genes for biosynthesis and assembly of CS3 pili of CFA/II enterotoxigenic *Escherichia coli*: Novel regulation of pilus production by bypassing an amber codon. *Mol Microbiol* 1989;3:1685-95.
12. Berman HM, Westbrook J, Feng Z, Gilliland G, Bhat TN, Weissig H, et al. The protein data bank. *Nucleic Acids Res* 2000;28:235-42.
13. Rost B, Sander C, Schneider R. PHD-An automatic mail server for protein secondary structure prediction. *Comput Appl Biosci* 1994;10:53-60.
14. Adamczak R, Porollo A, Meller J. The SABLE. Server, which was rigorously. *Bioinformatics*, 2004.
15. Bendtsen JD, Nielsen H, von Heijne G, Brunak S. Improved prediction of signal peptides: Signal P 3.0. *J Mol Biol* 2004;340:783-95.
16. Bennett-Lovsey RM, Herbert AD, Sternberg MJ, Kelley LA. Exploring the extremes of sequence/structure space with ensemble fold recognition in the program Phyre. *Proteins* 2008;70:611-25.
17. Krieger E, Nabuurs SB, Vriend G. Homology modeling. *Methods Biochem Anal* 2003;44:509-23.
18. Sujatha MS, Balaji PV. Fold-recognition and comparative modeling of human alpha2,3-sialyltransferases reveal their sequence and structural similarities to CstII from *Campylobacter jejuni*. *BMC Struct Biol* 2006;6:9.
19. Hardesty J. Protein Simulation, Function and Prediction. *Computational Structure Biology*, 2005.
20. Kaplan W, Littlejohn TG. Swiss-PDB Viewer (Deep View). *Brief Bioinform* 2001;2:195-7.
21. Niemeyer EA. Application Guide tools and methods used in Discovery Studio for the visualization. *Nucleic Acids Res* 2000;28:235-42.
22. Eskandari V, Yakhchali B, Minuchehr Z. In-silico prediction of permissive sites of CstH subunit of *E. coli* CS3 pili for insertion of foreign peptides. *Iran J Biol* 2010;1:73-84.
23. Xiong J. Protein Secondary Structure Prediction. In: Xiong J, editor. *Essential Bioinformatics*. 1<sup>st</sup> ed. New York: Cambridge University Press; 2006. p. 200-13.
24. Yamaoka Y, Alm RA. *Helicobacter pylori* outer membrane proteins. In: Yamaoka Y, editor. *Helicobacter pylori* Molecular Genetics and Cellular Biology. 1<sup>st</sup> ed. Norfolk: Caister Academic Press; 2008. p. 37-61.

**Cite this article as:** Teymournejad O, Mobarez AM, Hassan ZM, Moazzeni SM, Yakhchali B, Eskandari V. In silico prediction of exposure amino acid sequences of outer inflammatory protein A of *Helicobacter pylori* for surface display on *Escherichia coli*. *Indian J Hum Genet* 2012;18:83-6.

**Source of Support:** Nil, **Conflict of Interest:** None declared.