BclI RFLP for the human vimentin gene

E.M.Marcus, B.A.Smith, H.Telenius, R.M.Landsvater¹, C.H.C.M.Buys¹, S.Ferrari², B.A.J.Ponder and C.G.P.Mathew

Section of Human Cancer Genetics, Institute of Cancer Research, Haddow Laboratories, Sutton, Surrey SM2 5NG, UK, ¹Department of Human Genetics, State University of Groningen, Groningen, The Netherlands and ²Department of Pathology, Temple University Medical School, Philadelphia, PA 19140, USA

Source/Description : A 1.1 kb cDNA clone (hp4F1) encoding the human vimentin gene identified in a human library by screening with 4F1, a hamster vimentin cDNA (Ferrari et al, 1986).

Polymorphism : BclI (TGATCA) recognises a two allele
polymorphism :
 Bands Al : 8.1 kb
 A2 : 3.6 kb
 There is a faint constant band at 15.8 kb.

Frequency : in 47 unrelated Gaucesian individuals: Al : 0.70 A2 : 0.30

Not polymorphic for : AccI, AvaI, AvaII, BamHI, BanII, BglI, BstNI, BstXI, ClaI, EcoRI, HincII, HindIII, StuI.

<u>Chromosome location</u>: Mapped to chromosome 10pter-10q23 using somatic cell hybrids and to 10p13 by <u>in</u> <u>situ</u> hybridisation (Ferrari et al, 1987).

Mendelian inheritance : Codominant segregation was observed in 2 informative families.

Probe availability : Contact R. Baserga.

Acknowledgements : Supported by a programme grant to the Institute of Cancer Research by the Cancer Research Campaign. H.T. is supported by grants from several Swedish foundations. References :

Ferrari, S. et al; Mol Cell Biol, 6: 3614-3620 (1986). Ferrari, S. et al; Amer J Hum Genet, 41: 616-626 (1987).