

# CORRECTIONS

Vol. 159: 479–488, 2012

Lee S.H., Chung G.C., Jang J.Y., Ahn S.J., and Zwiazek J.J. Overexpression of PIP2;5 Aquaporin Alleviates Effects of Low Root Temperature on Cell Hydraulic Conductivity and Growth in Arabidopsis.

An error during the production process resulted in Table I in this article being replaced with a duplicate version of Table II. Table I appears below and has been added to the online version of this article.

**Table I.** Comparison of cell dimensions and water relations parameters in Arabidopsis seedlings subjected to 23°C root temperature

Water relations parameters, including P,  $\epsilon$ , and  $T_{1/2}$  of root cortical cells, were measured in Arabidopsis at a distance of about 25 to 30 mm from the root tip using the cell pressure probe. Different letters in each row for wild-type and transgenic plants indicate significant differences (unpaired *t* test, *P* = 0.05). Values are means  $\pm$  SE (*n* = 15–20 cells from 15–20 plants).

Plants	Cell Dimensions		Water Relations Parameters			
	Length	Diameter	P	$\epsilon$	$T_{1/2}$	$L_p \times 10^7$
	$\mu m$			MPa	s	$m s^{-1} MPa^{-1}$
Wild type	200 $\pm$ 3.6 <sup>a</sup>	32 $\pm$ 1.3 <sup>a</sup>	0.35 $\pm$ 0.03 <sup>a</sup>	3.9 $\pm$ 0.5 <sup>a</sup>	1.7 $\pm$ 0.1 <sup>a</sup>	7.2 $\pm$ 0.4 <sup>a</sup>
PIP1;4-1	200 $\pm$ 5.9 <sup>a</sup>	32 $\pm$ 1.7 <sup>a</sup>	0.34 $\pm$ 0.04 <sup>a</sup>	3.6 $\pm$ 0.7 <sup>a</sup>	1.8 $\pm$ 0.1 <sup>a</sup>	7.6 $\pm$ 0.4 <sup>a</sup>
PIP1;4-3	203 $\pm$ 6.5 <sup>a</sup>	32 $\pm$ 1.4 <sup>a</sup>	0.37 $\pm$ 0.04 <sup>a</sup>	3.5 $\pm$ 1.0 <sup>a</sup>	2.0 $\pm$ 0.4 <sup>a</sup>	6.2 $\pm$ 1.6 <sup>a</sup>
PIP2;5-1	198 $\pm$ 7.3 <sup>a</sup>	32 $\pm$ 1.1 <sup>a</sup>	0.35 $\pm$ 0.04 <sup>a</sup>	3.3 $\pm$ 0.7 <sup>a</sup>	1.7 $\pm$ 0.2 <sup>a</sup>	8.4 $\pm$ 0.9 <sup>a</sup>
PIP2;5-3	204 $\pm$ 7.7 <sup>a</sup>	32 $\pm$ 1.5 <sup>a</sup>	0.37 $\pm$ 0.03 <sup>a</sup>	4.1 $\pm$ 1.2 <sup>a</sup>	1.8 $\pm$ 0.3 <sup>a</sup>	6.6 $\pm$ 1.1 <sup>a</sup>