

Published in final edited form as:

Angew Chem Int Ed Engl. 2012 January 23; 51(4): 862–872. doi:10.1002/anie.201104807.

Transition-Metal-Catalyzed Denitrogenative Transannulation: Converting Triazoles into Other Heterocyclic Systems

Buddhadeb Chattopadhyay and **Vladimir Gevorgyan**

Department of Chemistry, University of Illinois at Chicago, 845 W Taylor Street, Rm 4500, Chicago, IL 60607 (USA)

Vladimir Gevorgyan: vlad@uic.edu

Abstract

Transition metal catalyzed denitrogenative transannulation of a triazole ring has recently received considerable attention as a new concept for the construction of diverse nitrogen-containing heterocyclic cores. This method allows a single-step synthesis of complex nitrogen heterocycles from easily available and cheap triazole precursors. In this Minireview, recent progress of the transition metal catalyzed denitrogenative transannulation of a triazole ring, which was discovered in 2007, is discussed.

Keywords

cyclization; nitrogen heterocycles; transition metals; transannulation; synthetic methods

1. Introduction

1,2,3-Triazoles are important heterocyclic units endowed with a broad spectrum of biological activities.^[1–5] They have been extensively used in medicinal chemistry,^[6, 7] biochemistry,^[8, 9] and in material science.^[10] According to the existing paradigm, the 1,2,3-triazole ring is a very robust heterocyclic unit, thus not surprisingly, its chemistry mostly involves functionalization of the core. However, recently, new chemistry involving ring opening of the 1,2,3-triazole ring in the presence of various transition-metal catalysts has been reported. Most importantly a new direction, a denitrogenative transannulation of triazoles into other N-containing heterocycles, has recently appeared. Although, several methods exist for the construction of various N-containing heterocycles, there is always a need for new, efficient, and general methods for the synthesis of these important classes of compounds. The denitrogenative transannulation approach offers obvious advantages over many existing methods, as it allows efficient, single-step interconversion of easily available 1,2,3-triazoles into a variety of other valuable N-containing heterocyclic systems.

This Minireview covers the transition metal catalyzed denitrogenative transannulation of 1,2,3-triazoles into highly functionalized five- and six-membered-ring heterocycles, as well as fused nitrogen-containing heterocycles, in a single step. The organization of this Minireview is based on the denitrogenative transannulation reaction of different types of triazoles with alkynes, nitriles, alkenes, allenes, and isocyanides. Both, the synthetic applications, as well as the mechanistic aspects of the described transannulation reactions are discussed.

2. Transannulation of Pyridotriazoles

In solution, the pyridotriazoles **1**^[11] exist in a closed/opened form equilibrium^[12] with the diazocompounds **2**^[13] (Scheme 1). Thus, not surprisingly, the former is sometimes capable of undergoing transformations that are characteristic for diazocompounds.^[14] It deserves mentioning that the position of this equilibrium depends upon several factors, such as temperature, solvent, and the nature of the substituent (R^1) at C7^[12b] of the triazole ring. It has been reported that the introduction of a halogen atom at C7 ($R^1 = Cl$) shifts the equilibrium to the right, which has been explained in terms of nonbonding repulsion between the lone pair of electrons on the halogen and nitrogen atom in the peri-position of **1** (Scheme 1).^[15]

Recently, Gevorgyan and co-workers demonstrated^[16] that the diazoform **2a,b** may be used as a source of a rhodium carbenoid species (Scheme 2). Thus, it was shown that the 7-halo-substituted pyridotriazole **3b**, in the presence of a rhodium catalyst, released dinitrogen from **2b** to produce the corresponding rhodium carbenoid **4b**. It was confirmed by its insertion into the Si–H bond of triethylsilane, a method developed by Doyle et al. for trapping the rhodium-stabilized carbenes originating from Rh^{II} complexes and diazoacetates.^[17] As expected, the pyridotriazoles **3a** and **3b** exhibited different reactivity towards triethylsilane under these reaction conditions. Thus, while **3a** remained unreactive, **3b** was smoothly converted into **5b**, the product of the rhodium carbenoid insertion into the Si–H bond. Thus, it became evident the 7-halo-substituted pyridotriazole **3b** can indeed serve as a convenient precursor for rhodium carbenoids (Scheme 2).

2.1. Transannulation with Alkynes and Nitriles

After revealing that the pyridotriazole **3b**, the surrogate of α -imino diazocompound **2b**, can be used as convenient precursors for the rhodium carbenoid **4b**, the reactivity of the latter in cycloaddition reactions with alkynes was explored (Scheme 3).^[16] It was found that treatment of the pyridotriazole **3b** with phenyl acetylene in the presence of $Rh_2(OAc)_4$, resulted in a mixture of the cyclopropene **6a** and indolizine **7a**, the products of the [2+1] and the formal [2+3] cycloaddition reactions, respectively. Interestingly, the cyclopropene **6a**, under the employed reaction conditions, did not undergo cycloisomerization into indolizine **7a**, thereby suggesting independent mechanistic paths for their formation. Selectivity of the transannulation reaction has been dramatically improved by employing a $[Rh_2(pfb)_4]$ catalyst to give **7a** as the sole reaction product in 78% yield. Under these reaction conditions, the 1-carbomethoxy-substituted pyridotriazole **3b** underwent smooth transannulation with terminal aryl and alkenyl alkynes to produce indolizines **7** in good to excellent yields (Scheme 3).

Next, the Gevorgyan group explored the possibility of a transannulation reaction of **3**, having various substitutions, with nitriles en route to N-fused imidazoles (Scheme 4). It was found that the pyridotriazoles **3** reacted smoothly with a variety of aryl, alkyl, and alkenyl nitriles **8** in the presence of $Rh_2(OAc)_4$ to afford the N-fused imidazopyridines **9** in good to high yields (Scheme 4). Importantly, 3-carbomethoxy-, 3-aryl-, as well as 7-bromo-, and 7-methoxy-substituted pyridotriazoles, proved to be equally efficient in this reaction.

It was proposed that this transannulation proceeds through the in situ generated rhodium carbenoid intermediate **10** (Scheme 5). A direct nucleophilic attack^[18] of the alkyne or nitrile at **10** produces the intermediate ylide **11** (path a, Scheme 5), which then cyclizes to form **7** or **9** via the cyclic zwitterion **12**. Alternatively (path b), [2+2] cycloaddition of the rhodium carbenoid **10** with an alkyne or nitrile produces the metallacyclobutene **13**, which can also arise from cyclization of **11**.^[19] Rhodacycle **13** then undergoes σ -bond metathesis

to produce the rhodium carbenoid **14**, which upon 6π electrocycloization and subsequent reductive elimination of rhodium furnished either the product **7** or **9**. The potential [2+1]/cycloisomerization sequence via **16** (path c) was ruled out since the cyclopropene **6a** did not cycloisomerize into **7a** under the reaction conditions (see Scheme 3).

3. Transannulation of N-Sulfonyl-1,2,3-triazoles

3.1. Transannulation with Alkynes, Nitriles, and Alkenes

N-Sulfonyl-1,2,3-triazole is an important heterocyclic unit that can easily be synthesized by the copper-catalyzed azide–alkyne cycloaddition reaction.^[20] This triazole is exceedingly resistant to thermal degradation and stays intact under harsh hydrolytic, reductive, and oxidative conditions.^[21, 22] In 2008, Gevorgyan, Fokin, and co-workers challenged the robustness of this heterocyclic unit. They demonstrated that N-sulfonyl-1,2,3-triazoles **17a** in the presence of 1 mol% $\text{Rh}_2(\text{OAc})_4$ smoothly reacts with styrene to quantitatively produce the *trans*-cyclopropane^[23] carboxaldehyde **18** after silica gel chromatography (Scheme 6).^[24] Apparently, the N-sulfonyl-1,2,3-triazole **17a** served as a surrogate for the diazoimine species **19**, which in turn was converted into the corresponding metal carbenoid **20**. A subsequent [2+1] cycloaddition of **20** with styrene, and hydrolysis of the formed iminocyclopropane **21** furnished the reaction product **18** (Scheme 6).

Inspired by this finding, they next attempted a transannulation reaction of the triazole **17a** with benzonitrile by employing two different protocols, namely, microwave-assisted and conventional heating. It was found that both methods were equally efficient in providing the transannulation product, imidazole **22a** in high yields (Scheme 7).

The developed methods have been applied to transannulation of differently C4-substituted N-sulfonyl-1,2,3-triazoles **17** with a number of nitriles (Scheme 8). The reactions appeared to be very general with respect to the triazole and nitrile components. Both the microwave and the conventional heating methods afforded high to excellent yields of the diversely substituted transannulation products **22** (Scheme 8).^[24] It deserves mentioning that in contrast to pyridotriazoles (see above), the triazoles **17** under these reaction conditions did not undergo transannulation reaction with terminal alkynes into pyrroles.

The proposed mechanistic rationale is related to the analogous transannulation of nitriles with diazoketones as reported by Helquist, Akermark, and co-workers.^[25] According to the path a in Scheme 9, a nucleophilic attack of the nitrile on the rhodium carbenoid **20**^[26] leads to the ylide **23**, which upon cyclization into the zwitterion **24** and subsequent loss of the metal, furnishes the imidazole **22**. Alternatively, ylide **23** may give rise to the rhodium carbenoid **25** through a [1,3] Rh shift. A subsequent cyclization of the intermediate **25** and then reductive elimination^[27] produces **22**. Also, a possible direct formation of **22** through a [3+2] cycloaddition of **20** with a nitrile was not ruled out (path b).

In 2009, Murakami and co-workers reported a nickel-catalyzed denitrogenative transannulation reaction of N-sulfonyl-1,2,3-triazoles with internal alkynes.^[28] They discovered that a combination of a $[\text{Ni}(\text{cod})_2]$ catalyst with the electron-rich and bulky phosphine ligand $\text{P}(n\text{Bu})\text{Ad}_2$, and AlPh_3 as a Lewis acid additive, was efficient for the transannulation of triazoles **17** with internal alkynes into tetrasubstituted pyrroles **26** (Scheme 10). It was found that the yields of the transannulation reaction with symmetrical alkynes were generally good, except for an *n*-hexyl-substituted triazole. Transannulation with unsymmetrical alkynes produced nearly equal amounts of regioisomers (last three structures in Scheme 10), whereas attempts on the employment of terminal alkynes were unsuccessful, presumably because of a facile self-oligimerization side process.^[28]

Mechanistically, it is believed that this reaction starts from a ring–chain tautomerization of the triazole **17a** into diazoimine **19** (Scheme 11), which is captured by nickel to give the nickel carbenoid **27**. The latter cyclizes into the azanickelacycle **28**. Subsequent insertion of the alkyne into the Ni–C bond leads to the corresponding six-membered nickelacycle **29**, which upon reductive elimination of the Ni⁰, furnishes the pyrrole **26**. It was hypothesized that the possible role of the Lewis acid in this transformation may involve a promotion of the ring–chain tautomerization of **17a** into **19**, or an acceleration of the reductive elimination of nickel^[29] from **29**.

Very recently, Gevorgyan et al. partially solved the problem of transannulation of monocyclic triazoles with terminal alkynes (see below) into pyrroles.^[30] It was reported, that employment of the [Rh₂(oct)₄]/AgOCOCF₃ binary catalyst system enables efficient transannulation of the N-sulfonyl-1,2,3-triazoles **17b** with arylalkynes to afford the corresponding transannulation products **30** in good to excellent yields (Scheme 12). Electron-rich alkynes were more efficient in this reaction than their electron-neutral counterparts, whereas electron-deficient arylalkynes did not undergo this transformation at all.

The following plausible mechanism for this transannulation reaction has been proposed (Scheme 13). Upon treatment with [Rh₂(oct)₄], the triazole **17b** transforms into the rhodium iminocarbene **20b**.^[24] A direct nucleophilic attack of the terminal alkyne at the latter produces ylide **31** (path a, Scheme 13),^[4, 24] which upon cyclization forms a cyclic zwitterionic species **32**. Elimination of the rhodium catalyst from **32** produces the reaction product **30**. In contrast, the in situ generated silver acetylide may attack **20b** to form the rhodium-containing propargylimine species **33** (path b). Alternatively, **33** may arise through a proton loss from **31** (path b'). Proton-assisted 5-*endo-dig* cyclization of **33** would afford cyclic intermediate **32**. However, a deuterium labeling experiment employing the deuterated alkyne resulted in formation of [D]-**30** with complete preservation of a deuterium label at C3, thus undoubtedly ruling out the possible involvement of the paths b and b', both of which would result in partial or complete deuterium scrambling. Although, the crucial role of silver trifluoroacetate in this transformation is not completely understood, this Lewis acid probably activates the electrophilic rhodium carbene moiety, through coordination to the imine, toward the nucleophilic attack by an alkyne. The higher reactivity of electron-rich alkynes in this transformation reasonably fits into the most plausible reaction being the ylide reaction (path a).

The synthetic usefulness of this transannulation reaction was showcased by an efficient three-component semi-one-pot synthesis of the pyrrole **35** from tosylazide **34** and two different terminal alkynes by a combined copper-catalyzed click/rhodium-catalyzed transannulation reaction sequence (Scheme 14).

Very recently, Fokin and co-workers showed^[31] that highly reactive Rh^{II} N-triflyl azavinyl carbenes can easily be produced from the MH-1,2,3-triazoles **36** by treatment with triflic anhydride in the presence of Rh^{II} complexes. These carbene intermediates efficiently engage olefins in highly enantio- and diastereoselective transformations, thus providing easy access to homochiral cyclopropane carboxaldehydes **37** and 2,3-dihydropyrroles **38** (Scheme 15). Although, the transannulation products were formed in high yield, the enantioselectivity of the transannulation products varied depending on the nature of the substituent (R¹) at C4 of the triazole ring.

4. Transannulation of N-Aroylbenzotriazoles

4.1. Transannulation with Alkynes

Nakamura et al. developed an interesting palladium-catalyzed transannulation reaction of N-arylbentotriazoles **39** with alkynes to give indoles **40** (Scheme 16).^[32] The authors took advantage of the closed/opened form equilibrium between the acyltriazole **39** and its diazonium isomer **41**, which serves as an equivalent of the haloanilide **42** that is employed in indole synthesis reported by Larock et al.^[33] From an environmental standpoint, the base-free conditions and benign by-product (N₂) of this transannulation reaction are the obvious advantages of this method over Larock's classical indole synthesis, which produces stoichiometric amounts of HX●base waste.

The reaction conditions for this transannulation require heating **39a** and the internal alkynes **43** in the presence of [Pd(PPh₃)₄] at 130°C without a solvent (Scheme 17). These reaction conditions allow the synthesis of the multisubstituted indoles **40a** in good yields. Performing the reaction using solvents, as well as employment of other palladium catalysts, were less efficient. The electronic nature of the substituents showed a pronounced effect on the efficiency of this reaction. Whereas triazoles possessing electron-withdrawing groups reacted well, those substituted with electron-donating groups reacted sluggishly, thus providing poor yields or no reaction even under prolonged reaction times. Reactions with unsymmetrical alkynes showed varied regioselectivity, thus favoring bulkier substituents (R⁶) at C2 of the indole; this trend is analogous to that observed in the Larock's indole synthesis. Expectedly, this Pd⁰-catalyzed method did not tolerate terminal alkynes (Scheme 17).

Mechanistically, this palladium-catalyzed transannulation reaction is quite similar to that of the nickel-catalyzed transannulation described above. First, Pd⁰ oxidatively inserts into the C–N bond of the diazonium moiety of the 2-iminobenzediazonium species **41a**, which is thermally generated from the benzotriazole **39a**.^[34] Insertion of the alkyne **43** into the Pd–C bond of the resulting intermediate **44** or **45** leads to the formation of the palladacycle **46**, which upon the reductive elimination yields indole **40a** and regenerates the Pd⁰ catalyst (Scheme 18).

5. Transannulation of 1,2,3-Benzotriazinones

5.1. Transannulation with Alkynes, allenes, and Alkenes

Murakami and co-workers found that the 1,2,3-benzotriazinones **47** are also good substrates for denitrogenative transannulation reactions. Thus, **47** in the presence of a nickel catalyst undergoes a facile reaction with alkynes to produce isoquinolones **48** (Scheme 19).^[35] The authors proposed that the reaction is initiated by the insertion of Ni⁰ into the N–N linkage of **47**, which upon loss of dinitrogen produces the azanickelacycle **49**.^[36] Insertion of the alkyne into the Ni–C bond leads to the formation of a seven-membered nickelacycle intermediate **50**,^[37] which after reductive elimination affords the final product **48** and regenerates the Ni⁰ catalyst (Scheme 19).

This reaction appeared to be very general in scope, as various symmetrical and unsymmetrical internal alkynes, as well as terminal alkynes, gave very high yields of the isoquinolones **48'** (Scheme 20). Unsymmetrical alkynes however, showed varied regioselectivity. Remarkably, terminal alkynes provided both excellent yields and regioselectivities in this reaction. It was found that the 1,2,3-benzotriazinones **47** possessing either electron-withdrawing or electron-donating aryl substituents at the nitrogen atom underwent smooth transannulation reactions at room temperature, whereas the reaction of

benzyl- and methyl-substituted substrates required higher temperatures. N-unsubstituted benzotriazinone failed to undergo this reaction.

Murakami et al. have also developed the nickel-catalyzed denitrogenative transannulation of 1,2,3-benzotriazinones with allenes.^[38] First, from the reaction of **47a** with stoichiometric amounts of $[\text{Ni}(\text{cod})_2]$ and dppbenz, the authors succeeded in isolating a five-membered azanickelacycle intermediate **49a**, the structure of which was confirmed by the single-crystal X-ray analysis (Scheme 21). Treatment of **49a** with an allene at 60°C in THF gave an isomeric mixture of the 3,4-dihydroisoquinolin-1(2*H*)-ones **51a** and **52a** (54:46) in 99% yield.

A catalytic version of this reaction (5 mol% $[\text{Ni}(\text{cod})_2]$ 20 mol% PMe_3 , THF, 60°C) was then applied for transannulation of different 1,2,3-benzotriazinones **47b** with a number of monosubstituted allenes (Scheme 22). Both the electron-withdrawing and electron-donating substituents at the N atom of the triazole moiety and at the aromatic ring of the benzotriazinone worked well, thus producing the differently substituted isoquinolones **51b** as a major regioisomer.^[39] Probably as a result of sterics, the regiochemistry was completely reversed in the reaction with *tert*-butyl- and trialkylsilyl-substituted allenes (Scheme 22).

Employment of the cyclic 1,3-disubstituted allene **53** resulted in an interesting outcome; the nature of the product varied depending upon the type of the phosphine ligand employed (Scheme 23). The use of PMe_3 in THF at 60°C produced the imino ester **54** in 75% yield, whereas employment of the bidentate phosphine ligand (*R,R*)-Me-duphos in toluene at 100°C afforded **55** as the sole product in 99% yield. The control experiment revealed that **54** in the presence of $[\text{Ni}(\text{cod})_2]$ and (*R,R*)-Me-duphos in toluene at 100°C was completely isomerized into **55**, thus confirming thermodynamic control in the formation of the latter in the reaction of **47a** and **53**.

The authors have also explored the asymmetric version of this transformation. It was shown that employment of bidentate phosphine ligands, such as (*R,R*)-Me-duphos and (*S,S,R,R*)-tangphos provided good enantioselectivities. Importantly, both the regio- and enantioselectivities were very high when the phosphinoxazoline ligand (*S,S*)-*i*Pr-foxap^[40] was employed (Scheme 24).^[38]

As an extension of this methodology, the same group has also developed the nickel-catalyzed transannulation of benzotriazinones with 1,3-dienes and activated alkenes.^[41] Interestingly, when the complex **49a** was mixed with the 1,3-diene **56** in the absence of a phosphine ligand, the formation of only trace amounts of **57** was detected (Scheme 25). However, addition of the dppf ligand provided **57** in 40% (Scheme 25).

Next, the generality of this approach was tested using a catalytic version of this reaction. Thus, employment of $[\text{Ni}(\text{cod})_2]$ (10 mol%) and dppf (10 mol%) in THF at 60°C allowed a facile reaction of differently substituted benzotriazinones **47b** with symmetrical 1,3-dienes **58** to form various N-protected isoquinolones **59** (Scheme 26). Except for the N-benzyl-substituted benzotriazinone (24%), the yields employing all other substrates were high. Benzotriazinones with both electron-donating and electron-withdrawing groups at the benzene ring were equally competent in this reaction. Employment of unsymmetrical dienes **58** was nearly as efficient in providing isoquinolones **59** as major regioisomers over **60**. For this transformation, the authors proposed a mechanism similar to that proposed for the nickel-catalyzed transannulation of benzotriazinones with allenes.^[38]

It was also shown that in the presence of $[\text{Ni}(\text{cod})_2]$ and $\text{P}(n\text{Bu})_3$, the benzotriazinones **47b** can undergo an efficient transannulation reaction with activated alkenes (Scheme 27). Thus,

electron-deficient alkenes, such as methyl acrylate, acrylonitrile, and acrylamide smoothly underwent transannulation with **47b** to give the dehydroisoquinolinones **61** in excellent yields. Pyridyl-containing alkenes were similarly efficient, whereas styrene gave low yield of the product. Electron-neutral and electron-rich alkenes did not participate in this reaction at all. (Scheme 27).^[41]

5.2. Transannulation with Isocyanides

The same research group has also shown^[42] that isocyanides can also be employed in this transannulation reaction. Thus, 1,2,3-benzotriazinone **47b** and benzothiazine dioxide **47c**, in the presence of a palladium catalyst and a phosphine ligand underwent smooth transannulation with isocyanides **62** to give the corresponding isocyanide incorporated products **63** in excellent yields (Scheme 28). Except for the N-alkyl-substituted triazinones, all other substrates tested exhibited excellent reactivity, thus giving rise to almost quantitative yields of the products. The reaction is also quite general with respect to the isocyanides **62**, as aryl, benzyl, cyclohexyl, and even aliphatic isocyanides were competent in this reaction for producing high yields of the transannulation products.

6. Transannulation of 1,2,3,4-Benzothiazinones

6.1. Transannulation with Allenes

In 2010, Murakami and co-workers reported^[43] the nickel-catalyzed enantioselective transannulation reaction of 1,2,3,4-benzothiazine-1,2(*2H*)-dioxide **64** with monosubstituted allenenes to produce 1,2,3,4-benzothiazine-1,1(*2H*)-dioxide derivatives **65** and **66**. It was proposed that this reaction is initiated by an oxidative addition of Ni⁰ into the N–N bond and subsequent elimination of the dinitrogen molecule to yield the five-membered intermediate **67**. A subsequent allene insertion into **67** generates the π -allylnickel intermediate **68**. An allylic amidation at the more-substituted carbon atom^[41b, 44] in the latter delivers the reaction product and regenerates the Ni⁰ catalyst (Scheme 29).

It was found that C₂-symmetric bidentate bisphosphine ligands such as (*S*)-binap,^[45] (*S,S'*, *R,R'*)-tangphos,^[46] and (*R,R*)-Me-duphos^[47] were not competent in this reaction. However, employment of unsymmetrical P,N-type bidentate ligands such as (*S,S*)-*i*Pr-foxap^[40] and quinap^[48] afforded both good yield and excellent enantioselectivity. This reaction was found to be quite general with respect to the alkyl substituent (R) at the N atom of the triazole moiety, thus producing **66** as the major regioisomer in good enantioselectivity. Reaction of the *tert*-butyl-containing substrate **64** (R = *t*Bu), probably resulting because of steric reasons, produced **65** as a major regioisomer. The *p*-tolyl-substituted substrate **64** (R = *p*-tolyl) was much less efficient in this reaction.

Various monosubstituted allenenes were equally effective in transannulation with **64a**, thus producing high yields and good enantioselectivities of the corresponding products. Allenenes possessing siloxy, benzyloxy, and N-phthalimidoyl groups at the alkyl chains also reacted well, although the enantioselectivities were found to be slightly lower (Scheme 30).^[43]

7. Conclusions

This Minireview highlights the increasing interest in the development of transition metal catalyzed transannulation reactions. This new approach may serve as a complimentary methodology for construction of heterocycles as it allows a general and highly efficient synthesis of complex and highly functionalized aromatic nitrogen heterocycles with diverse substitution patterns. Although additional development of novel, more general, and efficient

transannulation protocols is highly warranted, the progress achieved so far in this area holds promise for its extensive application in organic synthesis.

Acknowledgments

We thank the National Institutes of Health (Grant GM-64444) for financial support of this work.

Biographies



Vladimir Gevorgyan was born in Krasnodar, Russia. He received his B.S. in 1978 from Kuban State University and his Ph.D. in 1984 from the Latvian Institute of Organic Synthesis. He spent two years at Tohoku University, and was a Visiting Professor at Consiglio Nazionale delle Ricerche. He returned to Tohoku University in 1996 and was promoted to Associate Professor in 1997. In 1999 he moved to the University of Illinois at Chicago and was promoted to Professor in 2003. Prof. Gevorgyan's research includes the development of transition metal catalyzed synthetic methodologies.




Buddhadeb Chattopadhyay was born in Insura (Hooghly), West Bengal, India. He received his B.Sc. in 2001 from the Burdwan University, his M.Sc. at Visva-Bharati (Santiniketan), and then joined Prof. K. C. Majumdar at the University of Kalyani. His doctoral work focused on the development of common to medium-sized-ring heterocyclic compounds using transition-metal catalysts. He joined Prof. Gevorgyan's group in 2009 as a postdoctoral research associate to work on transition metal catalyzed transannulation and C-H activation chemistry.

References


1. Bourne Y, Kolb HC, Radic Z, Sharpless KB, Taylor P, Marchot P. *Proc. Natl. Acad. Sci. USA*. 2004; 101:1449. [PubMed: 14757816]
2. Whiting M, Muldoon J, Lin YC, Silverman SM, Lindstron W, Olson AJ, Kolb HC, Finn MG, Sharpless KB, Elder JH, Fokin VV. *Angew. Chem.* 2006; 118:1463. *Angew. Chem. Int. Ed.* 2006; 45:1435.
3. Tornøe CW, Sanderson SJ, Mottram JC, Coombs GH, Meldal M. *J. Comb. Chem.* 2004; 6:312. [PubMed: 15132590]
4. Pande V, Ramos MJ. *Bioorg. Med. Chem. Lett.* 2005; 15:5129. [PubMed: 16213715]
5. Holla BS, Mahalinga M, Karthikeyan MS, Poojary B, Akberathi PM, Kumari NS. *Eur. J. Med. Chem.* 2005; 40:1173. [PubMed: 15979767]
6. For medicinal chemistry, see: Kolb HC, Finn MG, Sharpless KB. *Angew. Chem.* 2001; 113:2056. *Angew. Chem. Int. Ed.* 2001; 40:2004. Norris P. *Curr. Top. Med. Chem.* 2008; 8:101. [PubMed: 18289080] Wilkinson BL, Bornaghi LF, Houston TA, Innocenti A, Vullo D, Supuran CT, Poulsen

- S-A. *J. Med. Chem.* 2007; 50:1651. [PubMed: 17343373] De SK, Stebbins JL, Chen L-H, Riel-Mehan M, Machleidt T, Dahl R, Yuan H, Emdadi A, Barile E, Chen V, Murphy R, Pellicchia M. *J. Med. Chem.* 2009; 52:1943. [PubMed: 19271755]
7. For drug discovery, see: Moorhouse AD, Santos AM, Gunaratnam M, Moore M, Neidle S, Moses JE. *J. Am. Chem. Soc.* 2006; 128:15972. [PubMed: 17165715] Lee LV, Mitchell ML, Huang S-J, Fokin VV, Sharpless KB, Wong C-H. *J. Am. Chem. Soc.* 2003; 125:9588. [PubMed: 12904015]
8. For peptidomimetic area, see: Pedersen DS, Abell A. *Eur. J. Org. Chem.* 2011:2399. Zanda M. *New J. Chem.* 2004; 28:1401. Volonterio A, Bellosa S, Bravin F, Bellucci MC, Bruche L, Colombo G, Malpezzi L, Mazzini S, Meille SV, Meli M, Ramirez de Arellano C, Zanda M. *Chem. Eur. J.* 2003; 9:4510. [PubMed: 14502637]
9. For conformational studies, see: Tornøe CW, Christensen C, Meldal M. *J. Org. Chem.* 2002; 67:3057. [PubMed: 11975567] Angelo NG, Arora PS. *J. Am. Chem. Soc.* 2005; 127:17134. [PubMed: 16332031] Angell Y, Burgess K. *J. Org. Chem.* 2005; 70:9595. [PubMed: 16268639] Oh K, Guan Z. *Chem. Commun.* 2006:3069.
10. For material science, see: Ostrovskii VA, Pevzner MS, Kofmna TP, Shcherbinin MB, Tselinskii IV. *Targets Heterocycl. Syst.* 1999; 3:476. Hiskey M, Chavez DE, Naud DL, Son SF, Berghout HL, Bome CA. *Proc. Int. Pyrotech. Semin.* 2000; 27:3. Helms B, Mynar JL, Hawker CJ, Frechet JMJ. *J. Am. Chem. Soc.* 2004; 126:15020. [PubMed: 15547986] Wu P, Feldman AK, Nugent AK, Hawker CJ, Scheel A, Voit B, Pyun J, Frechet MJ, Sharpless KB, Fokin VV. *Angew. Chem.* 2004; 116:4018. *Angew. Chem. Int. Ed.* 2004; 43:3928.
11. Regitz M. *Angew. Chem.* 1997; 109:786. *Angew. Chem. Int. Ed. Engl.* 1967; 6:733.
12. a) Regitz M, Arnold B, Danion D, Schubert H, Fusser G. *Bull. Soc. Chim. Belg.* 1981; 90:615.b) L'Abbé G. *Bull. Soc. Chim. Belg.* 1990; 99:281.c) L'Abbé G, Godts F, Toppet S. *J. Chem. Soc. Chem. Commun.* 1985:589.d) L'Abbé G, Luyten I, Toppet S. *J. Heterocycl. Chem.* 1992; 29:713.e) L'Abbé G, Godts F, Toppet S. *Bull. Soc. Chim. Belg.* 1986; 95:679.
13. For cyclopropanation with 2-pyridyl diazocompounds, see: Davies HML, Townsend RJ. *J. Org. Chem.* 2001; 66:6595. [PubMed: 11578209]
14. Abarca-González B. *J. Enzyme Inhib. Med. Chem.* 2002; 17:359. and references therein. [PubMed: 12683670]
15. Abarca-González B, Ballesteros R, Mojarred F, Jones G, Mouat DJ. *J. Chem. Soc. Perkin Trans.* 1987; 1:1865.
16. Chuprakov S, Frank, Hwang W, Gevorgyan V. *Angew. Chem.* 2007; 119:4841. *Angew. Chem. Int. Ed.* 2007; 46:4757.
17. Bagheri V, Doyle MP, Taunton J, Claxton EE. *J. Org. Chem.* 1988; 53:6158.
18. Padwa A, Austin DJ, Price AT, Semones MA, Doyle MP, Protopopova MN, Winchester WR, Tran A. *J. Am. Chem. Soc.* 1993; 115:8669.
19. Hoye TR, Dinsmore CJ, Johnson DS, Korkowski PF. *J. Org. Chem.* 1990; 55:4518.
20. a) Rostovtsev VV, Green LG, Fokin VV, Sharpless KB. *Angew. Chem.* 2002; 114:2708. *Angew. Chem. Int. Ed.* 2002; 41:2596.b) Tornøe CW, Christensen C, Meldal M. *J. Org. Chem.* 2002; 67:3057. [PubMed: 11975567] c) Raushel J, Fokin VV. *Org. Lett.* 2010; 12:4952. [PubMed: 20931987]
21. One notable exception: triazoles bearing a strong electron-withdrawing group, such as cyano, nitro, or sulfonyl groups at N1 are known to undergo facile ring opening to diazoimine tautomers. The ring-chain tautomerism manifests itself in various interconversions of triazoles and other heterocycles, collectively known as Dimroth rearrangements. See: Dimroth O. *Justus Liebigs Ann. Chem.* 1909; 364:183. Gilchrist TL, Gymer GE. *Adv. Heterocycl. Chem.* 1974; 16:33.
22. For other types of 5-lithiated triazoles that decompose already at $-78\text{ }^{\circ}\text{C}$, see: Whiting M, Fokin VV. *Angew. Chem.* 2006; 118:3229. *Angew. Chem. Int. Ed.* 2006; 45:3157. Bae I, Han H, Chang S. *J. Am. Chem. Soc.* 2005; 127:2038. [PubMed: 15713069] Cassidy MP, Raushel J, Fokin VV. *Angew. Chem.* 2006; 118:3226. *Angew. Chem. Int. Ed.* 2006; 45:3154. Cho SH, Yoo EJ, Bae I, Chang S. *J. Am. Chem. Soc.* 2005; 127:16046. [PubMed: 16287290] Yoo EJ, Bae I, Cho SH, Han H, Chang S. *Org. Lett.* 2006; 8:1347. [PubMed: 16562888]



23. For reviews on synthesis of cyclopropenes via metal-stabilized carbenes, see: Davies HML, Beckwith REJ. *Chem. Rev.* 2003; 103:2861. [PubMed: 12914484] Doyle MP, Forbes DC. *Chem. Rev.* 1998; 98:911. [PubMed: 11848918]
24. Horneff T, Chuprakov S, Chernyak N, Gevorgyan V, Fokin VV. *J. Am. Chem. Soc.* 2008; 130:14972. [PubMed: 18855475]
25. Connell R, Scavo F, Helquist P, Akermark B. *Tetrahedron Lett.* 1986; 27:5559.
26. Doyle KJ, Moody CJ. *Tetrahedron.* 1994; 50:3761.
27. Padwa A, Kassir JM, Xu SL. *J. Org. Chem.* 1997; 62:1642.
28. Miura T, Yamauchi M, Murakami M. *Chem. Commun.* 2009:1470.
29. Shen Q, Hartwig JF. *J. Am. Chem. Soc.* 2007; 129:7734. [PubMed: 17542591]
30. Chattopadhyay B, Gevorgyan V. *Org. Lett.* 2011; 13:3746. [PubMed: 21692488]
31. Grimster N, Zhang L, Fokin VV. *J. Am. Chem. Soc.* 2010; 132:2510. [PubMed: 20131915]
32. Nakamura I, Nemoto T, Shiraiwa N, Terada M. *Org. Lett.* 2009; 11:1055. [PubMed: 19243194]
33. a) Larock RC, Yum EK. *J. Am. Chem. Soc.* 1991; 113:6689. b) Larock RC, Yum EK, Refvik MD. *J. Org. Chem.* 1998; 63:7652. c) Kürti, L.; Czakó, B. *Strategic Applications of Named Reactions in Organic Synthesis.* New York: Elsevier; 2005. p. 260d) Leogane O, Lebel H. *Angew. Chem.* 2008; 120:356. *Angew. Chem. Int. Ed.* 2008; 47:350.
34. Solé D, Vallverdú L, Solans X, Font-Bardía M, Bonjoch J. *J. Am. Chem. Soc.* 2003; 125:1587. [PubMed: 12568619] For reviews of benzotriazoles, see: Katritzky AR, Suzuki K, Wang Z. *Synlett.* 2005:1656. Katritzky AR, Lan X, Yang JZ, Denisko OV. *Chem. Rev.* 1998; 98:409. [PubMed: 11848906]
35. Miura T, Yamauchi M, Murakami M. *Org. Lett.* 2008; 10:3085. [PubMed: 18558759]
36. For a similar type of the azanickelacyclic intermediacy, see: Takahashi T, Tsai F-Y, Li Y, Wang H, Kondo Y, Yamanaka M, Nakajima K, Kotora M. *J. Am. Chem. Soc.* 2002; 124:5059. [PubMed: 11982370] Duong HA, Louie J. *J. Organomet. Chem.* 2005; 690:5098. Duong HA, Louie J. *Tetrahedron.* 2006; 62:7552.
37. For an example of alkyne insertion into a seven-membered-ring nickelacyclic intermediate, see: Korivi RP, Cheng C-H. *Org. Lett.* 2005; 7:5179. [PubMed: 16268532]
38. Yamauchi M, Morimoto M, Miura T, Murakami M. *J. Am. Chem. Soc.* 2010; 132:54. [PubMed: 20000760]
39. A similar regioselectivity trend was observed in the insertion of allenes into palladacycles; see: Lu G, Malinakova HC. *J. Org. Chem.* 2004; 69:8266. [PubMed: 15549797]
40. Miyake Y, Nishibayashi Y, Uemura S. *Synlett.* 2008:1747.
41. Miura T, Morimoto M, Yamauchi M, Murakami M. *J. Org. Chem.* 2010; 75:5359. [PubMed: 20604510]
42. Miura T, Nishida Y, Morimoto M, Yamauchi M, Murakami M. *Org. Lett.* 2011; 13:1429. [PubMed: 21319832]
43. Miura T, Yamauchi M, Kosaka A, Murakami M. *Angew. Chem.* 2010; 122:5075. *Angew. Chem. Int. Ed.* 2010; 49:4955.
44. a) Pawlas J, Nakao Y, Kawatsura M, Hartwig JF. *J. Am. Chem. Soc.* 2002; 124:3669. [PubMed: 11929257] b) Grigg R, Liu A, Shaw D, Suganthan S, Woodall DE, Yoganathan G. *Tetrahedron Lett.* 2000; 41:7125.
45. Noyori R, Takaya H. *Acc. Chem. Res.* 1990; 23:345.
46. Tang W, Zhang X. *Angew. Chem.* 2002; 114:1682. *Angew. Chem. Int. Ed.* 2002; 41:1612.
47. Burk MJ. *J. Am. Chem. Soc.* 1991; 113:8518.
48. Alcock NW, Brown JM, Hulmes DI. *Tetrahedron: Asymmetry.* 1993; 4:743.



Scheme 1.
Closed/opened form equilibrium of pyridotriazoles.





Scheme 2.
Rhodium carbenoid insertion into a Si-H bond.




Scheme 3.


Transannulation of the pyridotriazole **3b** with alkynes. pfb = perfluorobutyrate.




Scheme 4.
Denitrogenative annulation of pyridotriazoles **3** with nitriles.



Scheme 5.
Proposed mechanism for transannulation of pyridotriazoles **3** with alkynes and nitriles.



Scheme 6.
Denitrogenative cyclopropanation of the N-sulfonyl-1,2,3-triazole **17a** with styrene. DCE = 1,2-dichloroethane, oct = octanoate.




Microwave:

[Rh₂(oct)₄], CHCl₃ } 82%
 140 °C, 15 min


Conventional heating:

[Rh₂(oct)₄], CHCl₃ } 85%
 80 °C, 15h



Scheme 7. Transannulation of N-sulfonyl-1,2,3-triazole **17a** with benzonitrile employing microwave-assisted and conventional heating methods.




Scheme 8.
Denitrogenative transannulation of N-sulfonyl-1,2,3-triazoles **17** with nitriles. TMS = trimethylsilyl.




Scheme 9.
Proposed mechanism for the transannulation of N-sulfonyl-1,2,3-triazoles **17** with nitriles.

**Scheme 10.**


Nickel-catalyzed transannulation of the N-sulfonyl-1,2,3-triazoles **17** with internal alkynes. The yields are those of the isolated product. [a] $[Ni(cod)_2]$ (15 mol%) and $P(nBu)Ad_2$ (30 mol%). [b] $110^\circ C$. Ad = adamantyl, cod = 1,5-cyclooctadiene.

**Scheme 11.**


Proposed mechanism of the nickel-catalyzed denitrogenative transannulation of N-tosonyl-1,2,3-triazole **17a** with internal alkynes. Ts = 4-toluenesulfonyl.

**Scheme 12.**


Rhodium-catalyzed denitrogenative transannulation of N-sulfonyl-1,2,3-triazoles **17b** with terminal alkynes.




Scheme 13.
Proposed mechanism for transannulation of N-sulfonyl-1,2,3-triazoles **17b** with terminal alkynes.




Scheme 14.
Semi-one-pot transannulation toward synthesis of pyrroles.

**Scheme 15.**


Transannulation of the NH-1,2,3-triazoles **36** with styrenes. NTTL = *N*-1,8-naphthoyl-*tert*-leucine.




Scheme 16.
Palladium-catalyzed transannulation versus the Larock indole synthesis.





Scheme 17.
Palladium-catalyzed denitrogenative transannulation of the N-arylbenzotriazoles **39a** with internal alkynes.




Scheme 18.
Proposed mechanism of the palladium-catalyzed transannulation of N-arylbenzotriazoles **39a** with internal alkynes.




Scheme 19. Nickel-catalyzed transannulation of the 1,2,3-benzotriazinones **47** with an internal symmetrical alkyne.

**Scheme 20.**

Nickel-catalyzed denitrogenative transannulation of 1,2,3-benzotriazinones **47** with alkynes.
pin = pinacol.

**Scheme 21.**


Nickel-catalyzed denitrogenative transannulation of the 1,2,3-benzotriazinones **47a** with allenes. dppbenz = 1,2-bis(diphenylphosphino) benzene.




R^1 = tolyl, aryl,
 R^2 = H, OMe
 R^3 = H, OMe, CO_2Me
 R = hexyl, $(\text{CH}_2)_2\text{OBn}$, $(\text{CH}_2)_2\text{OSi}t\text{BuMe}_2$,
 $(\text{CH}_2)_2\text{OH}$, $(\text{CH}_2)_2\text{CN}$, $\text{Si}t\text{BuMe}_2$

14 examples
 yield: 76-99%
 selectivity: up to 0:100

Scheme 22.
 Scope of the nickel-catalyzed transannulation of 1,2,3-benzotriazinones **47b** with allenes.

**Scheme 23.**

Nickel-catalyzed denitrogenative transannulation of 1,2,3-benzotriazinones **47a** with internal cyclic allene. See Scheme 29 for structure of $(R,R)\text{-Me-duphos}$.



R¹ = tolyl, aryl, CONPh₂


R² = H, OMe

R³ = H, OMe, CO₂Me


R = alkyl, protected alkyl

19 examples
yield: 76-99%
ee: 78-97%


Scheme 24.
Enantioselective synthesis of isoquinolones.

**Scheme 25.**


Transannulation of benzotriazinones with 1,3-dienes. dppf = 1,1'-bis(diphenylphosphino)ferrocene.





with symmetrical 1,3-dienes




with unsymmetrical 1,3-dienes




Scheme 26.
Scope of transannulation of the benzotriazinones **47b** with 1,3-dienes.




Scheme 27.
Nickel-catalyzed denitrogenative transannulation with alkenes.



Scheme 28. Palladium-catalyzed transannulation with isocyanides. Cp = cyclopentadienyl.

**Scheme 29.**

Transannulation of 1,2,3,4-benzothiazine-1,1(2*H*)-dioxide **64** with allenes in the presence of various chiral ligands.

**Scheme 30.**Transannulation of 1,2,3,4-benzothiazine-1,2(*2H*)-dioxide **64a** with various allenes.