

Age-dependent decline in renal function in patients with lung cancer

GEN OHARA^{1,2}, KOICHI KURISHIMA², KENSUKE NAKAZAWA², MIO KAWAGUCHI²,
KATSUNORI KAGOHASHI¹, HIROICHI ISHIKAWA³, NOBUYUKI HIZAWA² and HIROAKI SATOH¹

¹Division of Respiratory Medicine, Mito Medical Center, University of Tsukuba; ²Division of Respiratory Medicine, Institute of Clinical Medicine, University of Tsukuba; ³Division of Respiratory Medicine, Tsukuba Medical Center, Ibaraki, Japan

Received January 3, 2012; Accepted March 30, 2012

DOI: 10.3892/ol.2012.672

Abstract. The aim of this study was to clarify the effect of aging on renal function. Serum creatinine (SCr), blood urea nitrogen (BUN) and 24-h creatinine clearance (measured-CrCl) were examined in lung cancer patients and the measured-CrCl were compared with CrCl estimates by employing two commonly used equations. In total, 787 lung cancer patients who were diagnosed between 2001 and 2010 were retrospectively analyzed. SCr and urine creatinine, BUN and measured-CrCl were evaluated prior to treatment. The Cockcroft-Gault (CG) and modification of diet in renal disease (MDRD) formulae were also used to estimate CrCl. SCr, BUN and measured-CrCl showed a significant decline in the elderly. In the 787 lung cancer patients, a significant correlation coefficient was found between measured-CrCl and age. However, in patients aged 80 years or older, no significant correlation coefficient was found between measured-CrCl and age. In the comparison between the measured CrCl and the CrCls estimated by the two formulae, the CG-CrCl levels were lower than those of the measured-CrCl, whereas the MDRD-CrCl levels were higher. Age is a crucial factor influencing renal function in patients with lung cancer. Particularly in the elderly, a decline in CrCl and greater individual variability in CrCl, as well as discrepancies in measured-CrCl and estimated CrCls are significant factors.

Introduction

Lung cancer in the elderly is an increasingly common problem faced by the oncologist. Elderly patients have more co-morbid diseases and tend to tolerate toxic medical treatments more poorly than their younger counterparts. Recently, however,

clinical trials of platinum-based chemotherapy for selected elderly patients with good condition have been conducted, attracting attention to the utility of such therapy for the elderly (1,2). Assessment of renal function is a daily clinical task, particularly in geriatric practice. Impairment of renal function is common in elderly patients, and their glomerular filtration rate (GFR) should be considered prior to prescription of drugs that are eliminated through the renal route or performing investigations requiring iodine injection. In clinical practice, GFR is rarely measured, since it is expensive and requires the use of radioelements for isotopic clearance determination, therefore renal function is assessed by the determination of creatinine clearance (CrCl). Measured-CrCl has been used as the reference method since direct measurement of GFR using isotopic clearance is difficult to perform in elderly patients, and it is recognized to be a reliable estimate of GFR in older individuals (3). Nevertheless, the number of lung cancer patients older than 80 years evaluated for CrCl in previous studies has been too small (4,5). Therefore, additional studies focusing especially on renal function in patients aged over 80 years are required. To clarify the effect of aging on renal function, we measured serum creatinine (SCr), blood urea nitrogen (BUN), and 24-h creatinine clearance (measured-CrCl) in elderly patients with lung cancer and compared the measured-CrCl with CrCl estimates obtained from two commonly used formulae for CrCl.

Patients and methods

Patients. Data from 787 patients with newly diagnosed lung cancer who were admitted to the Division of Respiratory Medicine, Tsukuba University Hospital or Tsukuba Medical Center Hospital between January 2001 and April 2010 were retrospectively analyzed. For each patient, the diagnosis of lung cancer was confirmed with pathological and/or cytological specimens. Pathological and/or cytological diagnosis was defined by the WHO classification and patients were staged according to the seventh edition of the cancer staging system (6). Age, height, actual body weight and gender were recorded at the initial visit. The evaluation of clinical data for lung cancer patients was approved by the institutional ethics committee of University of Tsukuba Hospital.

Correspondence to: Dr Hiroaki Satoh, Division of Respiratory Medicine, Mito Medical Center, University of Tsukuba, Miya-machi 3-2-7, Mito, Ibaraki 310-0015, Japan
E-mail: hirosato@md.tsukuba.ac.jp

Key words: renal function, creatinine clearance, lung cancer, elderly

Table I. Characteristics of 787 patients with lung cancer.

Characteristic	
Age (years)	Median: 70, range: 21-87
≤49	30 (3.8%)
50-59	121 (15.4%)
60-69	251 (31.9%)
70-79	327 (41.6%)
≥80	58 (7.4%)
Gender	
Male	631 (80.2%)
Female	156 (19.8%)
Histology	
Adenocarcinoma	374 (47.5%)
Squamous cell carcinoma	191 (24.3%)
Small cell carcinoma	178 (22.6%)
Large cell carcinoma	20 (2.5%)
Others	24 (3.0%)
Clinical stage	
IA-III A	242 (30.7%)
IIIB	211 (26.8%)
IV	334 (42.4%)

Creatinine clearance (CrCl). Urinary and serum creatinine and BUN were measured using the enzymatic method. Prior to commencement of the lung cancer treatment, measured-CrCl in all patients was calculated three times and averaged. For each patient, the CrCl was also estimated using the Cockcroft and Gault (CG) and modification of diet in renal disease (MDRD) formulae as follows:

$$BSA = 0.007184 \times Ht^{0.725} \times Wt^{0.425}$$

$$\text{Measured-CrCl} = (1.73/BSA) \times (UCr \times Uvol)/(24 \times 60 \times SCr)$$

$$\text{CG-CrCl} = (1.73/BSA) \times \{(140-A)/(SCr+0.2)\} \times (Wt/72) \times 0.85 \text{ (in the case of a female patient) (7)}$$

$$\text{MDRD-CrCl} = (1/0.719) \times 194 \times SCr^{-1.094} \times A^{-0.287} \times 0.739 \text{ (in the case of a female patient) (8,9)}$$




Figure 1. The measured-CrCl among the decades of age. The measured-CrCl decreased in an age-dependent manner, with significant differences among the age decades ($P=0.001$, Kruskal-Wallis test).

where BSA indicates body surface area (m^2); Ht, height (cm); Wt, body weight (kg); UCr, urine creatinine concentration (mg/dl) (using the enzymatic method); Uvol, 24 h urine volume (ml/day); SCr, serum creatinine concentration (mg/dl) (using the enzymatic method); and A, age (years). We added 0.2 to a SCr value measured using an enzymatic peroxidase-antiperoxidase method in order to render it equivalent to the SCr value measured using the Jaffé method (10).

Statistical analysis. The Mann-Whitney U test or Kruskal-Wallis test was applied to elucidate the difference between two or more independent groups. Statistical significance between paired baseline and CrCl levels was evaluated using the Wilcoxon signed-rank test. To evaluate the correlation between two independent groups, the Spearman's rank correlation was used. Statistical analysis were performed using SPSS 10.1 for Windows (SPSS, Chicago, IL, USA). $P < 0.05$ was considered statistically significant.

Results

Patient characteristics. Of the 787 patients included in the present study, 80.2% ($n=631$) were men (Table I). The median

Table II. Serum levels of creatinine and blood urea nitrogen in each age decade.

Age (years)	Creatinine (mg/dl) (median, range)	P-value ^a	BUN (mg/dl) (median, range)	P-value ^a
≤49	0.60, 0.37-1.10	0.001	11.7, 5.5-19.8	0.001
50-59	0.70, 0.39-1.19		13.4, 5.3-27.1	
60-69	0.70, 0.30-2.20		14.9, 5.3-44.9	
70-79	0.80, 0.20-5.40		16.0, 3.7-86.4	
≥80	0.80, 0.79-2.60		17.1, 5.6-53.9	

^aData analysis was computed by Kruskal-Wallis test. BUN, blood urea nitrogen.

Table III. Value of measured-CrCl, CG-CrCl and MDRD-CrCl.

Patient age (years)	Measured CrCl (median, range)	CG-CrCl (median, range)	MDRD-CrCl (median, range)
All patients	89.4, 5.1-239.0	63.1, 6.3-141.2 ^a	104.4, 8.3-339.9 ^a
<70 years	102.1, 15.8-239.3	72.4, 22.5-141.2 ^a	114.1, 27.5-228.7 ^a
70-79 years	79.7, 5.1-159.3	54.9, 6.3-124.4 ^a	99.8, 8.3-339.9 ^a
≥80 years	72.6, 17.9-147.2	45.1, 17.2-71.8 ^a	85.2, 27.0-163.4 ^a

CrCl, Creatinine clearance; CG, Cockcroft-Gault; MDRD, modification diet in renal disease. ^aP=0.001 versus measured-CrCl.




Figure 2. Correlation between measured-CrCl and age in all 787 lung cancer patients. A statistically significant correlation coefficient was found between measured-CrCl and age (Spearman's p-value = -0.447, P=0.001).

age was 70 years (range, 21-87). The types of lung cancer comprised 609 non-small cell carcinomas and 178 small cell carcinomas. Stage IA-IIIa was noted in 242 patients, stage III-B in 211 patients, and stage IV disease in 334 patients.

Renal function with age. SCr and BUN levels were determined in each decade of age (Table II). Both SCr and BUN showed a statistically significant difference among the age decades (P=0.001, Kruskal-Wallis test). The measured-CrCl decreased in an age-dependent manner, with significant differences among the age decades (P=0.001, Kruskal-Wallis test) (Fig. 1). The CG-CrCl and MDRD-CrCl also decreased in an age-dependent manner (P=0.001 and P=0.001, respectively, data not shown).

Correlation between CrCl and age. In all the 787 lung cancer patients, a statistically significant correlation coefficient was found between measured-CrCl and age (Spearman's p-value = -0.447, P=0.001) (Fig. 2). In patients aged ≥70 years, a statistically significant correlation coefficient was also found between measured-CrCl and age, although Spearman's p-value was extremely low (Spearman's p-value = -0.172, P=0.001) (Fig. 3). However, in patients aged ≥80 years, a statistically significant correlation coefficient was not found (Spearman's p-value = -0.100, P=0.456, data not shown).




Figure 3. Correlation between measured-CrCl and age in patients aged ≥70 years. A statistically significant correlation coefficient was also found between measured-CrCl and age, although the Spearman's p-value was extremely low (Spearman's p-value = -0.172, P=0.001).

Comparison among measured-CrCl, CG-CrCl and MDRD-CrCl. The median values and ranges of the measured-CrCl, CG-CrCl, and MDRD-CrCl are shown in Table III. In all 787 patients, the CG-CrCl levels were significantly lower than those of the measured-CrCl (P=0.001, Wilcoxon signed-rank test). However, the MDRD-CrCl levels were significantly higher than those of the measured-CrCl (P=0.001). Particularly in patients aged 70-79 years, the CG-CrCl levels were lower than the measured-CrCl levels, and the MDRD-CrCl levels were significantly higher than the measured-CrCl levels (P=0.001 and P=0.001, respectively). These discrepancies were also observed in patients aged ≥80 years.

Discussion

With recent improvements in the medical management of lung cancer, the availability of many new drugs, better standards of medical care and more widespread health services, survival of elderly lung cancer patients has improved. In addition, Quiox *et al* recently reported that treatment with platinum-based doublet chemotherapy resulted in better survival for the very elderly with non-small cell lung cancer than did standard single-agent therapy (11). As the proportion of elderly patients

aged 80 years or older continues to increase, intensive chemotherapy is likely to bring a clinical benefit to elderly patients with clinically good conditions such as unimpaired renal function. However, few studies evaluating renal function in a large number of elderly patients with lung cancer have been published thus far.

Renal function is known to decrease with age and morphological changes, such as decrease of kidney weight, appearance of sclerotic glomeruli (12) and intimal proliferation in the renal artery, are some of the causes of renal dysfunction (13). SCr concentration remains the most widely used index of renal function in clinical practice (14). In the very elderly, however, serum creatinine is not always useful as a marker of renal function since creatinine production is low due to decreased muscle mass associated with aging (15,16). The GFR is generally used as an index of renal function and may be accurately measured through the renal clearance of either cold (inulin, iohexol) or radiolabeled (51Cr-EDTA, 99mTc-DTPA) exogenous filtration markers. Nonetheless, these methods are seldom used in clinical practice as they are invasive and expensive and require the use of radioelements for isotopic clearance determination. Instead, indirect methods are used for bedside renal function estimates, all of which are based on CrCl.

In oncological practice, estimation of GFR at the bedside is essential since renal insufficiency is directly associated with increased chemotherapeutic complications. The two most commonly used formulae, the CG and MDRD, were developed to estimate the CrCl (7,17,18). However, it is notable that in the original report in which the CG equation was validated, only 17 of a total of 249 patients were aged over 80 years (7). In addition, the MDRD formula was developed to estimate CrCl in patients with a variety of chronic renal diseases (19) and this population differs from those recruited for CrCl estimation with the CG formula as well as from elderly cancer patients such as those of the present study. Therefore, these two commonly used equations have not yet established an optimal accuracy in very elderly patients with lung cancer. Furthermore, a marked discrepancy between the CG and the MDRD formulae has been observed in a large population of over 65 years of age (20) and these formulae have been found to lack a consistent proportion of patients with renal failure within a population of octogenarian inpatients (21). Studies estimating renal function in solid cancer patients have previously been published (22-25). However, the number of the cancer patients aged 80 years or older evaluated for renal function was limited (24,25). Therefore, additional studies focusing specifically on renal function in very elderly lung cancer patients are required. To clarify the effect of aging on renal function, we evaluated SCr, BUN, and measured-CrCl in elderly patients with lung cancer. SCr concentration and BUN showed statistically significant increase among the decades of age, whereas measured CrCl showed a statistically significant decrease among the age decades. Analysis of the correlation between measured-CrCl and age showed a statistically significant correlation coefficient in all 787 lung cancer patients. A significant but weaker correlation coefficient was observed in patients aged 70 years or older. In patients aged 80 years or older, however, a statistically significant correlation coefficient was not found between measured-CrCl and age. We also compared

the measured-CrCl, CG-CrCl and MDRD-CrCl levels. In all 787 patients, the CG-CrCl levels were significantly lower than those of the measured-CrCl, whereas the MDRD-CrCl levels were significantly higher. Particularly in patients aged 70 and over, CG-CrCl was lower than the measured-CrCl, and the MDRD-CrCl levels were significantly higher than those of the measured-CrCl. Our results are consistent with those obtained by Péquignot *et al* from 121 patients aged 70 and older with an indwelling urinary catheter (4).

Despite the novel findings of the present study, there are several limitations. Firstly, GFR was not determined using the isotopic reference method. Such methods are difficult to use in elderly patients with lung cancer as they are invasive and seldom readily available in oncological practice. The use of the isotopic method in the elderly patients of our study would have introduced a major selection bias. Secondly, this study was retrospective and our findings were obtained from a limited number of patients that were neither population-based nor case-controlled. Nevertheless, we suggest that it is of interest to report the management of unselected groups of elderly lung cancer patients. In the present study, we confirmed the importance of age as a factor influencing renal function in patients with lung cancer. As an early detection method and curative therapy have yet to be established, all lung cancer patients have a high possibility of receiving chemotherapy at the time of the first presentation or recurrence. Advanced lung cancer patients are particularly likely to receive chemotherapy. In the present study, we revealed a decline in CrCl in the elderly. In addition, we showed greater individual variability in CrCl in elderly lung cancer patients. For elderly patients, therefore, careful evaluation of renal function is essential to avoid excessive dosage of chemotherapeutic drugs.

Acknowledgements

We are especially grateful to Ms. F. Miyamasu for her review of the manuscript.

References

1. Vamvakas L, Saloustros E, Karampeazis A and Georgoulis V: Advanced non-small-cell lung cancer in the elderly. *Clin Lung Cancer* 10: 158-167, 2009.
2. Lang K, Marciniak MD, Faries D, *et al*: Trends and predictors of first-line chemotherapy use among elderly patients with advanced non-small cell lung cancer in the United States. *Lung Cancer* 63: 264-270, 2009.
3. Lamb EJ, Webb MC, Simpson DE, *et al*: Estimation of glomerular filtration rate in older patients with chronic renal insufficiency: is the modification of diet in renal disease formula an improvement? *J Am Geriatr Soc* 51: 1012-1017, 2003.
4. Péquignot R, Belmin J, Chauvelier S, *et al*: Renal function in older hospital patients is more accurately estimated using the Cockcroft-Gault formula than the modification diet in renal disease formula. *J Am Geriatr Soc* 57: 1638-1643, 2009.
5. Verhave JC, Fesler P, Ribstein J, *et al*: Estimation of renal function in subjects with normal serum creatinine levels: influence of age and body mass index. *Am J Kidney Dis* 46: 233-241, 2005.
6. Lababede O, Meziane M and Rice T: Seventh edition of the cancer staging manual and stage grouping of lung cancer: quick reference chart and diagrams. *Chest* 139: 183-189, 2011.
7. Cockcroft DW and Gault MH: Prediction of creatinine clearance from plasma creatinine. *Nephron* 16: 31-41, 1976.
8. Matsuo S, Imai E, Horio M, *et al*: Revised equations for estimated GFR from serum creatinine in Japan. *Am J Kidney Dis* 53: 982-992, 2009.

9. Sato H, Kuroda T, Tanabe N, *et al*: Cystatin C is a sensitive marker for detecting a reduced glomerular filtration rate when assessing chronic kidney disease in patients with rheumatoid arthritis and secondary amyloidosis. *Scand J Rheumatol* 39: 33-37, 2010.
10. Horio M and Orita Y: Comparison of Jaffé rate assay and enzymatic method for the measurement of creatinine clearance. *Jap J Nephrol* 38: 296-299, 1996.
11. Quoiq E, Zalcmán G, Oster JP, *et al*: Carboplatin and weekly paclitaxel doublet chemotherapy compared with monotherapy in elderly patients with advanced non-small-cell lung cancer: IFCT-0501 randomised, phase 3 trial. *Lancet* 378: 1079-1088, 2011.
12. Kaplan C, Pasternack B, Shah H, *et al*: Age-related incidence of sclerotic glomeruli in human kidneys. *Am J Pathol* 80: 227-234, 1975.
13. Tauchi H, Tsuboi K and Sato K: Histology and experimental pathology of senile atrophy of the kidney. *Nagoya Med J* 4: 71-97, 1958.
14. Levey AS: Measurement of renal function in chronic renal disease. *Kidney Int* 38: 167-184, 1990.
15. Beck LH: Changes in renal function with aging. *Clin Geriatr Med* 11: 199-209, 1998.
16. Herig PJ and Carlson RE: Plasma creatinine and renal function in the elderly. *JAMA* 248: 31, 1982.
17. Levey AS, Bosch JP, Lewis JB, *et al*: A more accurate method to estimate glomerular filtration rate from plasma creatinine: a new prediction equation. Modification of Diet in Renal Disease study group. *Ann Intern Med* 130: 461-470, 1999.
18. Levey AS, Greene T, Kusek JW, *et al*: A simplified equation to predict glomerular filtration rate from plasma creatinine. *J Am Soc Nephrol* 11: A0828, 2000.
19. National Kidney Foundation: K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, stratification. *Am J Kidney Dis* 39: S1-S266, 2002.
20. Garg AX, Papaioannou A, Ferko N, *et al*: Estimating the prevalence of renal insufficiency in seniors requiring long-term care. *Kidney Int* 65: 649-653, 2004.
21. Rimon E, Kagansky N, Cojocarú L, Gindin J, Schattner A and Levy S: Can creatinine clearance be accurately predicted by formulae in octogenarian in-patients? *QJM* 97: 281-287, 2004.
22. Nishimura G, Tsukuda M, Horiuchi C, *et al*: Decrease of creatinine clearance rate with aging in patients with head and neck cancer in Japan. *Int J Clin Oncol* 12: 120-124, 2007.
23. Hurria A, Hurria A, Brogan K, *et al*: Effect of creatinine clearance on patterns of toxicity in older patients receiving adjuvant chemotherapy for breast cancer. *Drugs Aging* 22: 785-791, 2005.
24. Séronie-Vivien S, Toullec S, Malard L, *et al*: Contribution of the MDRD equation and of cystatin C for renal function estimates in cancer patients. *Med Oncol* 23: 63-73, 2006.
25. Thyss A, Saudes L, Otto J, *et al*: Renal tolerance of cisplatin in patients more than 80 years old. *J Clin Oncol* 12: 2121-2125, 1994.