

Quiescent keratocytes fail to repair MMC induced DNA damage leading to the long-term inhibition of myofibroblast differentiation and wound healing

James V. Jester,¹ Chyong Jy Nien,¹ Vasilis Vasilidou,² Donald J. Brown¹

¹The Gavin Herbert Eye Institute, University of California, Irvine, Irvine, CA; ²Department of Pharmaceutical Sciences, Skaggs School of Pharmacy and Pharmaceutical Sciences, University of Colorado Denver, Aurora, CO

Purpose: The purpose of this study was to determine the acute and long-term effects of mitomycin C (MMC) on quiescent rabbit corneal keratocytes regarding cell proliferation, myofibroblast differentiation and DNA repair.

Methods: Quiescent keratocytes cultured in serum-free media were exposed to various concentrations of MMC and then treated with transforming growth factor- β (TGF β). DNA damage was evaluated in both cultured keratocytes and live rabbit eyes following treatment with MMC. The long-term ability of quiescent keratocytes to repair MMC induced damage in vivo was evaluated in rabbits treated with MMC 2 months before 100 μ m deep lamellar keratectomy (LK) injury.

Results: MMC significantly blocked TGF β -induced cell proliferation and myofibroblast differentiation in cultured quiescent keratocytes and altered the transcriptional regulation of macrophage chemotactic protein-1 (MCP-1) and alpha smooth muscle actin (α SMA). MMC also induced phosphorylation of the nuclear histone marker of DNA damage, γ H2AX (a member of the H2A histone family), without induction of cell cycle entry or immediate DNA repair measured by Comet assay. In live rabbits, 0.2 mg/ml MMC significantly induced γ H2AX nuclear immunostaining ($p < 0.05$) throughout the cornea and corneas receiving 0.2 mg/ml MMC treatment 2 months before LK injury showed complete absence of any corneal scarring.

Conclusions: MMC induces DNA damage to quiescent corneal keratocytes, which remains unrepaired, resulting in abnormal cell replication and gene transcription that leads to long-term effects on corneal repair. Overall these findings suggest that there may be long-term and perhaps permanent consequences to the application of MMC as an anti-fibrotic therapy.

Mitomycin C (MMC) belongs to a family of anti-tumor quinolone antibiotics derived from *Streptomyces caespitosus*, and functions as a powerful bifunctional alkylating agent that induces DNA interstrand crosslinks (ICL). These changes lead to the inhibition of cell replication, altered gene transcription and cell death if not repaired [1,2]. While approved as an anti-cancer chemotherapeutic agent, MMC has recently gained widespread clinical acceptance in the ophthalmic community as an off-label prophylactic anti-fibrotic therapy to prevent corneal scarring and haze following corneal refractive surgery, including photorefractive keratectomy (PRK) and laser assisted in situ keratomileusis (LASIK) [3-5]. Treatment of patients immediately following PRK with 0.2 mg/ml (0.02%) MMC for 12 s to 2 min significantly decreases haze and results in better refractive predictability and improved efficacy for over a year and half after surgery [6-9].

Initial laboratory studies evaluating the effects of MMC in rabbits have shown that inhibition of corneal haze following PRK is associated with a significant reduction of keratocyte

cell density [10,11]. Later studies have shown that loss of keratocyte density after MMC treatment is associated with increased keratocyte apoptosis and the absence of myofibroblasts within the wound [12-14]. More recent studies have also shown that adjacent keratocytes fail to enter the cell cycle following PRK with MMC treatment suggesting that MMC blocks cell replication [15]. Furthermore, DNA damage as detected by phosphorylation of H2AX, a member of the H2A nuclear histone family that initiates recruitment of nuclear excision repair endonucleases [16-18], has been identified in the nuclei of corneal endothelial cells in eyes treated with MMC under ex vivo conditions [19]. These data indicate that the effect of MMC on corneal scarring and haze is related to the generation of DNA ICLs leading to defective cell replication and gene expression.

While the molecular mechanism underlying MMC's effect on corneal scarring and haze remains unknown, previous investigators have proposed that increased MMC-induced keratocyte apoptosis and the profound inhibition of keratocyte and myofibroblast progenitor cell proliferation may play a central role [15]. To more clearly understand the process by which MMC inhibits corneal scarring and haze after PRK, we have evaluated the effects of MMC on quiescent corneal keratocytes. Our findings indicate that

Correspondence to: James V. Jester, Ph.D., The Gavin Herbert Eye Institute, 843 Health Sciences Road, Irvine, CA; Phone: (949) 824-8047; FAX: (949) 824-9626; email: jjester@uci.edu

MMC induces DNA damage that leads to abnormal myofibroblast differentiation, proliferation and gene transcription. More disturbing is the finding that DNA damage in MMC treated keratocytes leads to long-term effects on myofibroblast differentiation and corneal repair in culture and animal wound healing models. Overall, the findings suggest that there may be long-term consequences to the use of MMC as an anti-fibrotic that have yet to be clinically manifested.

METHODS

Cell culture studies: Rabbit corneal keratocytes were isolated from rabbit eyes (Pel Freez, Rogers, AR) and cultured under serum-free conditions as previously described [20,21]. Cells were plated at a density of $1-5 \times 10^4$ cells/cm² in polystyrene coated tissue culture dishes (Falcon; Becton Dickinson, Franklin Lakes, NJ) and allowed to attach for 72 h. Cells were then treated with various concentrations of MMC (Calbiochem, La Jolla, CA) for 5 min and then rinsed in serum-free media. Cells were then cultured with and without transforming growth factor- β (TGF β , 5 ng/ml; Sigma-Aldrich, St. Louis, MO) for various times and evaluated by immunocytochemistry, western blotting, and RT-PCR.

Immunohistochemistry: Cells were fixed in 2% paraformaldehyde in PBS (pH=7.4), then treated with acetone (-20 °C) for 3 min, rinsed with PBS, blocked with goat serum (dilution 1:10 in PBS; Cappel; MP Biomedicals, Solon, OH) for 30 min at 37 °C and stained with monoclonal antibodies to α -smooth muscle actin (α SMA; Sigma, St. Louis, MO) to measure myofibroblast differentiation, the nuclear cell cycle protein Ki67 (clone MIB-2; Immunotech, Westbrook, ME) to measure cell cycle entry and γ H2AX (Abcam, Cambridge, MA) to assess DNA damage [16-18]. Sections were then stained with FITC conjugated goat anti-mouse IgG (Southern Bioech, Birmingham, AL) and counterstained with Alexa 543 phalloidin and DAPI. Sections were then evaluated using the Nikon Eclipse E600 epifluorescence microscope (Nikon Instruments Inc., Melville, NY).

Cell proliferation: To assess the effects of MMC on TGF β induced cell proliferation, keratocytes were cultured in 6-well plates (Falcon) and treated with various concentrations of MMC for 5 min. Cells were then cultured with and without TGF β and the change in cell number measured as previously describe using a Leica DMIRB Inverted microscope (Leica Microsystems, Wetzlar, Germany) and Optronics CCD camera (Goleta, CA) [22]. At least three wells/treatment condition were measured and the experiment repeated 3 times.

Western blotting: Total proteins were solubilized in buffer containing 25 mM Tris-HCl (pH 7.4), 1 mM EDTA, 1 mM EGTA, 10 mM dithiothreitol, 1% sodium dodecyl sulfate (SDS), 5 μ g/ml antipain, 5 μ g/ml pepstatin A, and 1 mM phenyl methyl sulfonyl fluoride (PMSF). Proteins were electrophoresed on precast 10% Tris-glycine sodium dodecyl sulfate polyacrylamide (SDS-PAGE) gels (Invitrogen,

Carlsbad, CA). Proteins were transferred to a polyvinylidene difluoride membranes (Bio-Rad, Hercules, CA), blocked for 2 h with 5% BSA in Tris saline and probed with antibodies to α SMA (Sigma-Aldrich) at 4 °C overnight. Blots were then washed with Tween-20 and Tris-buffered saline (50mM Tris, 150mM NaCl and 0.05% Tween-20) and incubated for 1 h with a Cy3 conjugated goat anti-mouse IgG (1:5000; Southern Biotech). Western blots were then scanned at an excitation of 532 nm and emission of 580 nm using a fluorescent imager (FMBIO III; Hitachi-MiraiBio Inc., Alameda, CA).

RNA extraction and PCR: Cells were washed 2 \times in cold PBS then directly lysed in RLT buffer (Qiagen, Valencia, CA) and the supernatant recovered. The samples were then passed over QiaShredder columns and then processed over RNeasy columns (both from Qiagen). The RNA was eluted from the silica column with 50 μ l water, and the product was analyzed using a NanoDrop Spectrophotometer (Thermo Scientific, Wilmington, DE) to estimate RNA concentrations. The RNA (500 μ g) obtained from cultured cells was converted to cDNA using the QuantiTect Reverse Transcription Kit (Qiagen). Reactions were performed for 30 min at 42 °C, and 5 min at 95 °C, followed by cooling to 4 °C. After synthesis, the cDNA from all sources was diluted to equivalent input RNA levels with TE (10 mM Tris/HCl, 1 mM EDTA [pH 7.5]; 10 ng input RNA/ μ l) and stored at -20 °C until use. cDNA samples were subjected to PCR using specific primers (Table 1). Primers were designed using Primer 3 Internet software program (The Whitehead Institute, Cambridge, MA), and their specificities were confirmed by a BLAST Internet software-assisted search of the non-redundant nucleotide sequence database (National Library of Medicine, Bethesda, MD). Polymerase chain reactions (PCR) were carried out as previously described with 5–25 ng reverse-transcribed RNA and 200 nM forward and reverse primers, in a total volume of 25 ml [23]. For quantitative real time PCR (qPCR), samples were normalized by b2-MG amplification and were amplified using the QuantiTect Syber Green reagents (Qiagen) in triplicate using an MJ Research Opticon Thermal Cycler (Waltham, MA). PCR controls without reverse transcriptase (water control) or with normal human genomic DNA as a template were routinely negative. PCR controls without reverse transcriptase (water control) were routinely negative. Product specificities were confirmed by DNA sequencing the products.

Comet assay: DNA repair by corneal fibroblasts and keratocytes was assessed by Comet assay using the protocol previously described by Cordelli et al. [24]. Briefly, slides were prepared by dipping in 1% agarose/PBS and allowing to air dry to a thin film. Cells were harvested 24 h post MMC treatment by scraping in cold PBS, centrifuged at 500 \times g and the cell pellet re-suspended in PBS to 10⁷ cells/ml. The cell suspension (10 μ l) was quickly mixed with 65 μ l 0.7% Low Melting Point agarose/PBS, coverslipped, and the agarose allowed to gel at 4 °C. After gelling, the coverslip was

TABLE 1. RT-PCR PRIMERS.

Product	Forward	Reverse	GenBank
α smooth muscle actin	TGTGCTATGTCGCTCTGGAC	CTTCTGCATACGGTCAGCAA	X60732
GAPDH	GAGCTGAACGGGAAACTCAC	CCCTGTTGCTGTAGCCAAAT	NM_001082253
β actin	ATCGTGATGGACTCCGGCGAC	AGGCCACGTAGCACAGC	NM_001101683
MCP-1	CACCCGGACACCTCTACTA	GATCCTTGCAAGAACCCTCA	M57440

removed and the gel overlaid with an additional 100 μ l of 0.7% LMP agarose/PBS and allowed to gel. Slides were then submerged overnight at 4 °C in lysis buffer containing: 2.5M NaCl, 100 mM EDTA, 10 mM Tris/HCl pH 10, 1% Triton X 100 and 10% DMSO. The following morning, the slides were placed in lysis buffer containing 10 mM DTT and allowed to incubate, for 30 min at 4 °C. The slides were then placed under freshly prepared electrophoresis buffer composed of: 300 mM NaOH, 1 mM EDTA, adjusted to pH 13 with HCl and held at 4 °C for 20 min. The samples were then electrophoresed at 4 °C, for 5 min at 25V. The slides were then immersed in 0.3 M sodium acetate in 70% alcohol for 30 min, then fixed in 100% alcohol for 2 h, followed by 70% alcohol for 5 min and then allowed to air dry. The slides were stained by immersion in 2 μ g/ml Ethidium Bromide and photographed under a 20 \times objective. Slides were scored using CometScore software v1.5 (TriTek Corporation, Summerduck, VA).

Rabbit studies: New Zealand albino rabbits were used in this study and all procedures were approved by the UCI IACUC and conducted in accordance with ARVO Statement for the Use of Animals in Ophthalmic and Vision Research. Prior to injury, all rabbits were anesthetized with intramuscular injections of xylazine (5 mg/Kg) (Anased, Lloyd Laboratories, Shenandoah, IA) and ketamine (22 mg/Kg) (Bioniche Pharma USA LLC, Lake Forest, IL). Tetracaine hydrochloride 0.5% eye drops (Alcon Laboratories Inc., Forth Worth, TX) were instilled into each right eye, and a speculum was used to open the eyelids. After surgery, eyes received topical gentamicin sulfate (Akorn, Buffalo Grove, IL) three times a day for 3 days and intramuscular buprenorphine (Buprenx; 0.1 mg/kg; Reckitt Benckiser Healthcare, Hull, UK) once and as needed for relief of pain.

DNA damage study: A total of 6 rabbits were used in this study. An 8-mm trephine (Bausch & Lomb, Rochester, NY) was used to mark the central cornea and the epithelium was then removed using a Bard Parker® #11 surgical blade (Becton Dickinson AcuteCare, Franklin Lakes, NJ). Both eyes of each rabbit were then randomly assigned to four groups of 3 eyes each and were treated with either 0.2 mg/ml or 0.02 mg/ml MMC for 15 s or 60 s. Animals were allowed to recover and then sacrificed 3 h after treatment.

MMC and delayed injury study: A total of 10 rabbits were used in this study. Initially rabbits received a unilateral epithelial scrape injury and then randomly assigned to two groups of 5 rabbits each and treated with 0.2 mg/ml MMC for 60 s or vehicle control applied using a surgical sponge as

previous described [25]. Animals were allowed to recover for 8 weeks and then received a lamellar keratectomy (LK) injury in the same treated eye using techniques previously reported [26,27]. Rabbits were then followed by in vivo confocal microscopy and then sacrificed 3 months after LK injury.

In vivo confocal microscopy (CM): Under anesthesia, animals received a baseline scan and were then followed at various times after injury by in vivo confocal microscopy (CM) to measure epithelial and stromal thickness and haze as described previously [28,29]. All in vivo CM examinations were performed using a tandem scanning confocal microscope (TSCM; Tandem Scanning Corporation, Reston, VA) with a 24 \times surface-contact objective (numerical aperture, 0.6; working distance, 1.5 mm). One drop of 2.5% hydromethylcellulose (Gonak, Akorn, Buffalo Grove, IL) was placed on the tip of the objective as a coupling gel. All camera settings were kept constant throughout the experiment. For each examination, repeated through focus data sets were obtained from the central corneal region. A total of three to five through-focus data sets for each eye at each time point were collected in the thinnest region of the cornea.

Ex vivo microscopy: After animals were sacrificed, corneas were fixed by anterior chamber perfusion for 4 min with 2% paraformaldehyde in phosphate-buffered saline, pH 7.4. After perfusion, corneas were dissected and left in 2% paraformaldehyde overnight. For assessment of DNA damage following MMC treatment, fixed corneal tissue was embedded in O.C.T Compound (Tissue-Teck, Sakura Finetek, Torrance, CA) and snap frozen in liquid nitrogen. Blocks were then cut on a Leica CM 1850 cryotome (Leica, Wetzlar, Germany) and stained with antibodies specific of γ H2AX for 1 h followed by staining with goat anti-mouse IgG conjugated to FITC for 1 h. Slides were then counter stained with DAPI (300 nM, Invitrogen, Carlsbad, CA) and imaged using a Nikon Eclipse E600 epifluorescent microscope equipped with a Photometrics CoolSnap FX camera (Roper Scientific, Tucson, AZ) and images digitized using Meta Imaging Series software (Molecular Devices, Downingtown, PA). All images were taken using the same camera settings and exposure time, and all slides were stained on the same day. To quantify the intensity of γ H2AX staining over keratocyte nuclei, the DAPI channel was used to define regions of interest over nuclei using the create region tool in the Meta Imaging software. These regions were then applied to the γ H2AX channel, and the average pixel intensity over the nuclei measured and recorded. To adjust for differences in staining, the average

background image intensity overlying non-nuclear areas was also measured and subtracted from the measured nuclear intensity for each slide. On average, a total of 20 nuclei from each corneal section from each region were measured.

To assess corneal fibrosis, collagen organization of the corneas was visualized by non linear optical imaging of second harmonic generated signals from collagen (SHG). SHG signals to detect collagen were generated using 800-nm infrared femtosecond laser pulses to obtain maximum SHG emission signal at 400 nm [25]. SHG forward-scattered signals passing through the tissue were collected using an 0.8 NA condenser lens with a narrow band-pass filter (400/50) placed in front of the transmission light detector. Backward-scattered SHG signals were detected using the Meta Detector set to collect 390 to 410 nm band width on the Zeiss 510 Meta Laser Scanning Confocal microscope (Carl Zeiss Microscopy GmbH, Jena, Germany). To assess cell density in the same optical plane, corneal blocks were also stained with Syto 59 (Invitrogen), and imaged using laser scanning confocal microscopy.

Statistical analysis: All results are reported as mean±standard deviation. Differences between treatment groups were assessed by two-way repeated-measures ANOVA and Bonferroni multiple comparisons (Sigma Stat version 3.11; Systat Software Inc., Point Richmond, CA). For the live rabbit studies we used a sample size of five eyes per group, with a power of 0.95 and $\alpha=0.05$ to detect a 50% reduction in haze. All tissue culture experiments were performed in triplicate and repeated at least three times.

RESULTS

Effects of MMC on TGF β -induced myofibroblast differentiation: To establish whether MMC blocked myofibroblast differentiation of quiescent keratocytes, cultures were stained with specific antibodies to α SMA, a known biomarker for myofibroblast differentiation [21], and the number of cells expressing α SMA quantified. By comparison, keratocytes cultured under serum-free conditions maintained a dendritic morphology and showed no staining for α SMA (Figure 1A), while cells exposed to TGF β alone showed cell spreading, increased actin filament assembly and marked expression of α SMA (Figure 1B). When cells were pre-treated with MMC for 5 min and then immediately treated with TGF β , MMC at concentrations of 0.02 mg/ml and 0.002 mg/ml blocked TGF β induced cell spreading, actin filament assembly and expression of α SMA at 72 h (Figure 1C,1D). Concentrations above 0.02 mg/ml induced marked keratocyte cell death. Quantifying the number of cells expressing α SMA showed that the pre-treatment of cells with MMC dose dependently and significantly reduced ($p<0.001$) the number of cells staining for α SMA. Western blots also confirmed the reduction in expression of α SMA after 72 h of TGF β treatment (Figure 1E, insert). Overall, these data

showed that MMC leads to abnormal TGF β -induced myofibroblast differentiation of rabbit corneal keratocytes.

Effects of MMC on TGF β -induced cell replication and gene transcription: Since DNA damage induced by MMC is known to block cell replication and gene transcription, we next evaluated the effect of MMC on TGF β -induced cell growth and gene expression of quiescent keratocytes (Figure 2). While keratocytes showed no increase in cell number over 7 days in culture with or without MMC treatment (data not shown), TGF β treatment alone resulted in a doubling in the cell number (Figure 2A). Pretreatment of cells with 0.02 mg/ml MMC, on the other hand, resulted in a loss of cells averaging only 77%±16% of the original cell numbers 7 days after treatment. Pretreatment with 0.002 mg/ml MMC resulted in only a modest increase in the number of cells over 7 days that remained significantly below TGF β alone treated cells ($p<0.001$). When treatment of cells with TGF β was delayed 7 days to allow for DNA repair, MMC treated keratocytes continued to show a dose dependent decrease in the ability to replicate and express α SMA in response to TGF β (data not shown) suggesting prolonged effects of MMC that are not repaired in culture.

To assess the effect of MMC on gene expression by quiescent keratocytes, the expression of MCP-1 and α SMA were evaluated by RT-PCR. MCP-1, is a small inducible cytokine belonging to the CC chemokine family that is involved in recruiting macrophages, memory T-cells and dendritic cells to sites of injury and is thought to play an important role in the development of fibrosis [30,31]. As shown in Figure 2B, TGF β treated quiescent keratocytes showed increased expression of mRNA for *MCP-1* and *α SMA*. Treatment with increasing concentrations of MMC showed a dose dependent loss of TGF β -induced MCP-1. Interestingly, expression of TGF β -induced *α SMA* mRNA showed increasing expression with higher concentrations of MMC. Overall, these findings are consistent with MMC induced DNA damage leading to abnormal cell replication and gene expression patterns.

MMC induced DNA damage and repair in culture: Since the effects of MMC on quiescent keratocytes were not reversed by extended cell culture, we next evaluated the MMC induced DNA damage and repair. As a bifunctional ankylosing agent, MMC induces DNA interstrand crosslinks (ICLs) that lead to phosphorylation of histone, H2AX, which recruits nuclear excision repair endonucleases that detach ICLs and repair DNA through a homologous recombination [16-18]. Antibodies specific for γ H2AX, the phosphorylated form of H2AX used as a molecular marker for DNA ICLs [16], stained keratocyte nuclei that had been treated with MMC (Figure 3A). Quantification of the number of cells stained by γ H2AX following MMC treatment showed a dose dependent increase with concentration of 0.07 mg/ml showing greater than 80% γ H2AX staining that peaked 2 days after treatment and




Figure 1. Concentration-dependent effects of MMC on keratocytes treated with TGFβ for 72 h. Cells were untreated (A) or treated with TGFβ alone (B) or after 0.02 mg/ml MMC (C) or 0.002 mg/ml MMC (D) treatment and then stained with phalloidin (red), anti-αSMA (green), and DAPI (blue). Cells treated with TGFβ alone showed a spread morphology and αSMA staining typical of myofibroblasts (B), while cells treated with MMC before TGFβ showed loss of cell spreading and decreased αSMA staining (C and D). Quantitation of αSMA staining (E) showed significantly reduced numbers of myofibroblasts following pre-treatment with MMC and decreased protein expression for αSMA (insert).

decreased to 40% by day 4 (data not shown). When MMC treated keratocytes were stained for Ki67, a marker of cell cycle entry and DNA repair, no Ki67 staining was detected (Figure 3B), suggesting that keratocytes did not go through a normal replication coupled DNA repair pathway. This is different from corneal fibroblasts that showed both γH2AX staining and Ki67 labeling following treatment with MMC (Figure 3C,3D, respectively).

To determine whether MMC treated keratocytes undergo DNA repair, a Comet assay was performed on MMC treated quiescent keratocytes and fibroblast cultures to detect interstrand breaks in nuclear DNA, necessary to uncouple and remove ICLs in preparation for DNA repair. In a Comet assay, DNA migrates from the nuclei, or head of the Comet, and forms a progressively longer tail depending on the number of breaks (Figure 3E, insert). Replicating cells, such as corneal





Figure 2. The effects of MMC on TGFβ-induced cell proliferation and gene expression. **A:** The effect of MMC and TGFβ on cell proliferation showed that 0.02 and 0.002 mg/ml significantly reduced ($p < 0.001$) cell growth compared to TGFβ alone. **B:** RT-PCR for *MCP-1* and *αSMA* showed that TGFβ alone induces the expression of both *MCP-1* and *αSMA*, but that MMC dose dependently blocks *MCP-1*, while increasing *αSMA* mRNA when normalized to glyceraldehyde 3-phosphate dehydrogenase (*GAPDH*).



fibroblasts, show a dose dependent increase in the Tail Moment of the Comet when treated with MMC and evaluated 24 h after treatment, indicating uncoupling of ICLs. By comparison, keratocytes showed a marked decrease in Tail moment 24 h after treatment with increasing doses of MMC, suggesting no DNA excision repair following MMC treatment. Overall, these data indicate that MMC dose dependently leads to DNA damage of quiescent corneal keratocytes in culture that does not undergo normal replicative DNA repair.

MMC induced DNA damage in the cornea: To determine whether MMC induced DNA damage in the cornea, rabbits were treated with MMC following scrape injury and then evaluated for the expression of γ H2AX 3 h after treatment (Figure 4). Sections stained with antibodies to γ H2AX (green)

and counter stained with nuclear dye (red) showed staining of nuclei both in the anterior stroma (Figure 4A, arrow), posterior stroma (Figure 4B), and corneal endothelium (Figure 4B, arrow). In the very anterior stroma, keratocyte nuclei show little or no staining, most likely related to the known keratocyte apoptosis and cell death following scrape injury (Figure 4A, arrowhead). When the intensity of γ H2AX staining over the nuclei was measured, very high intensity was detected over stromal keratocytes when exposed to 0.2 mg/ml MMC for 60 s (Figure 4C). There was a significant decrease in intensity ($p < 0.05$) when the exposure was reduced from 60 s to 15 s, but the intensity of staining remained significantly greater than that detected for corneas treated with 0.02 mg/ml MMC for either 60 s or 15 s, which was not significantly different from background. Regarding differences in staining




Figure 3. DNA repair in quiescent keratocytes compared to proliferating fibroblasts following MMC exposure. MMC treated Keratocytes (A and B) and corneal fibroblasts (C and D) were stained with γ H2AX (A and C, green), Ki67 (B and D, green) and DAPI (red) 24 h after treatment. Keratocytes showed only γ H2AX staining while fibroblasts showed γ H2AX and Ki67 staining. Comet assay (E) shows that MMC dose dependently increases the Comet tail in fibroblasts 24 h after treatment, while significantly decreasing the Comet tail in quiescent keratocytes.

between anterior and posterior keratocytes, no significant differences were detected for both 60 s and 15 s exposures to 0.2 mg/ml MMC, suggesting all cells in the corneal stroma are equally damaged by MMC treatment in the rabbit at a given dose and exposure. It should also be noted that expression of Ki67 was not detected in corneal sections, consistent with previous studies that have evaluated later time points (data not shown) [15]. Overall, these data indicate that 0.2 mg/ml MMC (either 60 s or 15 s exposure times) is the lowest dose that induces significant DNA damage.

Effect of delayed injury on MMC treated corneas: To access repair of MMC DNA damage to quiescent keratocytes in live rabbits, eyes were treated with 0.2 mg/ml MMC for 60 s following epithelial scrape injury. Rabbits were then allowed to recover for 2 months before 100 μ m deep LK injury followed by evaluation using in vivo confocal microscopy. As shown in previous studies [26,27], LK injury in control, vehicle treated eyes resulted in significant corneal scarring that was detected by in vivo confocal microscopy as intense light scattering (Scar) underlying the corneal epithelium (Epi) in XZ projections (Figure 5A). In regions of the scar, single planes (XY) showed a high density of scar fibroblasts embedded in highly light scattering extracellular matrix

(Figure 5B). By contrast, eyes treated 2 months previously with 0.2 mg/ml MMC showed no corneal scarring in XZ projections (Figure 5C) and in single planes (XY) underlying the corneal epithelium showed a sparse population of keratocytes embedded in low light scattering, transparent extracellular matrix (Figure 5D). Measurement of the light scattering from the corneas also showed significantly reduced ($p < 0.001$) corneal haze in eyes pretreated with 0.2 mg/ml MMC (Figure 5E).

The absence of corneal scarring and reduced keratocyte cell numbers was also confirmed by ex vivo confocal and nonlinear optical imaging of collagen using SHG imaging (Figure 6). In vehicle treated eyes, irregular collagen organization was detected by SHG (Figure 6A) that measured on average $84.2 \pm 27 \mu$ m in thickness. By comparison, eyes previously treated with 0.2 mg/ml MMC showed no irregular collagen deposition and only the presence of normal, broad collagen lamellae at the interface between the corneal epithelium and the stroma (Figure 6B). The nuclear density of the corneal scar in the vehicle treated eyes was also significantly greater ($p < 0.05$) averaging 382 ± 114 cells/mm² immediately underlying the corneal epithelium compared to the MMC treated eyes, which averaged 126 ± 67 cells/mm².




Figure 4. DNA damage in rabbit corneas following treatment with MMC. γ H2AX staining (green) and DAPI staining (red) of the anterior (A) and posterior (B) cornea following MMC treatment. Note that the most anterior keratocytes show only DAPI staining (arrowhead) suggesting apoptosis, while the remaining keratocytes (A, arrow) and posterior endothelial cells (B, arrow) show γ H2AX staining. Quantification of γ H2AX staining intensity (C) showed that 0.2 mg/ml MMC produced significantly greater staining compared to corneas treated with 0.02 mg/ml, and that 60 s exposure produced significantly greater staining compared to 15 s. It should also be noted that there was no significant difference (NS) in staining between cells in the anterior or posterior cornea.

Overall, this data suggests that MMC damage to quiescent corneal keratocytes goes unrepaired in rabbits for at least 2 months after treatment.

DISCUSSION

Previous studies have shown that MMC treatment following PRK results in the induction of keratocyte apoptosis, inhibition of keratocyte proliferation and reduced keratocyte repopulation of the anterior cornea [15,32]. While these

findings have lead some investigators to suggest that the apoptotic and anti-proliferative effects of MMC block activation of myofibroblast progenitor cells [33], the mechanism underlying the anti-fibrotic effect of MMC remains unclear. In this report, we show that MMC dose dependently blocks TGF β -induced proliferation and alters gene transcription of cultured quiescent rabbit keratocytes and that quiescent keratocytes do not recover a normal TGF β response for at least 7 days in culture. These observations




Figure 5. In vivo confocal microscopy of lamellar keratectomy injury in rabbit corneas 2 months after MMC treatment. In vivo confocal microscopy of rabbit corneas receiving vehicle control (A and B) or 0.2 mg/ml MMC (C and D) 2 months before lamellar keratectomy (LK). In control eyes, 3 months after LK injury, there is a dense scar underlying the epithelium (Epi) detected in the XZ projection (A) that contains densely populated scar fibroblasts seen in the XY plane underlying the epithelium (B). Eyes treated 2 months earlier with MMC showed no scar in the XZ projection (C) and decreased keratocytes underlying the epithelium in the XY plane (D) 3 months after LK injury. Quantification of light scattering showed that eyes treated with MMC 2 months before LK injury had significantly less haze (E).

support and extend previous observations in live rabbit eyes and cultured horse corneal fibroblasts and are consistent with the DNA interstrand crosslinking (ICL) effects of MMC [15,34]. More importantly, we show for the first time that the failure of quiescent keratocytes to recover from MMC treatment and respond to TGF β is associated with failed or greatly delayed DNA repair as evidenced by the absence of cell cycle entry following phosphorylation of H2AX in response to MMC induced ICLs, the prolonged expression of γ H2AX indicating the persistence of ICLs, and the lack of

formation of DNA interstrand breaks detected by Comet Assay. More troubling was the finding that the effect of MMC persisted for at least 2 months in live rabbits as evidenced by the absence of corneal fibrosis in response to LK injury. Together these findings suggest that the anti-fibrotic effect of MMC is due to persistent DNA damage that blocks the ability of quiescent keratocytes to respond to wound cytokines. While it is not known whether a similar mechanism underlies the anti-fibrotic effect of MMC in patients receiving refractive




Figure 6. Corneal scarring and keratocyte cell density in rabbit corneas 3 months after lamellar keratectomy in eyes treated with and without MMC. Second harmonic (A and B) and confocal imaging of the same plane stained with Syto59 (C and D) of control (A and C) and 0.2 mg/ml MMC treated corneas (B and D) 2 months before lamellar keratectomy injury. Control corneas 3 months after LK injury showed the presence of irregularly organized scar collagen (A) and a high density of scar fibroblast nuclei (B). By comparison, eyes treated with MMC 2 months before LK injury showed normal lamellar stromal collagen at the epithelial-stromal interface (C) and markedly decreased corneal keratocytes (D) 3 months after LK injury.

surgery, these findings suggest that there may be significant long-term consequences for MMC therapy.

Recently, studies evaluating the effects of MMC on the goat cornea have shown that topical treatment with therapeutic doses of MMC induced DNA damage detected by modified Comet assay as increased 'head' DNA, consistent with decreased Tail moment measured in this report [19]. Additionally, cultured goat corneal endothelial cells treated with various concentrations of MMC showed persistent γ H2AX staining suggesting markedly delayed or defective DNA repair. Importantly, these findings are similar to our findings in the rabbit keratocyte showing both prolonged γ H2AX staining and failed TGF β responsiveness, even after 7 days of recovery in culture. Our in vivo findings that γ H2AX uniformly and equally stains anterior and posterior stromal keratocytes as well as corneal endothelial cells, also support the previous observations that the corneal endothelium is a likely target for MMC-induced DNA damage even in thicker human corneas. Furthermore, the suspicion that DNA repair is lacking in the cornea based on prolonged γ H2AX staining is confirmed by our finding that 2 months recovery following

MMC treatment failed to restore a normal wound healing response in the rabbit.

While DNA repair mechanisms in the cornea are unknown, it is generally thought that cells enter the cell cycle following MMC treatment and that cycle progression at S-phase is blocked, where upon cells undergo DNA repair [35]. In cell cycling competent cells, nuclear excision repair endonucleases (NER) have been shown to 'unhook' and remove ICLs, followed by homologous recombination. However, studies of terminally differentiated cells, including muscle and nerve, have shown that this normal repair mechanism is turned off [36]. As an alternative, quiescent, serum-starved cells are thought to have transcription-coupled repair (TCR) mechanisms, acting only on the transcribed genes, and non-transcribed strand repair (NTSR) that act on non-transcribed genes, both through different mechanisms [37]. More recent studies of terminally differentiated muscle cells suggest that these alternative mechanisms are impaired or severely compromised and that accumulation of DNA damage may lead to restricted apoptosis and autophagy [36, 38]. Such a fate for MMC treated corneal keratocytes and endothelial cells is suggested by the finding of persistent

annexin V staining in goat corneal endothelial cells [19], and the decrease keratocyte density detected *in vivo* and in culture following TGF β treatment. Based on this data, additional investigation into the long-term effects of MMC in culture and live animals seems warranted. On a more fundamental level, a better understanding of DNA repair mechanisms of corneal keratocytes and endothelial cells is needed since these cells are exposed to DNA damaging UV and oxidative stress through out life. Importantly, these studies need to evaluate DNA repair mechanisms in quiescent and differentiated cells that critically lack the global genome repair mechanisms that have been extensively studied in serum cultured, cycling cells.

An additional finding in this study was the observation that the only dose of MMC to cause significant γ H2AX staining of corneal keratocytes and corneal endothelial cells was the currently accepted therapeutic dose. While there were significant differences in the intensity γ H2AX staining comparing 60 s exposure to 15 s exposure, a lower dose of MMC at either 60 s or 15 s exposure failed to show significantly detectable γ H2AX staining. Previous studies have extensively evaluated the effect of lower doses of MMC on corneal keratocyte density, cell cycling and haze, and observed some anti-fibrotic effects using a “prophylactic” dose of 0.002% MMC [15]. It should be noted, however, that this study also used a longer exposure interval of 2 min, and that this dose has been shown clinically to produce ‘breakout haze’ [5,15,39]. Studies performed in our laboratory that tested different doses of MMC with exposure times of 15 s found that the only dose that significantly reduced corneal fibrosis as measured by SHG imaging in rabbits was the recognized therapeutic dose, consistent with the γ H2AX staining and the clinical observations (data not shown).

In summary, this report provides evidence that DNA damage induced by MMC goes unrepaired in rabbit keratocytes. This finding suggests that DNA repair mechanisms in corneal keratocytes may be limited, and that further studies are needed to evaluate how keratocytes repair DNA damage in the cornea. Since accumulated, unrepaired DNA damage is thought to underlie many degenerative neurologic disorders [40], understanding DNA repair in the cornea may have wider implications than just for MMC-induced corneal injury. While it is widely recognized that the safety of using MMC as an off-label anti-fibrotic has yet to be established, our findings taken together with those of Roh et al. [19] suggest that the lack of DNA repair following MMC treatment could have important long-term consequences in those patients having refractive surgery. Besides the obvious long-term effects on wound healing following accidental injury or additional surgical intervention, the accumulative genomic and mitochondrial DNA damage induced by MMC may increase the susceptibility of patients to other degenerative corneal disorders. Clearly, caution in the use of MMC is warranted until a clearer understanding of the potential long-term side effects have been established.

ACKNOWLEDGMENTS

Supported in part by NEI grants, EY07348, EY016663, EY017963, The Discovery Eye Foundation, The Skirball Program in Molecular Ophthalmology, Support grant from Research to Prevent Blindness, Inc.

REFERENCES

1. Tomasz M. Mitomycin C: small, fast and deadly (but very selective). *Chem Biol* 1995; 2:575-9. [PMID: 9383461]
2. Galm U, Hager MH, Van Lanen SG, Ju J, Thorson JS, Shen B. Antitumor antibiotics: bleomycin, enediyne, and mitomycin. *Chem Rev* 2005; 105:739-58. [PMID: 15700963]
3. Azar DT, Jain S. Topical MMC for subepithelial fibrosis after refractive corneal surgery. *Ophthalmology* 2001; 108:239-40. [PMID: 11158781]
4. Majmudar PA, Forstot SL, Dennis RF, Nirankari VS, Damiano RE, Brenart R, Epstein RJ. Topical mitomycin-C for subepithelial fibrosis after refractive corneal surgery. *Ophthalmology* 2000; 107:89-94. [PMID: 10647725]
5. Shah RA, Wilson SE. Use of mitomycin-C for photorefractive keratectomy and photorefractive keratectomy surgery. *Curr Opin Ophthalmol* 2010; 21:269-73. [PMID: 20489622]
6. Argento C, Cosentino MJ, Ganly M. Comparison of laser epithelial keratomileusis with and without the use of mitomycin C. *J Refract Surg* 2006; 22:782-6. [PMID: 17061715]
7. Bedei A, Marabotti A, Giannechini I, Ferretti C, Montagnani M, Martinucci C, Barabesi L. Photorefractive keratectomy in high myopic defects with or without intraoperative mitomycin C: 1-year results. *Eur J Ophthalmol* 2006; 16:229-34. [PMID: 16703539]
8. de Benito-Llopis L, Teus MA, Sanchez-Pina JM, Fuentes I. Stability of laser epithelial keratomileusis with and without mitomycin C performed to correct myopia in thin corneas: a 15-month follow-up. *Am J Ophthalmol* 2008; 145:807-12. [PMID: 18342832]
9. Leccisotti A. Mitomycin-C in hyperopic photorefractive keratectomy. *J Cataract Refract Surg* 2009; 35:682-7. [PMID: 19304089]
10. Schipper I, Suppelt C, Gebbers JO. Mitomycin C reduces scar formation after excimer laser (193 nm) photorefractive keratectomy in rabbits. *Eye (Lond)* 1997; 11:649-55. [PMID: 9474312]
11. Xu H, Liu S, Xia X, Huang P, Wang P, Wu X. Mitomycin C reduces haze formation in rabbits after excimer laser photorefractive keratectomy. *J Refract Surg* 2001; 17:342-9. [PMID: 11383766]
12. Kim TI, Lee SY, Pak JH, Tchah H, Kook MS. Mitomycin C, ceramide, and 5-fluorouracil inhibit corneal haze and apoptosis after PRK. *Cornea* 2006; 25:55-60. [PMID: 16331043]
13. Kim TI, Pak JH, Lee SY, Tchah H. Mitomycin C-induced reduction of keratocytes and fibroblasts after photorefractive keratectomy. *Invest Ophthalmol Vis Sci* 2004; 45:2978-84. [PMID: 15326110]
14. Lai YH, Wang HZ, Lin CP, Chang SJ. Mitomycin C alters corneal stromal wound healing and corneal haze in rabbits after argon-fluoride excimer laser photorefractive

- keratectomy. *J Ocul Pharmacol Ther* 2004; 20:129-38. [PMID: 15117569]
15. Netto MV, Mohan RR, Sinha S, Sharma A, Gupta PC, Wilson SE. Effect of prophylactic and therapeutic mitomycin C on corneal apoptosis, cellular proliferation, haze, and long-term keratocyte density in rabbits. *J Refract Surg* 2006; 22:562-74. [PMID: 16805119]
 16. Clingen PH, Wu JY, Miller J, Mistry N, Chin F, Wynne P, Prise KM, Hartley JA. Histone H2AX phosphorylation as a molecular pharmacological marker for DNA interstrand crosslink cancer chemotherapy. *Biochem Pharmacol* 2008; 76:19-27. [PMID: 18508035]
 17. Mah LJ, El-Osta A, Karagiannis TC. gammaH2AX: a sensitive molecular marker of DNA damage and repair. *Leukemia* 2010; 24:679-86. [PMID: 20130602]
 18. van Attikum H, Gasser SM. Crosstalk between histone modifications during the DNA damage response. *Trends Cell Biol* 2009; 19:207-17. [PMID: 19342239]
 19. Roh DS, Cook AL, Rhee SS, Joshi A, Kowalski R, Dhaliwal DK, Funderburgh JL. DNA cross-linking, double-strand breaks, and apoptosis in corneal endothelial cells after a single exposure to mitomycin C. *Invest Ophthalmol Vis Sci* 2008; 49:4837-43. [PMID: 18658091]
 20. Beales MP, Funderburgh JL, Jester JV, Hassell JR. Proteoglycan synthesis by bovine keratocytes and corneal fibroblasts: maintenance of the keratocyte phenotype in culture. *Invest Ophthalmol Vis Sci* 1999; 40:1658-63. [PMID: 10393032]
 21. Jester JV, Barry-Lane PA, Cavanagh HD, Petroll WM. Induction of alpha-smooth muscle actin expression and myofibroblast transformation in cultured corneal keratocytes. *Cornea* 1996; 15:505-16. [PMID: 8862928]
 22. Jester JV, Ho-Chang J. Modulation of cultured corneal keratocyte phenotype by growth factors/cytokines control in vitro contractility and extracellular matrix contraction. *Exp Eye Res* 2003; 77:581-92. [PMID: 14550400]
 23. Brown DJ, Lin B, Holguin B. Expression of neuregulin 1, a member of the epidermal growth factor family, is expressed as multiple splice variants in the adult human cornea. *Invest Ophthalmol Vis Sci* 2004; 45:3021-9. [PMID: 15326116]
 24. Cordelli E, Fresegna AM, D'Alessio A, Eleuteri P, Spano M, Pacchierotti F, Villani P. ReProComet: a new in vitro method to assess DNA damage in mammalian sperm. *Toxicol Sci* 2007; 99:545-52. [PMID: 17675332]
 25. Farid M, Morishige N, Lam L, Wahlert A, Steinert RF, Jester JV. Detection of corneal fibrosis by imaging second harmonic-generated signals in rabbit corneas treated with mitomycin C after excimer laser surface ablation. *Invest Ophthalmol Vis Sci* 2008; 49:4377-83. [PMID: 18502995]
 26. Gatlin J, Melkus MW, Padgett A, Petroll WM, Cavanagh HD, Garcia JV, Jester JV. In vivo fluorescent labeling of corneal wound healing fibroblasts. *Exp Eye Res* 2003; 76:361-71. [PMID: 12573665]
 27. Jester JV, Barry-Lane PA, Petroll WM, Olsen DR, Cavanagh HD. Inhibition of corneal fibrosis by topical application of blocking antibodies to TGF beta in the rabbit. *Cornea* 1997; 16:177-87. [PMID: 9071531]
 28. Jester JV, Petroll WM, Cavanagh HD. Measurement of tissue thickness using confocal microscopy. *Methods Enzymol* 1999; 307:230-45. [PMID: 10506977]
 29. Li J, Jester JV, Cavanagh HD, Black TD, Petroll WM. On-line 3-dimensional confocal imaging in vivo. *Invest Ophthalmol Vis Sci* 2000; 41:2945-53. [PMID: 10967049]
 30. Ebihara N, Yamagami S, Yokoo S, Amano S, Murakami A. Involvement of C-C chemokine ligand 2-CCR2 interaction in monocyte-lineage cell recruitment of normal human corneal stroma. *J Immunol* 2007; 178:3288-92. [PMID: 17312179]
 31. Lee SK, Choi BK, Kang WJ, Kim YH, Park HY, Kim KH, Kwon BS. MCP-1 derived from stromal keratocyte induces corneal infiltration of CD4+ T cells in herpetic stromal keratitis. *Mol Cells* 2008; 26:67-73. [PMID: 18594181]
 32. Chang SW. Corneal keratocyte apoptosis following topical intraoperative mitomycin C in rabbits. *J Refract Surg* 2005; 21:446-53. [PMID: 16209441]
 33. Salomao MQ, Wilson SE. Corneal molecular and cellular biology update for the refractive surgeon. *J Refract Surg* 2009; 25:459-66. [PMID: 19507799]
 34. Buss DG, Sharma A, Giuliano EA, Mohan RR. Efficacy and safety of mitomycin C as an agent to treat corneal scarring in horses using an in vitro model. *Vet Ophthalmol* 2010; 13:211-8. [PMID: 20618797]
 35. Mladenov E, Tsaneva I, Anachkova B. Activation of the S phase DNA damage checkpoint by mitomycin C. *J Cell Physiol* 2007; 211:468-76. [PMID: 17167777]
 36. Fortini P, Dogliotti E. Mechanisms of dealing with DNA damage in terminally differentiated cells. *Mutat Res* 2010; 685:38-44. [PMID: 19941876]
 37. Bielas JH. Non-transcribed strand repair revealed in quiescent cells. *Mutagenesis* 2006; 21:49-53. [PMID: 16394029]
 38. Narciso L, Fortini P, Pajalunga D, Franchitto A, Liu P, Degan P, Frechet M, Demple B, Crescenzi M, Dogliotti E. Terminally differentiated muscle cells are defective in base excision DNA repair and hypersensitive to oxygen injury. *Proc Natl Acad Sci USA* 2007; 104:17010-5. [PMID: 17940040]
 39. Santhiago MR, Netto MV, Wilson SE. Mitomycin C: biological effects and use in refractive surgery. *Cornea* 2012; 31:311-21. [PMID: 22157595]
 40. Jeppesen DK, Bohr VA, Stevnsner T. DNA repair deficiency in neurodegeneration. *Prog Neurobiol* 2011; 94:166-200. [PMID: 21550379]

Articles are provided courtesy of Emory University and the Zhongshan Ophthalmic Center, Sun Yat-sen University, P.R. China. The print version of this article was created on 1 July 2012. This reflects all typographical corrections and errata to the article through that date. Details of any changes may be found in the online version of the article.