

OXA-163-Producing *Klebsiella pneumoniae* in Cairo, Egypt, in 2009 and 2010

Mohammed O. Abdelaziz,^a Celestino Bonura,^b Aurora Aleo,^b Ramadan A. El-Domany,^a Teresa Fasciana,^b and Caterina Mammina^b

Department of Microbiology and Immunology, Faculty of Pharmacy, Helwan University, Cairo, Egypt,^a and Department of Sciences for Health Promotion "G. D'Alessandro," University of Palermo, Palermo, Italy^b

Two genetically unrelated OXA-163-carrying *Klebsiella pneumoniae* strains were identified from two infection cases in June 2009 and May 2010 in Cairo, Egypt. OXA-163-producing *Enterobacteriaceae* had been previously reported in Argentina only. Both patients had no history of travel abroad. The emergence of this newly recognized OXA-48-related β -lactamase able to hydrolyze cephalosporins and carbapenems is especially worrying in a geographic area where OXA-48 is endemic and effective surveillance for antibiotic resistance is largely unaffordable.

Carbapenems are first-line drugs for severe infections caused by extended-spectrum β -lactamase (ESBL)-producing *Enterobacteriaceae*. However, the emergence and rapid diffusion of carbapenemase-producing Gram negatives in the recent years is compromising the therapeutic efficacy of this class of antimicrobial drugs (2, 11, 16).

Recently, two *bla*_{OXA-163}-producing isolates of *Klebsiella pneumoniae* and *Enterobacter cloacae* have been identified in Argentina (15). OXA-163 differs from OXA-48 by a single amino acid substitution and a 4-amino-acid deletion only, but it works as an extended-spectrum oxacillinase showing a weaker ability to hydrolyze carbapenems than OXA-48 but a concurrent significant ability to hydrolyze extended-spectrum cephalosporins (15). The *bla*_{OXA-163} gene has been putatively proposed as the equivalent in South America of the *bla*_{OXA-48} gene that is increasingly identified in the southern and eastern Mediterranean countries (1, 6, 8, 14).

Here, we report two autochthonous cases of infection caused by two genetically unrelated OXA-163-carrying *K. pneumoniae* strains in Cairo, Egypt, in 2009 and 2010.

Case 1 was a 31-year-old woman hospitalized in Cairo (Egypt) on June 5, 2009, due to acute myelocytic leukemia. She was on therapy by doxorubicin and cytarabine. After 16 days since admission, she was transferred to an intensive care unit because of heart and respiratory failure. Initial antibiotic therapy was started with vancomycin, azithromycin, and imipenem. Blood culture was positive for a carbapenem-resistant strain of *K. pneumoniae* (isolate 18). The patient died on June 23.

Case 2 was a 25-year-old man with an 18-month history of acute lymphocytic leukemia. He was in outpatient follow-up until May 2010, when he was admitted to the hospital due to a lower respiratory tract infection. He was started on cefepime and amikacin on May 14, but the clinical course was complicated. A sputum culture grew carbapenem-resistant *K. pneumoniae* (isolate 78). No further data about the treatment outcome were available for this patient.

Both patients had never traveled outside Egypt.

The two carbapenem-resistant *K. pneumoniae* strains were sent to the molecular epidemiology laboratory of the Department of Sciences for Health Promotion "G. D'Alessandro," University of Palermo, Italy, for confirmation and typing. The two isolates were initially tested for antimicrobial susceptibility by disk diffusion according to the Clinical and Laboratory Standards Institute

(CLSI) guidelines (3). Susceptibility testing for extended-spectrum cephalosporins and carbapenems was performed with Etest strips (bioMérieux, Marnes-La-Coquette, France). Results were interpreted according to the CLSI interpretative criteria (3).

PCR screening for carbapenemase-encoding genes of classes A (*bla*_{KPC}, *bla*_{GES}), B (*bla*_{VIM}, *bla*_{IMP}), and D (*bla*_{OXA-48}) and for ESBL-encoding genes *bla*_{TEM}, *bla*_{OXA}, *bla*_{SHV}, and *bla*_{CTX-M} was performed as described previously (4). PCR products were purified and sequenced using the BigDye Terminator version 1.1 cycle sequencing kit (Applied Biosystems, Warrington, United Kingdom) and the ABI Prism 310 Genetic Analyzer (Applied Biosystems, Foster City, CA). The nucleotide (nt) and deduced protein sequences were analyzed with the software available from the National Center for Biotechnology Information website (www.ncbi.nlm.nih.gov). Outer membrane protein (OMP) gene amplification was conducted using *ompK35*-F and -R and *ompK36*-F and -R, and PCR products were sequenced using OMP primers according to Kaczmarek et al. (8). The sequences were analyzed by comparison with reported nucleotide sequences in GenBank. Expression levels of the *ompK35* and *ompK36* genes were not investigated.

To evaluate their clonal relationship, the two *K. pneumoniae* isolates were submitted to pulsed-field gel electrophoresis (PFGE) after XbaI DNA digestion. Strain differentiation by PFGE was based on the criteria of Tenover et al. (17). The rep-PCR DiversiLab Microbial Typing System (bioMérieux, Marcy l'Étoile, France) was also used with the DiversiLab *Klebsiella* kit. DNA fragment separation and detection were done using the Agilent 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA), and results were analyzed and interpreted using the Kullback-Leibler method, as previously reported (13). Moreover, multilocus sequence typing (MLST) was performed on both isolates according to the protocol described on the *K. pneumoniae* MLST website (<http://www.pasteur.fr/recherche/genopole>

Received 15 December 2011 Returned for modification 23 January 2012

Accepted 6 April 2012

Published ahead of print 18 April 2012

Address correspondence to Caterina Mammina, caterina.mammina@unipa.it.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JCM.06710-11

TABLE 1 Main characteristics of two OXA-163-producing *K. pneumoniae* isolates detected in Egypt, June 2009 and May 2010

<i>K. pneumoniae</i> strain (patient)	MIC ($\mu\text{g/ml}$) ^a							<i>bla</i> genes	Sequence type
	Cefotaxime	Ceftriaxone	Ceftazidime	Aztreonam	Imipenem	Meropenem	Colistin		
18 (case 1)	>256	>256	>256	>256	6	8	0.25	TEM-1, SHV-1, CTX-M15, OXA-163	37
78 (case 2)	>256	>256	>256	>256	6	12	0.25	TEM-1, SHV-1, CTX-M15, OXA-163	20

^a MICs assessed by Etest.

/PF8/mlst/*Kpneumoniae*.html). MLST results were compared with the international *K. pneumoniae* MLST database at the Pasteur Institute in Paris, France.

The two *K. pneumoniae* isolates exhibited resistance to extended-spectrum cephalosporins (cefotaxime, ceftazidime, cefepime), aztreonam, and carbapenems (meropenem and imipenem) (Table 1). Additionally, both isolates were resistant to ciprofloxacin and gentamicin and susceptible to colistin.

For both isolates, positive PCR results were obtained for the *bla*_{TEM}, *bla*_{SHV}, and *bla*_{CTX-M} genes, and sequencing identified the narrow-spectrum β -lactamase TEM-1 and SHV-1 genes and the ESBL CTX-M-15. A *bla*_{OXA-48}-like gene was detected by PCR in both isolates that, by sequencing, was identified as *bla*_{OXA-163} (www.lahey.org).

The *ompK35* and *ompK36* genes of isolates 18 and 78 were amplified, sequenced, and aligned with those of OMP gene sequences of *K. pneumoniae* MGH 78578. The *ompK35* gene of both isolates showed a frameshift mutation due to a nucleotide deletion at nt 873. Moreover, in the *ompK35* gene of isolate 18, sequencing revealed one substitution, T→A, at nucleotide position 108, generating a TAA nonsense mutation which resulted in a premature stop codon at amino acid position 36. Silent point mutations were also found at nt 303, 420, 537, 648, and 786 on the nucleotide sequence of both isolates. Comparison of the *ompK36* gene sequence with the wild-type sequence revealed that both had an insertion of six nucleotides (5'-GGCGAC-3') at positions 405 to 410, due to a duplication of the adjacent region at nucleotide positions 399 to 404. Sequencing of the *ompK36* sequence of isolate 78 detected also a 9-nucleotide deletion at positions 556 to 564.

PFGE and rep-PCR showed that the two isolates were genetically unrelated to each other (Fig. 1). Additionally, MLST attributed two different sequence types, ST37 and ST20, respectively, to the strains 18 and 78.

The main characteristics of the two isolates of *K. pneumoniae* are summarized in Table 1.

The findings of this study are of concern for a number of reasons. First, the detection of OXA-163 outside the continent where it has been firstly identified was unexpected. This finding seems to stand against the hypothesis of OXA-163 as the equivalent in South America of OXA-48 in the Mediterranean area (15). The two patients had no history of travel abroad. Moreover, possible contacts with source patients from Argentina, though it was not possible to investigate them in detail, appear to be unlikely due also to the 1-year-long interval of time between the two cases. An autochthonous emergence in Egypt of the OXA-163 resistance determinant appears to be the more plausible explanation. Second, the two *K. pneumoniae* isolates were genetically unrelated based upon PFGE and MLST. This rules out the clonal expansion of an OXA-163-producing strain in Egypt and, alternatively, sup-

ports the hypothesis of horizontal transmission of this genetic determinant via plasmids, according with the previous report (14). Notably, strains belonging to ST20 and ST37 have been previously identified in different countries as ESBL or carbapenemase producers (11, 12). Moreover, it is of special interest that OXA-163 has been identified in Egypt, a country of North Africa, a geographical area where OXA-48-producing Gram negatives are considered endemic (6, 9, 10). Third, both strains had much higher MICs for carbapenems than those reported by Poirel et al. (15). Studies suggest that outer membrane permeability defects coupled with ESBL production can confer resistance to carbapenems (5, 7, 9). This is likely to account for the higher level of carbapenem resistance detected in our strains. It is also noteworthy that the presence of similar mutations altering the properties of the OmpK35 and OmpK36 porins have been previously described in an ST37 *K. pneumoniae* clone carrying *bla*_{CTX-M-15} and exhibiting reduced susceptibility to carbapenems (5). However, our study did not definitively attribute the carbapenem resistance phenotype to OXA-163, because transfer of antimicrobial agent resistance experiments and plasmid analysis were not performed. This is a substantial limitation of our report.

As previously reported about OXA-163-producing strains in Argentina, because of its unique hydrolysis ability mimicking a class A ESBL, OXA-163 could be able to spread unrecognized as a




FIG 1 XbaI pulsed-field gel electrophoresis patterns and rep-PCR patterns of two OXA-163-carrying *Klebsiella pneumoniae* strains, Cairo, Egypt, 2009 and 2010. (A) Lanes: MW, XbaI-digested DNA of *Salmonella* Braenderup H9812; 1, *K. pneumoniae* strain 18; 2, *K. pneumoniae* strain 78. (B) Lanes: 1, *K. pneumoniae* strain 18; 2, *K. pneumoniae* strain 78.

carbapenemase (15). Surveillance of emergence and dissemination of enterobacteria with complex carbapenem resistance mechanisms is a serious public health problem where the costs can be unaffordable, such as low-resource countries. Further studies are urgently needed to assess prevalence of the *bla*_{OXA-163} genetic determinant worldwide.

Nucleotide sequence accession numbers. The *ompK35* of isolate 78 was assigned accession number [JQ660370](#), and the *ompK36* sequences of isolates 18 and 78 were assigned accession numbers [JQ660371](#) and [JQ660372](#), respectively, in the GenBank database.

REFERENCES

1. Benouda A, Touzani O, Khairallah MT, Araj GF, Matar GM. 2010. First detection of oxacillinase-mediated resistance to carbapenems in *Klebsiella pneumoniae* from Morocco. *Ann. Trop. Med. Parasitol.* **104**:327–330.
2. Carmeli Y, et al. 2010. Controlling the spread of carbapenemase-producing Gram-negatives: therapeutic approach and infection control. *Clin. Microbiol. Infect.* **16**:102–111.
3. Clinical and Laboratory Standards Institute. 2011. Performance standards for antimicrobial susceptibility testing; 21st informational supplement. CLSI M100-S21. Clinical and Laboratory Standards Institute, Wayne, PA.
4. Dalenne C, Da Costa A, Decré D, Favier C, Arlet G. 2010. Development of a set of multiplex PCR assays for the detection of genes encoding important beta-lactamases in *Enterobacteriaceae*. *J. Antimicrob. Chemother.* **65**:490–495.
5. García-Fernández A, et al. 2010. An ertapenem-resistant ESBL-producing *Klebsiella pneumoniae* clone carries a novel *OmpK36* porin variant. *Antimicrob. Agents Chemother.* **54**:4178–4184.
6. Goren MG, Chmelnitsky I, Carmeli Y, Navon-Venezia S. 2011. Plasmid-encoded OXA-48 carbapenemase in *Escherichia coli* from Israel. *J. Antimicrob. Chemother.* **66**:672–673.
7. Jacoby GA, Mills DM, Chow N. 2004. Role of β -lactamases and porins in resistance to ertapenem and other β -lactams in *Klebsiella pneumoniae*. *Antimicrob. Agents Chemother.* **48**:3203–3206.
8. Kaczmarek FM, Dib-Hajj F, Shang W, Gootz TD. 2006. High-level carbapenem resistance in a *Klebsiella pneumoniae* clinical isolate is due to the combination of *bla*_{ACT-1} β -lactamase production, porin *OmpK35/36* insertional inactivation, and downregulation of the phosphate transport porin *phoE*. *Antimicrob. Agents Chemother.* **50**:3396–3406.
9. Ktari S, et al. 2011. Spread of *Klebsiella pneumoniae* isolates producing OXA-48 β -lactamase in a Tunisian university hospital. *J. Antimicrob. Chemother.* **66**:1644–1666.
10. Nordmann P, Naas T, Poirel L. 2011. Global spread of carbapenemase-producing *Enterobacteriaceae*. *Emerg. Infect. Dis.* **17**:1791–1798.
11. Orsi GB, et al. 2011. Risk factors and clinical significance of ertapenem-resistant *Klebsiella pneumoniae* in hospitalised patients. *J. Hosp. Infect.* **78**:54–58.
12. Oteo J, et al. 2009. Emergence of CTX-M-15-producing *Klebsiella pneumoniae* of multilocus sequence types 1, 11, 14, 17, 20, 35 and 36 as pathogens and colonizers in newborns and adults. *J. Antimicrob. Chemother.* **64**:524–528.
13. Perez F, et al. 2010. Carbapenem-resistant *Acinetobacter baumannii* and *Klebsiella pneumoniae* across a hospital system: impact of post-acute care facilities on dissemination. *J. Antimicrob. Chemother.* **65**:1807–1818.
14. Poirel L, Bonnin RA, Nordmann P. 2012. Genetic features of the wide-spread plasmid coding for the carbapenemase OXA-48. *Antimicrob. Agents Chemother.* **56**:559–562.
15. Poirel L, et al. 2011. OXA-163, an OXA-48-related class D β -lactamase with extended activity toward expanded-spectrum cephalosporins. *Antimicrob. Agents Chemother.* **55**:2546–2551.
16. Poirel L, Naas T, Nordmann P. 2010. Diversity, epidemiology, and genetics of class D β -lactamases. *Antimicrob. Agents Chemother.* **54**:24–38.
17. Tenover FC, et al. 1995. Interpreting chromosomal DNA restriction patterns produced by pulsed-field gel electrophoresis: criteria for bacterial strain typing. *J. Clin. Microbiol.* **33**:2233–2239.