

Hepatitis B Virus X Protein Upregulates mTOR Signaling through IKK β to Increase Cell Proliferation and VEGF Production in Hepatocellular Carcinoma

Chia-Jui Yen^{1,2}, Yih-Jyh Lin^{3,9}, Chia-Sheng Yen^{4,9}, Hung-Wen Tsai^{5,9}, Ting-Fen Tsai⁶, Kwang-Yu Chang^{1,7}, Wei-Chien Huang⁸, Pin-Wen Lin³, Chi-Wu Chiang^{9*}, Ting-Tsung Chang^{2,9*}

1 Institute of Clinical Medicine, National Cheng Kung University College of Medicine and Hospital, Tainan, Taiwan, **2** Department of Internal Medicine, National Cheng Kung University College of Medicine and Hospital, Tainan, Taiwan, **3** Department of Surgery, National Cheng Kung University College of Medicine and Hospital, Tainan, Taiwan, **4** Department of General Surgery, Chi-Mei Medical Center, Tainan, Taiwan, **5** Department of Pathology, National Cheng Kung University College of Medicine and Hospital, Tainan, Taiwan, **6** Department of Life Sciences and Institute of Genome Sciences, National Yang-Ming University, Taipei, Taiwan, **7** Institute of Cancer Research, National Health Research Institutes, Tainan, Taiwan, **8** Center for Molecular Medicine and Graduate Institute of Cancer Biology, China Medical University and Hospital, Taichung, Taiwan, **9** Institute of Molecular Medicine, College of Medicine, National Cheng Kung University, Tainan, Taiwan

Abstract

Hepatocellular carcinoma (HCC), a major cause of cancer-related death in Southeast Asia, is frequently associated with hepatitis B virus (HBV) infection. HBV X protein (HBx), encoded by a viral non-structural gene, is a multifunctional regulator in HBV-associated tumor development. We investigated novel signaling pathways underlying HBx-induced liver tumorigenesis and found that the signaling pathway involving I κ B kinase β (IKK β), tuberous sclerosis complex 1 (TSC1), and mammalian target of rapamycin (mTOR) downstream effector S6 kinase (S6K1), was upregulated when HBx was overexpressed in hepatoma cells. HBx-induced S6K1 activation was reversed by IKK β inhibitor Bay 11-7082 or silencing IKK β expression using siRNA. HBx upregulated cell proliferation and vascular endothelial growth factor (VEGF) production, and these HBx-upregulated phenotypes were abolished by treatment with IKK β inhibitor Bay 11-7082 or mTOR inhibitor rapamycin. The association of HBx-modulated IKK β /mTOR/S6K1 signaling with liver tumorigenesis was verified in a HBx transgenic mouse model in which pIKK β , pS6K1, and VEGF expression was found to be higher in cancerous than non-cancerous liver tissues. Furthermore, we also found that pIKK β levels were strongly correlated with pTSC1 and pS6K1 levels in HBV-associated hepatoma tissue specimens taken from 95 patients, and that higher pIKK β , pTSC1, and pS6K1 levels were correlated with a poor prognosis in these patients. Taken together, our findings demonstrate that HBx deregulates TSC1/mTOR signaling through IKK β , which is crucially linked to HBV-associated HCC development.

Citation: Yen C-J, Lin Y-J, Yen C-S, Tsai H-W, Tsai T-F, et al. (2012) Hepatitis B Virus X Protein Upregulates mTOR Signaling through IKK β to Increase Cell Proliferation and VEGF Production in Hepatocellular Carcinoma. *PLoS ONE* 7(7): e41931. doi:10.1371/journal.pone.0041931

Editor: Michael Bouchard, Drexel University College of Medicine, United States of America

Received: February 18, 2012; **Accepted:** June 27, 2012; **Published:** July 27, 2012

Copyright: © 2012 Yen et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This study was supported by grants from the National Science Council NSC 98-2314-B-006-018-MY2, National Cheng Kung University Comprehensive Cancer Center in Southern Taiwan (DOH99-TD-C-111-003), and the Ministry of Education's 'Five Year, 50 Billion' project, NCKU Infectious Diseases and Signaling Research Center. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: ttchang@mail.ncku.edu.tw (T-TC); chiangcw@mail.ncku.edu.tw (C-WC)

† These authors contributed equally to this work.

Introduction

Hepatocellular carcinoma (HCC), which occurs frequently in Southeast Asia, is one of the most important causes of cancer-related death in the world [1,2,3]. According to epidemiological studies [4,5,6,7], there is a strong correlation between chronic hepatitis B virus (HBV) infection and the occurrence of HCC. HBV X protein (HBx) is a well-known viral non-structural gene that operates as a multifunctional regulator by modulating activity of host cellular genes such as p53 [11,12,13] and transactivating some transcription factors including AP-1, NF- κ B, CREB, and TBP [14,15]. Moreover, HBx is involved in the activation of multiple signaling pathways linked to cell proliferation and survival, such as RAS/RAF/MAPK, MEKK1/JNK, and PI3K/Akt [16,17,18]. Chronic inflammation is one of the key conditions of persistent HBV infection and has been implicated in tumor development [19,20,21]. The proinflammatory cytokines and

chemokines, such as tumor necrosis factor α (TNF- α), IL-1, IL-6, and IL-8, produced in microenvironments, have been known to promote tumor development [22,23]. TNF- α is considered one of the most important factors involved in inflammation-mediated tumorigenesis [24,25,26], and the transcription factor NF- κ B, a downstream signaling transducer of TNF- α , has been implicated in oncogenesis by promoting expression of genes related to cell proliferation and survival [27]. Activation of the inhibitor of nuclear factor κ B (I κ B) kinase (IKK) by TNF- α phosphorylates I κ Bs and promotes degradation of I κ Bs, resulting in nuclear translocation of NF- κ B and induction of NF- κ B downstream genes [28,29]. The involvement of the IKK/NF- κ B pathway in HBV-induced hepatitis and HCC is well documented [30,31,32], whereas effects of IKKs independent of NF- κ B on tumorigenesis have also been found [33,34,35]. It was recently reported that IKK β increased tumor development and tumor angiogenesis by

activating the mTOR signaling pathway through inhibiting tuberous sclerosis 1 (TSC1) [36,37,38]. Moreover, aberrant activation of the mTOR/ribosomal protein S6 kinase 1 (S6K1) signaling pathway increased cell proliferation and angiogenesis in a rat HCC model [39,40]. In the present study, we investigated whether HBx can modulate IKK β to inactivate TSC1's inhibition on mTOR so that it contributes to HCC development. We found that HBx modulated IKK β /TSC1/mTOR signaling and up-regulated cell proliferation and VEGF production in both unstimulated and TNF- α -stimulated hepatoma cells. We further used an HBx transgenic mouse model to verify whether HBx upregulates IKK β /TSC1/mTOR signaling *in vivo*, and to examine the association of upregulated IKK β /TSC1/mTOR signaling with increased VEGF expression and angiogenesis in liver tumorigenesis. Furthermore, we investigated the status of IKK β /TSC1/mTOR signaling in specimens from HBV-associated human hepatomas, and analyzed the relationship between the status of IKK β /TSC1/mTOR signaling and the prognosis of HCC patients. We conclude that IKK β activates mTOR signaling through TSC1 suppression to contribute to one crucial mechanism underlying HBx-dependent pathogenesis of HCC.

Methods

Plasmid and Cell Lines

The pcDNA6.0-HBx plasmid was constructed by cloning the cDNA product of the HBx gene into the pcDNA6.0 expression vector. The cDNA of the HBx gene was obtained by preparing RNA from serum of an HBV (+) patient followed by a reverse transcription-polymerase chain reaction (RT-PCR) using reverse transcriptase (SuperScript III; Invitrogen), oligo(dT) primers, and HBx primers: Forward 5'-AAGCTTGCTGCTCGGGTGTGCTGCCAA-3' and Reverse 5'-GGTACCGG CAGAGGTGAAAAAGTTGCA-3'. The sequence of the HBx cDNA was confirmed by sequencing analysis. The expression vectors payw1.2WT and payw*7 [41] for wildtype and HBx-defective HBV genome, respectively, were kind gifts from Dr. Jack R. Wands (The Warren Alpert Medical School of Brown University).

Hep3Bx and HepG2x cells were derivatives of human hepatoma Hep 3B and Hep G2 cells (both were from ATCC), respectively, stably expressing the HBx gene, and were established by transfecting Hep3B and HepG2 cells with pcDNA6.0-HBx




Figure 1. The IKK β /TSC1/mTOR signaling pathway is activated by HBx. (A). Expression of HBx mRNA in Hep3Bx, HepG2x, and parental Hep3B and HepG2 cells was detected using semi-quantitative RT-PCR. Levels of GAPDH mRNA were used as an internal control. RNAs of a HBV-positive patient's serum (P) and RNAs of a control HBV-negative serum (N) were used as controls. (B). Levels of HBx protein were detected in lysates of Hep3Bx, HepG2x, and parental Hep3B and HepG2 cells using Western blotting by antibody specific against HBx protein and β -actin. (C). Levels of pIKK β (S181), pTSC1 (S511), pS6K1 (T389), total IKK β , total TSC1, total S6K1, and β -actin were assessed in lysates of Hep3Bx, HepG2x, and parental Hep3B and HepG2 cells using Western blotting by specific antibody as indicated. (D). Data shown are ratios of viable cells in Hep3Bx and HepG2x cells relative to that in Hep3B and HepG2 cells (set as 1), respectively, at 24 h after seeding using MTT assay. (E). Levels of pIKK β (S181), pS6K1 (T389), total IKK β , total S6K1, HBx, and β -actin were assessed in lysates of Huh7 cells transfected with empty vector alone, payw1.2WT, or payw*7. (F). Levels of pIKK β (S181), pTSC1 (S511), pS6K1 (T389), total IKK β , total TSC1, total S6K1, and β -actin were assessed in lysates of Hep3B and Hep3Bx with or without TNF- α treatment using Western blotting as described earlier. doi:10.1371/journal.pone.0041931.g001




Figure 2. TNF- α -stimulated increases of pTSC1 (S511), pS6K1 (T389) and cell proliferation in Hep3Bx and HepG2x cells are blocked by the IKK β inhibitor Bay 11-7082, siRNA specific for IKK β , and the mTOR inhibitor rapamycin. (A). Lysates of Hep3Bx and HepG2x cells treated with or without TNF- α in the presence or absence of Bay 11-7082 were analyzed for levels of pIKK β (S181), pTSC1 (S511), pS6K1 (T389), total IKK β , total TSC1, total S6K1, and β -actin using Western blotting as described earlier. (B). Lysates of Hep3Bx and HepG2x cells with transfection of IKK β siRNAs or control siRNAs were assessed for levels of pTSC1 (S511), pS6K1 (T389), total IKK β , total TSC1, total S6K1, and β -actin. (C). Lysates of Hep3Bx and HepG2x cells treated with or without TNF- α in the presence or absence of rapamycin were analyzed for levels of pIKK β (S181), pS6K1 (T389), total IKK β , and total S6K1. (D). Data shown are ratios of viable cells in Hep3B, Hep3Bx, HepG2, and HepG2x cells treated with or without TNF- α in the presence or absence of Bay11-7082 or rapamycin relative to that in Hep3B and HepG2 cells without any treatment (set as 1), at 24 h after seeding using MTT assay. Data are shown as means \pm S.D. of three experiments. Comparisons were made between different groups as indicated. *P<0.001 is determined by X test. doi:10.1371/journal.pone.0041931.g002

using Lipofectamine (Invitrogen) followed by drug selection [42]. These cells were maintained at 37°C in a 5% CO₂ incubator with Dulbecco's modified Eagle's/F12 medium plus 10% fetal bovine serum.

Antibodies and Western Blotting

The primary antibodies used in this study were anti-TSC1 (37-0400; Zymed Laboratories, Inc., San Francisco, CA), anti-phosphorylated S6 kinase (T389) (9205; Cell Signaling Technology, Inc., Beverly, MA), anti-S6 kinase (2215; Cell Signaling), anti-phosphorylated IKK β (S181) (2681; Cell Signaling), anti-IKK β (2684; Cell Signaling), anti-HBx (ab235; Abcam Co., Cambridge, UK), anti-VEGF-A (Santa Cruz Biotechnology, Inc., Santa Cruz,

CA), anti-CD31 (ab28364; Abcam Co., Cambridge, UK), and anti-actin (A2066; Sigma-Aldrich Co., St Louis, MO). The rabbit polyclonal antibody against the phospho-S511 of TSC1 was a kind gift from Dr Mien-Chie Hung (MD Anderson Cancer Center, Houston, TX). The expression of IKK β , pIKK β (S181), TSC1, pTSC1 (S511), S6K1 or pS6K1 (T389) was detected in cell lysates prepared from cells lysed as described previously [37]. Fifty micrograms of total protein lysates were resolved using sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE), transferred to polyvinylidene difluoride (PVDF) membranes, and then probed with specific antibodies and HRP-conjugated secondary antibodies. Immunoblots were then developed using enhanced chemiluminescence.




Figure 3. Expression of VEGF is increased in Hep3Bx and HepG2x cells and is further enhanced by TNF- α and blocked by IKK β inhibitor Bay 11-7082 or the mTOR inhibitor rapamycin. (A). The expression levels of secreted VEGF in the culture medium of Hep3B, Hep3Bx, HepG2, and HepG2x cells were measured by ELISA assay as described in the Methods. (B). The expression levels of VEGFA mRNA were assessed in Hep3B, Hep3Bx, HepG2, and HepG2x cells using semi-quantitative RT-PCR (left) or real-time RT-PCR (right) as described in the Methods. (C). The amounts of secreted VEGF in the culture medium of Hep3B, Hep3Bx, HepG2, or HepG2x cells treated with or without TNF- α in the presence or absence of Bay11-7082 or rapamycin were measured by ELISA assay. Data are shown as means \pm S.D. of three experiments. Comparisons were made between different groups as indicated. * P <0.001 is determined by X test. doi:10.1371/journal.pone.0041931.g003

Inhibitor Treatment and Knockdown of Gene Expression by siRNAs

BAY 11-7082 and rapamycin were purchased from Calbiochem (San Diego, CA), and recombinant human TNF- α was from Roche Applied Sciences (Indianapolis, IN). Hep3B, Hep3Bx, HepG2, and HepG2x cells were grown in complete medium and exposed to 20 μ g/ml TNF- α (Roche, Indianapolis, IN) with or without pretreatment of 40 μ M BAY 11-7082 for 45 min (Calbiochem, San Diego, CA) or 100 nM Rapamycin (Calbiochem, San Diego, CA) for 3 h.

The MTT assay was used to measure cell proliferation and viability in 5,000 cells seeded onto 96-well plates, treated with TNF- α with or without BAY 11-7082 or Rapamycin for 24 h. The amount of VEGF protein in the culture supernatants was assessed using ELISA (BioSource; Invitrogen Corp., Carlsbad, CA) according to the manufacturer’s instructions. IKK β siRNAs (Smartpool [M-003503]; Upstate Biotechnology, Charlottesville, VA) and control siRNAs (Smartpool [D-001206-13-05]; Upstate) were transfected into cells using Lipofectamine to knock down IKK β expression.


Real-time RT-PCR and Quantitative Real-time RT-PCR for mRNAs

Total RNA was extracted from cells 48 h after transfection using Rezol (ProTech Technology Enterprise Co., Ltd., Taipei, Taiwan) according to the manufacturer’s instructions. Subsequently, 4 μ g RNA was converted to cDNA using SuperScript III, oligo (dT)

primers, and PCR using the following primers: HBx Forward 5’–AAGCTTGCTGCTCGGGTGTGCTGCCAA–3’ and Reverse 5’–GGTACCGG CAGAGGTGAAAAAGTTGCA–3’; VEGFA Forward 5’–CATGAAC TTTCTGC TGTCTTGG–3’ and Reverse 5’–CATTGTGTGCTGTAGGAAGC–3’; GAPDH Forward 5’–TGAAGGTCGGAGTCAACGGATTGGT–3’ and Reverse 5’–CATG TGGGCCATGAGGTCACCAC–3’. For real time RT-PCR reactions, 1 μ g of total RNA from each sample were used in the RT reaction (M-MLV Reverse Transcriptase; Invitrogen). The TaqMan gene expression real time PCR assays (ABI PRISM 7900 HT Sequence Detection System; Applied Biosystems) were used to assess the mRNA expression levels of the endogenous VEGFA and GAPDH (Applied Biosystems, Foster City, CA, USA; assay ID: Hs00900055_m1 for VEGFA and Hs99999905_m1 for GAPDH). Expression analysis was done in triplicate for each sample. In each run, the endogenous control gene (GAPDH) and one no-template-control (NTC) were also run in triplicate. The fold difference for each sample was obtained using the following equation: $2^{-\Delta\Delta Ct}$. Ct is the threshold cycle.

HBx Transgenic Mice, Tissue Preparation, and Immunohistochemical Analysis

The lines of HBx transgenic mice used in this study were established and described elsewhere [43]. The HBx transgenic mice were bred in a specific pathogen-free environment and all mouse experiments complied with the guidelines in the “Guide for the Care and Use of Laboratory Animals” (NIH publications 86-23 revised 1985) and were approved by the Institutional Animal



compared to the normal liver tissues of the wild-type age-matched mouse. The HBx mRNA levels were also measured by RT-PCR and the GAPDH mRNA levels were used as an internal control. The relative levels of pIKK β (S181), pS6K1 (T389), and VEGF-A were quantified by densitometry and normalized with total IKK β , total S6K1, and actin. Results are shown as ratios of average levels of pIKK β (S181) pS6K1 (T389), and VEGF-A in non-tumor and tumor parts of liver tissues of three HBx transgenic mice (#824, #825, and #826) relative to that in the normal liver tissues of the wild-type age-matched mouse (set as 1). Data are shown as means \pm S.D. of measurements of three mice. (D). Immunohistochemistry analyses show expression levels of pIKK β (S181), pS6K1 (T389), VEGF-A, and CD31 in normal liver tissues of the wild-type mouse, and non-tumor and tumor parts of liver tissues of HBx transgenic mice. One representative data are shown. N = 3. doi:10.1371/journal.pone.0041931.g004

Care and Use Committee (IACUC) of College of Medicine, National Cheng Kung University (Approval Number: 98129). The HBx transgenic mice developed hepatic tumor after 13 to 16 months of age. To ensure the mice could develop hepatic tumor, the transgenic mice were bred for up to 21 months, and approximately 90% incidence of HCC was observed in HBx transgenic male mice at an age of 19–20 months. After the mice had developed liver cancer, the mice were sacrificed, and liver tissue was collected, extracted, fixed, and stained with hematoxylin and eosin as previously described [42]. pIKK β , pS6K1, VEGF-A, and CD31 expression was immunohistochemically detected on paraffin-embedded liver sections (3 μ m) using specific antibodies mentioned earlier.

Patients' Characteristics

Institutional Review Board of the Human Investigation Committee of College of Medicine, National Cheng Kung University approved the study. Written informed consent was obtained from patients participating in this study. A total of ninety-five patients admitted to National Cheng-Kung University Hospital (Tainan, Taiwan) with HBV-associated HCC who received curative surgery between 1 January 2003 and 31 December 2006 were enrolled, and samples of their resected liver tumor tissue were assembled in a tissue microarray. All 95 patients showed positive for serum HBV surface antigen and for HBx in tumor specimens analyzed using PCR, and negative for antibodies to the hepatitis C virus. The patients were regularly followed up at clinic visit every 1 to 3 months after curative surgery. The patients included 70 (73.7%) males and 25 (26.3%) females with age range of 44 to 77 years (mean age 60.7 years). The median follow-up time was 38 months (range, 6 to 53 months). At the end of the follow-up, 28 patients had died of disease. The 1-year disease specific survival rate was 92.1% and 3-year disease specific survival rate was 74.2%.

Immunohistochemistry and Clinical Association Study

Immunohistochemical staining for pIKK β , pTSC1, and pS6K1 protein expression was examined on adjacent 4- μ m formalin-fixed paraffin-embedded tissue sections. An experienced gastrointestinal pathologist reviewed all specimens. Amino-ethylcarbazole chromogen was used for visualization. The positive protein staining was only considered in cytoplasmic immunoreactivity of cancer cells on a semi-quantitative scale that combined staining intensity and percentage of positively stained cells. Staining intensity was evaluated as low (0–10% positive cancer cells) or high (>10% positive cancer cells). Statistical analyses were done using a χ^2 test, Fisher's exact test, or Kaplan-Meier survival test. Significance was set at $P < 0.05$.

Results

HBx Upregulates IKK β and mTOR Activity

We established human hepatoma Hep 3Bx and Hep G2x cell lines which stably express HBx to investigate the effects of HBx overexpression on IKK β /TSC1/mTOR signaling. The expres-

sion of HBx in Hep3Bx and Hep G2x was confirmed by both RT-PCR and Western blotting analysis (Fig. 1A, B). Next, we evaluated whether HBx overexpression affected levels of phospho-IKK β (pIKK β) (S181), phospho-TSC1 (pTSC1) (S511), and mTOR downstream phospho-S6K1 (pS6K1) (T389). While there were no significant differences in total protein levels of IKK β , TSC1, S6K1 in cells with or without overexpression of HBx (Fig. 1C), we found overexpression HBx enhanced levels of pIKK β , pTSC1, and pS6K1. Consistently, we observed significantly increased proliferation in cells stably expressing HBx compared with control cells expressing vector only (Fig. 1D). To verify the role of HBx in enhancing levels of pIKK β and pS6K1 in a more natural setting mimicking HBV infection, we transfected Huh7 cells with empty vector alone, wildtype HBV full genome expression vector payw1.2WT, or payw*7 which is a paywWT mutant vector harboring an ochre termination signal after codon 7 in the HBx open reading frame [41]. We found that levels of phosphorylation of IKK β and S6K1 in cells transfected with wildtype HBV expression vector payw1.2WT were higher than that in cells transfected with HBx defective mutant payw*7 or vector only (Fig. 1E). Moreover, when cells were treated with the proinflammatory cytokine TNF- α , overexpression of HBx synergistically enhanced TNF- α -stimulated pIKK β , pTSC1, and pS6K1 (Fig. 1F). These results demonstrate that HBx upregulates basal and TNF- α -induced IKK β and mTOR activity.

HBx Deregulates TSC1/mTOR Signaling and Increases Cell Proliferation through IKK β

We found that HBx enhanced basal and TNF- α -stimulated IKK β and mTOR activity, and concomitantly increased phosphorylation and inactivation of TSC1 (Fig. 1). We hypothesized that the effect of HBx on mTOR was mediated by IKK β -mediated phosphorylation and inactivation of TSC1. The Hep3Bx and HepG2x cells were treated with the IKK β inhibitor Bay11-7082 in the presence or absence of TNF- α , and the levels of pIKK β (S181), pTSC1 (S511), and pS6K1 (T389) were examined. In agreement with our hypothesis, inhibition of IKK β by Bay11-7082 abolished the TNF- α -stimulated phosphorylation of TSC1 and S6K1 in both Hep3Bx and HepG2x cells (Fig. 2A), the finding of HBx-upregulated mTOR activity mediated by IKK β was in parallel confirmed using a specific IKK β siRNA to knockdown IKK β expression (Fig. 2B). To confirm that the HBx-mediated increase in phosphorylation of S6K1 was through mTOR activation, cells were treated with mTOR inhibitor rapamycin when stimulated with TNF- α . As shown in Fig. 2C, basal and TNF- α -stimulated phosphorylation of S6K1 was abolished by rapamycin treatment in both Hep3Bx and HepG2x cells. Consistently, TNF- α -stimulated cell proliferation in Hep3Bx and HepG2x cells was completely blocked by treatment with Bay11-7082 or rapamycin (Fig. 2D). These results suggest that HBx activates mTOR activity and cell proliferation through IKK β -mediated inactivation of TSC1.




Figure 5. Positive association between pIKK β (S181), pTSC1(S511), and pS6K1(T389) in HBV-associated human HCC specimens. (A). Immunohistochemistry analysis of pIKK β (S181), pTSC1(S511) and pS6K1(T389) in tumor tissues of 95 human HBV-associated HCC specimens. Results of one representative specimens stained by specific antibodies are shown. (B). Upper graph shows percentages of specimens with low or high pIKK β (S181) expression in which pS6K1 (T389) expression was high or was not observed (low). Lower graph shows percentages of specimens with low or

high pIKK β (S181) expression in which pTSC1 (S511) expression was high or was not observed (low). Positive correlations were noted between pIKK β (S181) and pS6K1 (T389) (* P <0.01) and between pIKK β (S181) and pTSC1 (S511) (* P <0.01). (C) The Kaplan-Meier disease-free survival curves show that expression of pIKK β (S181) (p =0.003), pTSC1 (S511) (p =0.048), or pS6K1 (T389) (p =0.0027) is associated with early tumor recurrence. Co-expression of pIKK β (S181) and pS6K1 (T389) (p =0.0013) was a better predictor of patients' recurrence-free time survival in HCC patients who received curative surgery for up to 48-month investigation.
doi:10.1371/journal.pone.0041931.g005

HBx Increases VEGF Production through IKK β /TSC1/mTOR Signaling

Since activation of mTOR pathway could up-regulate the angiogenesis process [36,37], we used IKK β inhibitor Bay 11-7082 and mTOR inhibitor rapamycin to clarify whether HBx could induce VEGF production through IKK β /TSC1/mTOR signaling. The expression level of VEGF in cell culture supernatant was checked by ELISA, and we observed increased VEGF production in the Hep 3Bx and Hep G2x cells compared with control Hep 3B and Hep G2 cells (p <0.001) (Fig. 3A). Results of semi-quantitative and real-time RT-PCR assay showed that the expression of VEGF-A messenger RNA was also upregulated in Hep 3Bx and Hep G2x cancer cells (Fig. 3B). In order to further clarify whether HBx enhanced VEGF production was related to IKK β /TSC1/mTOR signaling, effects of TNF- α (20 ng/ml) combined with or without pretreatment of IKK β inhibitor Bay 11-7082 or mTOR inhibitor rapamycin were examined. We found that TNF- α -enhanced VEGF production in Hep 3Bx and Hep G2x was substantially suppressed by treatment with Bay 11-7082 or rapamycin, as compared with that in control Hep 3B and Hep G2 cells (p <0.001) (Fig. 3C). These observations indicate that HBx increases the VEGF production of the hepatoma cells due to modulation of the IKK β /TSC1/mTOR pathway.

HBx-mediated Upregulation of IKK β /TSC1/mTOR Signaling is Associated with Liver Tumorigenesis in HBx Transgenic Mice

To further understand the contribution of the HBx-modulated IKK β /TSC1/mTOR signaling pathway in liver tumorigenesis, we used an HBx transgenic mouse model which has been shown to develop liver tumors [42] (Fig. 4A, B). We found that the average expression levels of both pIKK β and pS6K1 were higher in both the non-tumor parts and liver tumor tissues of the HBx transgenic mice liver compared with the normal liver tissues of the wild-type mice (Fig. 4C). Concurrently, significantly higher levels of VEGF-A were also found in liver tumor tissues of HBx transgenic mice compared with the normal liver tissues of the wild-type mice (Fig. 4C). In agreement with the results of Western blottings, immunohistochemical staining revealed higher expression levels of pIKK β , pS6K1, VEGF-A, and CD31 in liver tumors of HBx transgenic mice compared with adjacent non-tumor parts and wild-type mice liver tissues (Fig. 4D). Thus, findings of the HBx transgenic mouse model support that HBx enhances the IKK β /mTOR signaling pathway and promotes VEGF-A production and new vessels formation.

Immunohistochemical Staining Reveals Positive Correlations between pIKK β and pTSC1, and between pIKK β and pS6K1 in HBV-associated Human HCC Tissue Specimens

To validate the relevance of the upregulation of TSC1/mTOR pathway via IKK β signaling in HBV-associated human HCC, we evaluated the expression of pIKK β , pTSC1, and pS6K1 in a tissue microarray of 95 HBV-associated human HCC tissue specimens by immunohistochemical staining. All 95 patients who received curative surgery were positive for both serum HBV surface antigen

and HBx, but negative for antibodies to hepatitis C virus (data not shown). Expression of pIKK β , pTSC1, and pS6K1 was detected in human HCC tissue specimens (Fig. 5A). Analysis of immunohistochemical staining of 95 HCC tissue specimens revealed that pS6K1 was detected in 35 (60%) of the 58 specimens with high pIKK β expression, but it was detected in only 10 (27%) of the 37 specimens with low pIKK β expression, indicating that pS6K1 expression was positively associated with pIKK β expression (P <0.01) (Fig. 5B, upper). More, pTSC1 expression was detected in 32 (55%) of the 58 specimens with high pIKK β expression, but it was detected in only 4 (11%) of the 37 specimens with low pIKK β expression (P <0.01) (Fig. 5B, lower). We next analyzed the correlation of the expression of pIKK β (S181), pTSC1 (S511), and pS6K1 (T389) in HCC specimens with patients' recurrence-free survival data. The Kaplan-Meier recurrence-free survival curves showed that high levels of pIKK β (S181), pTSC1 (S511) and pS6K1 (T389) were associated with early recurrence of HCC (Fig. 5C). Multivariate analysis indicated that expression of pIKK β (S181) (HR, 2.37; 95% CI, 1.33–4.2 [P =0.003]), pTSC1 (S511) (HR, 2.07; 95% CI, 1.01–4.26 [P =0.048]) and pS6K1 (T389) (HR, 2.56; 95% CI, 1.39–4.74 [P =0.0027]) were predictors of patients' recurrence-free time (Table 1). In addition, concomitant expression of pIKK β (S181) and pS6K1 (T389) was a better predictor of patients' recurrence-free survival (HR, 4.11; 95% CI, 1.65–7.92 [P =0.0013]) than each factor alone (Table 1). Taken together, these data suggest that HBx-modulated IKK β /TSC1/mTOR pathway may play a crucial role in HBV-associated human HCC development and progression.

Discussion

The link between HBV and the development of hepatocellular carcinoma has been well established, but the pathogenic mechanism responsible for the transformation of normal hepatocytes to HCC is still far from understood, especially in view of the contribution of HBx to cancer progression. We have provided evidence that HBx activates IKK β , which leads to inactivation of TSC1 and activation of mTOR/S6K1 and to the production of angiogenesis factor VEGF-A in HBx expressing hepatoma cells and liver tissues of HBx transgenic mice. The HBx-associated IKK β /TSC1/mTOR signaling pathway may play a molecular switch that allows HBV-related HCC tumor progression. This is likely to be clinically relevant to pathogenesis as we found a statistically significant correlation between the phosphorylation of IKK β and phosphorylation of TSC1, and between the phosphorylation of IKK β and phosphorylation of S6K1 in HBV-associated HCC specimens.

Chronic hepatitis B virus (HBV) infection causes an inflammation process in the normal liver tissue, resulting in liver damage that may subsequently evolve into liver cirrhosis and tumor development. The proinflammatory cytokine TNF- α has been shown to play a promoting role for tumor development [21,23,44], and activation of NF- κ B, a downstream signaling transducer of TNF- α , has long been implicated in the development of HBV-associated HCC [45,46]. HBx has been shown to activate NF- κ B by directly interacting with NF- κ B [47,48], up-regulating TNF- α expression [48], or promoting phosphorylation and degradation of I κ B [50]. HBx was also shown to activate NF- κ B and up-

Table 1. Recurrence free survival in multivariate analysis.

Var	Hazard Ratio (95% CI)	P-value
Gender		
Male	1.0	
Female	0.5	0.08
	(0.3–1.1)	
Age		
<50 years	1.0	
≥ 50 years	0.7	0.41
	(0.4–1.5)	
Albumin		
Albumin <3.5 g/dl	1.0	
Albumin ≥ 3.5 g/dl	0.6	0.17
	(0.3–1.2)	
AFP		
AFP<400 ng/ml	1.0	
AFP>400 ng/ml	0.5	0.08
	(0.2–1.1)	
Differentiation		
Well	1.0	
Moderate	0.9	0.77
	(0.5–1.7)	
Poor	1.3	0.55
	(0.6–3.1)	
Primary tumor T stage		
T1	1.0	
T2& T3&T4	1.2	0.59
	(0.6–2.4)	
BCLC		
A stage	1.0	
B stage	1.8	0.07
	(1.0–3.2)	
pIKKβ(S181)		
Low expression	1.0	
High expression	2.37	0.003
	(1.33–4.2)	
pTSC1(S511)		
Low expression	1.0	
High expression	2.07	0.048
	(1.01–4.26)	
pS6K1(T389)		
Low expression	1.0	
High expression	2.56	0.0027
	(1.39–4.74)	
pIKKβ(S181) & pS6K1(T389)		
Low expression & Low expression	1.0	
High expression & High expression	4.11	0.0013
	(1.65–7.92)	

doi:10.1371/journal.pone.0041931.t001

regulate genes involved in cell invasion by IKK β activation [51].

Recently, HBx was shown to activate NF- κ B through modulating TRAF2/TAK1 signaling cascade [52]. The mechanism by which HBx activates IKK β remains unclear, but it is conceivable that HBx may activate IKK β by up-regulating TNF- α receptor stimulation through increasing TNF- α expression [49] and/or by modulating the TRAF2/TAK1 signaling module to increase IKK β activation [52]. Additionally, HBx may activate IKK β through activation of Ras/Raf/MEK [53] or protein kinase C (PKC) pathway [54], which has been known to be associated with IKK activation. Nevertheless, HBx may use alternative mechanisms to activate IKK β . IKKs' activities independent of NF- κ B have been reported [33,34,35], and IKK β was found to phosphorylate TSC1 and block the inhibitory effect of TSC1 on mTOR activity, resulting in increased tumorigenesis and angiogenesis [36,37,38]. Consistent with these findings, our results demonstrate that HBx up-regulates IKK β to deregulate TSC1/mTOR signaling and to promote cell proliferation and VEGF production. Moreover, activation of the mTOR signaling pathway has been found in a significant portion of HCC examined, and mTOR inhibition showed antitumoral effects, although the relationship between the status of HBx and mTOR activation was not known [39,55,56]. HBx was demonstrated to enhance VEGF expression by up-regulation of HIF-1 α transcription [57,58]. Our results showed that HBx can modulate IKK β /TSC1/mTOR signaling to increase VEGF-A production and TNF- α further up-regulates VEGF-A production in Hep3B and HepG2 cells overexpressing HBx. Similarly, using an HBx transgenic mouse liver tumor model, we demonstrated higher expression of pIKK β , pS6K1, and VEGF-A, and neovascularization in tumor tissues when compared with the non-neoplastic area of the HBx transgenic liver.

We demonstrated that treatment of liver cancer cells with mTOR inhibitor rapamycin and IKK β inhibitor Bay11-7082 effectively blocked the HBx-induced cell proliferation and VEGF-A production. Currently available treatments for non-operable advanced HCC patients, including local ablation therapy (radio-frequency ablation and percutaneous ethanol injection), transarterial chemo-embolization (TACE), chemotherapy and radiotherapy, are ineffective, resulting in very poor survival rates of patients [2,3]. Furthermore, our observations in specimens of the clinical HBV-associated HCC patients demonstrated that the expression of pIKK β (S181), pTSC1 (S511), and pS6K1 (T389) in human HCC tissue samples was correlated with early tumor recurrence and poor patients' survival. Moreover, we showed that the combination of pIKK β (S181) and pS6K1 (T389) expression was a better predictor of survival. Our findings suggest that pIKK β (S181) and pTSC1 (S511) and pS6K1 (T389) might be used as prediction of poor treatment outcomes of HCC patients. Therefore, HCC patients who have a significantly elevated risk of poor treatment outcomes should receive more intensive therapy; for example, surgery followed by adjuvant chemotherapy or target therapy.

In conclusion, we demonstrate that HBx deregulates TSC1/mTOR signaling through IKK β and renders liver cancer cells more sensitive to TNF- α stimulation in activating mTOR downstream S6K1 activity through IKK β signaling. Activation of the IKK β /mTOR pathway occurs concomitantly with increased cell proliferation and angiogenesis, which may associate with the progression of the HCC. Consistently, blocking IKK β or mTOR signaling with Bay 11-7082 or rapamycin, respectively, inhibits the liver cancer cell growth and VEGF-A production, suggesting that inhibitors of IKK β or mTOR signaling may be useful as new therapeutics for the treatment of HBV-associated HCC.

Acknowledgments

We are grateful to Drs. Margherita Melegari and Jack R. Wands for providing the HBV expression constructs. We thank Ms. Ming-Yu Chang, Ms. Han-Yu Lin, and Mr. Chih-Chieh Tsao for their technical assistance.

References

- Mazzanti R, Gramantieri L, Bolondi L (2008) Hepatocellular carcinoma: epidemiology and clinical aspects. *Mol Aspects Med* 29, 130–143.
- Kulik LM, Mulcahy MF, Omary RA, Salem R (2007) Emerging approaches in hepatocellular carcinoma. *J Clin Gastroenterol* 41, 839–854.
- Rougier P, Mitry E, Barbare J-C, Taieb J (2007) Hepatocellular Carcinoma (HCC): An Update. *Semin Oncol* 34, Supplement 1: S12–S20.
- Marotta F, Vangieri B, Cecere A, Gattoni A (2004) The pathogenesis of hepatocellular carcinoma is multifactorial event. Novel immunological treatment in prospect. *Clin Ter* 155, 187–199.
- Cougot D, Neuveut C, Buendia MA (2005) HBV induced carcinogenesis. *J Clin Virol* 34, Suppl 1: S75–78.
- Anzola M (2004) Hepatocellular carcinoma: role of hepatitis B and hepatitis C viruses proteins in hepatocarcinogenesis. *J Viral Hepat* 11, 383–393.
- Barazani Y, Hiatt JR, Tong MJ, Busuttil RW (2007) Chronic viral hepatitis and hepatocellular carcinoma. *World J Surg* 31, 1243–1248.
- Kaido T, Uemoto S (2008) Recent evidence in the treatment of small hepatocellular carcinoma. *Hepatogastroenterology* 55, 1460–1462.
- Lau WY, Lai EC (2008) Hepatocellular carcinoma: current management and recent advances. *Hepatobiliary Pancreat Dis Int* 7, 237–257.
- Yamazaki S, Takayama T (2008) Surgical treatment of hepatocellular carcinoma: evidence-based outcomes. *World J Gastroenterol* 14, 685–692.
- Koskinas J, Petraki K, Kavantzis N, Rapti I, Kountouras D, et al. (2005) Hepatic expression of the proliferative marker Ki-67 and p53 protein in HBV or HCV cirrhosis in relation to dysplastic liver cell changes and hepatocellular carcinoma. *J Viral Hepat* 12, 635–641.
- Dewantoro O, Gani RA, Akbar N (2006) Hepatocarcinogenesis in viral Hepatitis B infection: the role of HBx and p53. *Acta Med Indones* 38, 154–159.
- Park US, Su JJ, Ban KC, Qin L, Lee EH, et al. (2000) Mutations in the p53 tumor suppressor gene in tree shrew hepatocellular carcinoma associated with hepatitis B virus infection and intake of aflatoxin B1. *Gene* 251, 73–80.
- Chaparro M, Sanz-Cameno P, Trapero-Marugan M, Garcia-Buey L, Moreno-Otero R (2007) Mechanisms of angiogenesis in chronic inflammatory liver disease. *Ann Hepatol* 6, 208–213.
- Edamoto Y, Hara A, Biernat W, Terracciano L, Cathomas G, et al. (2003) Alterations of RB1, p53 and Wnt pathways in hepatocellular carcinomas associated with hepatitis C, hepatitis B and alcoholic liver cirrhosis. *Int J Cancer* 106, 334–341.
- Tommasi S, Pinto R, Pilato B, Paradiso A (2007) Molecular pathways and related target therapies in liver carcinoma. *Curr Pharm Des* 13, 3279–3287.
- McCubrey JA, Steelman LS, Abrams SL, Lee JT, Chang F, et al. (2006) Roles of the RAF/MEK/ERK and PI3K/PTEN/AKT pathways in malignant transformation and drug resistance. *Adv Enzyme Regul* 46, 249–279.
- Kasai Y, Takeda S, Takagi H (1996) Pathogenesis of hepatocellular carcinoma: a review from the viewpoint of molecular analysis. *Semin Surg Oncol* 12, 155–159.
- Coussens LM, Werb Z (2002) Inflammation and cancer. *Nature* 420, 860–867.
- Balkwill F, Mantovani A (2001) Inflammation and cancer: back to Virchow? *Lancet* 357, 539–545.
- Lu H, Ouyang W, Huang C (2006) Inflammation, a key event in cancer development. *Mol Cancer Res* 4, 221–233.
- Rollins BJ (2006) Inflammatory chemokines in cancer growth and progression. *Eur J Cancer* 42, 760–767.
- Raman D, Baugher PJ, Thu YM, Richmond A (2007) Role of chemokines in tumor growth. *Cancer Lett* 256, 137–165.
- Luo JL, Maeda S, Hsu LC, Yagita H, Karin M (2004) Inhibition of NF-kappaB in cancer cells converts inflammation-induced tumor growth mediated by TNFalpha to TRAIL-mediated tumor regression. *Cancer Cell* 6, 297–305.
- Greten FR, Eckmann L, Gretchen TF, Park JM, Li ZW, et al. (2004) IKKbeta links inflammation and tumorigenesis in a mouse model of colitis-associated cancer. *Cell* 118, 285–296.
- Clement JF, Meloche S, Servant MJ (2008) The IKK-related kinases: from innate immunity to oncogenesis. *Cell Res* 18, 889–899.
- Basseres DS, Baldwin AS (2006) Nuclear factor-kappaB and inhibitor of kappaB kinase pathways in oncogenic initiation and progression. *Oncogene* 25, 6817–6830.
- Li H, Lin X (2008) Positive and negative signaling components involved in TNFalpha-induced NF-kappaB activation. *Cytokine* 41, 1–8.
- Scheidereit C (2006) I kappa B kinase complexes: gateways to NF-kappaB activation and transcription. *Oncogene* 25, 6685–6705.
- Budhu A, Wang XW (2006) The role of cytokines in hepatocellular carcinoma. *J Leukoc Biol* 80, 1197–1213.
- Cavin LG, Venkatraman M, Factor VM, Kaur S, Schroeder I, et al. (2004) Regulation of alpha-fetoprotein by nuclear factor-kappaB protects hepatocytes

Author Contributions

Conceived and designed the experiments: CJY CWC TTC. Performed the experiments: CJY YJL. Analyzed the data: HWT. Contributed reagents/materials/analysis tools: CSY TFT KYC WCH PWL. Wrote the paper: CJY CWC TTC.

- from tumor necrosis factor-alpha cytotoxicity during fetal liver development and hepatic oncogenesis. *Cancer Res* 64, 7030–7038.
- Biermer M, Puro R, Schneider RJ (2003) Tumor necrosis factor alpha inhibition of hepatitis B virus replication involves disruption of capsid Integrity through activation of NF-kappaB. *J Virol* 77, 4033–4042.
- Gish RG, Baron A (2008) Hepatocellular carcinoma (HCC): current and evolving therapies. *IDrugs* 11, 198–203.
- Pang RW, Poon RT (2007) From molecular biology to targeted therapies for hepatocellular carcinoma: the future is now. *Oncology* 72, Suppl 1: 30–44.
- Aravalli RN, Steer CJ, Cressman EN (2008) Molecular mechanisms of hepatocellular carcinoma. *Hepatology* 48, 2047–2063.
- Lee DF, Kuo HP, Chen CT, Hsu JM, Chou CK, et al. (2007) IKK beta suppression of TSC1 links inflammation and tumor angiogenesis via the mTOR pathway. *Cell* 130, 440–455.
- Yen CJ, Izzo JG, Lee DF, Guha S, Wei Y, et al. (2008) Bile acid exposure up-regulates tuberous sclerosis complex 1/mammalian target of rapamycin pathway in Barrett's-associated esophageal adenocarcinoma. *Cancer Res* 68, 2632–2640.
- Lee DF, Hung MC (2007) All roads lead to mTOR: integrating inflammation and tumor angiogenesis. *Cell Cycle* 6, 3011–3014.
- Semela D, Piguat AC, Kolev M, Schmitter K, Hlushchuk R, et al. (2007) Vascular remodeling and antitumoral effects of mTOR inhibition in a rat model of hepatocellular carcinoma. *J Hepatol* 46, 840–848.
- Bu X, Jia F, Wang W, Guo X, Wu M, et al. (2007) Coupled down-regulation of mTOR and telomerase activity during fluorouracil-induced apoptosis of hepatocarcinoma cells. *BMC Cancer* 7, 208.
- [41] Melegari M, Scaglioni PP, Wands JR (1998) Cloning and characterization of a novel hepatitis B virus x binding protein that inhibits viral replication. *J Virol* 72, 1737–1743.
- Ding Q, Xia W, Liu JC, Yang JY, Lee DF, et al. (2005) Erk associates with and primes GSK-3beta for its inactivation resulting in upregulation of beta-catenin. *Mol Cell* 19, 159–170.
- Wu YF, Fu SL, Kao CH, Yang CW, Lin CH, et al. (2008) Chemopreventive effect of silymarin on liver pathology in HBV X protein transgenic mice. *Cancer Res* 68, 2033–2042.
- Grivennikov SI, Karin M (2010) Inflammation and oncogenesis: a vicious connection. *Curr Opin Genet Dev* 20, 65–71.
- Guo SP, Wang WL, Zhai YQ, Zhao YL (2001) Expression of nuclear factor-kappa B in hepatocellular carcinoma and its relation with the X protein of hepatitis B virus. *World J Gastroenterol* 7, 340–344.
- Siddiqui A, Gaynor R, Srinivasan A, Mapoles J, Farr RW (1989) Transactivation of viral enhancers including long terminal repeat of the human immunodeficiency virus by the hepatitis B virus X protein. *Virology* 169, 479–484.
- Seto E, Mitchell PJ, Yen TS (1990) Transactivation by the hepatitis B virus X protein depends on AP-2 and other transcription factors. *Nature* 344, 72–74.
- Lucito R, Schneider RJ (1992) Hepatitis B virus X protein activates transcription factor NF-kappa B without a requirement for protein kinase C. *J Virol* 66, 983–991.
- Lara-Pezzi E, Majano PL, Gomez-Gonzalo M, Garcia-Monzon C, Moreno-Otero R, et al. (1998) The hepatitis B virus X protein up-regulates tumor necrosis factor alpha gene expression in hepatocytes. *Hepatology* 28, 1013–1021.
- Su F, Schneider RJ (1996) Hepatitis B virus HBx protein activates transcription factor NF-kappaB by acting on multiple cytoplasmic inhibitors of rel-related proteins. *J Virol* 70, 4558–4566.
- Chan CF, Yau TO, Jin DY, Wong CM, Fan ST, et al. (2004) Evaluation of nuclear factor-kappaB, urokinase-type plasminogen activator, and HBx and their clinicopathological significance in hepatocellular carcinoma. *Clin Cancer Res* 10, 4140–4149.
- Zhou Y, Wang S, Ma JW, Lei Z, Zhu HF, et al. (2010) Hepatitis B virus protein X-induced expression of the CXC chemokine IP-10 is mediated through activation of NF-kappaB and increases migration of leukocytes. *J Biol Chem* 285, 12159–12168.
- [53] Benn J, Schneider RJ (1994) Hepatitis B virus HBx protein activates Ras-GTP complex formation and establishes a Ras, Raf, MAP kinase signaling cascade. *Proc Natl Acad Sci U S A* 91, 10350–10354.
- [54] Kekulé AS, Lauer U, Weiss L, Lubber B, Hofschneider PH (1993) Hepatitis B virus transactivator HBx uses a tumour promoter signalling pathway. *Nature* 361, 742–745.
- Sahin F, Kannangai R, Adegbola O, Wang J, Su G, et al. (2004) mTOR and P70 S6 kinase expression in primary liver neoplasms. *Clin Cancer Res* 10, 8421–8425.

56. Sieghart W, Fuereder T, Schmid K, Cejka D, Werzowa J, et al. (2007). Mammalian target of rapamycin pathway activity in hepatocellular carcinomas of patients undergoing liver transplantation. *Transplantation* 83, 425–432.
57. Yoo YG, Cho S, Park S, Lee MO (2004) The carboxy-terminus of the hepatitis B virus X protein is necessary and sufficient for the activation of hypoxia-inducible factor-1alpha. *FEBS Lett* 577, 121–126.
58. Yoo YG, Oh SH, Park ES, Cho H, Lee N, et al. (2003) Hepatitis B virus X protein enhances transcriptional activity of hypoxia-inducible factor-1alpha through activation of mitogen-activated protein kinase pathway. *J Biol Chem* 278, 39076–39084.