

6'-Guanidinonaltrindole (6'-GNTI) Is a G Protein-biased κ -Opioid Receptor Agonist That Inhibits Arrestin Recruitment^{*§}

Received for publication, June 1, 2012, and in revised form, June 18, 2012
Published, JBC Papers in Press, June 26, 2012, DOI 10.1074/jbc.C112.387332

Marie-Laure Rives^{‡§}, Mary Rossillo[‡], Lee-Yuan Liu-Chen[¶],
and Jonathan A. Javitch^{‡§||}

From the Departments of [‡]Psychiatry and ^{||}Pharmacology, College of Physicians and Surgeons, Columbia University, New York, New York 10032, the [§]Division of Molecular Therapeutics, New York State Psychiatric Institute, New York, New York 10032, and the [¶]Department of Pharmacology and Center for Substance Abuse Research, School of Medicine, Temple University, Philadelphia, Pennsylvania 19140

Background: Arrestin recruitment to the κ -opioid receptor (KOR) has been linked to several adverse effects of analgesics, such as dysphoria and tolerance.

Results: We identified 6'-guanidinonaltrindole (6'-GNTI) as a potent KOR agonist for G protein activation that fails to recruit arrestin.

Conclusion: 6'-GNTI is an extreme G protein-biased KOR ligand.

Significance: 6'-GNTI is a lead toward analgesics with fewer arrestin-mediated adverse effects.

κ -Opioid receptor (KOR) agonists do not activate the reward pathway stimulated by morphine-like μ -opioid receptor (MOR) agonists and thus have been considered to be promising nonaddictive analgesics. However, KOR agonists produce other adverse effects, including dysphoria, diuresis, and constipation. The therapeutic promise of KOR agonists has nonetheless recently been revived by studies showing that their dysphoric effects require arrestin recruitment, whereas their analgesic effects do not. Moreover, KOR agonist-induced antinociceptive tolerance observed *in vivo* has also been proposed to be correlated to the ability to induce arrestin-dependent phosphorylation, desensitization, and internalization of the receptor. The discovery of functionally selective drugs that are therapeutically effective without the adverse effects triggered by the arrestin pathway is thus an important goal. We have identified such an extreme G protein-biased KOR compound, 6'-guanidinonaltrindole (6'-GNTI), a potent partial agonist at the KOR receptor for the G protein activation pathway that does not recruit arrestin. Indeed, 6'-GNTI functions as an antagonist to block the arrestin recruitment and KOR internalization induced by other nonbiased agonists. As an extremely G protein-biased KOR agonist, 6'-GNTI represents a promising lead compound in the

search for nonaddictive opioid analgesic as its signaling profile suggests that it will be without the dysphoria and other adverse effects promoted by arrestin recruitment and its downstream signaling.

κ -Opioid receptors (KOR)² are widely expressed in the periphery, the dorsal root ganglia, the spinal cord, and the supraspinal regions associated with pain modulation. KOR agonists have been shown to activate pain inhibitory pathways in the central nervous system, and peripherally restricted KOR agonists have been developed to target KOR located on visceral and somatic afferent nerves for relief of inflammatory, visceral, and neuropathic chronic pain (1, 2). The analgesic properties of KOR agonists are attributed to their ability to activate G proteins in the G_{i/o} family (3, 4) as the subsequent inhibition of cAMP production (5), as well as the activation of inward rectifier potassium channels (6) and blockade of calcium channels (7), has an inhibitory effect in neurons. In contrast to MOR agonists, KOR agonists are unable to activate the reward pathway and have therefore attracted considerable attention for their ability to exert potent analgesic effects without high abuse potential (1, 8).

Unfortunately, KOR agonists have been found to produce other significant adverse effects, such as dysphoria (9). Interestingly, their dysphoric effects require activation of the p38 MAPK pathway mediated by arrestin recruitment to the activated KOR (4, 10, 11). KOR-induced p38 MAPK activation has been demonstrated in heterologous expression systems, striatal neurons and astrocytes, spinal cord astrocytes, and *in vivo* (12–14) and requires KOR phosphorylation by the G protein-coupled receptor kinase 3 (GRK3) and subsequent arrestin3 recruitment (12). Based on such findings, Chavkin (11) has proposed that KOR-selective G protein-biased partial agonists that do not efficiently recruit arrestin would not cause dysphoria but would retain sufficient analgesic activity for the treatment of pain-related disorders.

As is evident from the above, KOR and many other G protein-coupled receptors (GPCRs) couple to a number of different downstream signaling cascades (4). It is now well accepted that ligands can display different efficacies and/or potencies for different signaling pathways. This phenomenon, referred to as functional selectivity, biased agonism, or ligand-directed signaling/receptor trafficking, has been widely observed (15, 16). For example, at the level of the DOR and MOR receptors, several compounds have been identified as agonists for G protein coupling but as antagonists (DOR) or partial agonists (MOR) for arrestin (17).

Interestingly, agonist-induced interactions with arrestin have also been shown to influence opioid-induced antinocice-

* This work was supported, in whole or in part, by National Institutes of Health Grants DA022413 and MH054137 (to J. A. J.).

§ This article contains supplemental Figs. 1 and 2.

¹ To whom correspondence should be addressed: Columbia University, P & S 11-401, 630 W. 168th St., Box 7, New York, NY 10032. Tel.: 212-305-7308; Fax: 212-305-5594; E-mail: jaj2@columbia.edu.

² The abbreviations used are: KOR, κ -opioid receptor; hKOR, human KOR; MOR, μ -opioid receptor; DOR, δ -opioid receptor; BRET, bioluminescence resonance energy transfer; 6'-GNTI, 6'-guanidinonaltrindole; EKC, ethylketocyclazocine; GPCR, G protein-coupled receptor; RLuc8, *Renilla* luciferase 8; CAMYEL, cAMP sensor using YFP-Epac-RLuc; GRK, G protein-coupled receptor kinase.

ption and tolerance *in vivo*. Although still an area of controversy (see also the "morphine paradox" (17, 18)), it has been proposed that the ability of an opioid agonist to promote arrestin-dependent desensitization and internalization *in vitro* is related to the degree of tolerance that develops after long term treatment *in vivo* (19, 20). Arrestin recruitment to KOR has also been proposed to play a role in antinociceptive tolerance (20, 21), making the development of a G protein-biased agonist an important goal for achieving analgesic efficacy without the adverse effects triggered by arrestin (4, 11).

In exploring the pharmacological properties of 6'-guanidinonaltrindole (6'-GNTI), previously proposed to be a DOR-KOR heteromer-selective ligand (22), we have identified it as a functionally selective G protein-biased KOR ligand, and thus, as a promising lead compound for treating pain without dysphoria and other arrestin-mediated side effects.

EXPERIMENTAL PROCEDURES

Constructs for Expression Vectors and Transfection—The cDNA for human KOR (hKOR) was obtained from the Missouri S&T cDNA Resource Center. For arrestin recruitment experiments, full-length *Renilla* luciferase 8 (RLuc8, provided by S. Gambhir) was fused in-frame to the C terminus of hKOR in the pcDNA3.1 vector. The following human G protein constructs used were provided by C. Gales (23, 24): untagged $G\alpha_{oA}$; $G\alpha_{oB}$ with RLuc8 inserted at position 91 ($G\alpha_{oB}$ -RLuc8); untagged $G\beta_1$ (β_1); untagged $G\gamma_2$ (γ_2). The human γ_2 subunit was fused to full-length mVenus at its N terminus (mVenus- γ_2), and we used the fusion construct human arrestin3-mVenus previously described (25). All the constructs were confirmed by sequencing analysis. A total of 20 μ g of plasmid cDNA (e.g. 0.2 μ g of hKOR-RLuc8, 15 μ g of arrestin3-mVenus, and 4.8 μ g of pcDNA3.1) was transfected into HEK-293T cells using polyethylenimine (Polysciences Inc.) in a 1:3 ratio in 10-cm dishes. Cells were maintained in culture with DMEM supplemented with 10% FBS. The transfected ratio among receptor, $G\alpha$, β_1 , and γ_2 , or arrestin was optimized by testing various ratios of plasmids encoding the different sensors. Experiments were performed 48 h after transfection.

BRET—BRET was performed as described (26). Briefly, cells were harvested, washed, and resuspended in a phosphate-buffered saline (PBS) solution. Approximately 200,000 cells/well were distributed in 96-well plates, and 5 μ M coelenterazine H (luciferase substrate) was added to each well. Five minutes after the addition of coelenterazine H, ligands were added to each well, and after 2 min for G protein activation or 5 min for arrestin recruitment, the BRET signal was determined by quantifying and calculating the ratio of the light emitted by mVenus, the energy acceptor (510–540 nm), over that emitted by RLuc8, the energy donor (485 nm). The drug-induced BRET signal was normalized, taking the E_{max} of the ethylketocyclazocine (EKC)-induced response as 100%. For measuring cAMP accumulation, we used a BRET-based cAMP sensor using the YFP-Epac-RLuc (CAMYEL) assay previously described (27). $G\alpha_{oA}$, β_1 , and γ_2 were co-expressed to enhance the signal-to-noise ratio, and the cells were treated for 5 min with 100 μ M forskolin prior to stimulation (27). The data were normalized and represented as

the percentage of forskolin-stimulated cAMP accumulation with 0 defined as the maximal inhibition triggered by EKC.

Surface Expression Determination by FACS Analysis—FACS was performed to determine the surface expression of FLAG-tagged hKOR stably expressed in CHO cells with a C6 flow cytometer (BD Accuri Cytometers). After stimulation of the cells with the corresponding drugs for 30 min at 37 °C, the cells were placed on ice and resuspended in PBS containing 0.5% BSA and 0.1% NaN₃. The cells were then incubated with a mouse FLAG-specific M2 (Sigma) antibody (dilution 1:500) for 30 min, washed, and incubated with secondary mouse-specific antibody labeled with Alexa Fluor 647 (Invitrogen). The fluorescence signal at 647 nm was determined using the appropriate filter set (675 \pm 25 nm) on the C6 flow cytometer.

Drugs—Pharmacological reagents were obtained from the National Institute on Drug Abuse Drug Supply Program (U50488), from Sigma (6'-GNTI), and from James Woods (EKC).

RESULTS

In a recent study, 6'-GNTI was shown to produce a prolonged antinociceptive response in a rat behavioral model of thermal allodynia (28). The effect was even more robust than that previously reported for another selective KOR agonist, U50488 (29). Although 6'-GNTI has been proposed to act as a potent DOR-KOR heteromer-selective ligand (22) and only as a weak agonist at KOR (22), *in vivo* studies showed only a small decrease in its potency in a radiant heat tail-flick assay of nociception in DOR knock-out mice (30). Thus, 6'-GNTI can activate KOR in the absence of DOR, consistent with its original description as a KOR agonist (31). We sought to investigate this controversy by assessing the ability of the compound to activate KOR when it is expressed alone.

6'-GNTI Is a Potent Partial Agonist at KOR for G Protein Activation—To measure G protein activation, we used a BRET-based assay (23). The hKOR receptor was co-expressed in HEK293T cells with $G\alpha_{oB}$ -RLuc8, β_1 , and mVenus- γ_2 (Fig. 1A). The drug-induced BRET signal corresponds to a dissociation of and/or conformational change within the $G\alpha\beta\gamma$ complex, and thus, to the activation of the co-expressed G protein. EKC and U50488 were found to be full agonists at KOR (Fig. 1A) with potencies of 2.5 \pm 1.6 and 43 \pm 24 nM, respectively ($n = 6$). Interestingly, 6'-GNTI robustly activated G protein with a potency of 1.6 \pm 1.3 nM and an E_{max} of 64 \pm 6% that of EKC ($n = 6$) (Fig. 1A). The BRET results with the G protein biosensor were essentially identical to our results measuring drug-induced inhibition of cAMP in a BRET-based CAMYEL sensor assay (27) (Fig. 1B). EKC and U50488 were similarly efficacious (IC₅₀ values of 1.4 \pm 1.1 and 38 \pm 30 nM, respectively), whereas 6'-GNTI had a relative E_{max} of 69 \pm 4% (IC₅₀ of 1.1 \pm 1.2 nM) ($n = 3$). Although the co-expression of DOR was previously reported to enhance the efficacy of 6'-GNTI for activation of a chimeric G_{q15} protein (22), we did not observe an increase in the efficacy or potency of 6'-GNTI for $G\alpha_o\beta\gamma$ activation or cAMP inhibition by KOR upon co-expression of DOR (supplemental Fig. 1, A and B).

6'-GNTI Is a Functional Antagonist for the Arrestin Pathway—We used a BRET-based assay to measure the recruitment of




FIGURE 1. 6'-GNTI is a potent partial agonist at hKOR for G protein activation and for inhibition of cAMP accumulation. *A*, the hKOR receptor was co-expressed with $G_{\alpha_{\text{O}}\beta}$ -RLuc8, $\beta 1$, and mVenus- $\gamma 2$ to assay G protein activation (23). *B*, the hKOR receptor was co-expressed with a BRET-based CAMYEL sensor to assay inhibition of forskolin-stimulated cAMP accumulation. Error bars in *A* and *B* indicate S.E.

arrestin3 fused to mVenus to KOR fused to RLuc8 (25). In contrast to EKC and U50488 (Fig. 2A), which both robustly recruited arrestin to KOR (EC_{50} of 17 ± 10 nM and 2.0 ± 1.2 μ M, respectively), 6'-GNTI was without significant effect (Fig. 2A), even after 30 min (supplemental Fig. 2A). To ensure that the absence of signal was not due to a sensitivity limit of the assay given that 6'-GNTI was shown to be a partial agonist for G protein activation, we co-expressed the G protein-coupled receptor kinase, GRK3, to enhance the arrestin BRET signal. GRKs, and more specifically GRK2 and GRK3, are known to enhance arrestin recruitment to various GPCRs, thereby promoting desensitization and internalization of the receptors (32). In the presence of GRK3, we observed a more than 2-fold increase in maximal EKC- and U50488-induced arrestin recruitment to KOR (Fig. 2B). In contrast, the signal triggered by 6'-GNTI was not significantly increased by GRK3, confirming the virtual absence of arrestin3 recruitment to KOR by this compound. 6'-GNTI was also unable to recruit arrestin2 (supplemental Fig. 2B), and the co-expression of DOR with KOR also did not lead to 6'-GNTI-induced recruitment of arrestin3 (supplemental Fig. 1B). These results suggested that 6'-GNTI, although a potent agonist for G protein activation and the cAMP pathway, has such low efficacy for arrestin recruitment that it functions as an antagonist for this pathway. Indeed, 6'-GNTI effectively inhibited EKC-induced arrestin3 recruitment to KOR (Fig. 2A). To confirm the lack of arrestin recruitment triggered by 6'-GNTI by probing its downstream effects, we measured drug-induced internalization. Both EKC and U50488 led to robust receptor internalization, whereas 6'-GNTI led to minimal internalization that did not reach significance (Fig. 2C). Consistent with our arrestin recruitment measurements, 6'-GNTI also potently inhibited EKC-induced KOR internalization (Fig. 2C).

DISCUSSION

We have found that the naltrindole derivative 6'-GNTI is a potent partial agonist for G protein activation at the KOR receptor. Moreover, 6'-GNTI is extremely G protein-biased as it failed to recruit arrestin and functioned as an antagonist for arrestin recruitment by other KOR agonists. This functional selectivity was manifested downstream as well because 6'-GNTI potently inhibited cAMP activation but functioned as an antagonist of agonist-mediated internalization. This extreme biased agonism is consistent with the potent and effi-

acious analgesic efficacy of 6'-GNTI observed *in vivo* (22, 28, 30) and also suggests that 6'-GNTI would not cause dysphoria because of its inability to activate the arrestin pathway. Thus, 6'-GNTI seems to be an ideal lead candidate that establishes the feasibility of an extremely G protein-biased KOR agonist satisfying the design features hypothesized by Chavkin (11). To our knowledge, no other such extreme G protein-biased KOR agonists have been identified to date, although there has been a report of an arrestin-biased KOR ligand (33). Interestingly, although not KOR-selective, etorphine and levorphanol were shown to reduce KOR internalization (34), and their potential functional selectivity at the level of arrestin recruitment merits further investigation.

It is also possible that the extreme G protein bias exhibited by 6'-GNTI would lead to less desensitization and tolerance at KOR. As mentioned above, although not without controversy (17, 18), the KOR-mediated arrestin pathway has been suggested to influence the development of antinociceptive tolerance *in vivo* (20, 21). It is noteworthy that arrestin3 KO mice display enhanced and prolonged antinociception after short term treatment with morphine and heroin in the hot plate test, a model of thermal antinociception (35). Thus, we hypothesize that after prolonged activation with 6'-GNTI, the KOR receptor would undergo less arrestin-mediated desensitization, which would result in enhanced rather than diminished efficacy over time through persistent G protein activation. In fact, the antiallo-dynic effect of U50488 was shown to decrease at high concentrations (29), and it was suggested that this might result from rapid desensitization of KOR. Interestingly, we found that EKC and U50488 were both more potent at G protein activation than at recruiting arrestin (7- and 47-fold, respectively) (see above). This could explain the dramatically blunted antinociceptive response to U50488 observed *in vivo* at high concentrations of the drug as at high concentrations of agonist, arrestin recruitment would be expected to severely limit G protein signaling and hence to limit the analgesic response. Based on our findings, we would predict the absence of such a U-shaped curve for 6'-GNTI analgesia *in vivo*. Nonetheless, further experiments investigating the desensitization properties of KOR will be necessary to test this hypothesis and to establish that KOR remains functional on the cell surface after prolonged activation by 6'-GNTI.




FIGURE 2. 6'-GNTI is a functional antagonist for the arrestin pathway. A and B, the hKOR receptor, fused to RLuc8, was co-expressed with arrestin3 (*Arr3*) fused to mVenus to assay arrestin recruitment to the receptor (25). A, 6'-GNTI does not induce arrestin3 recruitment to hKOR, in contrast to EKC and U50488, but can inhibit the recruitment induced by EKC. B, GRK3 dramatically enhanced arrestin recruitment to KOR induced by EKC and U50488 but did not affect the signal triggered by 6'-GNTI. A two-way analysis of variance test followed by Bonferroni's post tests was performed to determine the significance of the GRK3 effect on the E_{max} of each drug. C, after 30 min of incubation with the different drugs, the expression of the FLAG-tagged hKOR receptor, stably expressed in a CHO stable cell line, was measured by flow cytometry using an anti-FLAG antibody ($n = 3$). A one-way analysis of variance test with 99.9% confidence intervals ($p < 0.001$) followed by a Tukey's multiple test comparison was performed to determine the significance of each column as compared with vehicle (*Veh*). Error bars in A–C indicate S.E. ns, not significant.

Our data challenge the notion that 6'-GNTI is DOR-KOR heteromer-selective as we observe robust activation of G protein and inhibition of cAMP in HEK cells in which we only expressed KOR. This is consistent with the quite small alteration in the potency of 6'-GNTI described in a tail-flick assay in DOR KO mice (30). The original study suggesting decreased efficacy of 6'-GNTI in KOR relative to DOR-KOR used a chimeric G_{q15} aequorin signaling assay (22), which may read out differently for G protein activation. It is also possible that enhanced cell surface expression of KOR by co-expression of DOR impacted these results. Although it seems clear that DOR is not essential for 6'-GNTI function, either in HEK cells or *in vivo* (30), we do not rule out a role of DOR in modulating KOR signaling *in vivo*, through either a heteromer or a downstream cross-talk mechanism, as proposed by recent studies (28). Nonetheless, the profound functional selectivity of the compound seems likely to dominate its pharmacological properties.

Although functional selectivity/biased agonism was first described almost 20 years ago (36), it has emerged only recently as a key concept for the development of therapeutic drugs (37). GPCRs ligands can activate both G protein-dependent and G protein-independent signaling pathways, which differentially regulate their therapeutic actions and side effects. Understanding which signaling pathways contribute to therapeutic efficacy and which contribute to side effects will enable the design of better drug candidates and lead to safer and more effective therapies. For example, several novel compounds exhibiting antipsychotic-like activity *in vivo* were shown recently to be antagonists at dopamine D2 receptor of G_i -mediated inhibition of cAMP production but partial agonists for recruitment of arrestin3 to the dopamine D2 receptor (38). These functionally selective arrestin-biased ligands represent valuable chemical probes for investigation of the signaling pathways that are associated with antipsychotic efficacy and side effects. Similarly, the extreme G protein-biased functional selectivity of 6'-GNTI has great potential for effective analgesia with reduced liability for dysphoria and tolerance.

Acknowledgments—We thank Celine Gales and Sam Gambhir for providing cDNA constructs, James Woods for the EKC, and Marta Filizola and Prashant Donthamsetti for helpful comments on the manuscript.

REFERENCES

- Vanderah, T. W. (2010) δ - and κ -opioid receptors as suitable drug targets for pain. *Clin. J. Pain* **26**, (Suppl. 10) S10–S15
- Basbaum, A. I., and Jessell, T. M. (2000) The perception of pain in *Principles of Neural Science*, 4th Ed. (Kandel, E. R., Schwartz, J. H., and Jessell, T. M., eds) pp. 472–479, The McGraw-Hill Companies, New York, NY
- Taussig, R., Iniguez-Lluhi, J. A., and Gilman, A. G. (1993) Inhibition of adenylyl cyclase by $G_{i\alpha}$. *Science* **261**, 218–221
- Bruchas, M. R., and Chavkin, C. (2010) Kinase cascades and ligand-directed signaling at the κ -opioid receptor. *Psychopharmacology (Berl)* **210**, 137–147
- Schoffelmeer, A. N., Rice, K. C., Jacobson, A. E., Van Gelderen, J. G., Hogenboom, F., Heijna, M. H., and Mulder, A. H. (1988) μ -, δ -, and κ -Opioid receptor-mediated inhibition of neurotransmitter release and adenylyl cyclase activity in rat brain slices: studies with fentanyl isothiocyanate. *Eur. J. Pharmacol.* **154**, 169–178
- Henry, D. J., Grandy, D. K., Lester, H. A., Davidson, N., and Chavkin, C.

- (1995) κ -Opioid receptors couple to inwardly rectifying potassium channels when co-expressed by *Xenopus* oocytes. *Mol. Pharmacol.* **47**, 551–557
7. Tallent, M., Dichter, M. A., Bell, G. I., and Reisine, T. (1994) The cloned κ -opioid receptor couples to an *N*-type calcium current in undifferentiated PC-12 cells. *Neuroscience* **63**, 1033–1040
8. Wang, Y. H., Sun, J. F., Tao, Y. M., Chi, Z. Q., and Liu, J. G. (2010) The role of κ -opioid receptor activation in mediating antinociception and addiction. *Acta Pharmacol Sin* **31**, 1065–1070
9. Land, B. B., Bruchas, M. R., Lemos, J. C., Xu, M., Melief, E. J., and Chavkin, C. (2008) The dysphoric component of stress is encoded by activation of the dynorphin κ -opioid system. *J. Neurosci.* **28**, 407–414
10. Land, B. B., Bruchas, M. R., Schattauer, S., Giardino, W. J., Aita, M., Messinger, D., Hnasko, T. S., Palmiter, R. D., and Chavkin, C. (2009) Activation of the κ -opioid receptor in the dorsal raphe nucleus mediates the aversive effects of stress and reinstates drug seeking. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 19168–19173
11. Chavkin, C. (2011) The therapeutic potential of κ -opioids for treatment of pain and addiction. *Neuropsychopharmacology* **36**, 369–370
12. Bruchas, M. R., Macey, T. A., Lowe, J. D., and Chavkin, C. (2006) κ -Opioid receptor activation of p38 MAPK is GRK3- and arrestin-dependent in neurons and astrocytes. *J. Biol. Chem.* **281**, 18081–18089
13. Bruchas, M. R., Land, B. B., Aita, M., Xu, M., Barot, S. K., Li, S., and Chavkin, C. (2007) Stress-induced p38 mitogen-activated protein kinase activation mediates κ -opioid-dependent dysphoria. *J. Neurosci.* **27**, 11614–11623
14. Xu, M., Bruchas, M. R., Ippolito, D. L., Gendron, L., and Chavkin, C. (2007) Sciatic nerve ligation-induced proliferation of spinal cord astrocytes is mediated by κ -opioid activation of p38 mitogen-activated protein kinase. *J. Neurosci.* **27**, 2570–2581
15. Urban, J. D., Clarke, W. P., von Zastrow, M., Nichols, D. E., Kobilka, B., Weinstein, H., Javitch, J. A., Roth, B. L., Christopoulos, A., Sexton, P. M., Miller, K. J., Spedding, M., and Mailman, R. B. (2007) Functional selectivity and classical concepts of quantitative pharmacology. *J. Pharmacol. Exp. Ther.* **320**, 1–13
16. Lane, J. R., Powney, B., Wise, A., Rees, S., and Milligan, G. (2007) Protean agonism at the dopamine D2 receptor: (*S*)-3-(3-hydroxyphenyl)-*N*-propylpiperidine is an agonist for activation of G_{o1} but an antagonist/inverse agonist for G₁₁, G₁₂, and G₁₃. *Mol. Pharmacol.* **71**, 1349–1359
17. Molinari, P., Vezzi, V., Sbraccia, M., Grò, C., Riitano, D., Ambrosio, C., Casella, I., and Costa, T. (2010) Morphine-like opiates selectively antagonize receptor-arrestin interactions. *J. Biol. Chem.* **285**, 12522–12535
18. Bohn, L. M., Dykstra, L. A., Lefkowitz, R. J., Caron, M. G., and Barak, L. S. (2004) Relative opioid efficacy is determined by the complements of the G protein-coupled receptor desensitization machinery. *Mol. Pharmacol.* **66**, 106–112
19. Raehal, K. M., Schmid, C. L., Groer, C. E., and Bohn, L. M. (2011) Functional selectivity at the μ -opioid receptor: implications for understanding opioid analgesia and tolerance. *Pharmacol. Rev.* **63**, 1001–1019
20. Liu-Chen, L. Y. (2004) Agonist-induced regulation and trafficking of κ -opioid receptors. *Life Sci.* **75**, 511–536
21. McLaughlin, J. P., Myers, L. C., Zarek, P. E., Caron, M. G., Lefkowitz, R. J., Czyzyk, T. A., Pintar, J. E., and Chavkin, C. (2004) Prolonged κ -opioid receptor phosphorylation mediated by G protein receptor kinase underlies sustained analgesic tolerance. *J. Biol. Chem.* **279**, 1810–1818
22. Waldhoer, M., Fong, J., Jones, R. M., Lunzer, M. M., Sharma, S. K., Kostenis, E., Portoghese, P. S., and Whistler, J. L. (2005) A heterodimer-selective agonist shows *in vivo* relevance of G protein-coupled receptor dimers. *Proc. Natl. Acad. Sci. U.S.A.* **102**, 9050–9055
23. Galés, C., Van Durm, J. J., Schaak, S., Pontier, S., Percherancier, Y., Audet, M., Paris, H., and Bouvier, M. (2006) Probing the activation-promoted structural rearrangements in preassembled receptor-G protein complexes. *Nat. Struct. Mol. Biol.* **13**, 778–786
24. Saulière, A., Bellot, M., Paris, H., Denis, C., Finana, F., Hansen, J. T., Altié, M.-F., Seguelas, M.-H., Pathak, A., Hansen, J. L., Sénard, J. M., and Galés, C. (2012) Deciphering biased-agonism complexity reveals a new active AT1 receptor entity. *Nat. Chem. Biol.* **8**, 622–630
25. Klewe, I. V., Nielsen, S. M., Tarpø, L., Urizar, E., Dipace, C., Javitch, J. A., Gether, U., Egebjerg, J., and Christensen, K. V. (2008) Recruitment of β -arrestin2 to the dopamine D2 receptor: insights into antipsychotic and antiparkinsonian drug receptor signaling. *Neuropharmacology* **54**, 1215–1222
26. Guo, W., Urizar, E., Kralikova, M., Mobarec, J. C., Shi, L., Filizola, M., and Javitch, J. A. (2008) Dopamine D2 receptors form higher order oligomers at physiological expression levels. *EMBO J.* **27**, 2293–2304
27. Jiang, L. I., Collins, J., Davis, R., Lin, K. M., DeCamp, D., Roach, T., Hsueh, R., Rebres, R. A., Ross, E. M., Taussig, R., Fraser, I., and Sternweis, P. C. (2007) Use of a cAMP BRET sensor to characterize a novel regulation of cAMP by the sphingosine 1-phosphate/G₁₃ pathway. *J. Biol. Chem.* **282**, 10576–10584
28. Berg, K. A., Rowan, M. P., Gupta, A., Sanchez, T. A., Silva, M., Gomes, I., McGuire, B. A., Portoghese, P. S., Hargreaves, K. M., Devi, L. A., and Clarke, W. P. (2012) Allosteric interactions between δ - and κ -opioid receptors in peripheral sensory neurons. *Mol. Pharmacol.* **81**, 264–272
29. Berg, K. A., Rowan, M. P., Sanchez, T. A., Silva, M., Patwardhan, A. M., Milam, S. B., Hargreaves, K. M., and Clarke, W. P. (2011) Regulation of κ -opioid receptor signaling in peripheral sensory neurons *in vitro* and *in vivo*. *J. Pharmacol. Exp. Ther.* **338**, 92–99
30. Ansonoff, M. A., Portoghese, P. S., and Pintar, J. E. (2010) Consequences of opioid receptor mutation on actions of univalent and bivalent κ - and δ -ligands. *Psychopharmacology* **210**, 161–168
31. Sharma, S. K., Jones, R. M., Metzger, T. G., Ferguson, D. M., and Portoghese, P. S. (2001) Transformation of a κ -opioid receptor antagonist to a κ -agonist by transfer of a guanidinium group from the 5' - to 6' -position of naltrindole. *J. Med. Chem.* **44**, 2073–2079
32. Reiter, E., and Lefkowitz, R. J. (2006) GRKs and β -arrestins: roles in receptor silencing, trafficking, and signaling. *Trends Endocrinol Metab.* **17**, 159–165
33. Béguin, C., Potuzak, J., Xu, W., Liu-Chen, L. Y., Streicher, J. M., Groer, C. E., Bohn, L. M., Carlezon, W. A., Jr., and Cohen, B. M. (2012) Differential signaling properties at the κ -opioid receptor of 12-epi-salvinorin A and its analogues. *Bioorg. Med. Chem. Lett.* **22**, 1023–1026
34. Li, J. G., Zhang, F., Jin, X. L., and Liu-Chen, L. Y. (2003) Differential regulation of the human κ -opioid receptor by agonists: etorphine and levorphanol reduced dynorphin A- and U50,488H-induced internalization and phosphorylation. *J. Pharmacol. Exp. Ther.* **305**, 531–540
35. Raehal, K. M., and Bohn, L. M. (2011) The role of β -arrestin2 in the severity of antinociceptive tolerance and physical dependence induced by different opioid pain therapeutics. *Neuropharmacology* **60**, 58–65
36. Kenakin, T. (1995) Agonist-receptor efficacy. II. Agonist trafficking of receptor signals. *Trends Pharmacol. Sci.* **16**, 232–238
37. Whalen, E. J., Rajagopal, S., and Lefkowitz, R. J. (2011) Therapeutic potential of β -arrestin- and G protein-biased agonists. *Trends Mol. Med.* **17**, 126–139
38. Allen, J. A., Yost, J. M., Setola, V., Chen, X., Sassano, M. F., Chen, M., Peterson, S., Yadav, P. N., Huang, X. P., Feng, B., Jensen, N. H., Che, X., Bai, X., Frye, S. V., Wetsel, W. C., Caron, M. G., Javitch, J. A., Roth, B. L., and Jin, J. (2011) Discovery of β -arrestin-biased dopamine D2 ligands for probing signal transduction pathways essential for antipsychotic efficacy. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 18488–18493