

Michele Ferrari,^a Claudia Folli,^a
 Elisa Pincolini,^a Timothy S.
 McClintock,^b Manfred Rössle,^c
 Rodolfo Berni^a and Michele
 Cianci^{c*}

^aDepartment of Biochemistry and Molecular Biology, University of Parma, Parco Area delle Scienze 23/A, 43124 Parma, Italy, ^bUniversity of Kentucky College of Medicine, 800 Rose Street, Lexington, Kentucky, USA, and ^cEuropean Molecular Biology Laboratory, Hamburg Outstation, c/o DESY, Notkestrasse 85, 22603 Hamburg, Germany

Correspondence e-mail:
 michele.cianci@embl-hamburg.de

Received 5 March 2012
 Accepted 8 June 2012

PDB Reference: 4alo.

Structural characterization of recombinant crustacyanin subunits from the lobster *Homarus americanus*


Crustacean crustacyanin proteins are linked to the production and modification of carapace colour, with direct implications for fitness and survival. Here, the structural and functional properties of the two recombinant crustacyanin subunits H₁ and H₂ from the American lobster *Homarus americanus* are reported. The two subunits are structurally highly similar to the corresponding natural apo crustacyanin CRTC and CRTA subunits from the European lobster *H. gammarus*. Reconstitution studies of the recombinant crustacyanin proteins H₁ and H₂ with astaxanthin reproduced the bathochromic shift of 85–95 nm typical of the natural crustacyanin subunits from *H. gammarus* in complex with astaxanthin. Moreover, correlations between the presence of crustacyanin genes in crustacean species and the resulting carapace colours with the spectral properties of the subunits in complex with astaxanthin confirmed this genotype–phenotype linkage.

1. Introduction

The typical slate-grey/blue colour of the lobster carapace is generated by a number of distinct carotenoproteins, with the predominant one being the multimeric α -crustacyanin in complex with astaxanthin (3,3'-dihydroxy- β , β -carotene-4,4'-dione), which is characterized by a λ_{\max} of 632 nm (Zagalsky, 1985). In contrast, the absorption spectrum λ_{\max} in the visible region of astaxanthin in organic solvent is drastically blue-shifted (λ_{\max} of 478 nm in acetone). The change in colour of α -crustacyanin from blue to red upon protein denaturation and from red to blue upon astaxanthin-complex reconstitution has been a subject of investigation for over 60 years (Wald *et al.*, 1948; Zagalsky, 1985; Buchwald & Jenks, 1968).

The multimeric α -crustacyanin, a protein complex consisting of 16 crustacyanin protein subunits, dissociates reversibly in low ionic strength solution, initially forming α' -crustacyanin (λ_{\max} = 595 nm) and then with further incubation the heterodimeric β -crustacyanin (λ_{\max} = 580–590 nm) (Cheeseman *et al.*, 1966). Chemical removal of the carotenoid from β -crustacyanin results in reversible dissociation into two types of apoproteins with molecular weights of 21 kDa (type I; CRTC) and 19 kDa (type II; CRTA), respectively (Buchwald & Jenks, 1968; Quarmby *et al.*, 1977). β -Crustacyanin is in fact a heterodimer formed by one CRTC lipocalin subunit, one CRTA lipocalin subunit and two noncovalently bound astaxanthin molecules (Zagalsky, 1985; Cianci *et al.*, 2002). The crustacyanins of the lobsters *Homarus americanus* and *H. gammarus* exhibit identical absorption spectra but behave differently in ion-exchange chromatography and polyacrylamide gel electrophoresis (Zagalsky & Tidmarsh, 1985). In the case of *H. gammarus* five distinct subunits are evident on 6 M urea–PAGE gels, namely A₁, C₁ and C₂ (type I) and A₂ and A₃ (type II). The *H. americanus* crustacyanin subunits appear to consist of only two major subunits, namely H₁ (type I; CRTC) and H₂ (type II; CRTA), both of which behave like A_x subunits on a 6 M urea–PAGE gel (Fig. 1*a*). Only two genes encoding crustacyanin subunits have so far been identified in lobsters (Wade *et al.*, 2009), one for each group, suggesting that the differences between protein members of the same group could arise from post-translational modifications.

The presence of CRTC and/or CRTA might be linked to the ability to produce or modify carapace colour, with direct implications for the



fitness and survival of many crustaceans *via* camouflage or mate selection (Wade *et al.*, 2009).

Although the crystal structure of β -crustacyanin has been solved at 3.2 Å resolution (Cianci *et al.*, 2002) and the structural architecture of α -crustacyanin from *H. gammarus* has been investigated by EM/SAXS (Rhys *et al.*, 2011), there is still debate on the origin of the bathochromic shift of astaxanthin upon binding to carotenoproteins. The contributions of coplanarization, exciton interaction and polarization to the bathochromic shift of astaxanthin have also been investigated using chemical crystallography of astaxanthin and its derivatives (Bartalucci *et al.*, 2007, 2009; Helliwell, 2008) or by using halogenated canthaxanthins in crustacyanin-reconstitution studies (Liu *et al.*, 2002). These contributions have also been the subject of several theoretical reports (Durbeej & Eriksson, 2004; Ilagan *et al.*, 2005; Wijk *et al.*, 2005; Strambi & Durbeej, 2009; Helliwell, 2010; Polívka *et al.*, 2010; Neugebauer *et al.*, 2011).

The present study reports the cloning, heterologous expression and structural characterization of the two apo subunits H₁ and H₂ from *H. americanus* and the results of the reconstitution of their complexes with astaxanthin.

2. Materials and methods

2.1. Sequence analyses

A BLAST search (Altschul *et al.*, 1997) for sequences similar to the CRA1_HOMGA sequence (type I subunit from *H. gammarus*; GenBank accession No. P58989) identified EST sequence DV771534 (Fig. 1*b*) among 5184 *H. americanus* olfactory-organ 5'-end sequences

obtained from a subtracted directional cDNA library prepared in pBluescript (Stepanyan *et al.*, 2006). A second BLAST search obtained using the CRA2_HOMGA sequence (type II subunit from *H. gammarus*; GenBank accession No. P80007) as the query identified a second EST sequence of interest from the same source: DV774018 (Fig. 1*c*). Clones containing these putative crustacyanin sequences were rescued from frozen stocks of this library by amplicillin selection on agar plates.

The EST sequence DV771534 (H₁ from *H. americanus*) encoded 181 amino acids, the same as CRTC_HOMGA, with an overall protein sequence identity of 97%. The EST sequence DV774018 (H₂ from *H. americanus*) encoded 159 amino acids with a sequence identity of 95%. Based on this high degree of identity, the 15 N-terminal codons of A₃ from *H. gammarus* were added to cDNA clone DV774018 to produce what is predicted to be a full-length CRTA sequence (Fig. 2*b*).

2.2. Cloning, expression and purification of the H₁ subunit

The region of the cDNA clone DV771534 corresponding to the mature H₁ subunit was amplified by PCR using an *Nde*I-tailed upstream primer 5'-CATATGGACAAAATCCCAGACTTC-3' and the downstream primer 5'-CTAGAGTGTCTTCTGAGTATCGTA-3'. The product was then cloned into pGEM-T Easy (Promega) using the T/A cloning method and then transferred into the *Nde*I site of the expression plasmid pET11b (Novagen). *Escherichia coli* BL21 Rosetta-gami cells (Novagen) were transformed with the expression plasmid. The expression of the H₁ subunit was induced overnight at 310 K by adding 1 mM IPTG and the cells were lysed by sonication.



Figure 1 (a) 6 M urea-PAGE of crustacyanin pigments of *H. gammarus* (left) and *H. americanus* (right). Copyright Elsevier (1985), reproduced with permission from Zagalsky & Tidmarsh (1985). Sequence alignments of (b) CRTC and (c) CRTA crustacyanin proteins from *H. americanus* (HomAm) using the *H. gammarus* (HomGa) sequences as a reference. Amino-acid alignment was performed *via* the ClustalW2 algorithm (<http://www.ebi.ac.uk/Tools/msa/clustalw2/>; Altschul *et al.*, 1997).

The recombinant CRTC_HOMAM, which was present in the insoluble fraction, was dissolved in a denaturing medium [50 mM Tris-HCl pH 7.5, 2 mM EDTA, 2 mM PMSF, 0.1% (v/v) Triton X-100, 8 M urea] and then refolded by dialysis against 50 mM Tris-HCl pH 9.3. The soluble fraction at pH 7 was centrifuged and concentrated by ultrafiltration. The H₁ subunit was then purified to homogeneity using Ultrogel AcA54 gel-filtration chromatography. The final yield of purified protein was 2–2.5 mg per litre of culture.

2.3. Cloning, expression and purification of the H₂ subunit

The cDNA clone DV774018 lacked a complete N-terminus. To produce a full-length protein, albeit chimeric, for expression studies, three PCR reactions were performed sequentially to generate a cDNA encoding the first 15 N-terminal residues of CRA_HOMGA followed in-frame by the DV774018 sequence. For the 5' elongation of the coding sequence, three forward partially overlapping primers and a reverse primer were employed (Table 1). The sequence encoding the full-length chimeric protein was amplified using an *Nco*I-tailed upstream primer (Table 1), cloned into pGEM-T Easy

(Promega) using the T/A cloning method and transferred into the *Nco*I site of the expression plasmid pET28b (Novagen). *E. coli* BL21 Rosetta-gami cells (Novagen) were transformed with the expression plasmid. Protein expression and purification was carried out as previously described for the *H. americanus* H₁ subunit. The final yield of purified protein was 1.5–2 mg per litre of culture.

2.4. Crystallization, data collection and analysis of the H₁ subunit

Recombinant H₁ subunit was concentrated to a final concentration of 10.7 mg ml⁻¹ in 1 mM EDTA, 0.1 M Tris-HCl pH 7.0 by ultrafiltration using an Amicon cell with a 10 kDa cutoff membrane and used in crystallization experiments. Crystallization was performed at room temperature using the sitting-drop vapour-diffusion method at the EMBL Hamburg high-throughput crystallization facility (Mueller-Dieckmann, 2006). The best crystals were grown by mixing equal volumes (1 µl) of protein solution and 1 mM EDTA, 2.4 M ammonium sulfate, 5% (v/v) 2-methyl-2,4-pentanediol (MPD), 0.1 M Tris-HCl pH 8.0 buffer. Crystals grew to dimensions of 100 × 30 × 30 µm in a week.




Figure 2 Worm-type diagram of the H₁ subunit (red) superimposed on the A₁ subunit from β -crustacyanin (blue; PDB entry 1gka): (a) monomer A, (d) monomer B. Superimposition of the H₁ subunit binding pocket (red) superimposed on the A₁ subunit binding pocket from β -crustacyanin (blue), resulting in the astaxanthin molecule AXT1 from 1gka superimposed on the MPD-binding sites: (b) monomer A, (c) monomer A rotated by 90°, (e) monomer B, (f) monomer B rotated by 90°. $2F_o - F_c$ maps contoured at 1σ r.m.s. are shown.

X-ray diffraction data were collected from H_1 crystals using synchrotron radiation on the BW6-MPG beamline at the DORIS storage ring, *c/o* Deutsches Elektronen Synchrotron (DESY), Hamburg, Germany. Each single crystal was exposed to a cold nitrogen stream at 100 K without further cryoprotection. The data were processed with *HKL-2000* (Otwinowski & Minor, 1997). The crystals diffracted to 2.38 Å resolution and belonged to space group $P2_12_12_1$, with unit-cell parameters $a = 40.37$, $b = 78.42$, $c = 105.63$ Å.

The structure was solved by molecular replacement with *AMoRe* (Winn *et al.*, 2011) using the crystal structure of apo A_1 from *H. gammarus* (PDB entry 1h91; Cianci *et al.*, 2001) as the search model. The structure was refined using *REFMAC5* (Winn *et al.*, 2011; Murshudov *et al.*, 2011). Model building and water assignments were performed using *Coot* (Emsley *et al.*, 2010). General criteria for water assignment were B factor lower than 80 Å², $2F_o - F_c$ map σ level greater than 1 and interatomic contacts of between 2.3 and 3.5 Å.

Refinement using isotropic temperature factors converged to final R -factor and R_{free} (5% of data) values of 20.2% and 26.8%, respectively.

Table 1

Primers used for cloning of the *H. americanus* H_2 subunit.

CRTA1	Forward, 5'-AAATGTGCCTCCGTAGCAAACCAGGCCAAC-3'
CRTA2	Forward, 5'-TTTGTAACCTGCAGGAAAATGTGCCTCCGTAG-CAAAC-3'
CRTA3	Forward, 5'-GATGGAATTCCTTCATTGTAACTGCAG-GAAAATGTG-3'
CRTA	Reverse, 5'-TTAAGCTCTGTAGACACACTC-3'

The stereochemistry of the final model was checked using *PROCHECK* (Winn *et al.*, 2011). Structural superpositions were performed using *LSQKAB* (Winn *et al.*, 2011). Figures were prepared using *CCP4mg* (McNicholas *et al.*, 2011). Data-collection and refinement statistics are reported in Table 2.

2.5. SAXS data collection and analysis of the H_2 subunit from *H. americanus*

The synchrotron-radiation X-ray scattering data of the H_2 subunit were collected following standard procedures on SAXS beamline




Figure 3

(a) Ribbon diagram of an H_2 monomer, with cysteine residues highlighted. (b) H_2 trimer generated using the 1gka crystal packing, with point mutations shown in ball-and-stick representation. (c) 1gka crystal packing viewed down the threefold axis, showing the H_2 trimer. (d) Profile-fitting results of SAXS data using different models.

Table 2

Data-collection and model-refinement statistics.

Values in parentheses are for the highest resolution bin.

Data collection	
Wavelength (Å)	1.05
Detector	MAR 165 CCD
Oscillation angle (°)	0.35
No. of images	250
Space group	$P2_12_12_1$
Unit-cell parameters (Å)	$a = 40.38, b = 78.46, c = 105.63$
Resolution range (Å)	20.0–2.38 (2.46–2.38)
Total No. of reflections	48851
Unique reflections	13980
Multiplicity	3.5 (3.5)
Completeness (%)	98.9 (99.0)
$R_{\text{merge}}^{\dagger}$ (%)	7.5 (34.1)
Mean $I/\sigma(I)$	14.0 (2.5)
Refinement statistics	
No. of molecules in the asymmetric unit	2
No. of residues \ddagger	362
R factor \ddagger (%)	22.3
$R_{\text{free}}^{\ddagger}$ (%)	26.6
Cruickshank's DPI for coordinate error based on R factor \ddagger (Å)	0.98
B factors (Å ²)	
Average all-atom \S	31.5
Main-chain atoms \P	35.0
Side-chain atoms and waters \P	35.9
Average r.m.s. B factor (Å ²)	
Main-chain atoms \P	0.4
Side-chain atoms \P	0.9
Total No. of atoms	3083
Total No. of water molecules	97
Solvent content (%)	40.5
Matthews coefficient (Å ³ Da ⁻¹)	2.06
Ramachandran plot $\dagger\dagger$	
Most favoured region	286 [88.8%]
Additionally allowed region	34 [10.6%]
Generously allowed region	1 [0.3%]
Disallowed region	1 [0.3%]

\dagger $R_{\text{merge}} = \sum_{hkl} \sum_i |I_i(hkl) - \langle I(hkl) \rangle| / \sum_{hkl} \sum_i I_i(hkl)$, where $I_i(hkl)$ is the intensity of a reflection and $\langle I(hkl) \rangle$ is the mean intensity of all symmetry-related reflections i . \ddagger Taken from *REFMAC5* (Winn *et al.*, 2011; Murshudov *et al.*, 2011). \S DPI = $[N_{\text{atoms}} / (N_{\text{refl}} - N_{\text{params}})]^{1/2} \times R$ factor $\times D_{\text{max}} \times \text{compl}^{-1/3}$, where N_{atoms} is the number of atoms included in the refinement, N_{refl} is the number of reflections included in the refinement, D_{max} is the maximum resolution of the reflections included in the refinement, compl is the completeness of the observed data and, in isotropic refinement, $N_{\text{params}} \approx 4N_{\text{atoms}}$ (Cruickshank, 1999). \P Taken from *BAVERAGE* (CCP4; Winn *et al.*, 2011). $\dagger\dagger$ Taken from *PROCHECK* (CCP4; Winn *et al.*, 2011).

X33 (Roessle *et al.*, 2007) at the DORIS III storage ring, DESY, Hamburg. The scattering patterns from the H₂ subunit were measured using a sample-to-detector distance of 2.4 m, covering the momentum-transfer range $0.01 < s < 0.5 \text{ \AA}^{-1}$ [$s = 4\pi \sin(\theta)/\lambda$, where θ is the scattering angle and $\lambda = 1.5 \text{ \AA}$ is the X-ray wavelength]. In order to check for interprotein interactions, measurements were made at three protein concentrations: 15, 7.5 and 3.8 mg ml⁻¹. Repetitive measurements (120 s) of the same protein solution were performed in order to check for radiation damage, and no aggregation was found during the initial 120 s exposure. This initial exposure frame was taken as a reference for further analysis. The data were normalized to the intensity of the incident beam; the scattering of the buffer was subtracted and the difference curves were scaled for concentration. Data sets collected at different concentrations were merged to improve the data quality. All data-processing steps were performed using the *PRIMUS* package (Konarev *et al.*, 2003). The forward scattering $I(0)$ and the radius of gyration R_g were evaluated using the Guinier approximation (Guinier & Fournet, 1955) assuming that at very small angles ($s < 1.3/R_g$) the intensity was represented by $I(s) = I(0)\exp[-(sR_g)^2/3]$. These parameters were computed from the entire scattering patterns using the indirect transform package *GNOM* (Semenyuk & Svergun, 1991), which provides the distance

distribution function $p(r)$ of the particle. The molecular mass of the proteins was calculated by comparison with the forward scattering from a reference solution of bovine serum albumin (BSA).

Initial *ab initio* modelling trials using *DAMMIN* or *GASBOR* from the *ATSAS* program suite (Svergun, 1997; Svergun *et al.*, 2001) did not yield unique models (high χ values) and could not be used for structural interpretation. The multimeric state of the protein in solution was tested by the program *OLIGOMER* (Konarev *et al.*, 2003), which calculates the theoretical volume fractions of different protein oligomers, and this was compared with experimental SAXS data. For this analysis the A₃ model from the heterodimeric structure of β -crustacyanin (PDB entry 1gka; Cianci *et al.*, 2002) was used to create a pseudodimer of H₂. The trimeric H₂ model was derived from the crystal packing of β -crustacyanin (Fig. 3c).

2.6. Complex reconstitution with astaxanthin

The complex reconstitution of either H₁ or H₂ or a mixture of both subunits in the presence of astaxanthin was probed using the acetone method (Zagalsky, 1985). The protein (about 5 mM) dissolved in a mixture of equal volumes of 40 mM Tris-HCl pH 7, 0.2 M ammonium sulfate and acetone was incubated with astaxanthin dissolved in acetone (about 12 mM). The mixture was dialyzed overnight at 277 K in the dark against phosphate buffer pH 7. Protein solutions were analyzed for absorption spectra in the 240 and 700 nm range.

3. Results and discussion

3.1. H₁ subunit structure

The crystal structure of H₁ apo crustacyanin from *H. americanus* contains a homodimer in the crystallographic asymmetric unit.

The H₁ protein has 181 residues with the typical lipocalin folding of crustacyanins A₁ (Cianci *et al.*, 2001, 2002), C₁ (Gordon *et al.*, 2001) and C₂ (Habash *et al.*, 2004) (Figs. 2a–2d), with a β -barrel made up of two distinct β -sheets. Homodimerization is achieved *via* a close contact between the two β -strands of each subunit, similarly to the A₁ (Cianci *et al.*, 2001), C₁ (Gordon *et al.*, 2001) and C₂ (Habash *et al.*, 2004) crystal structures (see Fig. 4 of Gordon *et al.*, 2001). All the residues that differed between species were located on the protein surface in the crystal structure. Tyr16, Glu61 and Met99 are located on hairpins, Met30 on the first α -helix, Try66 on β -strand C and Gln145 on an α -helix.

Superimposition of the complete H₁ apo crustacyanin homodimer with its analogues leads to an r.m.s. deviation for C α atoms of 1.39 Å from the crystal structure of the apo A₁ subunit (PDB entry 1h91; Cianci *et al.*, 2001), of 1.41 Å from that of apo C₁ (PDB entry 14u1; Gordon *et al.*, 2001) and of 1.37 Å from that of apo C₂ (PDB code: 1sp2; Habash *et al.*, 2004). The comparison shows a good overall fit, in agreement with the high sequence homology.

The statistics of the Ramachandran plot for the apo H₁ subunit are reported in Table 2. The outlier residues are Tyr112 in both chains A and B, which are in a generously allowed region, similar to the previously reported apo CRTc subunits from *H. gammarus* (Cianci *et al.*, 2001). Other members of the lipocalin family have, as a feature, the torsion angles of this residue in a disallowed region of the Ramachandran plot (Cowan *et al.*, 1990; Zanotti *et al.*, 1993).

Apo subunits A₁ (Cianci *et al.*, 2001) and C₁ (Gordon *et al.*, 2001) from *H. gammarus* were crystallized in the presence of MPD, which could be found in the astaxanthin binding site of both proteins. Comparison of the MPD positions in the apo crustacyanin H₁ (this work), A₁ (Cianci *et al.*, 2001), C₁ (Gordon *et al.*, 2001) and C₂ (Habash *et al.*, 2004) subunits revealed that the MPD-binding

mode was preserved throughout the CRTC subunits from both *H. gammarus* and *H. americanus* given that no mutations were observed for the residues defining the MPD-binding pocket.

Moreover, the astaxanthin-binding site in the β -crustacyanin A_1 monomer corresponds to that of MPD in each H_1 monomer of the apo crustacyanin H_1 dimer. The superimposition of the three-dimensional structure of apo crustacyanin H_1 monomer *A*, for

instance, onto the structure of the A_1 monomer of β -crustacyanin (PDB entry 1gka; Cianci *et al.*, 2002) results in the superimposition of an astaxanthin molecule onto the positions of the pair of MPD molecules (Figs. 2*b*, 2*c*, 2*e* and 2*f*). MPD molecules are therefore mimicking the binding of parts of a carotenoid ring and the C14 atom. This is also evident for apo crustacyanin C_1 , C_2 and A_1 dimers. The superimposition of the residues of the apo H_1 crustacyanin that define the MPD-binding pocket (Gln37, Gln46, Val52, Asn54, Tyr56, Ser67, Ile95, Tyr97, Phe134 and Phe136) with their analogues in the A_1 monomer of β -crustacyanin (PDB entry 1gka; Cianci *et al.*, 2002) results in r.m.s. deviations of 0.9 Å for monomer *A* and 1.7 Å for monomer *B*. The larger r.m.s. deviation for monomer *B* is owing to Tyr56, which appears to be able to adopt a different conformation from monomer *A* (Fig. 2*e*) without affecting MPD binding (the r.m.s. deviation of Tyr56 in monomer *B* versus monomer *A* is 4.8 Å). The overall residue arrangement of the binding site of apo crustacyanin H_1 for the astaxanthin molecule is preserved.

3.2. H_2 subunit structure

The H_2 subunit shares 95% sequence identity with the A_3 subunit from *H. gammarus* (Cianci *et al.*, 2002). It is therefore highly likely that the H_2 subunit and the A_3 monomers adopt the same three-dimensional structure. The structure of the H_2 subunit from *H. americanus* was modelled on the known structure of the A_3 subunit from *H. gammarus* (PDB entry 1gka; Cianci *et al.*, 2002) and was thus predicted to have a lipocalin topology with a β -barrel made up of two distinct β -sheets (Fig. 3*a*).

The H_2 subunit structure model was then confirmed by SAXS. No conditions suitable for the production of protein crystals were found for this protein. A Kratky analysis of the SAXS data gave clear evidence for a folded H_2 subunit in solution. Further analysis of the SAXS data revealed that the H_2 subunit in solution is present in a dynamic equilibrium between monomeric and multimeric forms. Analysis using standard *ab initio* or rigid-body SAXS modelling did not give good results because of the polydispersity of the H_2 subunit in solution.

Several possible protein-oligomerization states were considered: monomer–dimer, monomer–dimer–trimer, monomer–trimer and dimer–trimer mixtures.

Optimum scattering curve fitting ($\chi = 0.84$) resulted from combination of the H_2 monomeric subunit (volume fractions of 65%) and its trimeric form (35%) predicted from the A_3 trimer in the crystal structure of β -crustacyanin (Cianci *et al.*, 2002), where it generates a crystallographic threefold axis (Figs. 3*b* and 3*c*).

Other combinations of monomeric and multimeric forms of the H_2 subunit tested with *OLIGOMER* resulted in no improvement in fit (Fig. 3*d*). The fact that only a mixture of the monomer and the trimer based on the crystallographic studies of the A_3 subunit could explain the scattering profile suggests not only that the apo crustacyanin H_2 subunit structure is conserved compared with apo crustacyanin A_3 , but also that the subunit preserves the same oligomerization properties in solution as observed in the crystal lattice. All of the residues that differed between species were located on the protein surface: Ala19, Pro165 and Gln166 are located on hairpins and Cys147 on an α -helix. They were not involved in generating intermolecular contacts (Fig. 3*b*). Within the putative apo H_2 subunit structure the Cys147 residue (associated with the T147C mutation) is directly opposite the Cys115 residue, thus potentially generating an intramolecular Cys115–Cys147 disulfide bridge, structurally similar to the observed Cys117–Cys150 disulfide bridge present in the H_1 subunit (Fig. 3*a*).




Figure 4
Reconstitution studies: (a) H_1 subunit incubated with astaxanthin, (b) H_2 subunit incubated with astaxanthin, (c) a solution containing both H_1 and H_2 subunits incubated with astaxanthin.

3.3. Complex reconstitution with astaxanthin

Recombination of apo H₁ protein singly with astaxanthin produced a bathochromic shift with λ_{max} = 580 nm (Fig. 4a), albeit with a low yield (OD₅₈₀/OD₂₈₅ = 0.6). A shoulder is also present with a peak at λ_{max} ≈ 545 nm (OD₅₄₅/OD₂₈₅ = 0.5), suggesting the presence of a second species in solution. For comparison, recombination of freshly prepared natural apo subunits from *H. gammarus* carapace singly with astaxanthin gives a dimeric-like product with λ_{max} = 585 nm in high yields when testing C₁ and C₂ apo proteins or a monomeric-like product with λ_{max} = 565 nm in low yields when testing A₁, A₂ and A₃ apo proteins (Quarmby *et al.*, 1977).

On the other hand, combination of astaxanthin with mixtures of the individual apo protein units separated by gel filtration or ion exchange gave only monomeric-like or dimeric-like products with λ_{max} = 565 nm (Zagalsky, 1985). The apo H₁ protein from *H. americanus* therefore has the ability to produce a bathochromic shift and to generate two recombination products with distinguishable λ_{max} at 580 and 545 nm, analogous to CRTC apo subunits from *H. gammarus*.

Recombination of apo H₂ subunit from *H. americanus* singly with astaxanthin gave a bathochromic shift with λ_{max} = 580 nm (Fig. 4b), albeit with a low yield. A shoulder is also present with a peak at λ_{max} ≈ 520 nm, suggesting the presence of a second species in solution. Previous reconstitution studies using natural apo H₂ subunit from *H. americanus* singly with astaxanthin showed a single peak at λ_{max} ≈ 540 nm after chromatofocusing (Milicua *et al.*, 1986).

The recombination of a mixture of H₁ and H₂ subunits with astaxanthin gave rise to a new single product attributable to β-crustacyanin and characterized by a single absorption maximum at λ_{max} = 570 nm (Fig. 4c) and with a yield (OD₅₇₀/OD₂₈₅ = 0.75) larger than those determined in the cases of the individual recombinant subunits in the presence of astaxanthin. For comparison, recombination of both types of freshly prepared natural apo subunits from *H. gammarus* with astaxanthin gives β-crustacyanin with variable but nevertheless good yields (OD₅₈₀/OD₂₈₅ > 2.5) and absorption maxima (565–585 nm) that differed according to the combination of apo proteins selected (Zagalsky, 1985; Quarmby *et al.*, 1977).

3.4. Genotype–phenotype correlation

Examination of the spectral properties of the subunits in complex with astaxanthin shed light on carapace coloration resulting from expression of CRTC and/or CRTA subunits (Table 3). In crustaceans the carapace colour is generated by subtractive mixing of the colours absorbed by the adducts. In crustaceans expressing only CRTC subunits the colours absorbed would be yellow (580 nm) and yellow–green (545 nm) similar to the H₁ subunit. By using Munsell’s colour wheel (Munsell, 1912; Supplementary Fig. 1¹) it can be determined that the complementary colours are violet and purple, respectively, thus generating a visible purple–violet colour. In the carapace of the shrimp *Alpheus alpheopsides* only CRTC subunits were found and the colour was reported to be blue (Wade *et al.*, 2009).

In crustaceans expressing only CRTA subunits the colours absorbed would be yellow (580 nm) and bluish–green (520 nm), similarly to the H₂ subunit, thus generating a visible purple–red colour. In the carapaces of *Palinurus cygnus* and *Dardanus megistos* only CRTA subunits are expressed and the reported colour is red (Wade *et al.*, 2009). Expression of the CRTA subunit alone may then

Table 3

Type of crustacyanin (CRTN) proteins present in Decapoda crustacea and their related colours.

Sample animal	Tissue	CRTN	Colour	Reference
<i>Panulirus cygnus</i>	Epithelium/carapace	A1/A2	Red	Wade <i>et al.</i> (2009)
<i>Panulirus ornatus</i>	Epithelium/carapace	—	Blue/orange	Wade <i>et al.</i> (2009)
<i>Panulirus versicolour</i>	Epithelium/carapace	A/C	Blue/green	Wade <i>et al.</i> (2009)
<i>Penaeus monodon</i>	Epithelium	A/C	Blue	Wade <i>et al.</i> (2009)
<i>Marsupenaeus japonicus</i>	Epithelium/EST	A/C	Red	Wade <i>et al.</i> (2009)
<i>Homarus americanus</i>	Carapace	A/C	Blue/green	Zagalsky (1985)
<i>Homarus gammarus</i>	Carapace	A/C	Blue/green	Zagalsky (1985)
<i>Cherax quadricarinatus</i>	Epithelium/carapace/EST	A/C	Blue	Wade <i>et al.</i> (2009)
<i>Alpheus</i> sp.	Whole animal	C	Blue	Wade <i>et al.</i> (2009)
<i>Macrobrachium rosenbergii</i>	Carapace	A(?) / C	Blue	Yang <i>et al.</i> (2011)
<i>Dardanus megistos</i>	Epithelium/carapace	A	Red	Wade <i>et al.</i> (2009)
<i>Gonadactylus smithii</i>	Epithelium/carapace	A	Green	Wade <i>et al.</i> (2009)

result in red hues and expression of the CRTC subunit alone results in blue hues.

In crustaceans expressing both CRTC and CRTA subunits the colours absorbed would be yellow (580 nm, β-crustacyanin-like complexes) to red (630 nm, α-crustacyanin-like complexes), similar to lobsters with visible blue to bluish–green colours. In the carapaces of *Palinurus versicolor*, *Penaeus monodon* and *Cherax quadricarinatus* both CRTC and CRTA subunits are expressed and their reported colours are blue or blue–green (Wade *et al.*, 2009). In *Macrobrachium rosenbergii* the MrLC gene encodes a lipocalin protein which has been shown to specifically bind astaxanthin (Yang *et al.*, 2011). Knockdown of the MrLC gene by RNA interference (RNAi) resulted in a shift in body colour from blue to orangish–red over the entire carapace (Yang *et al.*, 2011).

Apparent discrepancies in the correlation between genotype (the presence or absence of CRTC and/or CRTA crustacyanin genes) and phenotype (carapace colouration or hue) in crustaceans can potentially be explained either by different levels of protein expression, post-translational modifications, oligomerization (such as α-crustacyanin) or environmental and dietary factors (Wade *et al.*, 2009). The correlations between the presence of CRTC and/or CRTA crustacyanin genes in crustacean species and carapace colours, as reported in the literature, with the spectral properties of the subunit in complex with astaxanthin confirm this genotype–phenotype linkage.

Heterologous expression and purification of the two major recombinant crustacyanin subunits from *H. americanus* produce two apo subunits that are highly structurally similar to the corresponding subunits A₁, C₁ and C₂ from *H. gammarus* characterized in previous studies (Cianci *et al.*, 2002; Gordon *et al.*, 2001; Habash *et al.*, 2004). Moreover, complex-reconstitution studies of the recombinant proteins H₁ and H₂ with astaxanthin show the bathochromic shift of 85–95 nm typical of natural crustacyanin proteins from *H. gammarus*. Preparations of recombinant crustacyanin subunits provide several opportunities to further dissect the molecular basis of crustacean carapace coloration and its biotechnological applications.

The crystallographic coordinates (PDB entry 4alo) and structure-factor amplitudes for the H₁ subunit have been deposited in the Protein Data Bank. The theoretical coordinates for the H₂ subunit validated by SAXS are available from the corresponding author upon request.

MC thanks Dr G. Bourenkov for support during data collection. MPG is thanked for the provision of beam time at DORIS, DESY, Germany. Dr T. R. Schneider (EMBL Hamburg) is thanked for general support. MC is grateful to Dr P. Zagalsky, Professor J. R.

¹ Supplementary material has been deposited in the IUCr electronic archive (Reference: HV5216).

Helliwell (University of Manchester, England) and Dr I. Manolaridis (EMBL Hamburg) for helpful discussions.

References

- Altschul, S. F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W. & Lipman, D. J. (1997). *Nucleic Acids Res.* **25**, 3389–3402.
- Bartalucci, G., Coppin, J., Fisher, S., Hall, G., Helliwell, J. R., Helliwell, M. & Liaaen-Jensen, S. (2007). *Acta Cryst.* **B63**, 328–337.
- Bartalucci, G., Fisher, S., Helliwell, J. R., Helliwell, M., Liaaen-Jensen, S., Warren, J. E. & Wilkinson, J. (2009). *Acta Cryst.* **B65**, 238–247.
- Buchwald, M. & Jenks, W. (1968). *Biochemistry*, **7**, 844–859.
- Cheeseman, D. F., Zagalsky, P. F. & Ceccaldi, J. H. (1966). *Proc. R. Soc. Lond. B*, **164**, 130–151.
- Cianci, M., Rizkallah, P. J., Olczak, A., Raftery, J., Chayen, N. E., Zagalsky, P. F. & Helliwell, J. R. (2001). *Acta Cryst.* **D57**, 1219–1229.
- Cianci, M., Rizkallah, P. J., Olczak, A., Raftery, J., Chayen, N. E., Zagalsky, P. F. & Helliwell, J. R. (2002). *Proc. Natl Acad. Sci USA*, **99**, 9795–9800.
- Cowan, S. W., Newcomer, M. E. & Jones, T. A. (1990). *Proteins*, **8**, 44–61.
- Cruikshank, D. W. J. (1999). *Acta Cryst.* **D55**, 583–601.
- Durbeej, B. & Eriksson, L. A. (2004). *Phys. Chem. Chem. Phys.* **6**, 4190–4198.
- Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. (2010). *Acta Cryst.* **D66**, 486–501.
- Gordon, E. J., Leonard, G. A., McSweeney, S. & Zagalsky, P. F. (2001). *Acta Cryst.* **D57**, 1230–1237.
- Guinier, A. & Fournet, G. (1955). *Small Angle Scattering of X-rays*. New York: Wiley.
- Habash, J., Helliwell, J. R., Raftery, J., Cianci, M., Rizkallah, P. J., Chayen, N. E., Nneji, G. A. & Zagalsky, P. F. (2004). *Acta Cryst.* **D60**, 493–498.
- Helliwell, J. R. (2010). *Crystallogr. Rev.* **16**, 231–242.
- Helliwell, M. (2008). *Carotenoids*, Vol. 4, *Natural Functions*, edited by G. Britton, S. Liaaen-Jensen & H. Pfander, pp. 37–52. Basel: Birkhäuser Verlag.
- Ilagan, R. P., Christensen, R. L., Chapp, T. W., Gibson, G. N., Pascher, T., Polívka, T. & Frank, H. A. (2005). *J. Phys. Chem. A*, **109**, 3120–3127.
- Konarev, P. V., Volkov, V. V., Sokolova, A. V., Koch, M. H. J. & Svergun, D. I. (2003). *J. Appl. Cryst.* **36**, 1277–1282.
- Liu, J., Shelton, N. L. & Liu, R. S. (2002). *Org. Lett.* **4**, 2521–2524.
- McNicholas, S., Potterton, E., Wilson, K. S. & Noble, M. E. M. (2011). *Acta Cryst.* **D67**, 386–394.
- Millicua, J. C. G., Gárate, A. M., Barbon, P. G. & Gomez, R. (1986). *Comp. Biochem. Physiol. B Comp. Biol.* **85**, 621–626.
- Mueller-Dieckmann, J. (2006). *Acta Cryst.* **D62**, 1446–1452.
- Munsell, A. H. (1912). *Am. J. Psychol.* **23**, 236–244.
- Murshudov, G. N., Skubák, P., Lebedev, A. A., Pannu, N. S., Steiner, R. A., Nicholls, R. A., Winn, M. D., Long, F. & Vagin, A. A. (2011). *Acta Cryst.* **D67**, 355–367.
- Neugebauer, J., Veldstra, J. & Buda, F. (2011). *J. Phys. Chem. B*, **115**, 3216–3225.
- Otwinowski, Z. & Minor, W. (1997). *Methods Enzymol.* **276**, 307–326.
- Polívka, T., Frank, H. A., Enriquez, M. M., Niedzwiedzki, D. M., Liaaen-Jensen, S., Hemming, J., Helliwell, J. R. & Helliwell, M. (2010). *J. Phys. Chem. B*, **114**, 8760–8769.
- Quarby, R., Norden, D. A., Zagalsky, P. F., Ceccaldi, H. J. & Dumas, R. (1977). *Comp. Biochem. Physiol. B Comp. Biol.* **56**, 55–61.
- Rhys, N. H., Wang, M.-C., Jowitt, T. A., Helliwell, J. R., Grossmann, J. G. & Baldock, C. (2011). *J. Synchrotron Rad.* **18**, 79–83.
- Roessle, M. W., Klaering, R., Ristau, U., Robrahn, B., Jahn, D., Gehrman, T., Konarev, P., Round, A., Fiedler, S., Hermes, C. & Svergun, D. (2007). *J. Appl. Cryst.* **40**, s190–s194.
- Semenyuk, A. V. & Svergun, D. I. (1991). *J. Appl. Cryst.* **24**, 537–540.
- Stepanyan, R., Day, K., Urban, J., Hardin, D. L., Shetty, R. S., Derby, C. D., Ache, B. W. & McClintock, T. S. (2006). *Physiol. Genomics*, **25**, 224–233.
- Strambi, A. & Durbeej, B. (2009). *J. Phys. Chem. B*, **113**, 5311–5317.
- Svergun, D. I. (1997). *J. Appl. Cryst.* **30**, 792–797.
- Svergun, D. I., Petoukhov, M. V. & Koch, M. H. J. (2001). *Biophys. J.* **80**, 2946–2953.
- Wade, N. M., Tollenaere, A., Hall, M. R. & Degnan, B. M. (2009). *Mol. Biol. Evol.* **26**, 1851–1864.
- Wald, G., Nathanson, N., Jenks, W. & Tarr, E. (1948). *Biol. Bull. Mar. Biol.* **95**, 249–250.
- Wijk, A. A. van, Spaans, A., Uzunbajakava, N., Otto, C., de Groot, H. J., Lugtenburg, J. & Buda, F. (2005). *J. Am. Chem. Soc.* **127**, 1438–1445.
- Winn, M. D. *et al.* (2011). *Acta Cryst.* **D67**, 235–242.
- Yang, F., Wang, M.-R., Ma, Y.-G., Ma, W.-M. & Yang, W.-J. (2011). *J. Exp. Zool. A Ecol. Genet. Physiol.* **315**, 562–571.
- Zagalsky, P. F. (1985). *Methods Enzymol.* **111**, 216–247.
- Zagalsky, P. F. & Tidmarsh, M.-L. (1985). *Comp. Biochem. Physiol. B Comp. Biol.* **80**, 599–601.
- Zanotti, G., Ottonello, S., Berni, R. & Monaco, H. L. (1993). *J. Mol. Biol.* **230**, 613–624.