

Poor self-reported health and its association with biomarkers among Canadian Inuit

Helga Saudny¹, Zhirong Cao¹ and Grace M. Egeland^{1,2,3*}

¹Centre for Indigenous Peoples' Nutrition and Environment (CINE), School of Dietetics and Human Nutrition, McGill University, Ste. Anne de Bellevue, QC, Canada; ²Norwegian Institute of Public Health, Bergen, Norway; ³Department of Public Health and Primary Health Care, University of Bergen, Bergen, Norway

Objectives. To determine the extent to which demographic characteristics, clinical measurements and biomarkers were associated with poor self-reported health (SRH) among Inuit adults in the Canadian Arctic.

Study design. Cross-sectional survey was adopted as the study design.

Methods. The International Polar Year Inuit Health Survey carried out in 36 Canadian Arctic communities in 2007 and 2008 included Inuit men and women, aged 18 years or older, recruited from randomly selected households. The main outcome measure was SRH, which was dichotomized into good health (excellent, very good and good responses) and poor health (fair and poor responses).

Results. Of the 2,796 eligible households, 1,901 (68%) households and 2,595 participants took part in the survey. The weighted prevalence of poor SRH was 27.8%. Increasing age was significantly associated with poor SRH. The relative risk ratios for poor SRH was 2.0 (95% confidence interval [CI] 1.3–3.1) for men aged 50 years or older and 2.3 (95% CI 1.7–3.0) for women aged 50 years or older, compared with men and women aged 29 years or younger. After adjusting for age, gender and body mass index, poor SRH was significantly associated with smoking status (odds ratio [OR] = 1.5; CI 1.1–2.0), at-risk fasting glucose levels (≥ 6.1 mmol/L) (OR = 2.5; 95% CI 1.5–4.2) and elevated hs C-reactive protein levels ($> 3 - \leq 10$ mg/L) (OR = 2.1; 95% CI 1.4–3.1). Poor SRH was also significantly associated with a hypertriglyceridemic waist phenotype (high-risk waist circumference ≥ 102 cm for men and ≥ 88 cm for women with high triglyceride levels, ≥ 1.7 mmol/L), adjusted for age and gender, OR = 1.6; 95% CI 1.1–2.3.

Conclusions. Clinically relevant indicators of chronic disease risk were related to subjective assessment of SRH among Inuit.

Keywords: *Inuit; health research; self-reported health; biomarkers*

Received: 19 April 2012; Revised: 25 July 2012; Accepted: 26 July 2012; Published: 22 August 2012

Individuals participating in health research are often asked to rate their overall health by responding to a single question: “In general, would you say your health is excellent, very good, good, fair or poor”. This subjective and simple assessment of overall health is a widely used and recommended population health assessment tool (1–3) and has consistently been shown to predict mortality (4–6), functional decline (7,8), future morbidity (3,9), advanced cancer survival (10), clinical outcome measures (11,12) and other dimensions of health (13).

When asked to respond to this question, individuals are expected to quickly take into account their understanding of health and knowledge of their own health. The response would be a composite of known medical

diagnoses or conditions, cultural concepts of health, prescribed drugs, awareness of bodily sensations and symptoms, age, functional ability and positive or negative dispositions, among other traits (14–16).

However, only limited data are available explaining the relationship between biological processes and self-reported health (SRH). In a large community-based study, white blood cell count and hemoglobin were significantly associated with SRH even after adjusting for various diseases and socio demographic characteristics (17). In other recent studies, a strong association between inflammatory cytokines and poor SRH was noted (18–20), and a consistent association between HDL-cholesterol and good SRH was shown for men and women in different age groups in the Oslo Health Study (21).

The objective of this article is to investigate the association between demographic characteristics, clinical measurements and biomarkers with poor SRH and thus add to the understanding of how biological processes may play a part in the assessment of SRH. Further, by studying this relationship among Inuit, we can also examine if SRH has wider cross-cultural implications beyond the North American and European populations on which most previous research on SRH has been based.

Material and methods

Study design

The International Polar Year Inuit Health Survey (IPY-IHS) is a multifaceted, cross-sectional health survey conducted in the late summer and fall of 2007 and 2008 in 33 coastal and 3 inland communities in the Inuvialuit Settlement Region (ISR) of the Northwest Territories, Nunavut and the Nunatsiavut region of Labrador in Canada. All coastal communities were visited by the Canadian Coast Guard Ship (CCGS) *Amundsen*, which was equipped with research and laboratory facilities. The inland communities were visited by air by separate research crews. Communities provided the researchers with numbered housing lists from which a random sample of households was selected using a random numbers table or computer-generated random digits. All Inuit adults, aged 18 years or older, within the selected households were eligible to participate. Three attempts were made to contact the selected households. Households were visited by locally trained research assistants and survey team members, who explained the study protocol, obtained written consent from individuals and booked clinic appointments on the ship.

Participatory process

The IPY-IHS was developed using a participatory health research approach (22,23). Memoranda of Agreements were developed with steering committees in each of the 3 regions, consisting of representatives of Inuit communities and regional health officials, outlining responsibilities and processes to be followed. Similarly, separate community–university agreements were signed by all 36 local community councils, inviting the IPY-IHS teams into their communities.

Informed consent

In recognition of the strong oral traditions of the Inuit, a “visual” consent form, which duplicated the written informed consent form word-for-word and depicted all clinical and laboratory procedures, was created as a DVD in the appropriate Inuit languages. After watching the DVD, participants who consented to participating in the study signed the bilingual written consent form.

Questionnaires

Trained interviewers, who were bilingual in English and Inuit dialects, administered the questionnaires. One adult from each household completed a questionnaire concerning the environment at home, living conditions, access to country food and other questions about the household. All other eligible adults in the household then completed an individual questionnaire on current and past health, health-related behaviours, such as smoking, and socio-demographic information. Diet was assessed using both multiple-pass 24-hour food recalls and food frequency questionnaires. SRH was dichotomized by grouping excellent, very good and good responses into the “good” category, and fair and poor responses into the “poor” category.

Clinical measurements

Standing height was measured to the nearest 0.1 cm using a portable stadiometer (Road Rod 214 Portable Stadiometer, Seca, MD). Body weight was measured to the nearest 0.1 kg using the Tanita scale (Tanita TBF-300GS, Tanita Corporation of America, Inc., Arlington Heights, IL). Standard body type was assumed for all participants and 0.4 kg was subtracted for light clothing. Participants who had a pacemaker had their body weight measured with a Seca scale (Medical Scale Model 214, Seca Corp., Ontario, Canada). Body mass index (BMI) was calculated (kg/m^2) and classified into 3 categories: normal <25.0 , overweight 25.0–29.9 and obese ≥ 30 . Waist circumference was measured to the nearest 0.1 cm using a cloth retractable tape measure (ERP, Laval, Quebec, Canada) at the natural waist midway between the lowest rib margin and the lateral iliac crest. Waist circumference was dichotomized into “normal” (<102 cm for males and <88 cm for females) and “at risk” (≥ 102 cm for males and ≥ 88 cm for females) (24).

Blood pressure was measured, at rest, to the nearest 1 mm Hg using a BpTRU™ Vital Signs Monitor (VSM MedTech Ltd., Coquitlam, BC, Canada). Three readings, with a 2-minute rest period between readings, were recorded for each participant and the mean of the 3 readings was used in the analysis, and blood pressure was categorized as low-risk without diabetes, as at-risk with or without diabetes or on medication. Diabetes was characterized as having a fasting blood glucose level ≥ 7.0 mmol/L or OGTT ≥ 11.1 mmol/L or taking anti-diabetic medication.

Blood sampling

Participants were asked to fast (water was allowed) for at least 8 hours. Nurses carried out venipuncture and a maximum of 45 ml of blood was centrifuged on-site and decanted into appropriate tubes and stored at -80°C before being transported to the Center for Indigenous Peoples Nutrition and Environment (CINE) for laboratory analyses.

Fasting plasma and serum were used for the analyses of glucose, lipids and high sensitivity C-reactive protein (hsCRP), respectively. Analyses were carried out at Nutrasource Diagnostics, Guelph, ON (Life Labs/Gamma Dynacare and the Montreal General Hospital, Canada).

Fasting glucose results were classified into normal, <6.1 mmol/L; or at-risk, ≥6.1 mmol/L; or taking anti-diabetic medication. Blood lipid concentrations were categorized according to the Canadian Heart and Stroke Foundation’s cut-off values (24). HDL-cholesterol, normal >1.0 for males and >1.3 for females, abnormal ≤1.0 mmol/L for males and ≤1.3 mmol/L for females; at-risk cholesterol, normal LDL-cholesterol <5.0 or total cholesterol/HDL-cholesterol <6.0; and at-risk: LDL-cholesterol ≥5.0 or ratio of total cholesterol/HDL-cholesterol ≥6.0 mmol/L; triglycerides, normal <1.7 and abnormal ≥1.7 mmol/L. hsCRP concentrations were classified into 3 groups, low risk (<1.0 mg/L), average risk (≥1.0 – ≤3.0 mg/L) and high risk (>3.0– ≤10 mg/L), excluding 149 (6.75%) participants with values indicating an acute infection (>10 mg/L) (25). The hypertriglyceridemic waist phenotype was classified into 4 categories—normal waist with normal triglyceride levels, normal waist with high triglyceride levels, at-risk waist with normal triglyceride levels and at-risk waist with high triglyceride levels (26).

Statistical analyses

Data were analyzed using SAS version 9.2 (SAS Institute, Cary, NC). Weighted data were used in all analyses. Weight 1 was based on the total number of households in each community and the number of participating households per community; weight 2 was based on the total number of people in each household and the number of people sampled per household; weight 3 was used to correct for differences in the sex distribution between our sample and it’s population distribution. Risk ratios and 95% confidence intervals (CI) predicting poor SRH by age and gender were calculated and multivariate logistic regression was used to evaluate the relationship between demographic characteristics, clinical measurements and

biomarkers with poor SRH, adjusting for age, gender and BMI. Odds ratios (OR) and 95% CI were calculated.

Results

The logistics of the survey and the movement of the ship (depending on tide tables) varied across the Arctic. Generally, very small communities were overrepresented as the ship could handle approximately 40 participants per day. A total of 2,796 Inuit households were approached from which 1,901 households participated (68%), with an average of 1.38 participants per household (n = 2,595 participants, aged 18 years or older, 998 [38.5%] men and 1,597 [61.5%] women). The median number of participating households per community was 41, with a minimum of 17 and a maximum of 143. For individuals, the median was 54, with a minimum of 22 and a maximum of 172 individuals per community participating. The mean age of the survey population was 43 years (SD = 15) and the weighted prevalence of poor SRH was 27.8%. Poor SRH increased with advancing age, with a 2-fold higher risk for those aged ≥50 years compared to the 18–29 age group (Table I). The prevalence of poor SRH by categories of demographic characteristics and chronic disease risk factors identified no significant relationships with gender, primary language spoken in the home or in traditional food consumption, but identified a protective relationship for married individuals. In analyses adjusted for age and gender, poor SRH was related to a hypertriglyceridemic-waist phenotype. When further adjustments for age, gender and BMI were made, significant relationships persisted between chronic disease risk factors and poor SRH, where the prevalence of poor SRH was significantly higher among participants who had at-risk fasting blood glucose levels and high-risk hsCRP levels (Table II).

Discussion

The present study adds to the limited literature available on biomarkers as they relate to SRH and represents the first investigation, to our knowledge, among Inuit adults in the Canadian Arctic. The most notable findings

Table I. Weighted risk ratios (RR and 95% confidence intervals (CI) predicting self-rated poor health by age and gender; International Polar Year Inuit Health Survey, 2007–2008

Age group (years)	Men			Women		
	(%)	95% CI	RR (95% CI)	(%)	95% CI	RR (95% CI)
18–29	19.7	11.8–27.5	1.0	19.3	14.5–24.0	1.0
30–39	20.4	13.7–27.1	1.0 (0.6–1.7)	24.0	18.7–29.3	1.2 (0.9–1.7)
40–49	25.0	17.4–32.5	1.3 (0.8–2.1)	19.8	14.7–24.9	1.0 (0.7–1.5)
≥50	39.0	31.4–46.7	2.0** (1.3–3.1)	43.7	37.6–49.9	2.3*** (1.7–3.0)

p < 0.01; *p < 0.001.

Table II. Weighted prevalence and adjusted odds ratios^a (OR) and 95% confidence intervals (CI) of self-reported poor health by demographic and health indicators; International Polar Year Inuit Health Survey, 2007–2008

Variable	% Poor health (95% CI)	OR (95% CI)
Marital status		
Single (single, separated, divorced, widowed)	29.8 (25.5–34.1)	1.0
Married (married, common law partner)	26.9 (23.9–30.0)	0.8* (0.6–1.0)
Smoking at time of interview		
No	29.7 (25.2–34.2)	1.0
Yes	27.0 (24.0–30.0)	1.5* (1.1–2.0)
Hypertension (mm Hg)		
Low risk without diabetes	24.2 (21.6–26.9)	1.0
At risk with or without diabetes or taking antihypertensive medication ^b	36.0 (30.2–41.7)	1.0 (0.8–1.4)
Body mass index (BMI)^c		
Normal weight	22.8 (19.0–26.7)	1.0
Overweight (25.0–29.9)	25.8 (21.1–30.4)	1.1 (0.8–1.5)
Obese (≥ 30)	32.5 (28.1–36.9)	1.4* (1.0–1.9)
At-risk cholesterol (mmol/L)^d		
Normal	26.8 (24.2–29.4)	1.0
At-risk LDL ≥ 5.0 or total/HDL ≥ 6.0	36.9 (29.2–44.6)	1.1 (0.7–1.6)
At-risk fasting glucose (mmol/L)		
Normal	25.4 (22.8–27.9)	1.0
At-risk ≥ 6.1 or taking anti-diabetic medication	57.2 (46.4–68.0)	2.5* (1.5–4.2)
Hypertriglyceridemic-waist phenotype^c		
Normal waist with normal triglycerides < 1.7	23.8 (20.0–27.6)	1.0
Normal waist with high triglycerides ≥ 1.7	26.0 (16.2–35.7)	1.0 (0.6–1.7)
At-risk waist ^e with normal triglycerides < 1.7	26.8 (22.5–31.1)	1.0 (0.7–1.4)
At-risk waist ^e with high triglycerides ≥ 1.7	36.7 (29.3–44.1)	1.6* (1.1–2.3)
hsC-reactive protein (mg/L)		
Low risk	19.1 (15.7–22.4)	1.0
Average risk 1.0– ≤ 3.0	28.0 (23.9–32.2)	1.3 (0.9–1.7)
High risk > 3.0 – ≤ 10	40.4 (34.1–46.8)	2.1* (1.4–3.1)

^aAdjusted for age (years), gender and BMI.

^bAt risk: non-diabetics, SBP ≥ 140 , DBP ≥ 90 mm Hg; diabetics, SBP ≥ 130 , DBP ≥ 80 mm Hg.

^cAdjusted for age (years) and gender.

^dOr taking cholesterol lowering agents.

^eAt-risk waist was defined as ≥ 102 cm for men and ≥ 88 cm for women.

* $p < 0.05$.

include the significantly higher OR of poor SRH associated with at-risk fasting glucose levels, at-risk waist circumference with high triglyceride levels and elevated hsCRP levels. Our results contribute to the limited data suggesting that a physiologic basis may be attributed to SRH, as suggested by others (14,17,27).

This is the first study to report on a relationship between at-risk fasting blood glucose levels and poor SRH, where those with elevated fasting blood glucose levels were 2.5 times more likely to perceive their health as poor compared with normoglycemic individuals. The poorer assessment of health that was associated with elevated levels of fasting glucose may be the result of

physical sensations associated with disease progression, such as fatigue and poor sleep patterns, and important covariates, such as obesity. Further work is needed to explore the intuitive assessment of SRH and its relation to glucose homeostasis, especially in light of recently published results that suggest that Inuit ethnicity no longer prevents the development of diabetes, and the hypertriglyceridemic-waist phenotype is related to risk of fasting glycemia and type 2 diabetes (26). Our results of poorer SRH with a hypertriglyceridemic-waist phenotype provide additional information on the usefulness of SRH, beyond the more traditional objective health indicators.

Another interesting finding is the strong association between higher levels of hsCRP and poorer SRH, which is in agreement with other reports, indicating that increasing circulating levels of inflammatory intermediaries can influence health perception and relate to poorer health assessment (17–19). Worth citing is a recent study among relatively healthy older adults that demonstrated that poorer self-rated health was significantly associated with higher levels of C-reactive protein. These associations remained after controlling for age, BMI, gender, objective health conditions, depressive symptoms and other health behaviours (20). Similarly, our results support the aforementioned findings as individuals with high-risk hsCRP levels are twice more likely to report poor health than individuals with low or average risk levels, adjusted for age, gender and BMI. CRP levels have been related to obesity and smoking status (28). Though these associations warrant further investigation, they are beyond the scope of this paper.

Our results, like others, suggest that biochemical changes occurring at the cellular level, when the immune system is challenged, or covariates, such as obesity, result in associations with poorer SRH. Jylhä and colleagues have reached a similar conclusion based on results from a large community-based sample, where white blood cell count was significantly associated with poorer SRH, which remained after adjusting for a number of diseases, clinical measures, and socio-demographic conditions. The researchers concluded that SRH acted as a surrogate for disease and reflected subclinical physiological conditions (17). Likewise, a recent study of healthy Japanese men and women concluded that poor SRH was positively correlated with circulating immune markers even after adjusting for confounders (29).

Very few studies have investigated the relationship between blood lipid levels and SRH. The Oslo Health Study suggested that increasing levels of HDL-cholesterol increased the odds of reporting good health in most age groups and both sexes (21). In our study, at-risk cholesterol levels were not associated with poor SRH.

The view that older people are preoccupied with poor health may be exaggerated. While poor SRH and old age are clearly associated with mortality, (14) when overall health and not mortality is the outcome, the literature is not as convincing. Older subjects have been noted to report better health than younger subjects (30–32), but the literature provides contradictory results (33,34). Also, women have been noted to report poorer health than men (35), which is not supported by our study as both men and women in the ≥ 50 age group were 2 times more likely to rate their health as poor than their younger counterparts. The SRH response among those ≥ 50 years could reflect underlying subclinical or clinical diseases and/or disabilities that relate to the lower life expectancy

among Inuit, which is 67 years compared with 80 years for the general Canadian population (36).

Strengths

This study is the first to relate objectively measured biomarkers and clinical measurements to SRH in a large representative sample of Inuit adults from 3 regions across a vast geographic area in the Canadian Arctic. Nearly 20% of the adult Inuit population of the 3 regions participated in the survey (2,595 individuals) with 68% of households participating. A further strength of the study is its contribution of new evidence to the emerging body of research linking endocrine and immune-activated responses to SRH and the study's corroboration of existing information on clinical measurements and biomarkers. Furthermore, this study provides additional evidence of the transcultural nature of SRH and its validity among Inuit and possibly other Indigenous populations (37).

Limitations

The study is not without its limitations. The absolute value of the adjusted OR should be interpreted with caution as they tend to exaggerate the true relative risk for highly prevalent outcomes. Also, access to medical chart data was not available at the time of analyses, but nurses visually inspected and recorded medications taken and those taking medications for diabetes or dyslipidemia were excluded from analyses of fasting blood glucose and lipid-related biomarkers. Furthermore, we excluded individuals with hsCRP levels exceeding 10 mg/L from the hsCRP analysis, evidence of acute inflammation or other organic disease states that could have influenced SRH. However, only 1 marker of inflammation was available for analyses. Another limitation is that study participants received their anthropometric results during their clinic visit and prior to being interviewed, and knowledge of their obesity status may have influenced their responses to the SRH questions.

Finally, an inherent weakness is, of course, the subjective nature of SRH assessment, which is likely influenced by multiple and complex dimensions of health and perceptions, cultural conventions, willingness to present a positive or negative self-image, past and present health experiences and knowledge and, last but not least, an optimistic or pessimistic temperament (14). However, given these inherent complexities in SRH, the current findings are all the more notable.

Future research

More research linking endocrine and immune-activated response measures to SRH is needed. Also, more needs to be learned about the process that goes into the evaluation of self-perceived health. More studies linking SRH to biological processes and to self-reported and diagnosed

diseases will improve our understanding of the relationship between biological processes and SRH.

Conclusion and policy implications

The findings provide evidence of the trans-cultural validity of SRH for population-based health research. Further, the findings provide health care professionals with greater confidence in the usefulness of incorporating one simple question on SRH in their patient contacts.

Acknowledgements

We graciously thank all those who participated in the Canadian IPY-IHS. We acknowledge all the dedicated community research assistants, nurses and staff who made the survey possible. We also extend our warmest gratitude to the 3 steering committees representing 3 Inuit land claim regions who guided the survey from its inception to results dissemination.

Conflict of interest and funding

Funding was provided by the Government of Canada Federal Program for International Polar Year, Canadian Institutes of Health Research, Health Canada, Indian and Northern Affairs Canada, Government of Nunavut, Arctic-Net, Canada Research Chair Program and the Canadian Foundation for Innovation.

Ethical approval: Ethical approval for the research was obtained from Faculty of Medicine, Institutional Review Board, McGill University. Research licences were obtained from the Nunavut Research Institute and the Aurora Research Institute, regulatory agencies for scientific research in Canada's territories.

References

- de Bruin A, Picavet H, Nossikov A. Health interviews survey: towards international harmonization of methods and instruments. WHO Reg Pub Eur Ser. 1996;58:i-xiii-1-161.
- Robine JM, Jagger C, the Euro-Reves Group. Creating a coherent set of indicators to monitor health across Europe. The Euro-REVES 2 project. Eur J Pub Health. 2003;13 (Suppl):6-14.
- May M, Lawlor DA, Brindle P, Patel R, Ebrahim S. Cardiovascular disease risk assessment in older women: can we improve on Framingham? British Women's Heart and Health prospective cohort study. Heart. 2006;92:1396-401.
- Mossey JM, Shapiro E. Self-rated health: a predictor of mortality among the elderly. Am J Public Health. 1982;72: 800-8.
- Kaplan GA, Camacho T. Perceived health and mortality: a nine-year follow-up of the human population laboratory cohort. Am J Epidemiol. 1983;117:292-304.
- Nielsen AB, Siersma V, Hiort LC, Drivsholm T, Kreiner S, Hollnagel H. Self-rated general health among 40-year-old Danes and its association with all-cause mortality at 10-, 20-, and 29 years follow-up. Scand J Public Health. 2008;36: 3-11.
- Mor V, Wilcox V, Rakowski W, Hiris J. Functional transitions among the elderly: patterns, predictors, and related hospital use and clinical outcome. Am J Public Health. 1994;84:1274-80.
- Hubbard RA, Inoue LYT, Diehr P. Joint modeling of self-rated health and changes in physical functioning. J Am Stat Assoc. 2009;104:912-28.
- Moller L, Kristensen TS, Hollnagel H. Self-rated health as a predictor of coronary heart disease in Copenhagen, Denmark. J Epidemiol Community Health. 1996;50:423-8.
- Shadbolt B, Barresi J, Craft P. Self-rated health as a predictor of survival among patients with advanced cancer. J Clin Oncol. 2002;20:2514-9.
- Goldman N, Gleit DA, Chang MC. The role of clinical risk factors in understanding self-rated health. Ann Epidemiol. 2004;14:49-57.
- Lainscak M, Farkas J, Inkrot S, Gelbrich G, Neskovic AN, Rau T, et al. Self-rated health predicts adverse events during beta-blocker treatment: the CIBIS-ELD randomized trial analysis. Int J Cardiol. Available online 8 June 2011; doi:10.1016/j.ijcard.2011.05.037.
- Smith PM, Glazier RH, Sibley LM. The predictors of self-rated health and the relationship between self-rated health and health service needs are similar across socioeconomic groups in Canada. J Clin Epidemiol. 2010;63:412-21.
- Jylhä M. What is self-rated health and why does it predict mortality? Towards a unified conceptual model. Soc Sci Med. 2009;69:307-16.
- Mora PA, DiBonaventura MD, Idler E, Leventhal EA, Leventhal H. Psychological factors influencing self-assessments of health: toward an understanding of the mechanisms underlying how people rate their own health. Ann Behav Med. 2008;36:292-303.
- Nielsen ABS, Gannik D, Siersma V, de Fine Olivarius N. The relationship between HbA_{1c} level, symptoms and self-rated health in type 2 diabetic patients. Scand J Prim Health Care 2011;29:157-64.
- Jylhä M, Volpato S, Guralnik JM. Self-rated health showed a graded association with frequently used biomarkers in a large population sample. J Clin Epidemiol. 2006;59:465-71.
- Uden AL, Andreasson A, Elofsson S, Brismars K, Mathsson L, Ronnelid J, et al. Inflammatory cytokines, behavior and age as determinants of self-rated health in women. Clin Sci. 2007; 112:363-73.
- Lekander M, Elofsson S, Neve IM, Hansson LO, Uden AL. Self-rated health is related to levels of circulating cytokines. Psychosom Med. 2004;66:559-63.
- Christian LM, Glaser R, Porter K, Malarkey WB, Beversdorf D, Kiecolt-Glaser JK. Poorer self-rated health is associated with elevated inflammatory markers among older adults. Psychoneuroendocrinology. 2011;36:1495-504.
- Tomten SE, Hostmark AT. Self-rated health showed a consistent association with serum HDL-cholesterol in the cross-sectional Oslo Health Study. Int J Med Sci. 2007;4:278-87.
- WHO and CINE. Indigenous peoples and participatory health research. Geneva: WHO and Centre for Indigenous Peoples' Nutrition and Environment; 2003.
- ITK and NRI. Negotiating research relationships with Inuit communities: a guide for researchers. In: Nickels S, Shirley J, Laidler G, editors. Ottawa and Iqaluit: Inuit Tapiriit Kanatami and Nunavut Research Institute; 2007. 38 p.
- Heart and Stroke Foundation. Living with Cholesterol. Cholesterol and healthy living. p. 1-18. [cited 2012 February 13]. Available from: http://www.heartandstroke.qc.ca/site/c.pkI0L7MMJrE/b.5059373/k.FC5B/Heart_Disease__Living_with_Cholesterol.htm.
- Ridker PM, Cook N. Clinical usefulness of very high and very low levels of C-reactive protein across the full range of Framingham risk scores. Circulation. 2004;109:1955-9.

26. Egeland GM, Cao Z, Young TK. Hypertriglyceridemic-waist phenotype and glucose intolerance among Canadian Inuit: the International Polar Year Inuit Health Survey for Adults 2007–2008. *CMAJ*. 2011;183:1022.
27. Dantzer R, Kelley KW. Stress and immunity: an integrated view of relationships between the brain and the immune system. *Life Sci*. 1989;44:1995–2008.
28. Levitzky YS, Guo CY, Rong J, Larson MG, Walter RE, Keaney JF, et al. Relation of smoking status to a panel of inflammatory markers. The Framingham offspring. *Atherosclerosis*. 2008;201:217–24.
29. Akinori N, Takahashi M, Otsuka Y, Swanson NG. Is self-rated health associated with blood immune markers in healthy individuals? *Int J Behav Med*. 2010;17:234–42.
30. Cockerham WC, Sharp K, Wilcox JA. Aging and perceived health status. *J Gerontol*. 1983;38:349–55.
31. Idler EL. Age differences in self-assessments of health: age changes, cohort differences, or survivorship? *J Gerontol*. 1993;48:S289–300.
32. Ishizaki T, Yoshida H, Suzuki T, Shibata H. The association between self-rated health status and increasing age among older Japanese living in a rural community over a 6-year period: a longitudinal data analysis. *Gerontology*. 2009;55:344–52.
33. Kawada T, Wakayama Y, Katsumata M, Inagaki H, Otsuka T, Hirata Y, et al. Patterns in self-rated health according to age and sex in a Japanese national survey 1989–2004. *Gender Med*. 2009;6:329–34.
34. Chen H, Cohen P, Kasen S. Cohort differences in self-rated health: evidence from a three-decade, community based, longitudinal study of women. *Am J Epidemiol*. 2007;166:439–46.
35. Bath PA. Differences between older men and women in the self-rated health – mortality relationship. *The Gerontologist*. 2003;43:387–95.
36. Government of Canada. *Healthy Canadians. A federal report on comparable health indicators*. 2008. Ottawa, ON: Minister of Health; [cited 2012 February 13]. Available from: <http://www.hc-sc.gc.ca/hcs-sss/pubs/system-regime/index-eng.php>
37. Sibthorpe B, Anderson I, Cunningham J. Self-assessed health among indigenous Australians: how valid is a global question? *Am J Public Health*. 2001;91:1660–3.

***Grace Egeland**

Department of Public Health and Primary Health Care
 University of Bergen
 Kalfarveien 31
 NO-5018 Bergen
 Norway
 Email: g.egeland@isf.uib.no