

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

5-Amino-3-(4*H*-1,2,4-triazol-4-yl)-1*H*-1,2,4-triazoleBing Liu,^{a,b} João P. C. Tomé,^c Luís Cunha-Silva^b and Filipe A. Almeida Paz^{a*}

^aDepartment of Chemistry, University of Aveiro, CICECO, 3810-193 Aveiro, Portugal, ^bREQUIMTE, Departamento de Química e Bioquímica, Faculdade de Ciências, Universidade do Porto, 4169-007 Porto, Portugal, and ^cDepartment of Chemistry, University of Aveiro, QOPNA, 3810-193 Aveiro, Portugal
Correspondence e-mail: filipe.paz@ua.pt


Received 25 July 2012; accepted 4 August 2012

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{N}-\text{C}) = 0.007$ Å; R factor = 0.073; wR factor = 0.226; data-to-parameter ratio = 10.1.

The asymmetric unit of the title compound, $\text{C}_4\text{H}_5\text{N}_7$, comprises two independent but virtually superimposable molecules. Each molecule is planar with the dihedral angles between the five-membered rings being 2.8 (3) and 2.1 (3)°. The crystal structure is formed by an extensive network of relatively strong $\text{N}-\text{H}\cdots\text{N}$ hydrogen-bond interactions. Individual molecules are arranged into supramolecular zigzag chains running parallel to [001] by way of the strongest $\text{N}-\text{H}\cdots\text{N}$ interactions. Adjacent chains are interconnected by rather long ($D\cdots A$ distances range from *ca* 3.00 to 3.03 Å) but highly directional (interaction angles above *ca* 173°) hydrogen bonds forming a supramolecular layer in the *bc* plane.

Related literature

For the synthesis of 1-butyl-3-methylimidazolium bromide and 1,2-diformylhydrazine, see: Liu *et al.* (2007); Parnham & Morris (2006). For the use of triazole molecules, see: Wang *et al.* (2012); Zhang *et al.* (2009). For previous research studies on crystal engineering approaches, see: Fernandes *et al.* (2011); Silva *et al.* (2011); Amarante *et al.* (2009); Paz & Klinowski (2007). For graph-set notation, see: Grell *et al.* (1999). For a description of the Cambridge Structural Database, see: Allen (2002).



Experimental

Crystal data

$\text{C}_4\text{H}_5\text{N}_7$
 $M_r = 151.15$
Monoclinic, $P2_1/n$
 $a = 13.765$ (3) Å
 $b = 5.9378$ (12) Å
 $c = 16.635$ (4) Å
 $\beta = 112.914$ (12)°
 $V = 1252.4$ (5) Å³
 $Z = 8$
Mo $K\alpha$ radiation
 $\mu = 0.12$ mm⁻¹
 $T = 293$ K
 $0.10 \times 0.05 \times 0.03$ mm

Data collection

Bruker X8 Kappa CCD APEXII diffractometer
Absorption correction: multi-scan (SADABS; Sheldrick, 1998)
 $T_{\min} = 0.988$, $T_{\max} = 0.996$
8003 measured reflections
2192 independent reflections
975 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.079$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.073$
 $wR(F^2) = 0.226$
 $S = 0.96$
2192 reflections
217 parameters
8 restraints
H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.52$ e Å⁻³
 $\Delta\rho_{\min} = -0.36$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

| $D-\text{H}\cdots A$ | $D-\text{H}$ | $\text{H}\cdots A$ | $D\cdots A$ | $D-\text{H}\cdots A$ |
|---|--------------|--------------------|-------------|----------------------|
| $\text{N1}-\text{H1A}\cdots\text{N14}^{\text{i}}$ | 0.90 (1) | 2.43 (2) | 3.239 (6) | 150 (4) |
| $\text{N1}-\text{H1B}\cdots\text{N4}^{\text{ii}}$ | 0.90 (1) | 2.11 (1) | 3.002 (6) | 173 (5) |
| $\text{N2}-\text{H2}\cdots\text{N13}^{\text{i}}$ | 0.90 (1) | 1.93 (2) | 2.772 (6) | 155 (5) |
| $\text{N8}-\text{H8A}\cdots\text{N7}$ | 0.90 (1) | 2.43 (2) | 3.264 (6) | 154 (4) |
| $\text{N8}-\text{H8B}\cdots\text{N11}^{\text{iii}}$ | 0.90 (1) | 2.14 (1) | 3.034 (6) | 176 (5) |
| $\text{N10}-\text{H10}\cdots\text{N6}$ | 0.90 (1) | 1.91 (2) | 2.777 (6) | 161 (5) |

Symmetry codes: (i) $x, y, z - 1$; (ii) $x, y - 1, z$; (iii) $x, y + 1, z$.

Data collection: APEX2 (Bruker, 2006); cell refinement: SAINT-Plus (Bruker, 2005); data reduction: SAINT-Plus; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: DIAMOND (Brandenburg, 2009); software used to prepare material for publication: SHELXTL.

The authors gratefully acknowledge the Fundação para a Ciência e a Tecnologia (FCT, MEC, Portugal), the European Union, QREN, FEDER, COMPETE for financial support by the strategic projects PEst-C/CTM/LA0011/2011 (to CICECO) and Pest-C/EQB/LA0006/2011 (to REQUIMTE), the R&D projects PTDC/CTM/100357/2008 and PTDC/QUI-QUI/098098/2008 (FCOMP-01-0124-FEDER-010785), as well as the post-doctoral research grant SFRH/BPD/47566/2008 (to BL). They also wish to thank FCT for the specific funding towards the purchase of the single-crystal X-ray diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK5136).

References

- Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.
- Amarante, T. R., Gonçalves, I. S. & Almeida Paz, F. A. (2009). *Acta Cryst.* **E65**, o1962–o1963.
- Brandenburg, K. (2009). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2005). *SAINTE-Plus*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2006). *APEX2*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Fernandes, J. A., Liu, B., Tomé, J. C., Cunha-Silva, L. & Almeida Paz, F. A. (2011). *Acta Cryst.* **E67**, o2073–o2074.
- Grell, J., Bernstein, J. & Tinhofer, G. (1999). *Acta Cryst.* **B55**, 1030–1043.
- Liu, B., Zhang, X.-C. & Wang, Y.-F. (2007). *Inorg. Chem. Commun.* **10**, 199–203.
- Parnham, E. R. & Morris, R. E. (2006). *Chem. Mater.* **18**, 4882–4887.
- Paz, F. A. A. & Klinowski, J. (2007). *Pure Appl. Chem.* **79**, 1097–1110.
- Sheldrick, G. M. (1998). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Silva, P., Vieira, F., Gomes, A. C., Ananias, D., Fernandes, J. A., Bruno, S. M., Soares, R., Valente, A. A., Rocha, J. & Paz, F. A. A. (2011). *J. Am. Chem. Soc.* **133**, 15120–15138.
- Wang, N., Feng, Y.-C., Shi, W., Zhao, B., Cheng, P., Liao, D.-Z. & Yan, S.-P. (2012). *CrystEngComm*, **14**, 2769–2278.
- Zhang, X.-C. & Liu, B. (2009). *Inorg. Chem. Commun.* **12**, 808–810.

supplementary materials

Acta Cryst. (2012). E68, o2700–o2701 [doi:10.1107/S1600536812034691]

5-Amino-3-(4*H*-1,2,4-triazol-4-yl)-1*H*-1,2,4-triazole**Bing Liu, João P. C. Tomé, Luís Cunha-Silva and Filipe A. Almeida Paz****Comment**

In the context of our interest in Crystal Engineering approaches (Wang *et al.* 2012; Fernandes *et al.*; 2011; Silva *et al.*, 2011; Amarante, Gonçalves *et al.*, 2009; Amarante *et al.*, 2009; Paz & Klinowski, 2007), we are currently designing new triazole molecules which could be simultaneously employed in the construction of novel Metal-Organic Frameworks (MOFs) or organic crystals. We note that this type of molecule has received considerable interest in the synthesis of polynuclear complexes and in the preparation of MOFs (Zhang *et al.*, 2009). A survey in the Cambridge structural Database (Allen, 2002) and in the literature revealed that the use of the asymmetrical bridging bitriazole molecule 5-amino-3-(1,2,4-triazol-4-yl)-1*H*-1,2,4-triazole (HATrtr) has not been reported to date, either in MOFs nor in organic crystals. Furthermore, to the best of our knowledge, its crystal structure has not been reported. Following our recent efforts we were able to isolate good-quality single-crystals of the title compound as a minor secondary phase and here we wish to report its crystal structure at ambient temperature.

The asymmetric unit of the title compound (I) comprises two whole molecules (*i.e.*, $Z' = 2$) of HATrtr as depicted in Fig. 1. We note that these two individual moieties are almost perfectly overlaid by assuming a combination of both inversion and molecular flexibility (maximum distance of *ca* 0.021 Å with RMS of *ca* 0.014 Å). Nevertheless, this feature is not described by crystal symmetry as the "head-to-tail" orientation of the molecules can not be described by the screw-axis parallel to the *b*-axis of the unit cell. Both molecules are almost planar with the dihedral angles between the 1,2,4-triazole rings being only of *ca* 2.8 and 2.1°. In addition, the medium planes of all non-hydrogen atoms of each molecule subtend an angle of just *ca* 7.1°, indicating that the molecules can also be envisaged as coplanar.

HATrtr is rich in groups capable of forming strong N—H···N hydrogen bonding interactions (see Table 1 for further geometrical details), which direct the crystal packing features of the title compound. The most striking intermolecular interactions concern the double donation of hydrogen atoms from each 5-amino-3-(1,2,4-triazole moiety to the 1,2,4-triazole group of an adjacent molecule (N1—H1A···N14, N2—H2···N13, N8—H8A···N7 and N10—H10···N6), forming a $R_2^2(7)$ graph set motif (dashed yellow lines in Fig. 2) (Grell *et al.*, 1999). This supramolecular motif constitutes the basis of the formation of a supramolecular zigzag tape parallel to the *c* axis, for which the repeating motif are the two molecules composing the asymmetric unit. Tapes are interconnected by additional N—H···N interactions (N1—H1B···N4 and N8—H8B···N11) which, despite being slightly long ($d_{D···A}$ ranging from 3.002 (6) to 3.034 (6) Å), are nevertheless highly directional with the interaction angles approaching linearity (of *ca* 173 and 176°). Noteworthy, these inter-chain connections are further strengthened by the presence of two weak C—H···N interactions (not represented): C3—H3···N3ⁱ with $d_{C···N} = 3.389$ (6) Å and $\angle(\text{CHN}) = 174^\circ$; C8—H8···N9ⁱⁱ with $d_{C···N} = 3.435$ (6) Å and $\angle(\text{CHN}) = 167^\circ$ (symmetry codes: (i) $x, 1 + y, z$; (ii) $x, -1 + y, z$). The combination of these N—H···N and C—H···N contacts can be described by the $R_2^2(8)$ graph set motif.

The hydrogen bonding interactions connecting adjacent molecular units and summarized in the previous paragraph lead to the formation of a two-dimensional supramolecular layer placed in the *bc* plane as shown in Fig. 2. Individual layers close pack parallel to the *a*-axis of the unit cell mediated by offset π - π contacts between HATrtr molecules (Fig. 3 and Table 2 for intercentroid distances). Noteworthy, the inter-layer distance along the *a* axis alternates: layers are disposed into pairs so to promote the aforementioned strong offset π - π contacts within one pair; connections between adjacent pairs of supramolecular layers are weaker (*i.e.* longer inter-centroid distances - not shown).

Experimental

3,5-Di(amino)-1,2,4-triazole (98% purity) and $\text{Mn}(\text{OAc})_2 \cdot 4\text{H}_2\text{O}$ (99%+ purity) were purchased from Sigma-Aldrich and were used as received without further purification. 5-Amino-3-(1,2,4-triazol-4-yl)-1*H*-1,2,4-triazole and 1,2-diformylhydrazine were prepared by methods reported in the literature (Liu *et al.*, 2007). 1-Butyl-3-methylimidazolium bromide ([BMI]Br) was also prepared according to literature procedures (Parnham & Morris, 2006) (pale-yellow oil; yield: 78%).

1,2-Diformylhydrazine (0.1 mol; 8.8095 g) and 3,5-di(amino)-1,2,4-triazole (0.05 mol; 4.9564 g) were mixed in a 25 mL Teflon-lined stainless-steel reaction vessel and heated at 170 °C for 3 days in a furnace. After this time, the vessel was allowed to cool slowly to ambient temperature yielding 5-amino-3-(1,2,4-triazol-4-yl)-1*H*-1,2,4-triazole (HATrtr) as a white microcrystalline powder. The solid was then washed with copious amounts of water and ethanol and dried at *ca* 60 °C for 24 h. Yield: 6.85 g, 67.4%.

HATrtr (0.2 mmol, 0.0407 g) and $\text{Mn}(\text{OAc})_2 \cdot 4\text{H}_2\text{O}$ (1.0 mmol, 0.2451 g) were mixed with *ca* 0.50 g of [BMI]Br in a 25 mL teflon-lined stainless-steel reaction vessel. The resulting mixture was heated to 110 °C for 7 days. The vessel was then allowed to cool to ambient temperature at a rate of *ca* 1 °C/h. Small colourless platelets of HATrtr were formed as a minor secondary product, whose crystal structure is reported in this manuscript.


Refinement

Hydrogen atoms bound to carbon were placed at their idealized positions with $\text{C}-\text{H} = 0.93 \text{ \AA}$ and included in the final structural model in the riding-motion approximation with isotropic displacement parameters fixed at $1.2 \times U_{\text{eq}}(\text{C})$.


Hydrogen atoms associated with nitrogen have been directly located from difference Fourier maps and were included in the structural model with the $\text{N}-\text{H}$ and $\text{H} \cdots \text{H}$ (only for the $-\text{NH}_2$ moieties) distances restrained to 0.900 (5) and 1.560 (5) Å, respectively, in order to ensure a chemically reasonable environment. These hydrogen atoms were refined using a riding-motion approximation with isotropic displacement parameters fixed at $1.5 \times U_{\text{eq}}(\text{N})$.

Computing details


Data collection: *APEX2* (Bruker, 2006); cell refinement: *SAINTE-Plus* (Bruker, 2005); data reduction: *SAINTE-Plus* (Bruker, 2005); program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 2009); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

**Figure 1**

Asymmetric unit of the title compound showing all non-hydrogen atoms represented as anisotropic displacement ellipsoids drawn at the 50% probability level and hydrogen atoms as small spheres with arbitrary radius. The labeling scheme for all non-hydrogen atoms is also provided.

**Figure 2**

Two-dimensional supramolecular network placed in the bc plane formed by the N—H \cdots N hydrogen bonding interactions between adjacent molecular units. Zigzag supramolecular tapes running parallel to the c axis of the unit cell are based on $R_2^2(7)$ graph set motifs (yellow dashed lines) formed between the 5-amino-3-(1,2,4-triazole) moiety of one molecule and the 1,2,4-triazole group from the adjacent one. Connections between adjacent tapes (dashed green lines) are ensured by additional N—H \cdots N interactions along [010]. For geometrical details on the represented hydrogen bonding interactions see Table 1.

**Figure 3**

(a) Schematic representation of the supramolecular π - π interactions between HATrtr molecules belonging to adjacent supramolecular layers. For clarity one layer is represented as pink bonds and all hydrogen atoms have omitted. For inter-centroid distances see Table 2. **(b)** Crystal packing of the title compound viewed along [001] direction, clearly showing the planar nature of the two-dimensional supramolecular layer.

5-Amino-3-(4H-1,2,4-triazol-4-yl)-1H-1,2,4-triazole

Crystal data

$C_4H_5N_7$

$M_r = 151.15$

Monoclinic, $P2_1/n$

Hall symbol: $-P 2_1n$

$a = 13.765 (3) \text{ \AA}$

$b = 5.9378 (12) \text{ \AA}$

$c = 16.635 (4) \text{ \AA}$

$\beta = 112.914 (12)^\circ$

$V = 1252.4 (5) \text{ \AA}^3$

$Z = 8$

$F(000) = 624$

$D_x = 1.603 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
 Cell parameters from 823 reflections
 $\theta = 2.7\text{--}21.7^\circ$
 $\mu = 0.12 \text{ mm}^{-1}$

$T = 293 \text{ K}$
 Prism, yellow
 $0.10 \times 0.05 \times 0.03 \text{ mm}$

Data collection

Bruker X8 Kappa CCD APEXII
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 ω / φ scans
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1998)
 $T_{\min} = 0.988, T_{\max} = 0.996$

8003 measured reflections
 2192 independent reflections
 975 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.079$
 $\theta_{\max} = 25.0^\circ, \theta_{\min} = 3.7^\circ$
 $h = -16 \rightarrow 16$
 $k = -7 \rightarrow 5$
 $l = -19 \rightarrow 19$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.073$
 $wR(F^2) = 0.226$
 $S = 0.96$
 2192 reflections
 217 parameters
 8 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.1189P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.52 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.36 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | <i>x</i> | <i>y</i> | <i>z</i> | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|------------|-------------|--------------|----------------------------------|
| N1 | 0.6176 (4) | -0.3457 (6) | -0.1686 (3) | 0.0465 (13) |
| H1B | 0.625 (4) | -0.477 (4) | -0.140 (3) | 0.070* |
| H1A | 0.625 (4) | -0.341 (7) | -0.2198 (16) | 0.070* |
| N2 | 0.6244 (3) | 0.0473 (7) | -0.1487 (3) | 0.0431 (12) |
| H2 | 0.624 (4) | 0.106 (9) | -0.1987 (18) | 0.065* |
| N4 | 0.6274 (3) | 0.2015 (6) | -0.0855 (3) | 0.0443 (12) |
| C2 | 0.6268 (4) | 0.0647 (8) | -0.0238 (3) | 0.0346 (12) |
| N3 | 0.6228 (3) | -0.1582 (6) | -0.0393 (2) | 0.0326 (10) |
| C1 | 0.6217 (4) | -0.1618 (8) | -0.1201 (3) | 0.0368 (12) |
| N6 | 0.6346 (3) | 0.1685 (7) | 0.1884 (3) | 0.0477 (12) |
| N7 | 0.6317 (3) | 0.3877 (7) | 0.1576 (3) | 0.0493 (12) |
| C3 | 0.6290 (4) | 0.3720 (8) | 0.0783 (3) | 0.0449 (14) |

| | | | | |
|-----|------------|-------------|-------------|-------------|
| H3 | 0.6272 | 0.4936 | 0.0426 | 0.054* |
| N5 | 0.6292 (3) | 0.1502 (6) | 0.0556 (2) | 0.0339 (10) |
| C4 | 0.6328 (4) | 0.0325 (9) | 0.1264 (3) | 0.0420 (13) |
| H4 | 0.6338 | -0.1238 | 0.1302 | 0.050* |
| N10 | 0.6263 (3) | 0.2101 (6) | 0.3519 (3) | 0.0417 (12) |
| H10 | 0.636 (4) | 0.166 (9) | 0.3037 (19) | 0.063* |
| N11 | 0.6323 (3) | 0.0579 (6) | 0.4167 (3) | 0.0383 (11) |
| C6 | 0.6227 (3) | 0.1927 (7) | 0.4747 (3) | 0.0312 (11) |
| N9 | 0.6124 (3) | 0.4162 (6) | 0.4575 (3) | 0.0350 (10) |
| C5 | 0.6144 (4) | 0.4198 (8) | 0.3774 (3) | 0.0341 (12) |
| N14 | 0.6344 (3) | -0.1263 (7) | 0.6590 (3) | 0.0441 (11) |
| N13 | 0.6262 (3) | 0.0938 (7) | 0.6861 (3) | 0.0487 (12) |
| C7 | 0.6202 (4) | 0.2269 (8) | 0.6221 (3) | 0.0440 (14) |
| H7 | 0.6136 | 0.3827 | 0.6229 | 0.053* |
| N12 | 0.6250 (3) | 0.1088 (6) | 0.5547 (2) | 0.0342 (10) |
| C8 | 0.6339 (4) | -0.1094 (8) | 0.5814 (3) | 0.0411 (13) |
| H8 | 0.6390 | -0.2311 | 0.5480 | 0.049* |
| N8 | 0.6036 (4) | 0.6025 (7) | 0.3278 (3) | 0.0509 (13) |
| H8A | 0.605 (4) | 0.589 (7) | 0.2746 (15) | 0.076* |
| H8B | 0.615 (4) | 0.737 (4) | 0.355 (3) | 0.076* |

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|-----------|-----------|-----------|--------------|-----------|--------------|
| N1 | 0.083 (3) | 0.031 (2) | 0.035 (3) | 0.001 (2) | 0.033 (3) | -0.005 (2) |
| N2 | 0.084 (3) | 0.030 (2) | 0.026 (3) | -0.003 (2) | 0.032 (3) | -0.001 (2) |
| N4 | 0.083 (3) | 0.033 (2) | 0.028 (3) | -0.001 (2) | 0.033 (2) | 0.000 (2) |
| C2 | 0.045 (3) | 0.036 (3) | 0.026 (3) | 0.004 (2) | 0.017 (3) | -0.002 (2) |
| N3 | 0.051 (3) | 0.030 (2) | 0.022 (2) | 0.0005 (17) | 0.020 (2) | 0.0000 (17) |
| C1 | 0.050 (3) | 0.036 (3) | 0.030 (3) | 0.001 (2) | 0.022 (3) | -0.001 (2) |
| N6 | 0.075 (3) | 0.051 (3) | 0.025 (3) | 0.007 (2) | 0.028 (2) | -0.002 (2) |
| N7 | 0.066 (3) | 0.044 (3) | 0.044 (3) | -0.008 (2) | 0.028 (3) | -0.008 (2) |
| C3 | 0.067 (4) | 0.042 (3) | 0.030 (3) | -0.005 (3) | 0.022 (3) | -0.011 (2) |
| N5 | 0.052 (3) | 0.035 (2) | 0.021 (3) | 0.0012 (19) | 0.022 (2) | -0.0004 (18) |
| C4 | 0.059 (4) | 0.045 (3) | 0.027 (3) | 0.005 (2) | 0.022 (3) | 0.003 (3) |
| N10 | 0.073 (3) | 0.032 (3) | 0.029 (3) | 0.004 (2) | 0.029 (3) | 0.004 (2) |
| N11 | 0.067 (3) | 0.031 (2) | 0.023 (2) | -0.0016 (19) | 0.024 (2) | -0.0013 (19) |
| C6 | 0.043 (3) | 0.033 (3) | 0.022 (3) | 0.000 (2) | 0.017 (2) | 0.002 (2) |
| N9 | 0.055 (3) | 0.029 (2) | 0.026 (2) | 0.0003 (18) | 0.021 (2) | -0.0082 (18) |
| C5 | 0.050 (3) | 0.028 (3) | 0.028 (3) | 0.001 (2) | 0.020 (3) | 0.000 (2) |
| N14 | 0.061 (3) | 0.045 (3) | 0.030 (3) | 0.002 (2) | 0.022 (2) | 0.006 (2) |
| N13 | 0.073 (3) | 0.048 (3) | 0.029 (3) | -0.002 (2) | 0.024 (2) | 0.003 (2) |
| C7 | 0.069 (4) | 0.038 (3) | 0.033 (3) | -0.001 (2) | 0.029 (3) | 0.000 (3) |
| N12 | 0.052 (3) | 0.030 (2) | 0.024 (3) | -0.0001 (18) | 0.020 (2) | -0.0002 (18) |
| C8 | 0.059 (4) | 0.039 (3) | 0.032 (3) | 0.003 (2) | 0.026 (3) | 0.004 (2) |
| N8 | 0.097 (4) | 0.030 (2) | 0.036 (3) | 0.001 (2) | 0.037 (3) | 0.002 (2) |

Geometric parameters (Å, °)

| | | | |
|-------------|------------|----------------|------------|
| N1—C1 | 1.345 (6) | N10—C5 | 1.346 (6) |
| N1—H1B | 0.900 (5) | N10—N11 | 1.385 (5) |
| N1—H1A | 0.899 (5) | N10—H10 | 0.901 (5) |
| N2—C1 | 1.335 (6) | N11—C6 | 1.299 (5) |
| N2—N4 | 1.382 (5) | C6—N9 | 1.353 (5) |
| N2—H2 | 0.899 (5) | C6—N12 | 1.410 (5) |
| N4—C2 | 1.312 (6) | N9—C5 | 1.344 (6) |
| C2—N3 | 1.346 (5) | C5—N8 | 1.336 (6) |
| C2—N5 | 1.403 (5) | N14—C8 | 1.293 (6) |
| N3—C1 | 1.337 (6) | N14—N13 | 1.401 (5) |
| N6—C4 | 1.303 (6) | N13—C7 | 1.302 (6) |
| N6—N7 | 1.394 (6) | C7—N12 | 1.346 (6) |
| N7—C3 | 1.307 (6) | C7—H7 | 0.9300 |
| C3—N5 | 1.370 (6) | N12—C8 | 1.360 (6) |
| C3—H3 | 0.9300 | C8—H8 | 0.9300 |
| N5—C4 | 1.353 (6) | N8—H8A | 0.899 (5) |
| C4—H4 | 0.9300 | N8—H8B | 0.900 (5) |
| | | | |
| C1—N1—H1B | 114 (3) | C5—N10—N11 | 109.6 (4) |
| C1—N1—H1A | 123 (3) | C5—N10—H10 | 129 (4) |
| H1B—N1—H1A | 120 (4) | N11—N10—H10 | 121 (4) |
| C1—N2—N4 | 110.0 (4) | C6—N11—N10 | 100.6 (4) |
| C1—N2—H2 | 134 (4) | N11—C6—N9 | 118.7 (4) |
| N4—N2—H2 | 116 (4) | N11—C6—N12 | 120.8 (4) |
| C2—N4—N2 | 100.2 (4) | N9—C6—N12 | 120.5 (4) |
| N4—C2—N3 | 118.1 (4) | C5—N9—C6 | 100.6 (4) |
| N4—C2—N5 | 120.5 (4) | N8—C5—N9 | 125.8 (4) |
| N3—C2—N5 | 121.4 (4) | N8—C5—N10 | 123.7 (4) |
| C1—N3—C2 | 101.1 (4) | N9—C5—N10 | 110.5 (4) |
| N2—C1—N3 | 110.6 (4) | C8—N14—N13 | 106.2 (4) |
| N2—C1—N1 | 122.8 (5) | C7—N13—N14 | 106.9 (4) |
| N3—C1—N1 | 126.6 (5) | N13—C7—N12 | 110.9 (4) |
| C4—N6—N7 | 107.4 (4) | N13—C7—H7 | 124.5 |
| C3—N7—N6 | 106.8 (4) | N12—C7—H7 | 124.5 |
| N7—C3—N5 | 110.1 (5) | C7—N12—C8 | 104.6 (4) |
| N7—C3—H3 | 124.9 | C7—N12—C6 | 127.8 (4) |
| N5—C3—H3 | 124.9 | C8—N12—C6 | 127.6 (4) |
| C4—N5—C3 | 105.1 (4) | N14—C8—N12 | 111.4 (4) |
| C4—N5—C2 | 127.7 (4) | N14—C8—H8 | 124.3 |
| C3—N5—C2 | 127.2 (4) | N12—C8—H8 | 124.3 |
| N6—C4—N5 | 110.6 (4) | C5—N8—H8A | 120 (3) |
| N6—C4—H4 | 124.7 | C5—N8—H8B | 117 (3) |
| N5—C4—H4 | 124.7 | H8A—N8—H8B | 120 (4) |
| | | | |
| C1—N2—N4—C2 | 0.4 (5) | C5—N10—N11—C6 | 0.3 (5) |
| N2—N4—C2—N3 | -0.7 (6) | N10—N11—C6—N9 | -0.9 (6) |
| N2—N4—C2—N5 | -179.9 (4) | N10—N11—C6—N12 | -179.6 (4) |
| N4—C2—N3—C1 | 0.7 (6) | N11—C6—N9—C5 | 1.0 (6) |

| | | | |
|-------------|------------|----------------|------------|
| N5—C2—N3—C1 | 180.0 (4) | N12—C6—N9—C5 | 179.8 (4) |
| N4—N2—C1—N3 | 0.0 (6) | C6—N9—C5—N8 | 177.7 (5) |
| N4—N2—C1—N1 | 179.4 (4) | C6—N9—C5—N10 | -0.7 (5) |
| C2—N3—C1—N2 | -0.4 (5) | N11—N10—C5—N8 | -178.2 (5) |
| C2—N3—C1—N1 | -179.8 (5) | N11—N10—C5—N9 | 0.3 (5) |
| C4—N6—N7—C3 | -0.4 (5) | C8—N14—N13—C7 | 0.7 (5) |
| N6—N7—C3—N5 | 0.4 (6) | N14—N13—C7—N12 | -0.6 (6) |
| N7—C3—N5—C4 | -0.3 (5) | N13—C7—N12—C8 | 0.3 (6) |
| N7—C3—N5—C2 | 180.0 (4) | N13—C7—N12—C6 | -178.9 (4) |
| N4—C2—N5—C4 | -177.5 (5) | N11—C6—N12—C7 | 177.2 (4) |
| N3—C2—N5—C4 | 3.3 (7) | N9—C6—N12—C7 | -1.5 (7) |
| N4—C2—N5—C3 | 2.1 (7) | N11—C6—N12—C8 | -1.8 (7) |
| N3—C2—N5—C3 | -177.1 (4) | N9—C6—N12—C8 | 179.4 (4) |
| N7—N6—C4—N5 | 0.2 (5) | N13—N14—C8—N12 | -0.5 (5) |
| C3—N5—C4—N6 | 0.0 (5) | C7—N12—C8—N14 | 0.2 (5) |
| C2—N5—C4—N6 | 179.7 (4) | C6—N12—C8—N14 | 179.4 (4) |

Hydrogen-bond geometry (\AA , $^\circ$)

| $D-H\cdots A$ | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|------------------------------------|----------|-------------|-------------|---------------|
| N1—H1A \cdots N14 ⁱ | 0.90 (1) | 2.43 (2) | 3.239 (6) | 150 (4) |
| N1—H1B \cdots N4 ⁱⁱ | 0.90 (1) | 2.11 (1) | 3.002 (6) | 173 (5) |
| N2—H2 \cdots N13 ⁱ | 0.90 (1) | 1.93 (2) | 2.772 (6) | 155 (5) |
| N8—H8A \cdots N7 | 0.90 (1) | 2.43 (2) | 3.264 (6) | 154 (4) |
| N8—H8B \cdots N11 ⁱⁱⁱ | 0.90 (1) | 2.14 (1) | 3.034 (6) | 176 (5) |
| N10—H10 \cdots N6 | 0.90 (1) | 1.91 (2) | 2.777 (6) | 161 (5) |

Symmetry codes: (i) $x, y, z-1$; (ii) $x, y-1, z$; (iii) $x, y+1, z$.

Selected $\pi-\pi$ contacts

| $\pi-\pi$ interaction | $d(\text{Cg}\cdots\text{Cg})$ |
|--|-------------------------------|
| $\text{Cg}(1)\cdots\text{Cg}(2)$ ⁱ | 3.580 (3) |
| $\text{Cg}(3)\cdots\text{Cg}(4)$ ⁱⁱ | 3.700 (3) |

Symmetry codes: (i) $1-x, -y, -z$; (ii) $1-x, -y, 1-z$; Cg(1): centroid of the ring formed by N2, N3, N4, C1 and C2; Cg(2): centroid of the ring formed by N5, N6, N7, C3 and C4; Cg(3): centroid of the ring formed by N9, N10, N11, C5 and C6; Cg(4): centroid of the ring formed by N12, N13, N14, C7 and C8.