

The Binding Site of the V-ATPase Inhibitor Apicularen Is in the Vicinity of Those for Bafilomycin and Archazolid

Received for publication, April 13, 2012, and in revised form, July 11, 2012. Published, JBC Papers in Press, July 19, 2012, DOI 10.1074/jbc.M112.372169

Christin Osteresch[‡], Tobias Bender^{§¶}, Stephanie Grond[¶], Paultheo von Zezschwitz[¶], Brigitte Kunze^{||}, Rolf Jansen^{||}, Markus Huss^{‡¶}, and Helmut Wieczorek[‡]

From the [‡]Fachbereich Biologie/Chemie, Abteilung Tierphysiologie, Universität Osnabrück, Barbarastrasse 11, 49069 Osnabrück, the [§]Fachbereich Chemie, Philipps Universität Marburg, Hans-Meerwein Strasse, 35032 Marburg, the [¶]Institut für Organische Chemie, Eberhard Karls Universität Tübingen, Auf der Morgenstelle 18, 72076 Tübingen, and the ^{||}Helmholtz Zentrum für Infektionsforschung, Inhoffenstrasse 7, 38124 Braunschweig, Germany

Background: Apicularen is a specific V-ATPase inhibitor that binds to the V_O complex of the holoenzyme.

Results: Apicularen binds at the interface of the V_O subunits a and c.

Conclusion: The binding site for apicularen is in the vicinity of those for bafilomycin and archazolid.

Significance: We propose the first model of binding site arrangement for these three classes of V-ATPase inhibitors.

The investigation of V-ATPases as potential therapeutic drug targets and hence of their specific inhibitors is a promising approach in osteoporosis and cancer treatment because the occurrence of these diseases is interrelated to the function of the V-ATPase. Apicularen belongs to the novel inhibitor family of the benzolactone enamides, which are highly potent but feature the unique characteristic of not inhibiting V-ATPases from fungal sources. In this study we specify, for the first time, the binding site of apicularen within the membrane spanning V_O complex. By photoaffinity labeling using derivatives of apicularen and of the plecomacrolides bafilomycin and concanamycin, each coupled to ¹⁴C-labeled 4-(3-trifluoromethyl-diazirine-3-yl)benzoic acid, we verified that apicularen binds at the interface of the V_O subunits a and c. The binding site is in the vicinity to those of the plecomacrolides and of the archazolids, a third family of V-ATPase inhibitors. Expression of subunit c homologues from *Homo sapiens* and *Manduca sexta*, both species sensitive to benzolactone enamides, in a *Saccharomyces cerevisiae* strain lacking the corresponding intrinsic gene did not transfer this sensitivity to yeast. Therefore, the binding site of benzolactone enamides cannot be formed exclusively by subunit c. Apparently, subunit a substantially contributes to the binding of the benzolactone enamides.

Vacuolar-type ATPases (V-ATPases)² are found in the endomembrane system of all eukaryotic cells and in the plasma membranes of many animal cells. They energize multiple transport processes and regulate the pH in cells and organelles by coupling ATP hydrolysis to proton pumping (1, 2). Their heteromultimeric structure includes two main complexes, the catalytic V₁ complex with a subunit composition of A₃B₃CDE₃FG₃H and the proton translocating V_O complex with the subunits a, c_n, c'' (in fungi additional subunit c'), d, and

e (2, 3). ATP hydrolysis takes place at the hexameric headpiece built by three A catalytic subunits and three B regulatory subunits. The emerging free energy is converted into a rotation of the central stalk subunits D, F, and d and the ring formed by the proteolipid subunit c (2). For transmembrane transport, protons enter the proteolipid ring via the cytosolic hemichannel of subunit a and bind reversibly to the conserved essential glutamate in each c subunit. After rotation of the c-ring, protons leave via the luminal hemichannel of subunit a (2, 4).

By regulating the intracellular or intraorganellar pH, organellar V-ATPases operate in diverse processes such as receptor-mediated endocytosis, protein processing, and degradation as well as intracellular transport. Plasma membrane V-ATPases are involved in bone resorption, extracellular acidification, or energization of secondary active transport processes (2, 5). The connection of V-ATPase function to diseases such as osteoporosis or cancer and the verification of the V-ATPase as a suitable therapeutic target certainly require a comprehensive investigation of the V-ATPase, its inhibitors, and the intermolecular interactions. In the 1980s the plecomacrolide bafilomycin was identified as the first specific V-ATPase inhibitor, exhibiting nanomolar IC₅₀ values (6). Until now, a reasonable number of other inhibitory compounds from different sources have been discovered (7). Among them are the macrolactone archazolid and the benzolactone amide apicularen, both of which show a similar inhibitory efficacy as the popular plecomacrolides bafilomycin and concanamycin (8–10). However, the benzolactone enamides exhibit a unique feature as they do not, in contrast to the other inhibitors, affect V-ATPases from fungi, and therefore they are the first source specific V-ATPase inhibitors (7, 11). In consequence, this class of compounds may be one of the most promising groups of candidates for a therapeutic use of V-ATPase inhibitors.

Concerning the inhibitor-binding sites and the mechanisms of V-ATPase inhibition, two discrete approaches using plecomacrolides have led to first insights as follows. Labeling experiments with the purified V₁V_O holoenzyme of *Manduca sexta* and a ¹²⁵I-labeled derivative of concanamycin (Fig. 1) revealed the binding of plecomacrolides to the V_O subunit c (12). Simul-

¹ To whom correspondence should be addressed. Tel.: 49-541-9693502; Fax: 49-541-9693503; E-mail: huss@biologie.uni-osnabrueck.de.

² The abbreviations used are: V-ATPase, vacuolar-type ATPase; PAL, photoaffinity labeling; NCD-4, N-cyclohexyl-N'(4-(dimethylamino)naphthyl)-carbodiimide; D-, diazirinylbenzoyl-labeled inhibitor.

taneously, mutational analysis of the V_O subunit *c* in *Neurospora crassa* disclosed that certain single amino acid exchanges in the sequence of this subunit altered the affinity of bafilomycin to the V-ATPase (13). Later on, additional single amino acid exchanges in subunit *c* of the V-ATPases from *N. crassa* and *Saccharomyces cerevisiae* resulted in a more precise localization of the plecomacrolide-binding site, which accordingly resides at the interface between helices 1 and 2 of one subunit *c* and helix 4 of an adjacent subunit *c* in the ring (14, 15). Interestingly, the *c*-ring did not appear to contain the whole plecomacrolide-binding site because mutations in subunit *a* of the yeast V-ATPase also conferred resistance to bafilomycin (16). In our previous photoaffinity labeling (PAL) studies using the concanamycin derivative mentioned above, the photoactivatable cross-linking diazirinyl group was bound to the macrocyclic ring of the inhibitor that led to an exclusive label at subunit *c* (12). Labeling of merely subunit *c* was surprising regarding the length (6.4 Å) and flexibility of the attached diazirinyl. However, with respect to the mutational analysis and modeling of the binding site within the *c*-ring, this was a strong indication that position C9 of concanamycin may be deeply buried between two adjacent *c* subunits. In this study, we used derivatives of bafilomycin and concanamycin modified with the newly developed ^{14}C -labeled 4-(3-trifluoromethyl-diazirin-3-yl)benzoic acid (17). By repositioning the diazirinyl moiety to the opposite side of the plecomacrolide structures (Fig. 1), we anticipated labeling not only of subunit *c* but now also of subunit *a*. For the modification at C23, we did not expect strong influence on the inhibitory efficacy, as in previous studies it had already been shown that this position has only a negligible effect and that it does not seem to belong to the major pharmacophore (18–20).

The binding site of the archazolids originally had been presumed to overlap to a large extent with that of the plecomacrolides as archazolid prevented binding of a concanamycin derivative (10). However, the binding site for archazolids is relocated to the equatorial region of the *c*-ring and therefore overlaps with the plecomacrolide-binding site to a minor extent than previously thought (21). This revision has been derived from recent site-directed mutagenesis of the yeast V-ATPase subunit *c* and labeling of the *M. sexta* V-ATPase using a radioactive derivative of archazolid A as well as the fluorescent dicyclohexylcarbodiimide derivative NCD-4.

Up to now, information concerning the binding site of the benzolactone enamides is rare. For the benzolactone enamide salicylhalamide A, it has been reported that it binds to a different site than the plecomacrolides, although it inhibits proton translocation through the V_O complex (12, 22). Recent labeling experiments in the presence of apicularen revealed no interference of plecomacrolide, archazolid, or NCD-4 binding to subunit *c* (10, 21). Yet it was not possible to elucidate where apicularen binds within the V_O complex. The development of the ^{14}C -labeled 4-(3-trifluoromethyl-diazirin-3-yl)benzoic acid mentioned above now provided a convenient way to prepare an apicularen derivative that irreversibly cross-links to the protein upon UV exposure and therefore could be used to identify the interacting V-ATPase subunit(s) (17).

Furthermore, we used the radioactive derivatives of apicularen, bafilomycin, and concanamycin as well as nonradioactive compounds in competition assays to gain new insights into the interaction of the inhibitors. Considering the fact that the fungal V-ATPases are insensitive to benzolactone enamides, we used yeast deletion mutants deficient in subunit *Vma3*, *Vph1*, or *Stv1* for the heterologous expression of their human or insect homologues to prove whether it is possible to transfer sensitivity against apicularen to the yeast-human or yeast-insect hybrid V-ATPase. Summing up all available information, we provide a model predicting the arrangement of the binding sites for plecomacrolides, archazolids, and benzolactone enamides within the V_O complex.

EXPERIMENTAL PROCEDURES

Inhibitors—Bafilomycin A_1 , concanamycin A, 21-deoxyconcanolide A, apicularen A, archazolid A, and saliphenylhalamide were isolated or prepared as published (8, 9, 18, 23). 21-*O*-[4-(3-Trifluoromethyl-diazirin-3-yl)benzoyl]bafilomycin A_1 (D-bafilomycin), 23-*O*-[4-(3-trifluoromethyl-diazirin-3-yl)benzoyl]-21-deoxyconcanolide A (D-concanolide), and their ^{14}C -labeled derivatives were prepared according to Ref. 17. Based on previous evaluation of apicularen analogues, we favored position C3 to attach the diazirinyl moiety (7, 24). For synthesis of 3-*O*-[4-(3-trifluoromethyl-diazirin-3-yl)benzoyl]apicularen (D-apicularen), to a solution of apicularen A (1.5 mg, 3.4 μmol) in CH_2Cl_2 (5 ml) was added 4-(dimethylamino)-pyridine (0.9 mg, 7 μmol), 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide (1.1 mg, 5.7 μmol), and 4-(3-trifluoromethyl-diazirin-3-yl)benzoic acid (1.1 mg, 4.8 μmol), and the solution was stirred for 2.5 h in the dark at 25 °C. Column chromatography of the mixture on silica gel ($\text{CHCl}_3/\text{MeOH}$ 25:1) furnished 1.4 mg (63%) of D-apicularen ($R_f = 0.17$, $\text{CHCl}_3/\text{MeOH}$ 50:1). NMR data of D-apicularen is as follows: ^1H NMR (600 MHz, $[\text{D}_6]$ -acetone): $\delta = 0.99$ (t, $J = 7.5$ Hz, 3H, 25- H_3), 1.46–1.56 (m, 3H, 14- H_b , 10- H_b , 12- H_b), 1.65 (ddd, $J = 5.3, 7.8, 13.1$ Hz, 1H, 12- H_a), 1.78 (dt, $J = 10.9, 14.7$ Hz, 1H, 14- H_a), 1.95 (dt, $J = 4.6, 13.0$ Hz, 1H, 10- H_a), 2.00–2.06 (m, 2H, 16- H_2), 2.27 (dq, $J = 1.6, 7.5$ Hz, 2H, 24- H_2), 2.60 (dd, $J = 1.5, 14.6$ Hz, 1H, 8- H_b), 3.44 (dd, $J = 10.4, 14.6$ Hz, 1H, 8- H_a), 3.83–3.89 (m, 2H, 9- H , 11- OH), 3.98 (m, 1H, 11- H), 4.23 (m, 1H, 13- H), 4.97 (dt, $J = 7.4, 14.5$ Hz, 1H, 17- H), 5.39 (m, 1H, 15- H), 5.69 (d, $J = 11.5$ Hz, 1H, 20- H), 5.79 (m, 1H, 23- H), 6.71 (dd, $J = 10.4, 14.5$ Hz, 1H, 18- H), 6.85 (dt, $J = 1.1, 11.5$ Hz, 1H, 21- H), 7.23 (m, 2H, 4- H , 6- H), 7.44 (dd, $J = 7.7, 8.2$ Hz, 1H, 5- H), 7.50 (d, $J = 8.5$ Hz, 2H, Ar- H), 7.51 (m, 1H, 22- H), 8.23 (d, $J = 8.7$ Hz, 2H, Ar- H), 8.92 (d, $J = 10.4$ Hz, 1H, NH). ^{13}C NMR (150.8 MHz, $[\text{D}_6]$ -acetone) is as follows: $\delta = 14.3$ (C-25), 21.0 (C-24), 29.9 (CN $_2$, hidden), 35.8 (C-16), 38.5 (C-14), 39.0 (C-10), 39.8 (C-8), 40.1 (C-12), 64.7 (C-11), 67.5 (C-13), 73.7 (C-9), 74.6 (C-15), 106.9 (C-17), 120.7 (C-20), 121.6 (C-6), 123.0 (q, $J_{\text{C-F}} = 275$ Hz, CF $_3$), 125.4 (C-22), 126.4 (C-18), 127.8 (C-Ar), 129.0 (C-4), 130.5 (C-5), 130.6 (C-7), 131.4 (C-Ar), 131.7 (C-2) 134.7 (C-Ar), 136.9 (C-21), 140.8 (C-Ar), 141.5 (C-23), 147.7 (C-3), 163.5 (C-19), 164.1 (C-1'), 167.9 (C-1).

Using ^{14}C -labeled 4-(3-trifluoromethyl-diazirin-3-yl)benzoic acid, the ^{14}C -labeled apicularen derivative was prepared accordingly with a specific activity of 37.9 mCi/mmol as determined with a

Binding Site of the V-ATPase Inhibitor Apicularen

liquid scintillation counter. All inhibitors were dissolved in DMSO and stored in stock solutions (10 mM) at -80°C .

Photoaffinity Labeling—Thirty μg of *M. sexta* V_1V_O holoenzyme, 20 μg of V_1 complex, or 10 μg of V_O complex, respectively, were incubated with 52 μM 21-*O*-[4-(3-trifluoromethyl-diazirin-3-yl)-[1- ^{14}C]benzoyl]bafilomycin A_1 (^{14}C -D-bafilomycin), 52 μM 23-*O*-[4-(3-trifluoromethyl-diazirin-3-yl)-[1- ^{14}C]benzoyl]-21-deoxyconcanolide A (^{14}C -D-concanolide), or 100 μM 3-*O*-[4-(3-trifluoromethyl-diazirin-3-yl)-[1- ^{14}C]benzoyl]apicularen (^{14}C -D-apicularen) for 5 min at 25°C . In initial experiments, the presence of ATP apparently led to a slightly better label efficacy (data not shown). This may be the consequence of an increased number of V-ATPase inhibitor complexes forced into these specific conformations in which the inhibitor is, as Bowman and co-workers suggested, bound “like a stone in the gears” (14). Therefore, 1 mM ATP in 1.5 mM MgCl_2 was added to a final volume of 40 μl . For competition assays, 52 μM of the ^{14}C -diazirinylbenzoyl-labeled inhibitors were used after a preincubation with a 10-fold excess of a non-radioactive inhibitor for 5 min at 25°C . Cross-linking was induced by irradiating the samples for 1 min with UV light (366 nm) on ice. Subsequently, the samples were separated by SDS-PAGE (T 17%, C 0.4%) and stained with Coomassie. The gels were then dried on Whatman paper, exposed to a phosphor screen for 72 h, and analyzed with a phosphorimager (GE Healthcare).

Minimum Energy Calculation for D-Concanolide—For a theoretical insight from MM2-force field calculations on isolated systems in the gas phase the diazirinyl-labeled concanolides were investigated as example for nanomolar V-ATPase inhibitors. We calculated in respect to the possible positions which the flexible diazirinyl group may occupy in comparison to the pharmacophore using CS Chem3D Pro 12.0.2.1076, Gaussian 98, (CambridgeSoft, Cambridge, MA) and the energy minimization with the MM2-algorithm.

Yeast Strains and Growth Conditions—The yeast strains used in this study were BMA64-1B (*MAT α* , *ura3-52*; *trp1 Δ 2*; *leu2-3-112*; *his3-11*; *ade2-1*; *can1-100*) (Euroscarf, Frankfurt, Germany) and BMA64-1B Δ *vma3* (*MAT α* , *ura3-52*, *trp1 Δ 2*, *leu2-3-112*, *his3-11*, *ade2-1*, *can1-100*, *vma3::HIS3*) (21). The complemented strain BMA64-1B Δ *vma3*+*VMA3* was obtained by transformation of the plasmid carrying the *VMA3* gene pRS415/*VMA3* (21) into the *vma3* deletion mutant by electroporation.

The mutant strain BY4741 Δ *vph1* Δ *stv1* was obtained by crossing the haploid strains BY4741 Δ *vph1* and BY4742 Δ *stv1* (both Euroscarf, Frankfurt, Germany). Each 500 μl of the appropriate overnight cultures were mixed and pelleted and afterward incubated overnight at 30°C on agar plates containing YPDA, pH 5.5. Small amounts of the resulting zygotes were resuspended in 100 μl of H_2O , applied on a new agar plate, separated by using a micromanipulator (Singer MSM), and again incubated at 30°C . To obtain haploid double deletion mutants, the diploid strains were sporulated on agar plates containing 2% potassium acetate for 8 days at 30°C . Again, a small amount of spores were resuspended in 100 μl of H_2O , treated with 100 μg of zymolyase (100 units/mg, MP Biomedicals) for 10 min, and applied on an agar plate (YPDA, pH 5.5), and the

tetrads were separated by using a micromanipulator (Singer MSM). By selection on plates containing 250 $\mu\text{g}/\mu\text{l}$ G418, those tetrads of which two of the colonies grew on media with G418 and two colonies that did not grow on G418 were used for a PCR analysis of the genotype. By using two different primer combinations, (a) TACTCACGGCCGGCTAGTAATCCAGTTGCCGAGC and ATCGCGAGCCCATTATACC and (b) GACATAGGCCACGAAGGTG and ATCGCGAGCCCATTATACC, as well as the isolated genomic DNA of the four tetrad colonies as templates, the deletions of the genes *vph1* and *stv1* were confirmed.

Cells were usually grown on YPD, pH 5.5, additionally containing 0.02% adenine (YPDA, pH 5.5). The medium was buffered to pH 5.5 with 50 mM MES, 50 mM MOPS. For YPDA, pH 7.5, the pH was adjusted using NaOH. YPDA, pH 5.5, medium with 0.1 M CaCl_2 was obtained by adding 10% of a sterile 1 M CaCl_2 stock solution after autoclaving the medium. For plates, 2% agar was added. Selection of cells was carried out by using plates with SD medium containing 0.67% yeast nitrogen base without amino acids, 2% glucose, 1.5% agar, and 10% drop out amino acid solution (21) lacking the respective amino acid. The medium was buffered with 50 mM MES, 50 mM MOPS and adjusted to pH 5.5 using HCl.

Plasmid Construction—The plasmid pXJ40-KKO-16k containing the coding sequence of the gene *ATP6VOC* of the human V-ATPase subunit c was kindly provided by M. A. Skinner (25). The *ATP6VOC* gene was amplified by PCR using the plasmid pXJ40-KKO-16k as template and the forward primer TACTCAAAGCTTATGTCCGAGTCCAAGAGC and the reverse primer TACTCAAAGCTTCTAAGCGTAATCTGGTACGTCGTATGGGTACTTTGTGGAGAGGATGAG. Notably, the second primer additionally contains the coding sequence for a single C-terminal HA tag. The PCR product was cloned into the yeast expression vector pGADT7 by using the HindIII restriction sites resulting in a plasmid called pGADT7/*ATP6VOC*. The gene of the *M. sexta* proteolipid subunit c was amplified by PCR using a pBluescript plasmid containing the accordant coding sequence (26) as template as well as the forward primer TACTCAAAGCTTATGGCCGAAAATCCAAATC and the reverse primer TACTCAAAGCTTTTACTGTTCGTGTACAG. The PCR product was cloned into the vector pGADT7 by using the HindIII restriction sites. The resulting plasmid pGADT7/*Msc* and the plasmid pGADT7/*ATP6VOC* were transformed into the *vma3* deletion mutant BMA64-1B Δ *vma3* by electroporation, and the cells were selected on SD medium without leucine.

The construction of a plasmid carrying the coding sequence for a hybrid subunit containing the N-terminal half of yeast *VPH1* and the C-terminal half of the human a4 subunit was performed by homologous recombination in the yeast wild type strain BY4741 (Euroscarf, Frankfurt, Germany). For this purpose, the C-terminal half of the human a4 subunit encoded by the amino acids 396–840 and additional homologous regions were amplified by PCR using the primers CGGTATTGCTCAGTACAGAGAAATCAATGCTGGTTTACCCTACACCATCATCACCTTCCCCTTCC and GGTGGATTGGATTGCAAGTCTAACGTTTTTCATGAGATAAGCTACTCCTCGGCTGTGCCATCCA and the plasmid pJV97ha4 (a gift of Fiona

Binding Site of the V-ATPase Inhibitor Apicularen




FIGURE 1. Structures of the PAL inhibitor derivatives D-bafilomycin, D-concanolide, ¹²⁵I-concanolide, D-apicularen, saliphenylhalamide, and parent compounds.

Karet) as template. For the plasmid pRS415/*VPH1*, the yeast *VPH1* gene and in addition 400 bp upstream and downstream implying its native promoter and terminator was cloned into the yeast CEN vector pRS415. Afterward, the plasmid was linearized by the restriction enzyme BspI. This open pRS415/*VPH1* as well as the C-terminal half of the human $\alpha 4$ subunit were transformed into the yeast wild type strain BY4741. By homologous recombination, the C-terminal half of *VPH1* encoded by the amino acids 411–840 was replaced by the C-terminal half of the human $\alpha 4$ subunit resulting in a yeast human hybrid subunit a. The plasmid was called pRS415/*VPH1/a4*.

Serial Drop Dilution Assay—To analyze the V-ATPase function, serial drop dilution assays were carried out. Yeast strains were grown overnight in 5 ml of YPDA, pH 5.5, and diluted to 10⁵ cells/ml in distilled water. Serial 10-fold dilutions from 10⁴ cells/ml to 10⁰ cells/ml were prepared, and each 5 μ l was dropped onto plates with different media. After 3 days of incubation at 30 °C, pictures were taken.

Other Methods—The V₁V_O-ATPase holoenzyme, the V_O complex, and the V₁ complex were isolated from *M. sexta* midgut as published previously (12, 27). Yeast vacuolar membranes were purified as described by Bockelmann *et al.* (21).

V-ATPase assays using the *M. sexta* V₁V_O holoenzyme or V₁ complex or yeast vacuolar membranes were carried out as already reported (21, 27, 28). Protein content was determined using Amido Black 10B (28). SDS-PAGE, Western blotting on nitrocellulose, and immunostaining were performed as described previously (12, 29). For immunodetection, the primary antibodies against the V-ATPase subunits A (30) and e (31) from *M. sexta*, Vma1 (A6422, Invitrogen), and Vma6 (gift from Christian Ungermann, University of Osnabrück) from yeast, the HA tag (Y-11) (sc-805, Santa Cruz Biotechnology), and the secondary antibodies against rabbit (A3687, Sigma) and mouse (A3562, Sigma) were used.

RESULTS AND DISCUSSION

Labeling of the V-ATPase with Radioactive PAL Inhibitors—Before the diazirinyl-labeled derivatives of bafilomycin, concanamide, and apicularen (for structures see Fig. 1) were used for photoaffinity labeling, we tested in enzyme activity assays whether the modifications had an influence on V-ATPase inhibition. Compared with the natural products, the labeled derivatives exhibited reductions of their inhibitor potential by a factor of only 5–100 (Fig. 2) and therefore appeared to be highly suitable for the labeling assays.

Binding Site of the V-ATPase Inhibitor Apicularen

To determine which subunits of the V-ATPase interact with the inhibitors, the V_1V_O holoenzyme, the V_O complex, or as a control, the V_1 complex were incubated with the ^{14}C -diazirinyllabeled derivatives of the inhibitors, and the cross-link reactions were induced by the exposition to UV light. The results of autoradiography after SDS-PAGE are shown in Fig. 3. Generally speaking, labeling patterns appeared to be rather similar for all three inhibitors. The comparison of the exposed (lanes with even numbers, 4–20) and unexposed samples (lanes with odd numbers, 5–21) revealed a clear labeling of the V_O subunits a and c in the V_1V_O holoenzyme as well as in the V_O complex.

To date, subunit c was the only polypeptide of the V_O complex that could be labeled successfully by inhibitors such as concanamycin, archazolid, or NCD-4 (12, 21). Yet the appearance of a label at subunit a was not unexpected as its participation in plecomacrolide binding had already been shown by site-directed mutagenesis studies in the yeast V-ATPase (16). Additionally, it was anticipated that placement of the diazirinyl in the concanamycin derivative at position C23 instead of position C9 as in the previous ^{125}I -concanolide used in 2002 (12, 17)




FIGURE 2. Inhibition of the ATPase activity of the *M. sexta* V_1V_O holoenzyme by bafilomycin, concanamycin, apicularen, and their derivatives. Values represent the means \pm S.D. of experiments with three different preparations. The specific enzyme activity of the controls without inhibitors was $1.6 \pm 0.4 \mu\text{mol}\cdot\text{mg}^{-1}\cdot\text{min}^{-1}$.

would lead to a different labeling pattern. This concept is based on the assumption that the diazirinyl attached to the C9 of the inhibitor is, as already mentioned above, buried in the binding site between two adjacent c subunits, whereas the diazirinyl at position C23 probably is localized at the interface between subunits a and c. For a rough estimation of the space-compassing manner of the diazirinyl derivatives and how they label two opposing subunits, we exemplarily performed a minimum energy calculation for D-concanolide. The model provides a view on the theoretical positions and spanned distances of the diazirinyl group, which is, with respect to the pharmacophore, flexible around its single bonds of the ester moiety. The results of the calculations pointed to two preferred conformations, one with the diazirinyl moiety above and one below the macrolactone ring close to the molecule (Fig. 4) and a small distance of the label. These theoretical data support the hypothesis derived from the functional assays that labeling of both subunits a and c is conclusive even though the inhibitor is bound in one certain orientation in the binding site.

The irradiated samples of the V_O complex exhibited a further band below subunit a, which was most probably a degradation product of subunit a and also visible in the Coomassie staining of the V_O complex. Labeling of the V_1 subunits in the V_1V_O holoenzyme and in the V_1 complex are considered unspecific as it occurred in both the irradiated and nonirradiated samples. In addition, we may exclude specific inhibitory interactions with the V_1 complex as it was already shown previously that the ATPase activity of the V_1 complex of *M. sexta* and yeast is not affected by the plecomacrolide concanamycin and that salicylhalamide, another benzolactone enamide, did not inhibit the V_1 complex of bovine clathrin-coated vesicles (22, 27, 32). In this study, we confirmed these results for the V_1 complex from *M. sexta* using either $30 \mu\text{M}$ of D-bafilomycin, D-concanolide, D-apicularen, or their parent compounds. None of these inhibitors had an effect, neither on the Ca^{2+} -ATPase nor Mg^{2+} -ATPase activity of the isolated V_1 complex (data not shown).




FIGURE 3. Photoaffinity labeling of *M. sexta* V_1V_O holoenzyme, V_1 complex, and V_O complex with the ^{14}C -labeled derivatives of D-bafilomycin, D-concanolide, and D-apicularen. For the labeling assays, $30 \mu\text{g}$ of V_1V_O holoenzyme, $20 \mu\text{g}$ of V_1 complex, or $10 \mu\text{g}$ of V_O complex were first incubated for 5 min at 25°C with $52 \mu\text{M}$ ^{14}C -D-bafilomycin, $52 \mu\text{M}$ ^{14}C -D-concanolide, or $100 \mu\text{M}$ ^{14}C -D-apicularen, respectively. Before the samples were exposed to UV light (366 nm) for 1 min (+) or kept in the dark (-), 1 mM Mg-ATP was added. Afterward, the protein subunits were separated by SDS-PAGE, stained with Coomassie Blue, and exposed to a phosphor screen. Lanes 1–3, typical staining with Coomassie Blue; lanes 4–21, readout of the phosphor screen.




FIGURE 4. Minimum energy calculation for D-concanolide. For 21-deoxyconcanolide A, the space of the connected diazirinyl moiety is placed preferably at the east hemisphere. The length of the benzoyl-diaziranyl label can be calculated to 6.4 Å and its distance from the macrolactone ring to 5.4 Å (A). Minimum energy calculations suggest that the diazirinyl group is situated preferably above and below the macrolactone ring very close to the molecule (A and B). A, benzoyl diazirinyl system is on the other side of the carbonyl group of the macrolactone ester. B, benzoyl diazirinyl system is on the same side of the carbonyl group of the macrolactone ester.

Additional labeling of V_1 subunits may thus be interpreted as a consequence of unspecific binding of the derivatives due to the high protein and inhibitor concentrations in the assays, as they appeared in both the exposed and unexposed samples and even in the purified V_1 complex. A further label close to the V_1 subunits F and G was found in the V_1V_O holoenzyme and in the V_O complex but not in the V_1 complex. To check the origin of this unexpected label, pieces containing subunits F and G as well as the areas above, in between, and below these subunits were excised from a gel for further analysis. Scintillation counting revealed no radioactivity in pieces containing subunits F and G, but in contrast, radioactivity was detected at the interspace between the subunits and below subunit F (data not shown). In addition, mass spectrometric analysis (ESI-MS) verified subunits F and G, but no protein was detected in the areas above, in between, and below the subunits. Therefore, we suppose that these labels result from cross-links of the diazirinyl-derivatives with lipid or detergent molecules that remain bound specifically to the V_O complex during purification. This assumption is supported by the results of von Ballmoos *et al.* (33) who observed cross-links between lipids and diazirinyl-labeled carbodiimide inhibitors interacting with the F_O subunit c of the ATP synthase.




FIGURE 5. Western blot analysis of the V-ATPase with ^{14}C -D-apicularen. For the identification of the ^{14}C -D-apicularen-labeled band with the approximate molecular mass of subunit e 30 μg of the V_1V_O holoenzyme were incubated with 100 μM ^{14}C -D-apicularen, and before the samples were exposed to UV light (366 nm) or kept in the dark, 1 mM Mg-ATP was added. Afterward, the protein subunits were separated by SDS-PAGE. Lane 1, staining of the SDS-gel with Coomassie Blue; lane 2, staining of the proteins with Ponceau S after electro-transfer onto a nitrocellulose membrane; lane 3, immunostaining with monoclonal anti-A (221-9) and anti-e (224-3) antibodies; lane 4, autoradiography of the irradiated sample; lane 5, autoradiography of the nonirradiated sample.

Treatment of the V_1V_O holoenzyme as well as the V_O complex with ^{14}C -D-apicularen exhibited an additional radioactively labeled band with an approximate molecular mass of 20 kDa (Fig. 3, lanes 16 and 20). This size suggests that two polypeptides of the V_O complex could be labeled subunit c'' and/or subunit e. Although a gene encoding a putative subunit c'' was identified in the *M. sexta* genome project with a calculated molecular mass of 27 kDa, we have no indication that a product of this gene is an integral part of the purified V-ATPase derived from the plasma membrane of the midgut from *M. sexta*. Extensive analysis of all proteins of the purified V_1V_O holoenzyme, the purified V_O complex, and chloroform/methanol extracts of both by MALDI-MS and ESI-MS supplemented by N-terminal sequencing revealed so far no evidence for the presence of a subunit c'' (12, 34).³ Based on this state of knowledge, we so far do not see any evidence for the presence of subunit c'' in the plasma membrane V-ATPase of *M. sexta*. However, Western blot analysis of the ^{14}C -D-apicularen-labeled V_1V_O holoenzyme using monoclonal antibodies against subunit e clearly identified this labeled band as subunit e (Fig. 5, lane 3). Being a part of the V_O complex probably associated with subunit a (35), it is very likely that subunit e gets labeled due to its close proximity to subunit a, its position opposite to the c-ring, or even due to a direct contribution to the binding site. Anyhow, the

³ M. Huss and H. Wiczorek, unpublished results.

Binding Site of the V-ATPase Inhibitor Apicularen




FIGURE 6. Photoaffinity labeling of the V-ATPase with ^{14}C -D-bafilomycin after preincubation with different inhibitors. Samples with 30 μg of V_1V_0 holoenzyme were preincubated for 5 min at 25 $^\circ\text{C}$ with DMSO, 625 μM bafilomycin (*baf*), concanamycin (*con*), archazolid (*arch*), apicularen (*api*), or saliphenylhalamide (*saliphe*), respectively. Then the samples were incubated for 5 min at 25 $^\circ\text{C}$ with 52 μM ^{14}C -D-bafilomycin. Before the samples were exposed to UV light (366 nm) for 1 min (+) or kept in the dark (-), 1 mM Mg-ATP was added. Then protein subunits were separated by SDS-PAGE, stained with Coomassie Blue, and exposed to a phosphor screen. *A*, lane 1, typical staining with Coomassie Blue; lanes 2–15, readout of the phosphor screen. *B*, analysis of the pixel intensity of the labeled bands with ImageQuant. Values show the means \pm S.D. of two independent preparations, except for DMSO ($n = 6$) and apicularen ($n = 5$).

result that D-apicularen is the first inhibitor also labeling subunit e indicates again that the binding site of the benzolactone enamides is indeed different from the binding sites for the plecomacrolides and the archazolids and that the mechanism by which the benzolactone enamides inhibit V-ATPases may also be a different one. Unfortunately, nearly nothing is known about the exact localization or the function of subunit e in the V-ATPase, except that it is an integral and essential part of the V_0 complex (12, 31, 36, 37). Although subunit e is positioned next to subunit a in some models of the V_0 complex, until now there are no experimental data supporting a direct interaction between subunit e and any other subunit of the V_0 complex (36, 38). Therefore, it appears difficult to integrate subunit e into a concise model of the localization of the inhibitor-binding sites for which reason we will draw our model without subunit e. Anyway, taken together, the cross-linking studies with all ^{14}C -diazirinyllabeled inhibitors clearly identified both the V_0 subunits a and c to harbor parts of the binding sites.




FIGURE 7. Photoaffinity labeling of the V-ATPase with ^{14}C -D-concanolide after preincubation with different inhibitors. Samples with 30 μg of V_1V_0 holoenzyme were preincubated for 5 min at 25 $^\circ\text{C}$ with DMSO, 625 μM bafilomycin (*baf*), concanamycin (*con*), archazolid (*arch*), apicularen (*api*), or saliphenylhalamide (*saliphe*), respectively. Then samples were incubated for 5 min at 25 $^\circ\text{C}$ with 52 μM ^{14}C -D-concanolide. Before the samples were exposed to UV light (366 nm) for 1 min (+) or kept in the dark (-), 1 mM Mg-ATP was added. Afterward protein subunits were separated by SDS-PAGE, stained with Coomassie Blue, and exposed to a phosphor screen. *A*, lane 1, typical staining with Coomassie Blue; lanes 2–13, readout of the phosphor screen. *B*, analysis of the pixel intensity of the labeled bands with ImageQuant. Values show the means \pm S.D. of two independent preparations, except for DMSO ($n = 8$) and apicularen ($n = 5$).

Specification of the Arrangement of the Binding Sites for Bafilomycin, Concanamycin, and Apicularen by Competitive Labeling—To investigate to what extent the binding sites for apicularen, concanamycin, and bafilomycin overlap within the V_0 complex, we performed photoaffinity labeling assays with the ^{14}C -diazirinyllabeled derivatives in the presence of a 10-fold excess of the natural compounds, *i.e.* the V_1V_0 holoenzyme was preincubated with DMSO, bafilomycin, concanamycin, archazolid, apicularen, or saliphenylhalamide, respectively. For ^{14}C -D-bafilomycin, preincubation with the plecomacrolides bafilomycin and concanamycin as well as with the macrolide archazolid led to a significant reduction of the label in the V_0 subunit c (Fig. 6, lanes 4, 6, and 8). This outcome confirmed the results obtained in analogous experiments performed with bafilomycin, concanamycin, and archazolid where labeling of the V_0 subunit c by ^{125}I -concanolide A or ^{14}C -(bis-diazirinyll-




FIGURE 8. Photoaffinity labeling of the V-ATPase with ^{14}C -D-apicularen after preincubation with different inhibitors. Samples with $30\ \mu\text{g}$ of V_1V_0 holoenzyme were preincubated for 5 min at 25°C with DMSO, $625\ \mu\text{M}$ bafilomycin (*baf*), concanamycin (*con*), archazolid (*arch*), apicularen (*api*), or saliphenylhalamide (*saliphe*). Then the samples were incubated for 5 min at 25°C with $52\ \mu\text{M}$ ^{14}C -D-apicularen. Before the samples were exposed to UV light ($366\ \text{nm}$) for 1 min (+) or kept in the dark (-), $1\ \text{mM}$ Mg-ATP was added. Then protein subunits were separated by SDS-PAGE, stained with Coomassie Blue, and exposed to a phosphor screen. *A*, lane 1, typical staining with Coomassie Blue; lanes 2–15, readout of the phosphor screen. *B*, analysis of the pixel intensity of the labeled bands with ImageQuant. Values show the means \pm S.D. of two independent preparations, except for DMSO ($n = 7$), apicularen ($n = 5$), and D-apicularen ($n = 3$).

archazolid was suppressed in a similar manner (10, 12, 21). Evidently, the interaction with the V_0 subunit a also was significantly reduced by an excess of these compounds. The same effects can be seen in the ^{14}C -D-concanolide series, where bafilomycin, concanamycin, and archazolid impeded labeling of both V_0 subunits a and c (Fig. 7, lanes 4, 6, and 8). Notably, in neither series did the benzolactone enamides apicularen and saliphenylhalamide prevent labeling by ^{14}C -D-plecomacrolides (Figs. 6, lanes 10 and 12, and 7, lanes 10 and 12). This result supports the previous assumption that the binding site for the benzolactone enamides is significantly different from that for the plecomacrolides (10, 12, 22).

Competition assays using ^{14}C -D-apicularen led to the unexpected result that the native apicularen itself did nearly not prevent the labeling (Fig. 8, lane 10). A more effective reduction of labeling was achieved by preincubation with the nonradioactive D-apicularen and with saliphenylhalamide, another benzolactone enamide (Fig. 8, lanes 12 and 14). Therefore, we suggest that the less bulky native apicularen might be too small to cover




FIGURE 9. A, growth of the human Vma3 homologue *ATP6VOC*-expressing yeast strain. Serial 10-fold dilutions of the yeast wild type strain BMA64-1B, the deletion mutant BMA64-1B Δ vma3, and the deletion mutant BMA64-1B Δ vma3 complemented with pRS415/*VMA3* were dropped on different media. After 3 days of incubation at 30°C , pictures of the plates were taken. *B*, assembly of the V-ATPase in isolated vacuolar membranes of the *ATP6VOC*-expressing strain. Each $10\ \mu\text{g}$ of isolated vacuolar membranes of the wild type, the deletion mutant, and the strains complemented with *ATP6VOC* or *VMA3* were separated by SDS-PAGE, transferred to nitrocellulose, and analyzed by immunostaining with anti-A, anti-d, and anti-HA antibodies.

the complete binding site of ^{14}C -D-apicularen. By contrast, the nonradioactive D-apicularen has the same size, and saliphenylhalamide has a comparable size (Fig. 1), and therefore both compounds are more suitable to displace ^{14}C -D-apicularen. Furthermore, we may exclude binding of D-apicularen to an alternative binding site due to the sheer presence of the attached diazirinyl group, because a control compound modified with a diazirinyl was, in contrast to D-apicularen, not able to inhibit the V-ATPase (39). Surprisingly, an excess of the plecomacrolides bafilomycin or concanamycin also reduced labeling by ^{14}C -D-apicularen (Fig. 8, lanes 4 and 6), whereas apicularen did not reduce labeling by ^{14}C -D-bafilomycin as described above. Yet, in experiments using D-apicularen and ^{14}C -D-bafilomycin, a partial reduction of the radioactive label was detectable (Fig. 6, lane 14). Based on these results, we suspect the diazirinyl group of D-apicularen to extend into the plecomacrolide-binding site leading to a displacement by the plecomacrolides or the other way around. Yet the binding site for the native apicularen is obviously different from that for the plecomacrolides as they were not displaced by apicularen in labeling experiments (see above).

The macrolactone archazolid did not prevent labeling of the V-ATPase by ^{14}C -D-apicularen (Fig. 8, lane 8). This was expected from our previously published results with apicularen, NCD-4, and archazolid that had elucidated that archazolid binds to subunit c of the V_1V_0 holoenzyme and has an overlapping but different binding site as bafilomycin (21). Taken together, our results now show that apicularen has a binding site within the subunit a or c of the V_0 complex, which differs

Binding Site of the V-ATPase Inhibitor Apicularen

from those for archazolid and for the plecomacrolides, but a slight overlap of the binding sites for the plecomacrolides and apicularen cannot be excluded.

Exploration of the Apicularen-binding Site by Genetic Manipulation of the Yeast V-ATPase—To find out whether subunit a or c harbors the apicularen-binding site, we took advantage of the facts that bakers' yeast can easily be genetically manipulated and furthermore that apicularen does not inhibit the fungal V-ATPases. The expression of the apicularen-sensitive homologues of either human V-ATPase subunit a or c in appropriate yeast deletion mutants should result in a hybrid V-ATPase, which is active and sensitive to apicularen. This would elegantly prove to which of these two subunits apicularen binds. The possibility to complement a yeast *vma3* deletion mutant strain with c subunits of other organisms has already been reported, e.g. for the 16-kDa proteolipid from *Nephrops norvegicus* or *Drosophila melanogaster* (40, 41). In our approach, we expressed the human *ATP6VOC* in a yeast

$\Delta vma3$ mutant strain that resulted in a partially restored *vma*⁻ phenotype as this strain grew on media with an elevated CaCl₂ concentration, whereas growth on media with alkaline pH failed (Fig. 9A) (42, 43). This moderately recovered V-ATPase activity was also confirmed by Western blot analysis of purified vacuolar membranes that revealed an appropriate amount of assembled V₁V_O holoenzyme by immunodetection of the V₁ subunit A as well as the V_O subunits d and human c (Fig. 9B). From ATPase activity assays with isolated vacuolar membranes, a specific enzyme activity of $0.064 \pm 0.005 \mu\text{mol}\cdot\text{mg}^{-1}\cdot\text{min}^{-1}$ was determined for the *ATP6VOC*-expressing strain, which is about 16% of the wild type activity and, according to Leng *et al.* (44), nearly enough activity to exhibit wild type growth. Even though this hybrid V-ATPase containing *ATP6VOC* was highly sensitive to bafilomycin A₁, it still was not sensitive to apicularen (Fig. 10). In parallel, we also expressed the proteolipid subunit c of *M. sexta* in a yeast $\Delta vma3$ mutant strain. The analysis of the growth of the resulting strain on media with an elevated CaCl₂ concentration or an alkaline pH as well as the Western blot analysis of the V-ATPase assembly in isolated vacuolar membranes and ATPase activity assays revealed the same results (data not shown) as for the expression of the human *ATP6VOC*. Similarly, this hybrid V-ATPase expressing the *M. sexta* V_O subunit c was not sensitive to apicularen. Therefore, subunit c of the V_O complex obviously does not harbor the binding site of apicularen.

The next step was to express the human isoform a4 of the V-ATPase in a $\Delta vph1\Delta stv1$ mutant strain to check whether this hybrid V-ATPase is sensitive to apicularen, and therefore subunit a may host its binding site. Unfortunately, this approach did not lead to an active V-ATPase (data not shown). Furthermore, the expression of a yeast-human hybrid subunit a containing the N-terminal half of yeast Vph1, which is important for targeting and regulation of the V-ATPase (45), connected with the C-terminal half of the human a4 subunit did not lead to the restoration of V-ATPase activity (data not shown). Nevertheless, we can conclude that subunit c does not exclusively form the binding site for apicularen and that subunit a contributes substantially to the binding of apicularen.




FIGURE 10. Inhibition of the V-ATPase activity of vacuolar membranes from the wild type strain BMA64-1B (WT) and the *ATP6VOC*-expressing strain (hybrid) by bafilomycin A₁ and apicularen A. Values represent the means of two independent preparations \pm S.D. The specific enzyme activity of the controls was $0.064 \pm 0.005 \mu\text{mol}\cdot\text{mg}^{-1}\cdot\text{min}^{-1}$ (*ATP6VOC*-expressing strain) and $0.41 \pm 0.1 \mu\text{mol}\cdot\text{mg}^{-1}\cdot\text{min}^{-1}$ (wild type strain BMA64-1B).




FIGURE 11. Model of the inhibitor-binding site arrangement in the membrane. *Left*, side view; *right*, top view from cytosol. For simplicity, the N-terminal part of subunit a is drawn as a single sphere, and only two c subunits of the c-ring are shown. The four transmembrane helices of the c subunits are numbered (1–4). The suggested locations of the inhibitor-binding sites of apicularen, bafilomycin, and archazolid are colored in red, blue, and yellow, respectively. D, the position of the diazirinyl group attached to apicularen. E, the position of the essential glutamate in helix 4.

Modeling of the Binding Site for Apicularen in the Vicinity of Those for Plecomacrolides and Archazolid—Taken together, the information from our results presented here and from previously published data on the binding sites for archazolid and for bafilomycin can be integrated in a concise model arranging the binding sites for three different classes of V-ATPase inhibitors within the V_O complex (Fig. 11). Based on mutagenesis studies of the c subunit performed in *N. crassa* and *S. cerevisiae*, the binding site for the plecomacrolides is localized between two adjacent c subunits in the cytosolic half of the membrane bilayer (15, 21), indicated in blue in our model. The archazolid-binding site, displayed in yellow, has been shown to reside in the equatorial region of one single subunit c covering the essential glutamate (21). Now we suggest that the binding site for apicularen, indicated in red, is located at the interface of subunits a and c, above the equatorial region and in the cytosolic half of the membrane bilayer. Our assumption is based on the following results: (i) apicularen labels both subunits a and c; (ii) it does not interfere with archazolid binding, and (iii) it does not interfere with bafilomycin binding but may interact in the vicinity of bafilomycin, because D-apicularen, with its attached diazirinyl group, obviously interferes with the binding of the competing inhibitor (Fig. 6, lane 14). As already mentioned above, we decide to not include subunit e into our model as there is no reliable information on its localization available.

To gain further insights into the binding site of apicularen, it would be illuminative to construct a benzolactone enamide-sensitive subunit a in yeast by exchanging amino acids using the sequences of the human and insect subunits a as a guideline. This effort should be accompanied by elucidating the peptides cross-linked to D-apicularen by mass spectrometric analysis. In this context, modified derivatives with the diazirinyl group placed at another position or one of the improved diazirinylbenzoic acids with perfluorobutyl and perfluorooctyl chains (F-PAL) (39) could also be applied.

Acknowledgments—We thank Florenz Sasse (Braunschweig, Germany) for providing the inhibitor archazolid A; Xiao-Song Xie (Dallas, TX) for the inhibitor saliphenylhalamide, and Fiona E. Karet (Cambridge, UK), Mhairi A. Skinner (Guelph, Canada), and Julian A. S. Dow (Glasgow, UK) for supplying the plasmids carrying the coding sequences for subunit a of the human V-ATPase and subunit c from the V-ATPases of *H. sapiens* and *M. sexta*, respectively. We also thank Florian Zubeil (Eberhard Karls Universität Tübingen, Germany) for the minimum energy calculations and Martin Dransmann for excellent technical assistance.

REFERENCES

- Beyenbach, K. W., and Wieczorek, H. (2006) The V-type H^+ ATPase. Molecular structure and function, physiological roles, and regulation. *J. Exp. Biol.* **209**, 577–589
- Forgac, M. (2007) Vacuolar ATPases. Rotary proton pumps in physiology and pathophysiology. *Nat. Rev. Mol. Cell Biol.* **8**, 917–929
- Muench, S. P., Huss, M., Song, C. F., Phillips, C., Wieczorek, H., Trinick, J., and Harrison, M. A. (2009) Cryoelectron microscopy of the vacuolar ATPase motor reveals its mechanical and regulatory complexity. *J. Mol. Biol.* **386**, 989–999
- Toei, M., Toei, S., and Forgac, M. (2011) Definition of membrane topology and identification of residues important for transport in subunit a of the vacuolar ATPase. *J. Biol. Chem.* **286**, 35176–35186
- Wieczorek, H., Beyenbach, K. W., Huss, M., and Vitavska, O. (2009) Vacuolar-type proton pumps in insect epithelia. *J. Exp. Biol.* **212**, 1611–1619
- Bowman, E. J., Siebers, A., and Altendorf, K. (1988) Bafilomycins. A class of inhibitors of membrane ATPases from microorganisms, animal cells, and plant cells. *Proc. Natl. Acad. Sci. U.S.A.* **85**, 7972–7976
- Huss, M., and Wieczorek, H. (2009) Inhibitors of V-ATPases. Old and new players. *J. Exp. Biol.* **212**, 341–346
- Sasse, F., Steinmetz, H., Höfle, G., and Reichenbach, H. (2003) Archazolids, new cytotoxic macrolactones from *Archangium gephyra* (myxobacteria). Production, isolation, physicochemical, and biological properties. *J. Antibiot.* **56**, 520–525
- Kunze, B., Jansen, R., Sasse, F., Höfle, G., and Reichenbach, H. (1998) Apicularens A and B, new cytostatic macrolides from *Chondromyces* species (Myxobacteria). Production, physicochemical, and biological properties. *J. Antibiot.* **51**, 1075–1080
- Huss, M., Sasse, F., Kunze, B., Jansen, R., Steinmetz, H., Ingenhorst, G., Zeeck, A., and Wieczorek, H. (2005) Archazolid and apicularen. Novel specific V-ATPase inhibitors. *BMC Biochem.* **6**, 13
- Boyd, M. R., Farina, C., Belfiore, P., Gagliardi, S., Kim, J. W., Hayakawa, Y., Beutler, J. A., McKee, T. C., Bowman, B. J., and Bowman, E. J. (2001) Discovery of a novel antitumor benzolactone enamide class that selectively inhibits mammalian vacuolar-type (H^+)-ATPases. *J. Pharmacol. Exp. Ther.* **297**, 114–120
- Huss, M., Ingenhorst, G., König, S., Gassel, M., Dröse, S., Zeeck, A., Altendorf, K., and Wieczorek, H. (2002) Concanamycin A, the specific inhibitor of V-ATPases, binds to the V_o subunit c. *J. Biol. Chem.* **277**, 40544–40548
- Bowman, B. J., and Bowman, E. J. (2002) Mutations in subunit c of the vacuolar ATPase confer resistance to bafilomycin and identify a conserved antibiotic-binding site. *J. Biol. Chem.* **277**, 3965–3972
- Bowman, E. J., Graham, L. A., Stevens, T. H., and Bowman, B. J. (2004) The bafilomycin/concanamycin-binding site in subunit c of the V-ATPases from *Neurospora crassa* and *Saccharomyces cerevisiae*. *J. Biol. Chem.* **279**, 33131–33138
- Bowman, B. J., McCall, M. E., Baertsch, R., and Bowman, E. J. (2006) A model for the proteolipid ring and bafilomycin/concanamycin-binding site in the vacuolar ATPase of *Neurospora crassa*. *J. Biol. Chem.* **281**, 31885–31893
- Wang, Y., Inoue, T., and Forgac, M. (2005) Subunit a of the yeast V-ATPase participates in binding of bafilomycin. *J. Biol. Chem.* **280**, 40481–40488
- Bender, T., Huss, M., Wieczorek, H., Grond, S., and von Zezschwitz, P. (2007) Convenient synthesis of a [$1-^{14}C$]diazirinylbenzoic acid as a photoaffinity label for binding studies of V-ATPase inhibitors. *Eur. J. Org. Chem.* **2007**, 3870–3878
- Dröse, S., Boddien, C., Gassel, M., Ingenhorst, G., Zeeck, A., and Altendorf, K. (2001) Semisynthetic derivatives of concanamycin A and C, as inhibitors of V- and P-type ATPases. Structure-activity investigations and developments of photoaffinity probes. *Biochemistry* **40**, 2816–2825
- Gagliardi, S., Gatti, P. A., Belfiore, P., Zocchetti, A., Clarke, G. D., and Farina, C. (1998) Synthesis and structure-activity relationships of bafilomycin A₁ derivatives as inhibitors of vacuolar H^+ -ATPase. *J. Med. Chem.* **41**, 1883–1893
- Ingenhorst, G., Bindseil, K. U., Boddien, C., Dröse, S., Gassel, M., Altendorf, K., and Zeeck, A. (2001) Synthesis of a doubly labeled concanamycin derivative for ATPase binding studies. *Eur. J. Org. Chem.* **2001**, 4525–4532
- Bockelmann, S., Menche, D., Rudolph, S., Bender, T., Grond, S., von Zezschwitz, P., Muench, S. P., Wieczorek, H., and Huss, M. (2010) Archazolid A binds to the equatorial region of the c-ring of the vacuolar H^+ -ATPase. *J. Biol. Chem.* **285**, 38304–38314
- Xie, X. S., Padron, D., Liao, X., Wang, J., Roth, M. G., and De Brabander, J. K. (2004) Salicylhalamide A inhibits the V_O sector of the V-ATPase through a mechanism distinct from bafilomycin A₁. *J. Biol. Chem.* **279**, 19755–19763
- Jansen, R., Kunze, B., Reichenbach, H., and Höfle, G. (2000) Apicularen A and B, cytotoxic 10-membered lactones with a novel mechanism of action from *Chondromyces* species (Myxobacteria). Isolation, structure elucidation

Binding Site of the V-ATPase Inhibitor Apicularen

- tion, and biosynthesis. *Eur. J. Org. Chem.* **2000**, 913–919
24. Petri, A. F., Sasse, F., and Maier, M. E. (2005) Synthesis and biological evaluation of apicularen A analogues. *Eur. J. Org. Chem.* **2005**, 1865–1875
25. Skinner, M. A., and Wildeman, A. G. (2001) Suppression of tumor-related glycosylation of cell surface receptors by the 16-kDa membrane subunit of vacuolar H⁺-ATPase. *J. Biol. Chem.* **276**, 48451–48457
26. Dow, J. A., Goodwin, S. F., and Kaiser, K. (1992) Analysis of the gene encoding a 16-kDa proteolipid subunit of the vacuolar H⁺-ATPase from *Manduca sexta* midgut and tubules. *Gene* **122**, 355–360
27. Gräf, R., Harvey, W. R., and Wiczorek, H. (1996) Purification and properties of a cytosolic V₁-ATPase. *J. Biol. Chem.* **271**, 20908–20913
28. Wiczorek, H., Cioffi, M., Klein, U., Harvey, W. R., Schweikl, H., and Wolfersberger, M. G. (1990) Isolation of goblet cell apical membrane from tobacco hornworm midgut and purification of its vacuolar-type ATPase. *Methods Enzymol.* **192**, 608–616
29. Wiczorek, H., Putzenlechner, M., Zeiske, W., and Klein, U. (1991) A vacuolar-type proton pump energizes K⁺/H⁺ antiport in an animal plasma membrane. *J. Biol. Chem.* **266**, 15340–15347
30. Sumner, J. P., Dow, J. A., Earley, F. G., Klein, U., Jäger, D., and Wiczorek, H. (1995) Regulation of plasma membrane V-ATPase activity by dissociation of peripheral subunits. *J. Biol. Chem.* **270**, 5649–5653
31. Merzendorfer, H., Huss, M., Schmid, R., Harvey, W. R., and Wiczorek, H. (1999) A novel insect V-ATPase subunit M9.7 is glycosylated extensively. *J. Biol. Chem.* **274**, 17372–17378
32. Parra, K. J., Keenan, K. L., and Kane, P. M. (2000) The H subunit (Vma13p) of the yeast V-ATPase inhibits the ATPase activity of cytosolic V₁ complexes. *J. Biol. Chem.* **275**, 21761–21767
33. von Ballmoos, C., Appoldt, Y., Brunner, J., Granier, T., Vasella, A., and Dimroth, P. (2002) Membrane topography of the coupling ion binding site in Na⁺-translocating F₁F₀-ATP synthase. *J. Biol. Chem.* **277**, 3504–3510
34. Huss, M., and Wiczorek, H. (2007) Influence of ATP and ADP on dissociation of the V-ATPase into its V₁ and V_O complexes. *FEBS Lett.* **581**, 5566–5572
35. Muench, S. P., Trinick, J., and Harrison, M. A. (2011) Structural divergence of the rotary ATPases. *Q. Rev. Biophys.* **44**, 311–356
36. Compton, M. A., Graham, L. A., and Stevens, T. H. (2006) Vma9p (subunit e) is an integral membrane V_O subunit of the yeast V-ATPase. *J. Biol. Chem.* **281**, 15312–15319
37. Sambade, M., and Kane, P. M. (2004) The yeast vacuolar proton-translocating ATPase contains a subunit homologous to the *Manduca sexta* and Bovine e subunits that is essential for function. *J. Biol. Chem.* **279**, 17361–17365
38. Toei, M., Saum, R., and Forgac, M. (2010) Regulation and isoform function of the V-ATPases. *Biochemistry* **49**, 4715–4723
39. Burkard, N., Bender, T., Westmeier, J., Nardmann, C., Huss, M., Wiczorek, H., Grond, S., and von Zezschwitz, P. (2010) New fluororous photoaffinity labels (F-PAL) and their application in V-ATPase inhibition studies. *Eur. J. Org. Chem.* **2010**, 2176–2181
40. Finbow, M. E., Goodwin, S. F., Meagher, L., Lane, N. J., Keen, J., Findlay, J. B., and Kaiser, K. (1994) Evidence that the 16-kDa proteolipid (subunit c) of the vacuolar H⁺-ATPase and ductin from gap junctions are the same polypeptide in *Drosophila* and *Manduca*. Molecular cloning of the *Vha16k* gene from *Drosophila*. *J. Cell Sci.* **107**, 1817–1824
41. Harrison, M. A., Jones, P. C., Kim, Y. I., Finbow, M. E., and Findlay, J. B. (1994) Functional properties of a hybrid vacuolar H⁺-ATPase in *Saccharomyces* cells expressing the *Nephrops* 16-kDa proteolipid. *Eur. J. Biochem.* **221**, 111–120
42. Nelson, H., and Nelson, N. (1990) Disruption of genes encoding subunits of yeast vacuolar H⁺-ATPase causes conditional lethality. *Proc. Natl. Acad. Sci. U.S.A.* **87**, 3503–3507
43. Ohya, Y., Umemoto, N., Tanida, I., Ohta, A., Iida, H., and Anraku, Y. (1991) Calcium-sensitive *cls* mutants of *Saccharomyces cerevisiae* showing a pet⁻ phenotype are ascribable to defects of vacuolar membrane H⁺-ATPase activity. *J. Biol. Chem.* **266**, 13971–13977
44. Leng, X. H., Manolson, M. F., Liu, Q., and Forgac, M. (1996) Site-directed mutagenesis of the 100-kDa subunit (Vph1p) of the yeast vacuolar (H⁺)-ATPase. *J. Biol. Chem.* **271**, 22487–22493
45. Kawasaki-Nishi, S., Bowers, K., Nishi, T., Forgac, M., and Stevens, T. H. (2001) The N-terminal domain of the vacuolar proton-translocating ATPase a subunit controls targeting and *in vivo* dissociation, and the C-terminal domain affects coupling of proton transport and ATP hydrolysis. *J. Biol. Chem.* **276**, 47411–47420