

Functional and inducible expression of a transfected murine class II major histocompatibility complex gene

(antigen presentation/cosmid clones/ γ -interferon/protoplast fusion/T-helper hybridoma)

VIRGINIA FOLSOM*, DANIEL P. GOLD*, JANICE WHITE†, PHILIPPA MARRACK†, JOHN KAPPLER†, AND SUSUMU TONEGAWA*

*Center for Cancer Research and Department of Biology, Massachusetts Institute of Technology, Cambridge, MA 02139; and †National Jewish Hospital and Research Center, National Asthma Center, Denver, CO 80206

Communicated by S. E. Luria, December 30, 1983

ABSTRACT Using the spheroplast fusion technique, we have introduced the cloned E_{β}^b gene into two *d* haplotype cell lines, the B lymphoma line A20-2J and the macrophage tumor line P388D₁. Analysis with a monoclonal antibody indicates that the product of the transfected E_{β}^b gene associates with the endogenous E_{α} chain to form an $E_{\alpha}^d E_{\beta}^b$ complex. While expression of $E_{\alpha}^d E_{\beta}^b$ is constitutive in A20-2J cells transfected with the E_{β}^b gene, surface expression of $E_{\alpha}^d E_{\beta}^b$ is detected in transfected macrophage cells only after treatment of cells with culture supernatants from concanavalin A (Con A)-stimulated T cells. Transfected B lymphoma cells and transfected Con A supernatant-treated macrophage cells have acquired the ability to present antigen to $E_{\alpha}^d E_{\beta}^b$ -restricted T-cell hybridomas. The observed inducible expression of the transfected gene in the macrophage host indicates that sequences responsible for regulated expression of the E_{β}^b gene may be associated with the transfected gene. In combination with directed mutagenesis, the system described here provides a means to study (i) E_{β}^b sequences that are important in determining the restriction specificity of the *E* molecule and (ii) sequences associated with the E_{β} gene that may be important in the regulation of E_{β} chain expression.

The class II major histocompatibility antigens have been implicated in cell–cell interactions of the immune system (for review, see refs. 1 and 2). These polymorphic cell-surface glycoproteins control the level of immune response to certain antigens, probably through their role during presentation of antigen to class II-restricted antigen-specific T cells: a class II-restricted T cell recognizes specific antigen only when it is presented on the surface of a cell that expresses the appropriate allelic form of a class II molecule. Although the mechanism of restricted antigen presentation is not understood at a molecular level, it is clear that models for T-cell recognition of antigen should incorporate both a T-cell receptor (or receptors) and antigen in association with class II molecules on the surface of the presenting cell.

Expression of class II antigens is limited to certain cells of the immune system, including B cells and macrophages (3–5). A relatively high level of constitutive expression of class II antigens is observed in B cells. This can be increased by certain treatments but not by treatment with γ -interferon (unpublished observation). In macrophages, expression of class II antigens is modulated. Resting macrophages express low levels of surface class II antigen, and expression of these molecules can be induced with γ -interferon or with supernatants from Con A-stimulated T cells containing γ -interferon (5–7). Although induction by γ -interferon appears to operate at the level of transcriptional control (8, 9), virtually nothing is known about the mechanisms that regulate expression of class II molecules.

In mouse, two class II molecules, each consisting of an α and a β protein chain, have been defined (2, 10). The *A* molecule is designated as $A_{\alpha}A_{\beta}$ and the *E* molecule as $E_{\alpha}E_{\beta}$. Structural analysis of these molecules has been limited by a lack of protein sequence data. Recently, however, efforts from several laboratories have led to the isolation and characterization of cDNA and genomic clones for the A_{α} , A_{β} , E_{α} , and E_{β} chains (11–16). Protein sequences have been deduced and exon/intron organizations defined. Although sequence analysis provides some understanding of the organization of class II molecules and the genes encoding them, identification of functionally important regions remains tentative.

We have chosen to use a DNA transfection system, in combination with directed mutagenesis, to study the structure–function relationship of class II molecules and the regulation of their expression. Our initial goal has been to introduce a cloned class II gene into a host cell that has been shown (i) to express class II molecules and (ii) to be capable of presenting antigen to specific T-cell hybridomas in culture. We reasoned that a cell line that expresses endogenous class II genes would also express a transfected gene, provided appropriate DNA control elements were included in association with the transfecting gene. Once expression of a transfected gene has been established, we can begin to modify the gene prior to transfection and to determine the effect of defined modification on the expression and function of the class II molecule.

In this paper we report structural and functional evidence for the expression of the E_{β}^b gene after introduction into a B-cell lymphoma or a macrophage tumor. We also show that surface expression of the transfected E_{β}^b gene is inducible by Con A-activated T-cell supernatants after introduction into macrophage tumor cells.

MATERIALS AND METHODS

All cells were cultured in medium containing 10% fetal calf serum and 50 μ M 2-mercaptoethanol. The BALB/c B lymphoma line A20-2J was previously designated L10.A2J but has been recently reidentified (and renamed) as a derivative of the A20 tumor (17). The P388D₁ macrophage tumor line (18) was obtained from the American Type Culture Collection. The helper T-cell hybridomas BDK-38.2, BDK-44.5, and DO-11.10 have been described (19, 20). Two additional T-cell hybridomas, BDO-23.2 and BDO-34.1, were derived by established methods (21) and represent fusion products of normal T cells from ovalbumin-primed (C57BL/6 \times DBA/2) F₁ mice and the AKR thymoma BW5147. Hybrids were screened for their ability to recognize ovalbumin presented by the A20- E_{β}^b -1 transfected cell line (see *Results*).

The Y-17 monoclonal antibody (22) that recognizes a variety of *E* molecules, including $E_{\alpha}^d E_{\beta}^b$ but not $E_{\alpha}^d E_{\beta}^d$, was received from Donal Murphy (Yale University, New Haven,




FIG. 1. (A) Restriction enzyme map of the cosmid 7-01- E_{β}^b . Only the structure of the cosmid insert is presented. Six exons, which are indicated by dark boxes, have been placed relative to conserved restriction enzyme sites by analogy to the E_{β}^d gene (16). The striped box indicates the d haplotype $EcoRI$ - Pvu II fragment from phage λ 11.1. (16), which was used as a probe for the isolation of cosmid 7-01- E_{β}^b . (B) Map of the recombinant plasmid pSV2- E_{β}^b . The 22-kb Hpa I fragment from cosmid 7-01- E_{β}^b is indicated as a thin line with dark boxes representing exons of the E_{β}^b gene. Exons are indicated as follows: L, leader; β_1 , first external domain; β_2 , second external domain; TM, transmembrane; CP, cytoplasmic; 3'UT, 3' untranslated (see ref. 16). The Eco gpt gene (striped region), Simian virus 40 DNA sequences (dotted region), and ampicillin gene (shaded area) of the pSV2gpt vector are as indicated (see ref. 27). Modified Bam HI sites are labeled as (Bam). Transcriptional orientations of E_{β}^b and Eco gpt genes are indicated by arrows.

CT). The anti- $I-A^d$ specific MKD6 antibody is described by Kappler *et al.* (21). Hybridoma culture supernatants were used for immunofluorescence and immunoprecipitation. For immunofluorescence, the detection antibody was a fluoresceinated goat anti-mouse Fab antisera (a gift from Henry Wortis, Tufts University Medical School, Boston, MA). Fluorescence was measured on an Ortho fluorescence-activated cell sorter, and immunoprecipitation of [35 S]methionine-labeled cell extracts was done as described by P. Jones (23).

Parental and transfected cell lines were tested for their ability to present antigen as described elsewhere (24). Briefly, 1×10^5 presenting cells were cultured with 1×10^5 responder T cells in the presence of 200 μ g of antigen for 24 hr in 200 μ l of culture medium. Interleukin 2 (IL-2) production by the T-cell hybridoma was assayed by the ability of the culture supernatant to support the growth of an IL-2-dependent T-cell line, HT-2 (25). Units of IL-2 were determined as described (24). For induction of P388D₁ and derived lines, cells were incubated for 48 hr with culture supernatants from the T-cell hybridoma FS7-20 at 1:8 dilution (26).

The C57BL/6 cosmid library was constructed and screened by established methods (27). To construct the pSV2- E_{β}^b plasmid, the pSV2gpt vector (28) was linearized with Bam HI, treated with the Klenow fragment of *Escherichia coli* DNA polymerase I to create flush ends, and digested with calf alkaline phosphatase. A 22-kilobase (kb) Hpa I fragment isolated from cosmid 7-01- E_{β}^b was fused by blunt-end ligation to the pSV2gpt vector. Preparation of high molecular weight DNA was by standard methods (27). Southern blotting and hybridization were as described (29).

For transfection, bacteria containing pSV2- E_{β}^b or pSV2gpt plasmids were converted to protoplasts as described by Sandri-Goldin *et al.* (30). Protoplasts were fused to A20-2J or P388D₁ cells with 50% polyethylene glycol (British Drug House) as described by Gillies *et al.* (31). After fusion, cells were plated and allowed to recover in normal growth medium for 48–72 hr, at which time an equal volume of selective medium was added. Selective medium contained xanthine at 250 μ g/ml, hypoxanthine at 15 μ g/ml, and mycophenolic acid at 2 μ g/ml (for P388D₁) or 6 μ g/ml (for A20-2J). Xanthine and hypoxanthine were from Sigma. Mycophenolic acid was a gift from Eli Lilly. Plates were refed every 2–3 days with fresh selective medium. Colonies were picked for expansion after 2–3 wk.

RESULTS

As transfection hosts we have chosen the BALB/c B lymphoma cell line A20-2J and the DBA/2 macrophage tumor line P388D₁. The A20-2J line constitutively expresses both A and E molecules of the d haplotype and presents antigen in a restricted fashion to T-cell hybridomas in culture (17). In normal growth medium, P388D₁ expresses d haplotype class II molecules at low levels and presents antigen only poorly. However, when P388D₁ is cultured with supernatant from Con A-stimulated T-cells, class II antigens are expressed at high levels and the induced culture presents antigen efficiently to restricted T-cell hybridomas (32). The E_{β} gene of b haplotype (E_{β}^b) was chosen as a transfecting gene, as specific reagents were available to assay expression and function of $E_{\alpha}^d E_{\beta}^d$ hybrid molecules.

Isolation of the E_{β}^b Gene. The C57BL/6 cosmid library was screened with a probe derived from the first intron of the E_{β}^d gene (see Fig. 1A and ref. 16). Three overlapping clones were isolated, and cosmid clone 7-01- E_{β}^b was selected for restriction enzyme analysis (Fig. 1A).

Transfection of the E_{β}^b Gene into BALB/c Hosts. The pSV2- E_{β}^b plasmid was introduced into A20-2J or P388D₁ cells by spheroplast fusion. For A20-2J cells, the frequency of stable transformation was 5×10^{-5} with pSV2- E_{β}^b and 1×10^{-4} with pSV2gpt. After transfection, A20-2J cells were




FIG. 2. Southern blot analysis of E_{β}^b -transfected cell lines. High molecular weight DNA was digested with Bam HI, separated by electrophoresis on 0.8% agarose gels, and blotted to nitrocellulose. Blots were probed with a nick-translated 1.2-kb Eco RI- Pvu II fragment isolated from a recombinant phage containing part of the E_{β}^d gene (see legend to Fig. 1A). The sizes of DNA bands are indicated in kilobases. (A) Data for A20-2J and A20- E_{β}^b -transfected lines: lane 1, A20-2J; lane 2, A20- E_{β}^b -1; lane 3, A20- E_{β}^b -2; lane 4, A20- E_{β}^b -5. (B) Results for P388D₁ (lane 1) and the P388D₁- E_{β}^b -transfected population (lane 2).




FIG. 3. Surface expression of $E_{\alpha}^d E_{\beta}^b$ in an E_{β}^b -transfected A20 clone. A20-2J or A20- E_{β}^b -5 cells were incubated with Y-17 culture supernatant, followed by incubation with fluoresceinated goat anti-mouse Fab antisera. Fluorescence was measured on an Ortho fluorescence-activated cell sorter; 30,000 cells were analyzed per sample. The vertical axis indicates cell number and the horizontal axis indicates fluorescence intensity.

plated in 96-well culture plates, such that colonies were obtained in 1 of ≈ 10 wells. Colonies picked for analysis were therefore assumed to represent individual transfection events.

P388D₁ transfectants were selected on 60-mm tissue culture plates. Numerous colonies were visible after transfection with pSV2- E_{β}^b or pSV2gpt. Plates were grown to confluency so that our transfected P388D₁ lines represent populations of many individual transfectants. Frequencies of transfection, subsequently determined for a subline of P388D₁, were 3×10^{-5} and 1.5×10^{-4} with pSV2- E_{β}^b and pSV2gpt, respectively.

E_{β}^b DNA sequences are present in several A20- E_{β}^b transfectants and in the P388D₁- E_{β}^b population, as determined by Southern blot analysis (Fig. 2). Digestion is with *Bam*HI endonuclease. The endogenous E_{β}^d gene is seen as a 4.8-kb band, and the exogenous E_{β}^b gene is represented by an 8.7-kb band. Clearly, the E_{β}^b band is seen in all A20- E_{β}^b transfectants (Fig. 2A) and in the P388D₁- E_{β}^b population (Fig. 2B), whereas it is absent in DNA from the parental cell lines. Each of the A20- E_{β}^b transfectants contains an estimated 50–100 copies of the E_{β}^b gene per haploid genome, and the P388D₁- E_{β}^b population contains an average of 3–4 copies per haploid genome. A third restriction enzyme fragment is seen at 6.5 kb in A20-2J and in A20- E_{β}^b transfectants. The origin of this fragment is unclear, as a fragment of this size is not expected from the restriction map of the E_{β}^d gene (16).

Expression of E_{β}^b in A20-2J Transfectants. Four A20- E_{β}^b transfectants and two A20-pSV2gpt transfectants were examined for reactivity to the Y-17 monoclonal antibody by radioimmunoassay (data not shown). Surface expression of




FIG. 4. Immunoprecipitation of $E_{\alpha}^d E_{\beta}^b$ from A20- E_{β}^b -5 cells. Nonidet P-40 extracts of [³⁵S]methionine-labeled cells were immunoprecipitated with Y-17 antibody and inactivated *Staphylococcus aureus* bacteria. Extracts from 1×10^6 cells were used for the A20-2J (lane 1) and A20- E_{β}^b -5 (lane 2) samples. Extracts from 2.5×10^6 spleen cells were analyzed for the BALB/c (lane 3) and (BALB/c \times C57BL/6) F₁ (lane 4) samples. Immunoprecipitates were analyzed by electrophoresis on a 12% polyacrylamide gel in NaDodSO₄ buffer. The gel was fluorographed, dried, and exposed to Kodak XAR-5 film with an intensifying screen for 24 hr (lanes 1 and 2) or 2 wk (lanes 3 and 4). Molecular weight markers ($\times 10^{-3}$) are indicated (lane M).

$E_{\alpha}^d E_{\beta}^b$ was indicated for all A20- E_{β}^b transfectants tested, whereas the A20-2J parent and pSV2gpt transfectants failed to react with Y-17. Total surface expression of *E* molecules was increased in E_{β}^b -transfected lines, as measured with the monoclonal antibody 14-4-4S, which recognizes *E* molecules of all haplotypes (33). Reaction with the anti-*I-A*^d monoclonal antibody MKD6 indicated that surface expression of the *A* molecule was not altered by transfection.

The A20- E_{β}^b -5 transfectant cell line was further analyzed by fluorescence-activated cell sorting after reaction with Y-17. The results, as shown in Fig. 3, indicate that the vast majority of cells in the A20- E_{β}^b population are expressing surface $E_{\alpha}^d E_{\beta}^b$. The expression of surface $E_{\alpha}^d E_{\beta}^b$ is similar to that of $A_{\alpha}^d A_{\beta}^d$, as seen by analysis with the MKD6 antibody (not shown).

To further characterize the product of the transfected E_{β}^b gene, Nonidet P-40 extracts of [³⁵S]methionine-labeled A20-2J or A20- E_{β}^b -5 cells were analyzed by NaDodSO₄/polyacrylamide gel electrophoresis after immunoprecipitation with the Y-17 antibody (Fig. 4). Labeled extracts of spleen cells from BALB/c or (BALB/c \times C57BL/6) F₁ mice expressing $E_{\alpha}^d E_{\beta}^b$ were used as controls. Specific products of the expected sizes are seen in the A20- E_{β}^b -5 and F₁ spleen samples. E_{α} and E_{β} chains exist in several glycosylated




FIG. 5. Surface expression of $E_{\alpha}^d E_{\beta}^b$ is induced in the P388D₁- E_{β}^b population by treatment with T-cell supernatants. P388D₁ or P388D₁- E_{β}^b cells were incubated with the Y-17 or MKD6 (anti-*I-A*^d) antibody followed by incubation with fluoresceinated goat anti-mouse Fab antisera. Fluorescence was measured on an Ortho fluorescence-activated cell sorter; 15,000 cells were analyzed per sample. Profiles are shown for uninduced (–IFN) and induced (+IFN) cultures as indicated (IFN, interferon). Cell number is indicated on the vertical axis and fluorescence intensity is along the horizontal axis. (A) P388D₁ \times anti-*I-A*^d; (B) P388D₁ \times Y-17; (C) P388D₁- E_{β}^b \times anti-*I-A*^d; and (D) P388D₁- E_{β}^b \times Y-17.

Table 1. Antigen presentation by A20-2J and transfected A20-2J lines

T-cell hybridoma	Ag/H-2 specificity	Ag in culture	Units of IL-2 per ml produced in response to					
			LB-15.13*	A20-2J	A20-pSV-1	A20-E β^b -1 [†]	A20-E β^b -2	A20-E β^b -5
DO-11.10	cOVA/I-A ^d	cOVA	ND	320	80	80	80	80
BDK-38.2	KLH/I-E ^{d/b} ‡	None	<10	<10	<10	<10	<10	<10
BDK-38.2	KLH/I-E ^{d/b}	KLH	640	<10	<10	>640	320	640
BDK-44.5	KLH/I-A ^{b/d}	KLH	320	<10	<10	<10	<10	<10
BDO-23.2	cOVA/I-E ^{d/b}	cOVA	320	<10	ND	320 (<10)	ND	ND
BDO-34.1	cOVA/I-E ^{d/b}	cOVA	80	<10	ND	40 (<10)	ND	ND

Ag, antigen; ND, not done; cOVA, chicken ovalbumin; KLH, keyhole limpet hemocyanin. Ten units of IL-2 per ml is the minimum detectable by our assay.

*LB-15.13 is the I^b-, I^d-bearing fusion product of A20-2J to C57BL/10 B cells (17).

[†]Numbers in parentheses indicate units of IL-2 per ml produced when competing Y-17 antibody was added to the presentation assay.

‡I-E^{d/b} refers to the hybrid molecule E α^d E β^b . I-A^{b/d} refers to A α^b A β^d .

forms and cannot be fully distinguished from each other by one-dimensional gel electrophoresis. Regions of the gel containing E α^d or E β^b species (or both) are therefore indicated with parentheses. The invariant chain I_i, which is found in the cytoplasm in association with A α A β and E α E β complexes (10), migrates as a distinct band. A20-E β^b -5 and F₁ spleen immunoprecipitates were also analyzed by two-dimensional gel electrophoresis (not shown). Identification of E α^d , E β^b , and I_i chains (as designated in Fig. 4) was confirmed by a comparison of our two-dimensional gels with published data for F₁ spleen (34).

Inducible Expression of E β^b in a P388D₁ Transfectant Cell Line. The P388D₁ and P388D₁-E β^b cell lines were analyzed by indirect immunofluorescence for expression of E α^d E β^b and A α^d A β^d , both before and after treatment with supernatants from Con A-activated T cells. Fluorescence-activated cell sorting data for this experiment are presented in Fig. 5. Expression of the A α^d A β^d molecule is induced by treatment with T-cell supernatant in both P388D₁ and P388D₁-E β^b (Fig. 5 A and C), whereas the E α^d E β^b heterodimer is expressed only in the P388D₁-E β^b -transfected line after induction (Fig. 5D). Similar results have recently been obtained with γ -interferon (data not shown).

Presentation of Antigens to Restricted T-Cell Hybridomas. The data presented in Table 1 indicate that E β^b -transfected A20-2J lines have specifically acquired the ability to present antigens to E α^d E β^b -restricted T-cell hybridomas. Although E β^b -transfected lines can present ovalbumin or keyhole limpet hemocyanin to appropriate T-cell lines, the A20-2J parent and a pSV2gpt-transfected A20-2J derivative are not capable of functional interaction with E α^d E β^b -restricted T-cell hybridomas.

P388D₁ and transfected P388D₁ lines have been assayed for their ability to present antigen to class II-restricted T-cell hybridomas (Table 2). After incubation with supernatants from Con A-stimulated T cells, all P388D₁-derived lines can present ovalbumin to the I-A^d-restricted DO-11.10. However, only the P388D₁-E β^b line can present antigen to the E α^d E β^b -restricted lines BDO-23.2 and BDO-34.1. Presentation by P388D₁-E β^b to BDO-23.2 and BDO-34.1 is weak before induction and is increased 16-fold for BDO-23.2 and 4-fold for BDO-34.1 after incubation of P388D₁-E β^b cells with supernatants from activated T cells.

DISCUSSION

The class II antigens of the major histocompatibility complex affect the immune responsiveness to many antigens. To understand this phenomenon, it is important to determine at the molecular level (i) how class II molecules function in the presentation of antigen to restricted T cells and (ii) how the expression of class II molecules is regulated in antigen-presenting cells. These questions may be addressed through the use of directed mutagenesis and a defined DNA transfection system. To this end, we have introduced the cloned mouse E β^b gene into two *d* haplotype host cell lines—the B lymphoma line A20-2J and the macrophage tumor line P388D₁. Using a monoclonal alloantibody that recognizes the hybrid E α^d E β^b molecule, we have shown the transfected gene to be constitutively expressed in the B lymphoma host. In contrast, surface expression of E α^d E β^b could be detected in the transfected P388D₁-E β^b macrophage line after, but not before, incubation of cells with supernatants from Con A-stimulated T cells. In addition, we have shown that transfected B lymphoma cells or transfected, induced macrophage cells can functionally present antigens to appropriate T-helper cell hybridomas. The ability of a transfected line to present antigen was consistent with the pattern of cell-surface expression of E molecules measured by antibody: A20-2J cells transfected with the E β^b gene effectively presented antigen to E α^d E β^b -restricted T-cell hybridomas, whereas the P388D₁ cells transfected with the E β^b gene presented antigen optimally only after being cultured in T-cell-derived conditioned medium (with known inducing activity).

Functional expression of a transfected A β gene has recently been reported (35). However, it is not known whether A and E molecules play identical roles in the generation of immune response to antigens, and it will be important to understand how both A and E molecules function in cell-cell interactions of the immune system. The transfection system described here has the advantage that function may be assayed after introduction of a single class II gene into a recipient cell. In our system, transfected cell lines, which express E α^d E β^b (in addition to endogenous E α^d E β^b), are assayed against a panel of E α^d E β^b -restricted responder T-cell hybridomas. Because the polymorphism observed in E molecules is contributed principally by the E β chain (E α is the most highly conserved of the four class II polypeptides, whereas E β is

Table 2. Inducible antigen presentation by P388D₁ and transfected P388D₁ lines

T-cell hybridoma	Ag/H-2 specificity	LB-15.13	Units of IL-2 per ml produced in response to					
			P388D ₁		P388D ₁ -pSV		P388D ₁ -E β^b -1	
			Uninduced	Induced	Uninduced	Induced	Uninduced	Induced
DO-11.10	cOVA/I-A ^d	>640	<10	640	<10	>640	10	640
BDO-23.2	cOVA/I-E ^{d/b}	>640	<10	<10	<10	<10	40	640
BDO-34.1	cOVA/I-E ^{d/b}	320	<10	<10	<10	<10	20	80

See legend to Table 1 for abbreviations.

highly polymorphic; see ref. 36), it is expected that analysis of *E* restricting elements can be carried out by modifying (and transfecting) the *E_β* gene alone. By contrast, both the α and β chains of the *A* molecule are polymorphic and both chains contribute to the determination of restriction specificity. Analysis of *I-A*-restricted cell-cell interactions may therefore be more complex. The restricting function of hybrid *E_β* chains can be initially investigated by shuffling exons between *E_{β^b}* and *E_{β^d}* genes. The hybrid gene approach has been successfully applied to the study of class I major histocompatibility complex molecules (37, 38) and should allow localization of those regions of the *E_β* chain that are important in determining the restriction specificity of the *E* molecule.

Expression of class II major histocompatibility complex molecules is both positively and negatively regulated in macrophages. Positive regulation has been demonstrated in response to soluble products of stimulated T cells (6, 7), whereas negative regulation has been observed in response to prostaglandins (5). Recent data indicate that the T-cell factor responsible for induction is γ -interferon (7, 39). Results from human (8) and murine (9) systems suggest that regulation by γ -interferon may operate at the level of transcriptional control, in that increased levels of mRNA are observed after γ -interferon induction. In fact, HLA-DR $_{\alpha}$ and HLA-DR $_{\beta}$ (the human equivalents of *E_α* and *E_β*, respectively) mRNA levels are simultaneously increased in response to γ -interferon treatment, indicating that transcriptional regulation by γ -interferon may operate on both α and β chain genes (8).

The data presented here indicate that the 22-kb DNA fragment containing the *E_β* gene and its flanking regions may contain all of the sequence information required to make this gene susceptible to regulation by γ -interferon in macrophage cells. Premanipulation of this DNA fragment followed by transfection of P388D₁ cells may lead to localization of one or more specific sequence element(s) required for the induction. In this context, it is interesting that an apparently tissue-specific transcriptional enhancer element is associated with the *E_β* gene (unpublished observations). It is possible that the enhancer element itself is an indirect target for the inducing action of γ -interferon. A somewhat analogous involvement of an enhancer element in the modulation of gene expression by an externally provided agent has been seen in the glucocorticoid hormone-mediated induction of mouse mammary tumor virus genes (40).

We thank Dr. Donal Murphy for the Y-17 hybridoma cell line and Dr. Charles Wood for his advice on construction of the cosmid library. We also thank Ms. Lauren Ormsby for technical assistance and Mrs. Eleanor Basel for her patient preparation of this manuscript. V.F. is supported by a fellowship from the Cancer Research Institute. D.P.G. is supported by a National Institutes of Health training grant. This work was supported by Grants AI-17879 (S.T.) and CA-14051 (a core grant to S. Luria) from the National Institutes of Health.

- Benacerraf, B. & McDevitt, H. O. (1972) *Science* **175**, 273–278.
- Klein, J., Juretic, A., Baxevanis, C. N. & Nagy, Z. A. (1981) *Nature (London)* **291**, 455–460.
- Hammerling, G., Deak, B., Mauve, G., Hammerling, U. & McDevitt, H. O. (1974) *Immunogenetics* **1**, 68–74.
- Steeg, P., Moore, R. & Oppenheim, J. (1980) *J. Exp. Med.* **152**, 1734–1744.
- Snyder, D. S., Beller, D. I. & Unanue, E. R. (1982) *Nature (London)* **299**, 163–165.
- McNicholas, J. M., King, D. P. & Jones, P. P. (1983) *J. Immunol.* **130**, 449–455.
- King, D. P. & Jones, P. P. (1983) *J. Immunol.* **131**, 315–318.
- Rosa, F., Hatat, D., Abadie, A., Wallach, D., Revel, M. & Fellous, M. (1983) *EMBO J.* **2**, 1585–1589.
- Nakamura, M., Manser, T., Pearson, G. D. N., Daley, M. J. & Geftter, M. L. (1984) *Nature (London)* **307**, 381–382.
- McDevitt, H. O. (1980) *Immunology* **80**, 503–512.
- Steinmetz, M., Minard, K., Horvath, S., McNicholas, J., Frelinger, J., Wake, C., Long, E., Mach, B. & Hood, L. (1982) *Nature (London)* **300**, 35–43.
- Benoist, C. O., Mathis, D. J., Kanter, M. R., Williams, V. E., II, & McDevitt, H. O. (1983) *Proc. Natl. Acad. Sci. USA* **80**, 534–538.
- Choi, E., McIntyre, K., Germain, R. N. & Seidman, J. G. (1983) *Science* **221**, 283–286.
- Mallisen, M., Hunkapiller, T. & Hood, L. (1983) *Science* **221**, 750–753.
- Mathis, D. J., Benoist, C. O., Williams, V. E., II, Kanter, M. R. & McDevitt, H. O. (1983) *Cell* **32**, 745–754.
- Saito, H., Maki, R. A., Clayton, L. K. & Tonegawa, S. (1983) *Proc. Natl. Acad. Sci. USA* **80**, 5520–5524.
- Kappler, J., White, J., Wegmann, D., Mustain, E. & Marrack, P. (1982) *Proc. Natl. Acad. Sci. USA* **79**, 3604–3607.
- Koren, H., Handwerker, B. & Wunderlich, J. (1975) *J. Immunol.* **114**, 894–897.
- Marrack, P., Graham, S., Leibson, H. J., Roehm, N., Wegmann, D. & Kappler, J. (1982) in *Isolation, Characterization, and Utilization of T Lymphocyte Clones*, eds. Fathman, C. G. & Fitch, F. (Academic, New York), pp. 119–126.
- White, J., Haskins, K. M., Marrack, P. & Kappler, J. (1983) *J. Immunol.* **130**, 1033–1037.
- Kappler, J. W., Skidmore, B., White, J. & Marrack, P. (1980) *J. Exp. Med.* **153**, 1198–1214.
- Lerner, E., Matis, L., Janeway, C., Jones, P., Schwartz, R. & Murphy, D. (1980) *J. Exp. Med.* **152**, 1085–1101.
- Jones, P. P. (1980) in *Selected Methods in Cellular Immunology*, eds. Mishell, B. & Shiigi, S. (Freeman, San Francisco), pp. 398–440.
- Walker, E., Warner, N. L., Chesnut, R., Kappler, J. & Marrack, P. (1982) *J. Immunol.* **128**, 2164–2169.
- Watson, J. (1979) *J. Exp. Med.* **150**, 1510–1519.
- Zlotnik, A., Roberts, W. K., Vasil, A., Blumenthal, E., Larosa F., Leibson, H. J., Endres, R. O., Graham, S. D., Jr., White, J., Hill, J., Henson, P., Klein, J. R., Bevan, M. J., Marrack, P. & Kappler, J. (1983) *J. Immunol.* **131**, 794–800.
- Maniatis, T., Fritsch, E. F. & Sambrook, J. (1982) *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Laboratory, Cold Spring Harbor, NY), pp. 295–305.
- Mulligan, R. C. & Berg, P. (1980) *Science* **209**, 1422–1427.
- Southern, E. (1975) *J. Mol. Biol.* **98**, 503–515.
- Sandri-Goldin, R. M., Goldin, A. L., Levine, M. & Glorioso, J. C. (1981) *Mol. Cell. Biol.* **1**, 743–752.
- Gillies, S. D., Morrison, S. L., Oi, V. T. & Tonegawa, S. (1983) *Cell* **33**, 717–728.
- Birmingham, J. R., Chestnut, R. W., Kappler, J. W., Marrack, P., Kubo, R. & Grey, H. M. (1982) *J. Immunol.* **128**, 1491–1492.
- Ozato, K., Mayer, N. & Sachs, D. H. (1980) *J. Immunol.* **124**, 533–540.
- McNicholas, J. M., Murphy, D. A., Matis, L. A., Schwartz, R. H., Lerner, E. A., Janeway, C. A., Jr., & Jones, P. P. (1982) *J. Exp. Med.* **155**, 490–507.
- Germain, R. N., Norcross, M. A. & Margulies, D. H. (1983) *Nature (London)* **306**, 190–194.
- Steinmetz, M. & Hood, L. (1983) *Science* **222**, 727–733.
- Evans, G. A., Margulies, D. H., Shykind, B. & Seidman, J. G. (1982) *Nature (London)* **300**, 755–757.
- Ozato, K., Evans, G. A., Shykind, B., Margulies, D. H. & Seidman, J. G. (1983) *Proc. Natl. Acad. Sci. USA* **80**, 2040–2043.
- Wong, G. H. W., Clark-Lewis, I., McKimm-Breschkin, J. L., Harris, A. W. & Schrader, J. W. (1983) *J. Immunol.* **131**, 788–793.
- Chandler, V. L., Maler, B. A. & Yamamoto, K. R. (1983) *Cell* **23**, 489–499.