

Erratum to: Practical consensus guidelines for the management of enuresis

Johan Vande Walle · Soren Rittig · Stuart Bauer ·
Paul Eggert · Daniela Marschall-Kehrel · Serdar Tekgul

Published online: 1 May 2012
© Springer-Verlag 2012

Erratum to: Eur J Pediatr
DOI 10.1007/s00431-012-1687-7

There is an error in the section entitled Desmopressin therapy.

The text currently says:

Since tablets require up to 200 ml of fluid intake, which is ~25 % of a 7-year-old's bladder capacity, the Melt formulation is more suited to the antidiuretic indication of desmopressin [40].

This should in fact read:

Since tablets require up to 60 ml of fluid intake, which is ~25 % of a 7-year-old's bladder capacity, the Melt formulation is more suited to the antidiuretic indication of desmopressin [40].

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00431-012-1687-7>.

J. Vande Walle
Pediatric Nephrology Unit, Ghent University Hospital,
Ghent, Belgium

S. Rittig
Department of Pediatrics, Aarhus University Hospital,
Skejby, Denmark
e-mail: soren.rittig@skejby.rm.dk

S. Bauer
Department of Urology, Children's Hospital,
Harvard Medical School,
Boston, MA, USA
e-mail: stuart.bauer@childrens.harvard.edu

P. Eggert
University Children's Hospital,
Kiel, Germany
e-mail: p.eggert@pediatrics.uni-kiel.de

D. Marschall-Kehrel
Urology Consultancy,
Frankfurt/Main, Germany
e-mail: kehrel.daniela@t-online.de

S. Tekgul
Department of Urology, Hacettepe University School of Medicine,
Ankara, Turkey
e-mail: tekgul-k@tr.net

J. Vande Walle (✉)
Pediatric Nephrology UZ Ghent,
De Pintelaan 185,
9000 Ghent, Belgium
e-mail: johan.vandewalle@uzgent.be