

Symptoms of depression, anxiety, and somatization in female victims and perpetrators of intimate partner violence in Maputo City, Mozambique

Antonio Eugenio Zacarias^{1,2}

Gloria Macassa³

Joaquim JF Soares¹

Leif Svanström¹

Diddy Antai^{1,4}

¹Karolinska Institutet, Department of Public Health Sciences, Division of Social Medicine, Stockholm, Sweden;

²Eduardo Mondlane University, Faculty of Medicine, Maputo, Mozambique;

³University of Gävle, Department of Occupational and Health Sciences, Gävle, Sweden;

⁴Division of Global Health and Inequalities, The Angels Trust – Nigeria, Abuja, Nigeria

Background: Little knowledge exists in Mozambique and sub-Saharan Africa about the mental health (symptoms of depression, anxiety, and somatization) of women victims and perpetrators of intimate partner violence (IPV) by type of abuse (psychological aggression, physical assault without/with injury, and sexual coercion). This study scrutinizes factors associated with mental health among women victims and perpetrators of IPV over the 12 months prior to the study.

Methods and materials: Mental health data were analyzed with bivariate and multiple regression methods for 1442 women aged 15–49 years who contacted Forensic Services at Maputo Central Hospital (Maputo City, Mozambique) for IPV victimization between April 1, 2007 and March 31, 2008.

Results: In bivariate analyses, victims and perpetrators of IPV's scored higher on symptoms of mental health than their unaffected counterparts. Multiple regressions revealed that controlling behaviors, mental health comorbidity, social support, smoking, childhood abuse, sleep difficulties, age, and lack of education were more important in explaining symptoms of mental health than demographics/socioeconomics or life-style factors. Victimization and perpetration across all types of IPV were not associated with symptoms of mental health.

Conclusion: In our sample, victimization and perpetration were not important factors in explaining mental ill health, contrary to previous findings. More research into the relationship between women's IPV victimization and perpetration and mental health is warranted as well as the influence of controlling behaviors on mental health.

Keywords: women, depression, anxiety, somatization, victims, perpetrators, controlling behaviors, child abuse, social support

Introduction

Intimate partner violence (IPV) toward women has received increased global attention for its serious violation of women's health, legal rights, and quality of life.¹ The prevalence and magnitude of IPV makes it a public health problem and an important cause of morbidity and mortality,² with physical,^{2,3} sexual,^{4,5} and mental health consequences.^{6–10} Most studies investigating the mental health effects of IPV against women have focused predominantly on physical assault with or without injury and sexual coercion; however, the effect of psychological aggression on mental health within violent relationships may not have been sufficiently investigated.^{6,8,9,11–14} Existing studies show adverse effects of IPV on women's health, such as chronic pain,^{2,11,15} chronic stress-related symptoms, such as high blood pressure, abdominal pain, loss of appetite,¹⁶ and central nervous system problems, such as headaches, fainting, back pain, and seizures.^{2,11}

Correspondence: Antonio Eugenio Zacarias

Karolinska Institutet, Department of Public Health Sciences, Division of Social Medicine, 171 77 Stockholm, Sweden
Tel +258 823 087 990
Fax +46 08 334 693
Email antonio.zacarias@ki.se

Several studies have also suggested a relationship between psychological aggression and poor mental health effects, such as depression and anxiety.^{6,8,9,11,13,17–20} However, the frequent co-occurrence of psychological aggression with physical violence, as well as measurement difficulties, combine to reduce certainty regarding the independent effects of psychological aggression on mental health.²¹ Although knowledge about the comparative mental health effects of IPV by type is limited, some studies have indicated that psychological aggression may be more strongly linked to depression, anxiety, somatization, and post-traumatic stress disorder (PTSD)^{6,11,18–20,22} than other types of IPV.

Most studies reporting poor mental health among victimized women derive from high-income countries, and this relation may not be generalizable to low income countries, particularly in sub-Saharan Africa (SSA). Evidence from Botswana, Ethiopia, Ghana, Namibia, South Africa, and Tanzania points to a positive relation between IPV victimization and poor mental health, eg, depression.^{12,13,23–27}

Traditionally, women have been considered the predominant victims of IPV and men the perpetrators of IPV; however, the picture may be more complex than previously thought, as indicated by increasing evidence reporting women as perpetrators of violence,^{28–30} particularly in response to prior abuse, and often with high prevalence rates. Although most studies of women as perpetrators of IPV have focused mainly on physical assault, it is not uncommon for women to employ other forms of abuse against men, eg, psychological aggression.^{31–34} However, not much research attention has been given to the relation between women's IPV perpetration and their own health status. Existing studies, which are mainly from Western countries, have revealed that female perpetrators of IPV can have mental health problems, such as depression and anxiety.^{18,24,26,35–38} As far as we know, only one study has addressed the comparative effects of women's use of various IPV types on their own mental health, with results showing that psychological aggression was more strongly associated with depression, anxiety, and somatization than with other IPV types, eg, physical assault.¹⁸

Studies addressing the mental health of female victims and perpetrators of IPV are frequently limited to demographic/socioeconomic and life-style factors, particularly among perpetrators, whereas the effects of important factors, such as controlling behaviors, are not often considered. Omission of what has been shown to be a potentially crucial factor in explaining mental health consequences

could skew the relationship between IPV and mental health consequences towards confirmation. Controlling behaviors have been associated with women as victims^{40,41} and perpetrators of IPV,^{24,26,32,34,42–44} as well as with poor mental health (eg, depression) among female victims and perpetrators of IPV.^{18,24,26,45} The effect of controlling behavior on mental health is reported to be more salient than the effect of victimization.⁴⁵

Therefore, the aim of this study was to investigate the association between women as victims and perpetrators of IPV in relation to mental health consequences (depression, anxiety, and somatization) and associated factors in order to provide a better understanding of mental health in women victims and perpetrators of IPV and its management in Mozambique and similar contexts.

Methods and materials

Setting

Cross-sectional data gathered during the 12 months between April 1, 2007 and March 31, 2008 (consecutive cases) among women visiting the Forensic Services' unit at Maputo Central Hospital for abuse by a partner were used for this study. The interviews were conducted by selected trained females: four medical students from the Faculty of Medicine, and two nurses from Forensic Services. These interviewers were informed in detail about the research and various facets of IPV. They were also thoroughly informed about each and every scale included in the questionnaire, and they were trained to use it. Upon their arrival at the Forensic Services unit, the participant women were approached by the interviewers and carefully informed about all aspects of the study and the extent of their participation. Voluntariness and confidentiality were strongly emphasized. The participants were also informed that refusal to participate would not result in any negative consequences. If the women agreed to participate, consent was obtained (verbally and/or in writing). All the participants provided verbal consent prior to the commencement of the interviews, which were on average 1 hour and conducted in an isolated room by means of a questionnaire. Inclusion criteria included: (a) age 15–49 years; (b) victims of IPV; (c) acceptance to participate; and (d) resident in Maputo city, Mozambique. Data processing and preservation followed the usual anonymous and confidentiality rules, rendering public only results from aggregated data. The results (aggregate data) were available for participants upon request.

Participants

A total of 1500 women aged 15–49 years who had been exposed to abuse by a partner were approached. Of this number, 1442 women agreed to participate in the study (response rate of 96.1%), and 58 declined. Responses to the questions about violence were received from 1429 to 1433 participants depending on the type of violence. These women were referred by female nongovernmental organizations or the police, or they were self-referred.

Measures

Outcome

Mental health was assessed with the Symptom Check List-Revised (SCL-90-R),⁴⁶ which consists of 90 items divided into nine symptom dimensions. This study used dimensions of symptoms of depression, anxiety, and somatization. Depression was assessed using 13 items covering areas such as thoughts of suicide and loss of vital energy. Anxiety was assessed using 10 items that reflect feelings such as nervousness and fear. Somatization was assessed using 12 items covering areas such as cardiovascular and gastrointestinal complaints. The item scores ranged from 0 to 4 (not at all, to very much). The scores ranged from 0–52 for symptoms of depression, 0–40 for anxiety, and 0–48 for somatization, with high scores corresponding to high symptom reporting. Among women as victims, the Cronbach's α were 0.83 for symptoms of depression, 0.81 for anxiety, and 0.82 for somatization. The corresponding Cronbach's α among women as perpetrators were 0.83, 0.81, and 0.83, respectively.

Exposure

Intimate partner violence (IPV) was measured with the Conflict Tactic Scales-Version 2 (CTS2),⁴⁷ consisting of 39 items, aimed at respondents both as victims and as perpetrators (a total of 78 items). The CTS2 contains questions assessing the following: cognitive (eg, I suggested a compromise to a disagreement) and emotional (eg, I showed my partner I cared even though we disagreed)-oriented negotiation; minor (eg, insulted or swore at my partner) and severe (eg, called my partner fat or ugly) psychological aggression; minor (eg, pushed or shoved my partner) and severe (eg, beat up my partner) physical assault; minor (eg, made my partner have sex without a condom) and severe sexual coercion (eg, used threats to make my partner have sex); minor (eg, had a sprain) and severe (eg, had a broken bone from a fight with my partner) physical assault with injury; as well as chronicity of abusive acts (how often the acts happened),

which may have occurred once, twice, 3–5, 6–10, 11–20, or >20 times during the past year or did not occur in the past year but occurred before or never happened.^a The CTS2 is a previously validated and reliable instrument.⁴⁷ Cronbach's α among women victims was 0.89 for physical assault, 0.82 for psychological aggression, 0.73 for sexual coercion, and 0.65 for physical assault with injury. The correspondent Cronbach's α among women perpetrators were 0.79, 0.73, 0.70, and 0.63, respectively.

Abuse in childhood was measured with four items, one each for psychological abuse (eg, shouted or yelled at); physical abuse (eg, beaten up); sexual abuse (eg, forced to have sex); injury (eg, bruised); and chronicity (how often the acts happened). The acts may have happened once, twice, 3–5, 6–10, 11–20 or >20 times or never happened.^b The items obtained data about the respondent's exposure to abuse before 15 years of age. Cronbach's α was 0.72 for physical abuse, 0.70 for psychological abuse, 0.68 for sexual abuse, and 0.71 for injury.

Controlling behaviors were measured with the Controlling Behaviors Scale – Revised (CBS-R), consisting of 24 items^{43,44} that do not include items of physical assault. The CBS-R has good discriminative ability and can be scored to derive five subscores, each representing a particular type of control tactic or a total controlling behavior score (eg, “try to restrict time individual spent with family or friends”). The response format ranges from 0 to 4 (never to always), with a total range of 0–96. The women reported how often their partner used each act of controlling behavior toward them, as well as how often they used each act of controlling behavior toward their partner in the past year. This study used the total controlling behavior, and high scores corresponded to high controlling behaviors. Cronbach's α was 0.91 for women using control and 0.93 for women being controlled.

Social support was measured with the 12-item Schedule for Social Interaction.⁴⁸ Six items assessed “social attachment” (availability of deep emotional relationships) and the other six items assessed “social integration” (availability of peripheral social networks). Item scores ranged from 1 to 6 (ie, not available to available), with social attachment scores ranging from 0 to 6, social integration ranging from

^aThe CTS2 are scored by adding the midpoints of the response categories. For the categories 0, 1 and 2 the midpoints are the same. For category 3 (3–5 times) the midpoint is 4; for category 4 (6–10 times) the midpoint is 8; for category 5 (11–20 times) the midpoint is 15; and for category 6 (>20 times) the midpoint is 25.

^bThe scaling was based on CTS2. See also footnote a.

6 to 36, and social interaction (total) ranging from 6 to 42. High scores (components, total) corresponded to high social support. For this study, social attachment and social integration were used. Cronbach's α for social attachment among women as victims and perpetrators were 0.75 and 0.80, respectively, and corresponding Cronbach's α for social integration were 0.87 and 0.88, respectively.

Use of alcohol/cigarettes and sleep difficulties were assessed as dichotomous yes/no variables. Body mass index (BMI) was based on self-reported height and weight and calculated for each woman with the formula, kg/m^2 . Finally, demographic and socioeconomic characteristics were assessed with a classification system used in Mozambique (Ministry Council–Ministry of Finance), and included the following: age (in years), marital status (categorized as single, married/cohabitant, divorced/separated, and widow), children at home (assessed as yes or no variable), housing (categorized as conventional, and nonconventional), education (assessed as no education, low, intermediate, and high), occupational status (assessed as blue collar-worker, low white-collar-worker, middle-high white collar-worker and student/other), socioeconomic status (assessed as work for other, liberal/own business, student and domestic/other), salary/financial resources (categorized yes or no), and financial strain, which reflected how respondents make ends meet, was assessed as “no,” “sometimes,” “often,” and “always” format. A woman was regarded as experiencing financial strain if her response was anything but “no.”

Statistical analyses

Cross-tabulations/means and standard deviations (SD), and Pearson's chi-square analyses were used to assess the demographic, socioeconomic, and lifestyle factors associated with the IPV types. The significance level was set at $P < 0.05$. Using analysis of variance, descriptive statistics of mental health scores (symptoms of depression, anxiety, and somatization) were examined among women who were victims of IPV types compared with those who were not victims of IPV, as well as women who were perpetrators of IPV types (psychological aggression, physical assault without/with injury, sexual coercion) compared with those who were not perpetrators of IPV during the past 12 months.

Multiple linear regression analyses were conducted to examine the explanatory factors in the association between women's IPV victimization and perpetration during the past 12 months independent of type and the outcome variables (symptoms of depression, anxiety, and somatization). The selected factors (exposures) were variables significantly

associated with the IPV types in bivariate analyses conducted in a previous study.⁴⁹ These included married/cohabitant, secondary education, blue-collar worker and middle/high white-collar (occupational status), work for others (socioeconomic status), salary/financial resources, financial strain, children at home, living in nonconventional housing, BMI, use of cigarettes and alcohol, abuse as a child, and controlling behaviors (controlling behaviors over/by partner). In addition, sleep difficulties, being a victim and a perpetrator of IPV across all types of abuse, as well as social support variables, were added. Finally, depending on the outcome being analyzed, depression, anxiety, or somatization were also used as exposure variables to assess for influence of comorbidity with other mental health consequences. For example, in the analysis of depression and associated factors, anxiety and somatization were added as exposure variables.^c All variables were entered into the multiple linear regression models in a single block to control for possible confounding between these variables. Results were expressed as standardized betas and P -values. Significance levels for bivariate and multiple regression analyses were set at $P < 0.05$.

Ethical consideration

Ethical authorization was given by The National Ethical Committee at the Ministry of Health of Mozambique (122/CNBS/06).

Results

Characteristics of the women by IPV types

Differences in demographic, socioeconomic, and lifestyle factors associated with IPV types are presented in Table 1.

Psychological aggression

Significantly higher proportions of women who reported psychological aggression were single (53%, $n = 489$; $P < 0.0001$), had intermediate educational level (56%, $n = 51$; $P < 0.001$), and did not use alcohol (60%, $n = 558$; $P = 0.018$). However, there were no significant associations between psychological aggression and age, children at home, housing, occupational status, socioeconomic status, salary/financial resources, financial strain, BMI and smoking.

^cOmitting control, abuse as a child and the mental dimensions as exposure factors in the models did not affect the importance of victimization and perpetration across types (data not shown here). Replacing violence across types by each and every type did not change either the picture (data not shown here).

Table 1 Demographics, socioeconomics, and life-style characteristics of women

Characteristics	Psychological (n = 1433)		Physical (n = 1432)		Sexual (n = 1433)		Injury (n = 1432)	
	Yes (%)	No (%)	Yes (%)	No (%)	Yes (%)	No (%)	Yes (%)	No (%)
Age (years)	<i>P</i> = 0.398		<i>P</i> = 0.120		<i>P</i> = 0.626		<i>P</i> < 0.0001	
Mean ± SD	28.7 ± 8.1	28.8 ± 8.7	28.7 ± 7.9	29.5 ± 8.8	28.6 ± 7.9	29.6 ± 8.7	29.6 ± 7.7	28.8 ± 8.7
Marital status	(n = 1405)	<i>P</i> < 0.0001	(n = 1404)	<i>P</i> < 0.0001	(n = 1405)	<i>P</i> < 0.0001	(n = 1404)	<i>P</i> < 0.0001
Single	489 (53)	251 (51)	388 (51)	352 (55)	390 (54)	350 (51)	209 (43)	531 (57)
Married/cohabitant	302 (33)	103 (21)	253 (34)	151 (23)	230 (32)	175 (25)	174 (36)	230 (25)
Divorced/separated	88 (10)	90 (18)	86 (11)	92 (14)	68 (10)	110 (16)	75 (16)	103 (11)
Widow	36 (4)	46 (10)	32 (4)	50 (8)	26 (4)	56 (8)	22 (5)	60 (7)
Children at home	(n = 1348)	<i>P</i> = 0.171	(n = 1347)	<i>P</i> = 0.501	(n = 1348)	<i>P</i> = 0.988	(n = 1347)	<i>P</i> = 0.252
Yes	649 (73)	352 (77)	554 (75)	446 (73)	513 (74)	488 (74)	397 (85)	603 (68)
No	239 (27)	108 (13)	185 (25)	162 (27)	178 (26)	169 (26)	71 (15)	276 (32)
Housing	(n = 1408)	<i>P</i> = 0.667	(n = 1407)	<i>P</i> = 0.897	(n = 1408)	<i>P</i> = 0.337	(n = 1407)	<i>P</i> = 0.01
Conventional ^a	787 (86)	411 (84)	650 (85)	547 (85)	615 (86)	583 (85)	398 (82)	799 (87)
Nonconventional ^b	134 (14)	76 (16)	116 (15)	94 (15)	104 (14)	106 (15)	89 (18)	121 (13)
Education	(n = 1433)	<i>P</i> = 0.001	(n = 1432)	<i>P</i> = 0.397	(n = 1433)	<i>P</i> = 0.090	(n = 1432)	<i>P</i> = 0.356
No ^c	45 (5)	46 (9)	43 (6)	48 (7)	37 (5)	54 (7)	34 (7)	57 (6)
Low ^d	236 (25)	148 (30)	206 (26)	178 (28)	185 (26)	199 (29)	145 (30)	239 (26)
Inter ^e	519 (56)	241 (49)	427 (55)	333 (51)	399 (55)	361 (52)	252 (51)	508 (54)
High ^f	131 (14)	57 (12)	98 (13)	89 (14)	105 (14)	83 (12)	59 (12)	128 (14)
Occupational status	(n = 1433)	<i>P</i> = 0.314	(n = 1432)	<i>P</i> = 0.744	(n = 1433)	<i>P</i> = 0.901	(n = 1432)	<i>P</i> < 0.0001
Blue-collar worker	504 (56)	294 (62)	440 (59)	358 (57)	403 (57)	395 (58)	300 (64)	498 (55)
Low white-collar worker	47 (5)	18 (4)	35 (5)	30 (5)	35 (5)	30 (4)	23 (5)	42 (5)
Middle/high white-collar worker	103 (11)	50 (10)	86 (12)	67 (10)	82 (12)	71 (11)	67 (14)	86 (9)
Student/other	249 (28)	116 (24)	187 (25)	177 (28)	182 (26)	183 (27)	78 (17)	286 (31)
Socioeconomic status	(n = 1417)	<i>P</i> = 0.114	(n = 1416)	<i>P</i> = 0.117	(n = 1417)	<i>P</i> = 0.111	(n = 1416)	<i>P</i> < 0.0001
Work for others ^g	320 (34)	145 (30)	277 (36)	188 (29)	264 (37)	201 (29)	197 (40)	268 (29)
Liberal/own business ^h	149 (16)	99 (20)	124 (16)	124 (20)	118 (16)	130 (19)	87 (18)	161 (17)
Student	251 (27)	115 (24)	190 (24)	175 (27)	184 (25)	182 (26)	80 (16)	285 (31)
Domestic/other ⁱ	210 (23)	128 (26)	183 (24)	155 (24)	159 (22)	179 (26)	126 (26)	212 (23)
Salary/financial resources	(n = 1386)	<i>P</i> = 0.655	(n = 1386)	<i>P</i> = 0.202	(n = 1386)	<i>P</i> = 0.063	(n = 1386)	<i>P</i> < 0.0001
Yes	424 (46)	213 (45)	362 (48)	275 (44)	344 (48)	293 (43)	256 (53)	381 (42)
No	490 (54)	259 (55)	400 (52)	349 (56)	376 (52)	382 (57)	224 (47)	525 (58)
Financial strain	(n = 1405)	<i>P</i> = 0.102	(n = 1404)	<i>P</i> = 0.812	(n = 1405)	<i>P</i> = 0.368	(n = 1404)	<i>P</i> = 0.012
Yes	774 (84)	420 (87)	651 (85)	542 (85)	605 (84)	589 (86)	429 (88)	764 (83)
No	149 (16)	62 (13)	117 (15)	94 (15)	114 (16)	97 (14)	57 (12)	154 (17)
BMI (n)	(n = 1316)	<i>P</i> = 0.734	(n = 1315)	<i>P</i> = 0.120	(n = 1316)	<i>P</i> = 0.569	(n = 1315)	<i>P</i> = 0.185
Mean ± SD	25 ± 4.6	24.5 ± 4.8	25.1 ± 4.6	24.5 ± 4.8	24.9 ± 4.5	24.7 ± 4.9	25.4 ± 4.7	24.5 ± 4.7
Alcohol use	(n = 1412)	<i>P</i> = 0.018	(n = 1432)	<i>P</i> = 0.009	(n = 1433)	<i>P</i> < 0.001	(n = 1432)	<i>P</i> = 0.009
Yes	364 (40)	156 (31)	310 (41)	210 (33)	300 (42)	220 (32)	205 (42)	315 (34)
No	558 (60)	334 (69)	456 (59)	435 (67)	419 (58)	473 (68)	281 (58)	610 (66)
Cigarette use	(n = 1402)	<i>P</i> = 0.587	(n = 1401)	<i>P</i> = 0.066	(n = 1402)	<i>P</i> = 0.003	(n = 1398)	<i>P</i> < 0.0001
Yes	71 (7)	34 (9)	66 (8)	39 (6)	68 (9)	37 (5)	57 (12)	48 (5)
No	843 (93)	454 (91)	721 (92)	575 (94)	645 (91)	652 (95)	426 (88)	870 (95)

Notes: ^aHousing in cement; ^bHousing in local material, eg, mud, wood; ^cCannot read/write; ^dPrimary school/similar; ^eSecondary school/similar; ^fUniversity/similar; ^geg, clerks; ^heg, accountants; ⁱAt home/unemployed.

Abbreviation: SD, standard deviation.

Physical assault

Among women who reported physical assault, a significantly higher proportion were single (51%, *n* = 388; *P* < 0.0001) and did not drink alcohol (59%, *n* = 456; *P* = 0.009). However, there were no significant associations between physical assault and age, children at home, housing, educational level,

occupational status, socioeconomic status, salary/financial resources, financial strain, BMI, and smoking.

Sexual coercion

A significantly higher proportion of women who reported sexual coercion were single (54%, *n* = 390; *P* < 0.0001),

did not drink alcohol (58%, $n = 419$; $P < 0.0001$), and did not smoke cigarettes (91%, $n = 645$; $P = 0.003$). However, there were no significant associations between sexual coercion and age, children at home, housing, educational level, occupational status, socioeconomic status, salary/financial resources, financial strain and BMI.

Physical assault with injury

The number of women reporting physical assault with injury was significantly associated with age (Mean/SD = 29.6 ± 7.7 ; $P < 0.0001$) and included a higher proportion of women who were single (43%, $n = 209$; $P < 0.0001$), resident in conventional housing (82%, $n = 398$; $P < 0.01$), were blue collar workers (64%, $n = 300$; $P < 0.0001$), worked for others (40%, $n = 197$; $P < 0.0001$), received salary/financial resources (53%, $n = 256$; $P = 0.0001$), were experiencing financial strain (88%, $n = 429$; $P = 0.012$), did not drink (58%, $n = 281$; $P = 0.009$), and did not smoke (88%, $n = 426$; $P < 0.0001$). However, there were no significant associations among physical assault with injury and children at home, educational level, and BMI.

Descriptive statistics of mental health scores by IPV type for victims and perpetrators

Results of the descriptive statistics are presented in Table 2.

Victims

Compared to women who were not psychologically abused, women who were psychologically abused scored higher on symptoms of depression ($F[1,1419] = 27.9$; $P < 0.0001$), anxiety ($F[1,1419] = 10.6$; $P = 0.0012$) and somatization ($F[1,1419] = 14.3$; $P = 0.002$). Similarly, the following symptom scores were exhibited among women who were physically assaulted: depression ($F[1,1419] = 54.8$; $P < 0.0001$), anxiety ($F[1,1419] = 24.6$; $P < 0.0001$), and somatization ($F[1,1419] = 23.1$; $P < 0.0001$); women who experienced sexual coercion: depression ($F[1,1419] = 21.3$; $P = 0.0001$), anxiety ($F[1,1420] = 18.5$; $P < 0.0001$), and somatization ($F[1,1419] = 15.7$; $P = 0.0001$); and women who experienced physical assault with injury: depression ($F[1,1419] = 45.5$; $P < 0.0001$), anxiety ($F[1,1419] = 27.1$; $P < 0.0001$), and somatization ($F[1,1419] = 26.6$; $P < 0.0001$), compared to women who were not physically assaulted, did not experience sexual coercion, and did not experience physical assault with injury, respectively (Table 2).

Table 2 Mean/SD for SCL-90-R (symptoms of depression, anxiety, and somatization) among women victims and perpetrators of intimate partner violence by type (psychological aggression, physical assault, sexual coercion, and injury) within/above psychiatric outpatient norms (POPn)

Variables	Psychological (n = 1431)		Physical (n = 1429)		Sexual (n = 1431)		Injury (n = 1431)		POPn (n = 577)	
	Yes ^a n	No ^b n	Yes ^a n	No ^b n	Yes ^a n	No ^b n	Yes ^a n	No ^b n	Yes ^a n	No ^b n
Victims (n)	(1421)		(1421)		(1421)		(1421)		(1421)	
Depression	930; $P < 0.0001$	491	773; $P < 0.0001$	648	725; $P < 0.0001$	696	489; $P < 0.0001$	932	1.94 ± 0.93	
Mean (SD)	1.3 (0.875)	1.1 (0.900)	1.4 (0.847)	1.0 (0.909)	1.3 (0.889)	1.1 (0.878)	1.4 (0.878)	1.1 (0.879)		
Anxiety	930; $P = 0.0012$	491	773; $P = 0.0001$	648	725; $P < 0.0001$	696	489; $P < 0.0001$	932	1.59 ± 0.90	
Mean (SD)	1.0 (0.877)	0.856 (0.898)	1.1 (0.879)	0.835 (0.893)	1.1 (0.897)	0.851 (0.864)	1.1 (0.896)	0.874 (0.870)		
Somatization	930; $P = 0.002$	491	773; $P = 0.0001$	648	725; $P = 0.0001$	696	489; $P < 0.0001$	932	0.99 ± 0.78	
Mean (SD)	0.854 (0.688)	0.709 (0.691)	0.884 (0.681)	0.708 (0.694)	0.875 (0.689)	0.730 (0.686)	0.933 (0.694)	0.736 (0.682)		
Perpetrators (n)	(1421)		(1420)		(1421)		(1421)			
Depression	910; $P = 0.0269$	511	542; $P = 0.0028$	878	557; $P = 0.00816$	864	321; $P < 0.0001$	1100	1.94 ± 0.93	
Mean (SD)	1.2 (0.894)	1.1 (0.936)	1.3 (0.872)	1.1 (0.928)	1.2 (0.875)	1.2 (0.933)	1.3 (0.799)	1.0 (0.824)		
Anxiety	910; $P = 0.0138$	511	542; $P < 0.0001$	878	557; $P = 0.0004$	864	321; $P < 0.0001$	1100	1.59 ± 0.90	
Mean (SD)	1.0 (0.864)	0.880 (0.933)	1.1 (0.846)	0.879 (0.908)	1.1 (0.870)	0.890 (0.899)	1.3 (0.840)	0.868 (0.886)		
Somatization	910; $P = 0.0076$	511	542; $P = 0.0008$	878	557; $P = 0.0017$	864	321; $P < 0.0001$	1100	0.99 ± 0.78	
Mean (SD)	0.837 (0.674)	0.734 (0.738)	0.879 (0.667)	0.751 (0.715)	0.859 (0.665)	0.768 (0.708)	0.909 (0.604)	0.740 (0.683)		

Notes: ^aSustained or inflicted; ^bNot sustained or not inflicted.

Abbreviation: SD, standard deviation.

Perpetrators

Compared to women who did not perpetrate any IPV type, women who were perpetrators of psychological aggression scored higher on symptoms of depression ($F[1,1419] = 4.9$; $P = 0.0269$), anxiety ($F[1,1419] = 6.1$; $P = 0.0138$), and somatization ($F[1,1419] = 7.1$; $P = 0.0076$). Similar higher scores were observed among perpetrators of physical assault for symptoms of depression ($F[1,1418] = 8.9$; $P = 0.0028$), anxiety ($F[1,1418] = 17.4$; $P < 0.0001$), and somatization ($F[1,1418] = 11.3$; $P = 0.0008$), compared to women who did not perpetrate physical assault; among perpetrators of sexual coercion for symptoms of depression ($F[1,1419] = 3.04$; $P = 0.0081$), anxiety ($F[1,1419] = 12.7$; $P = 0.0004$), and somatization ($F[1,1419] = 5.75$; $P = 0.0017$), compared to women who did not perpetrate sexual coercion. Scores among women who perpetrated physical assault with injury were symptoms of depression ($F[1,1419] = 44.0$; $P < 0.0001$), anxiety ($F[1,1419] = 51.1$; $P < 0.0001$), and somatization ($F[1,1419] = 32.1$; $P < 0.0001$), compared to women who perpetrated physical assault (Table 2).

Multiple regression analyses of mental health

Results of the multiple linear regression analyses conducted to examine factors associated with the different outcome variables are presented in Table 3.

Symptoms of depression

There were significantly negative associations between symptoms of depression and being married/cohabitant ($\beta = -0.053$; $P = 0.016$), and social attachment ($\beta = -0.055$; $P = 0.011$). In contrast, symptoms of depression were positively associated with no education ($\beta = 0.047$; $P = 0.042$), controlling behaviors by partner ($\beta = 0.174$; $P < 0.0001$), sleep difficulties ($\beta = 0.067$; $P = 0.001$), as well as comorbidity with symptoms of anxiety ($\beta = 0.600$; $P < 0.0001$), and somatization ($\beta = 0.165$; $P < 0.0001$).

Symptoms of anxiety

The associations between symptoms of anxiety and smoking ($\beta = 0.051$; $P < 0.012$), control over partner ($\beta = 0.086$; $P < 0.0001$), and comorbidity with depression ($\beta = 0.619$; $P < 0.0001$), as well as somatization ($\beta = 0.276$; $P < 0.0001$) were positive and significant, whereas symptoms of anxiety were negatively associated with control by partner ($\beta = -0.102$; $P < 0.0001$), and social attachment ($\beta = -0.047$; $P = 0.030$).

Symptoms of somatization

Age ($\beta = 0.082$; $P = 0.004$), sleep difficulties ($\beta = 0.107$; $P < 0.0001$), control by partner ($\beta = 0.082$; $P = 0.005$), social attachment ($\beta = 0.056$; $P = 0.034$), social integration ($\beta = 0.058$; $P = 0.021$), and abuse as a child ($\beta = 0.097$; $P < 0.0001$), as well as comorbidity with symptoms of depression ($\beta = 0.253$; $P < 0.0001$), and anxiety ($\beta = 0.409$; $P < 0.0001$) were significantly and positively associated with reporting symptoms of somatization (Table 3).

Discussion

Characteristics of the women by IPV types

This study examined the associations between IPV types during the past 12 months, the mental health consequences of IPV, and explanatory factors among women resident in Maputo City, Mozambique who visited the Forensic Services unit at Maputo Central Hospital to get legal assistance by completing a clinical forensic report. Consistent with previous studies,^{50,51} our results indicated varying significant differences between exposure to IPV and several factors, such as age, marital status, housing, educational level, occupational and socioeconomic status, financial resources and financial strain, and alcohol and cigarette consumption. However, as shown in the multiple regressions analyses, the explanatory factors in our sample population were not important factors in elucidating symptoms of depression, anxiety and somatization, which is consistent with previous studies.^{11,12,19,23}

Descriptive analyses of IPV by type and mental health

In general, women's IPV victimization and perpetration during the past 12 months were related to symptoms of depression, anxiety, and somatization. However, significant effects were more apparent in relation to being victimized by abuse than by perpetrating abuse. This is consistent with previous findings concerning IPV victimization,^{7,8,17,23,24,26,36,52} as well as IPV perpetration.^{35,37-39,53,54}

Among victims, symptoms of depression, anxiety, and somatization were somewhat less evident in conjunction with psychological aggression than with the other violence types (as indicated by the level of significance of the associations), which is not similar to findings in other studies.^{11,18,19} Similarly, among perpetrators, symptoms of mental health consequences were somewhat more significantly associated with other types of IPV compared to psychological aggression, which is contrary to a recent study showing a stronger

Table 3 Factors associated with symptoms of depression, anxiety and somatization expressed as betas (β) and *P*-values among all women

Variables	Depression		Anxiety		Somatization	
	β	<i>P</i> -values	β	<i>P</i> -values	β	<i>P</i> -values
Age	-0.027	<i>P</i> = 0.249	-0.008	<i>P</i> = 0.727	0.082	<i>P</i> = 0.004
Marital status						
Single						
Married/cohabitant	-0.053	<i>P</i> = 0.016	-0.004	<i>P</i> = 0.847	0.023	<i>P</i> = 0.407
Divorced/separated	-0.037	<i>P</i> = 0.081	-0.036	<i>P</i> = 0.092	-0.005	<i>P</i> = 0.843
Widow	0.017	<i>P</i> = 0.404	0.002	<i>P</i> = 0.903	-0.020	<i>P</i> = 0.429
Children at home	0.041	<i>P</i> = 0.095	-0.019	<i>P</i> = 0.432	-0.003	<i>P</i> = 0.913
Housing						
Conventional						
Nonconventional	0.008	<i>P</i> = 0.661	0.024	<i>P</i> = 0.195	-0.020	<i>P</i> = 0.390
Education level						
No	0.047	<i>P</i> = 0.042	-0.043	<i>P</i> = 0.070	0.037	<i>P</i> = 0.200
Low	0.019	<i>P</i> = 0.571	-0.042	<i>P</i> = 0.212	0.058	<i>P</i> = 0.151
Inter	-0.010	0.741	-0.022	<i>P</i> = 0.483	0.070	<i>P</i> = 0.063
High						
Occupational status						
Blue-collar worker						
Low white-collar worker	0.003	<i>P</i> = 0.895	-0.033	<i>P</i> = 0.105	0.037	<i>P</i> = 0.133
Middle/high white-collar worker	0.001	<i>P</i> = 0.976	0.024	<i>P</i> = 0.308	0.001	<i>P</i> = 0.965
Student/other	0.033	<i>P</i> = 0.819	-0.078	<i>P</i> = 0.597	-0.028	<i>P</i> = 0.857
Socioeconomic status						
Worker for others						
Liberal/own business	-0.019	<i>P</i> = 0.394	0.015	<i>P</i> = 0.527	0.050	<i>P</i> = 0.075
Student	-0.018	<i>P</i> = 0.904	0.063	<i>P</i> = 0.672	0.088	<i>P</i> = 0.627
Domestic/other	0.002	<i>P</i> = 0.944	0.001	<i>P</i> = 0.971	0.024	<i>P</i> = 0.521
Salary/financial resources	0.004	<i>P</i> = 0.896	-0.022	<i>P</i> = 0.434	0.013	<i>P</i> = 0.707
Financial strain	0.010	<i>P</i> = 0.609	-0.002	<i>P</i> = 0.931	-0.004	<i>P</i> = 0.862
BMI	0.027	<i>P</i> = 0.169	-0.031	<i>P</i> = 0.111	0.028	<i>P</i> = 0.248
Drinking	0.026	<i>P</i> = 0.170	-0.011	<i>P</i> = 0.576	0.005	<i>P</i> = 0.824
Smoking	-0.029	<i>P</i> = 0.141	0.051	<i>P</i> = 0.012	-0.031	<i>P</i> = 0.212
Sleep difficulties	0.067	<i>P</i> = 0.001	-0.006	<i>P</i> = 0.759	0.107	<i>P</i> < 0.0001
Depression	-	-	0.619	<i>P</i> < 0.0001	0.253	<i>P</i> < 0.0001
Anxiety	0.600	<i>P</i> < 0.0001	-	-	0.409	<i>P</i> < 0.0001
Somatization	0.165	<i>P</i> < 0.0001	0.276	<i>P</i> < 0.0001	-	-
Control by partner	0.174	<i>P</i> < 0.0001	-0.102	<i>P</i> < 0.0001	0.082	<i>P</i> = 0.005
Control over partner	-0.020	<i>P</i> = 0.414	0.086	<i>P</i> < 0.0001	-0.027	<i>P</i> = 0.371
Social attachment	-0.055	<i>P</i> = 0.011	-0.047	<i>P</i> = 0.030	0.056	<i>P</i> = 0.034
Social integration	-0.003	<i>P</i> = 0.887	0.001	<i>P</i> = 0.972	0.058	<i>P</i> = 0.021
Victimization	-0.032	<i>P</i> = 0.336	0.052	<i>P</i> = 0.126	-0.033	<i>P</i> = 0.430
Perpetration	0.028	<i>P</i> = 0.407	-0.065	<i>P</i> = 0.062	0.053	<i>P</i> = 0.210
Abuse as a child	0.010	<i>P</i> = 0.605	-0.025	<i>P</i> = 0.196	0.097	<i>P</i> < 0.0001

effect of psychological aggression on mental health than other forms of IPV.¹⁸

Multiple regression analyses of mental health

Victimization and perpetration during the past 12 months were not independently associated with symptoms of depression, anxiety, and somatization. Explanations may be

twofold: the first is methodological, in that the analysis was first based on women's self-report of these symptoms rather than on clinical evidence; the second is the possibility that other factors, such as controlling behaviors, may have been more crucial in explaining the mental health of the women in our study than the abusive acts were. As suggested elsewhere,⁴⁹ these women were involved in relationships within which the occurrence of mutual abuse (ie, being both abused

and abusing) was the norm rather than the exception. Thus, the effect of abuse on their mental health may not have been significantly “visible” because these women may have adapted to abuse as a “normal” part of their lives. It is also possible that mental health problems may have existed prior to the abuse (the past 12 months), and therefore did not have a major influence on them. Our findings therefore stress the need for more research using longitudinal study designs to investigate the relation between IPV victimization, perpetration, and mental health in the Sub-Saharan African (including Mozambique) context.

Controlling behaviors by partner was positively associated with symptoms of depression and somatization and might be attributed to the effects of men’s controlling behaviors on women to instill fear and limit their independence and physical and social freedoms, which in turn might lead to feelings of hopelessness, reduced self-esteem, and psychopathology in the form of symptoms of depression and somatization. However, controlling behaviors by partner was also negatively associated with anxiety. The explanation for this result might be that women who have adapted to and accepted the social norms imposed by the patriarchal society in which they reside (characterized by tolerance of gender roles, economic dependency, and IPV) were less likely to exhibit symptoms of anxiety. Interestingly, women’s controlling behaviors over their partner was also associated with anxiety, which may indicate fear of the partner’s reactions or that the women were ashamed of their coercive behaviors. These findings therefore warrant further investigation. However, our findings are in agreement with those of recent studies, which indicate a relation between controlling behaviors and poor mental health,^{18,24,26,45,55} and the effects of controlling behaviors upon mental health were more marked than those of victimization.⁴⁵

The association of abuse as a child with symptoms of somatization is consistent with previous findings linking this abuse to poor mental health, including somatization.^{15,26,56–58} Although the mechanisms explaining the link between childhood trauma and symptoms of somatization in adulthood remain unclear, psychoanalytic explanations remain outside the scope of this study. However, it is plausible that a two-way social learning process is involved; on the one hand, as children the women may have been punished for going against social norms and thereby learned to suppress emotional expression. On the other hand, the effects of abuse in adulthood may have resulted in remorse and positive attention from significant others (ie, the perpetrator). These women may then have acquired an increased awareness of,

and sensitivity to, bodily reactions and a distorted perception of, eg, the experienced emotional reactions.

Sleep difficulties were associated with symptoms of depression and somatization, which is consistent with previous studies that found an association between sleep difficulties and depression and somatization.^{59,60} Contradictory findings have found that depression may lead to insomnia and that sleep difficulties may contribute to depression.⁶¹ It is possible that the women’s problematic situations (eg, history of child abuse) may have led to sleep difficulties, which over time resulted in a range of sadness, hopelessness, and body sensations characteristics of symptoms of depression and somatization. Depression and somatization involve various sensations (eg, loss of interest and musculoskeletal pains) which, if persistent, might over time result in sleep difficulties. However, due to the cross-sectional character of our data, a causal inference could not be made.

The association between smoking and symptoms of anxiety is consistent with previous studies.^{62–64} As we did not assess in depth the participant’s smoking patterns, it is difficult to fully assess the role of smoking in this study. However, one could hypothesize that these women smoked in order to alleviate the symptoms of anxiety. Social attachment was negatively associated with symptoms of depression and anxiety, suggesting a protective effect against mental health consequences. Women with persons (eg, family members) with whom they could share, eg, deep emotions and preoccupations tend to be less likely to experience intense and persistent depressive and anxious symptoms, which is consistent with previous observations that such ties have a protective effect in depression and anxiety.^{65–67} Possible explanations may be that the women developed self-esteem and gained much needed support that helped them to cope with the strains of depressive and anxious symptoms. Nevertheless, social attachment and social integration were positively associated with symptoms of somatization. Although the role of social support in somatization appears to be complex, it may be that persons (eg, family members and work colleagues) with whom the women could share, talk freely about various matters, and turn to for moral support may also have been more likely to experience somatic symptoms themselves. However, low social support has been associated with increased levels of somatic complaints among elderly in seven European countries,⁶⁸ whereas another study with younger participants revealed no relation between social support and somatization.⁶⁹ Studies on abused women reported that social support has moderating effects upon or reduces the risk of somatic complaints and poor physical

health.⁶⁷ The findings in the present study are inconsistent with these previous studies, and it is possible that the complaints of the women in our study may have led to support, which in turn led to more frequent complaints and perhaps their exacerbation. However, more research into the relation of social support and somatization is warranted, particularly in the African context.

Consistent with findings in other studies, not having an education was positively associated with symptoms of depression, which highlights the links among lower educational attainment, IPV, and higher levels of depressive symptoms.^{70,71} Without an education, abused women are economically dependent on their partners, often face repeated or long periods of abuse, and are either uninformed about social services or too scared to access these services. The inherent lack of social support further worsens their social isolation and ultimately leads to depression.

Being married/cohabitant was negatively associated with symptoms of depression, which is in accord with previous data, indicating that married/cohabitant people in comparison with divorced, separated or widowed people show less numerous, intense, and persistent depressive symptoms.⁷² Explanations for this phenomenon include, eg, that married/cohabitant persons may have more social resources and support, which could serve as protective factors against depression.⁷²

Age was associated with symptoms of somatization. In our sample, younger adult women were more likely to exhibit symptoms of somatization, which might be the result of a combination of factors, including economic dependence on their intimate partners, the main effects of IPV, possible experience of IPV in childhood, as well as biological factors related to sex and puberty, which may intensify physical sensations of symptoms somatization. Our findings are consistent with prior studies that indicated a close link between younger age and somatization.⁷³

Finally, the strong association observed between comorbidity and symptoms of depression, anxiety, and somatization in this study is consistent with findings from previous epidemiological studies in the general population,^{74–77} and may only mirror an already well known comorbidity among these conditions, which occur frequently among women.⁷⁸

Limitations

This study has several limitations. First, firm temporal relations and causal links cannot be established with a cross-sectional study and therefore require another type of

study design. Second, the sample of women was limited to women who sought legal aid at the Forensic Services unit of Maputo Central Hospital, Maputo City, and they all had prior experiences of IPV. Conclusions cannot therefore be generalized to the general population since the sample may not be representative of women in the rest of the country, and their IPV experiences could either differ or be similar to those of women in general (though some of our results were consistent with those of other studies). Third, data for this study relied on women's self-report and subjective assessment of their mental health. Hence, the accuracy of data could not be objectively verified. However, the SCL-90-R is sensitive to depressive disorders.⁴⁶ Fourth, in-depth assessment of certain variables (eg, how much alcohol was consumed in a typical drinking day) could not be performed. Fifth, all instruments were translated into Portuguese, back-translated, and culturally adapted, which may result in human error. However, some of our findings are congruent with other data, and the Cronbach's α were generally high. Sixth, CTS2 covers a limited number of psychological abuse dimensions. However, despite its limitations, CTS2 is one of the most widely used tools for identifying IPV and indeed captures central aspects of IPV in a reliable and valid manner.⁴⁷ Seventh, many of the women were both victims and perpetrators of IPV, which could have influenced our results. Thus, caution should be used in interpreting the results although they are in some areas congruent with observations from other studies. In spite of these limitations, this study may have provided new insights into the relation between women's sustained and inflicted IPV and mental health, at least in the context of Mozambique.

Conclusion

A large number of our women who sustained or inflicted the various types of IPV reported relatively high levels of symptoms of depression, anxiety, and somatization. Moreover, 25.9% and 25.5% of women as victims and perpetrators, respectively, showed psychopathological levels across these mental dimensions within or above those reported by psychiatric out patients. Important factors in explaining mental health problems included controlling behaviors, mental comorbidity, social support, not having any education, age, smoking, abuse as a child, and sleep difficulties. The findings contribute to the literature on the relation between women's IPV victimization and/or perpetration and mental health within the context of Mozambique. More research into the relation between

women's IPV victimization and/or perpetration and mental health, and between controlling behaviors and mental health is warranted.

Disclosure

The authors report no conflicts of interest in this work.

References

- Ellsberg M, Jansen HA, Heise L, et al. Intimate partner violence and women's physical and mental health in the WHO multi-country study on women's health and domestic violence: an observational study. *Lancet*. 2008;371(9619):1165–1172.
- Campbell J. Health consequences of intimate partner violence. *Lancet*. 2002;359:1331–1336.
- Devries K, Watts C, Yoshihama M, et al. Violence against women is strongly associated with suicide attempts: evidence from the WHO multi-country study on women's health and domestic violence against women. *Soc Sci Med*. 2011;73(1):79–86.
- Borwankar R, Diallo R, Sommerfelt AE. *Gender-Based Violence in Sub-Saharan Africa: A Review of Demographic and Health Survey Findings and Their Use in National Planning*. Washington, DC: USAID; 2008.
- Vives-Cases C, Ruiz-Cantero MT, Escribà-Agüir V, et al. The effect of intimate partner forms violence and other of violence against women on health. *J Public Health*. 2011;33(1):15–21.
- Mechanic MB, Weaver TL, Resick PA. Mental health consequences of intimate partner abuse: A multidimensional assessment of four different forms of abuse. *Violence Against Women*. 2008;14(6):634–654.
- Houry D, Kaslow NJ, Thompson MP. Depressive symptoms in women experiencing intimate partner violence. *J Interpers Violence*. 2005;20:1467–1477.
- Pico-Alfonso MA, Garcia-Linares MI, Celda-Navarro N, et al. The impact of physical, psychological, and sexual intimate partner violence on women's mental health: depressive symptoms, posttraumatic stress disorder, state anxiety, and suicide. *J Womens Health (Larchmt)*. 2006;15:599–611.
- Romito P, Turan JM, de Marchi M. The impact of current and past interpersonal violence on women's mental health. *Soc Sci Med*. 2005;60:1717–1727.
- Ansara DL, Hindin MJ. Psychosocial consequences of intimate partner violence for women and men in Canada. *J Interpers Violence*. 2011;26(8):1628–1645.
- Coker AL, Davis KE, Arias J, et al. Physical and mental health effects of intimate partner violence for men and women. *Am J Prev Med*. 2002;23:260–268.
- Deyessa N, Berhane Y, Alem A, et al. Intimate partner violence and depression among women in rural Ethiopia: a cross-sectional study. *Clin Pract Epidemiol Ment Health*. 2009;5:8.
- Tadegge AD. The mental health consequences of intimate partner violence against women in Agaro Town, southwest Ethiopia. *Trop Doct*. 2008;38:228–229.
- Zlotnick C, Johnson DM, Kohon R. Intimate partner violence and long-term psychosocial functioning in national sample of American women. *J Interpers Violence*. 2006;21:262–275.
- Pico-Alfonso MA, Garcia-Linares N, Celda-Navarro C, Hebert J, Martinez M. Changes in cortisol and dehydroepiandrosterone in women victims of physical and psychological intimate partner violence. *Biol Psychiatry*. 2004;56:233–240.
- Bonomi AE, Anderson ML, Rivara FP, Thompson RS. Health outcomes in women with physical and sexual intimate partner violence exposure. *J Womens Health (Larchmt)*. 2007;16(7):987–997.
- Hegarty K, Gunn J, Chondros P, Small R. Association between depression and abuse partner of women attending general practice: descriptive, cross-sectional survey. *Br Med J*. 2004;328:621–624.
- Próspero M, Kim M. Mutual partner violence: mental health symptoms among female and male victims in four racial/ethnic groups. *J Interpers Violence*. 2009;24:2039–2056.
- Ruiz-Perez I, Plazaola-Castano J. Intimate partner violence and mental health consequences in women attending family practice in Spain. *Psychosom Med*. 2005;67:791–797.
- Taft CT, Murphy CM, King LA, Dedejn JM, Musser PH. Post-traumatic stress disorder symptomatology among partners of men in treatment for relationship abuse. *J Abnorm Psychol*. 2005;114:259–268.
- Follingstad DR. Rethinking current approaches to psychological abuse: conceptual and methodological issues. *Aggress Violent Behav*. 2007;12:439–458.
- Jordan CE, Campbell R, Follingstad D. Violence and women's mental health: the impact of physical, sexual, and psychological aggression. *Annu Rev Clin Psychol*. 2010;6:607–628.
- Gelaye B, Arnold D, Williams MA, Goshu M, Berhane Y. Depressive symptoms among female college students experiencing gender-based violence in Awassa, Ethiopia. *J Interpers Violence*. 2009;24:464–481.
- Jankey O, Próspero M, Fawson P. Mutually violent attitudes: effects on intimate partner violence and mental health symptoms among couples in Botswana, Africa. *J Aggress Confl Peace Res*. 2011;3:4–11.
- Kaminer D, Grimsrud A, Myer L, Stein D, Williams D. Risk for posttraumatic stress disorder associated with different forms of interpersonal violence in South Africa. *Soc Sci Med*. 2008;67:1589–1595.
- Próspero M, Dwumah P, Ofori-Dua K. Violent attitudes and mental health symptoms among mutually violent Ghanaian couples. *J Aggress Confl Peace Res*. 2009;1:16–23.
- Williams SL, Williams DR, Stein DJ, et al. Multiple traumatic events and psychological distress: the South Africa stress and health study. *J Trauma Stress*. 2007;20:845–855.
- Lipsky S, Caetano R, Field CA, Bazargan S. Violence-related injury and intimate partner violence in an urban emergency department. *J Trauma*. 2004;57:352–359.
- Kernsmith P. Exerting power or striking back: A gendered comparison of motivations for domestic violence perpetration. *Violence Vict*. 2005;20:173–185.
- Houry D, Rhodes KV, Kemball RS, et al. Differences in female and male victims and perpetrators of partner violence with respect to WEB scores. *J Interpers Violence*. 2008;23:1041–1055.
- Archer J. Sex differences in physically aggressive acts between heterosexual partners: a meta-analytic review. *Psychol Bull*. 2000;126:651–680.
- Hines DA. Predictors of sexual coercion against women and men: a multilevel, multinational study of university students. *Arch Sex Behav*. 2007;36:403–422.
- Koenig MA, Lutalo T, Zhao F, Nalugoda F, et al. Domestic violence in rural Uganda: evidence from a community-based study. *Bull World Health Organ*. 2003;81:53–60.
- Laroche D. *Aspects of the Context and Consequences of Domestic Violence: Situational Couple Violence and Intimate Terrorism in Canada in 1999*. Quebec City, Canada: Quebec Institute de la Statistique du Quebec; 2005.
- Carney M, Buttell F. Exploring the relevance of attachment theory as dependent variable in treatment of women mandated into treatment for domestic violence offenses. *J Offender Rehabil*. 2006;41:33–62.
- Fergusson DM, Horwood LJ, Ridder EM. Partner violence and mental health outcomes in a New Zealand birth cohort. *J Marriage Fam*. 2005;67:1103–1119.
- Henning K, Renauber B, Holdford R. Victim or offender? Heterogeneity among women arrested for intimate partner violence. *J Fam Violenc*. 2006;21:351–368.
- Orcutt H, Garcia M, Pickett S. Female-perpetrated intimate partner violence and romantic attachment style in a college student sample. *Violence Vict*. 2005;20:287–302.

39. Stuart GL, Moore TM, Gordon KC, Ramsey SE, Kahler CW. Psychopathology in women arrested for domestic violence. *J Interpers Violence*. 2006;21:376–389.
40. Antai D. Controlling behavior, power relations within intimate relationships and intimate partner physical and sexual violence against women in Nigeria. *BMC Public Health*. 2011;11:511.
41. Krantz G, Nguyen DV. The role of controlling behaviour in intimate partner violence and its health effects: a population based study from rural Vietnam. *BMC Public Health*. 2009;9:143.
42. Capaldi DM, Kim HK, Shortt JW. Observed initiation and reciprocity for physical aggression in young at-risk couples. *J Fam Violence*. 2007;22:101–111.
43. Graham-Kevan N, Archer J. Physical aggression and control in heterosexual relationships: the effect of sampling procedure. *Violence Vict*. 2003;18:181–198.
44. Graham-Kevan N, Archer J. Investigating three explanations of women's relationship aggression. *Psychol Women Quart*. 2005;29:270–277.
45. Próspero M. Sex-symmetric effects of coercive behaviors on mental health? *J Interpers Violence*. 2009; 24:128–146.
46. Derogatis LR. *SCL-90-R Administration, Scoring, and Procedures Manual for the Revision Version*. Baltimore, MD: 1994.
47. Straus MA, Hamby SL, Boney-McCoy S, Sugarman DB. The revised conflict tactics scales (CTS2): development and preliminary psychometric data. *J Fam Issues*. 1996;17:283–316.
48. Undén AL, Orth-Gomér K. Development of social support instrument for use in population surveys. *Soc Sci Med*. 1989;29:387–392.
49. Zacarias AE, Macassa G, Soares JJ. Women as perpetrators of IPV: the experience of Mozambique. *J Aggress Confl Peace Res*. 2012;4:5–27.
50. Dunkle KL, Jewkes RK, Brown HC, et al. Prevalence and partners of gender-based violence and revictimization among women attending antenatal clinics in Soweto, South Africa. *Am J Epidemiol*. 2004;160:230–239.
51. Jewkes R, Levi J, Penn-Kekana L. Risk factors for domestic violence: findings from a South African cross-sectional study. *Soc Sci Med*. 2002;55:1603–1617.
52. Avdibegovic E, Sinanovic O. Consequences of domestic violence on women's mental health in Bosnia and Herzegovina. *Croat Med J*. 2006;47:730–741.
53. Abel EM. Comparing the social service utilization, exposure to violence, and trauma symptomatology of domestic violence female "victims" and female "batterers". *J Fam Violence*. 2001;16:401–420.
54. Goldenson J, Geffner R, Foster S, Clipson C. Female domestic violence offenders: their attachment security, trauma symptoms, and personality organization. *Violence Vict*. 2007;22:530–543.
55. Próspero M. The effect of coercion on aggression and mental health among reciprocally violent couples. *J Fam Violence*. 2008;23:195–202.
56. Carbone-López K, Kruttschnitt C, MacMillan R. Patterns of intimate partner violence and their associations with physical health, psychological distress, and substance use. *Public Health Rep*. 2006;121:382–392.
57. Nicolaidis C, Curry M, McFarland B, Gerrity M. Violence, mental health, and physical symptoms in an academic internal medicine practice. *J Gen Intern Med*. 2004;19:819–827.
58. Waldinger RJ, Schulz MS, Barsky AJ, Ahern DK. Mapping the road from childhood trauma to adult somatization: the role of attachment. *Psychosom Med*. 2006;68:129–135.
59. Kim K, Uchiyama M, Liu X, et al. Somatic and psychological complaints and their correlates with insomnia in the Japanese general population. *Psychosom Med*. 2001;63:441–446.
60. Üstün TB, Privett M, Lecrubier Y, et al. Form, frequency and burden of sleep problems in general health care: a report from the WHO collaborative study on psychological problems in general health care. *Eur Psychiatry*. 1996;11:5–10.
61. Riemann D. Insomnia and comorbid psychiatric disorders. *Sleep Med*. 2007;8(4):S15–S20.
62. Feldner MT, Babson KA, Zvolensky MJ. Smoking, traumatic event exposure, and post-traumatic stress: a critical review of the empirical literature. *Clin Psychol Rev*. 2007;27:14–45.
63. Johnson JG, Choen P, Pine DS, Klein DF, Kasen S, Brook JS. Association between cigarette smoking and anxiety disorders during adolescence and early adulthood. *JAMA*. 2000;284:2348–2351.
64. Mykletun A, Overland S, Aaro LE, Liabo H-M, Stewart R. Smoking in relation to anxiety and depression: evidence from a large population survey: the HUNT study. *Eur Psychiatry*. 2008;23:77–84.
65. Almeida J, Subramanian SV, Kawachi I, Molnar BE. Is blood thicker than water? Social support, depression and modifying role of ethnicity/nativity status. *J Epidemiol*. 2011;65:51–56.
66. Cacioppo JT, Hawkley LC, Crawford LE, et al. Loneliness and health: potential mechanisms. *Psychosom Med*. 2002;64:407–417.
67. Coker AL, Smith PH, Thompson MP, et al. Social support protects against the negative effects of partner violence on mental health. *J Womens Health Gen Gen Med*. 2002;11(5):465–476.
68. Soares JFF, Torres-Gonzales HBF, Ioannidi-Kapoulou E, et al. *Abuse and Health in Europe*. Kaunas, Lithuania: Lithuanian University of Health Sciences Press; 2010.
69. Ladwig KH, Marten-Mittag B, Erazo N, Gündel H. Identifying somatization disorder in a population-based health examination survey: psychosocial burden and gender differences. *Psychosomatics*. 2001;42:511–518.
70. Winter LB, Steer RA, Jones-Hicks L, Beck AT. Screening for major depression disorders in adolescent medical outpatients with the Beck Depression Inventory for Primary Care. *J Adolesc Health*. 1999;24:389–394.
71. Rodriguez MA, Heilemann MV, Fielder E, Ang A, Nevarez F, Mangione CM. Intimate partner violence, depression, and PTSD among pregnant Latina women. *Ann Fam Med*. 2008;6:44–52.
72. Whisman MA, Wistock LM, Tolejko N. Marriage and depression. In: Keyes CLM, Goodman SH, editors. *Women and Depression*. Cambridge, MA: Cambridge University Press; 2006:219–240.
73. Spitzer C, Barnow S, Gau K, Freyberger HJ, Grabe HJ. Childhood maltreatment in patients with somatization disorder. *Aust NZ J Psychiatry*. 2008;42:335–341.
74. Jacobi F, Wittchen H-U, Höltling C, et al. Prevalence, co-morbidity and correlates of mental disorders in general population: results from the German health interview and examination survey (GHS). *Psychol Med*. 2004;34:597–611.
75. Kessler RC, Angermeyer M, Anthony JC, et al. Lifetime prevalence and age-of-onset distributions of mental disorders in the World Health Organization's World Mental Health Survey Initiative. *World Psychiatry*. 2007;6:168–176.
76. Seedat S, Scott KM, Angermeyer MC, et al. Cross-national associations between gender and mental health disorders in the World Health Organization World Mental Health Surveys. *Arch Gen Psychiatry*. 2009;66:785–795.
77. Williams DR, Herman A, Stein DJ, et al. Twelve-month mental disorders in South Africa: prevalence, service use and demographic correlates in population-based South African stress and health study. *Psychol Med*. 2008;38:211–220.
78. Hamilton JA, Russo NF. Women and depression: research, theory, and social policy. In: Keyes CLM, Goodman SH, editors. *Women and Depression*. Cambridge, MA: Cambridge University Press; 2006:479–522.

International Journal of Women's Health

Dovepress

Publish your work in this journal

The International Journal of Women's Health is an international, peer-reviewed open-access journal publishing original research, reports, reviews and commentaries on all aspects of women's healthcare including gynecology, obstetrics, and breast cancer. Subject areas include: Chronic conditions (migraine headaches, arthritis, osteoporosis);

Endocrine and autoimmune syndromes; Sexual and reproductive health; Psychological and psychosocial conditions. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/international-journal-of-womens-health-journal>