

Preinspiratory calcium rise in putative pre-Bötzinger complex astrocytes

Yasumasa Okada^{1,2}, Takuya Sasaki³, Yoshitaka Oku⁴, Naoya Takahashi³, Megumi Seki³, Sakiko Ujita³, Kenji F. Tanaka⁵, Norio Matsuki³ and Yuji Ikegaya³

¹Division of Internal Medicine and Laboratory of Electrophysiology, Murayama Medical Center, Musashimurayama, Tokyo 208-0011, Japan

²Division of Pulmonary Medicine, Department of Internal Medicine, School of Medicine, Keio University, Tokyo 160-8582, Japan

³Laboratory of Chemical Pharmacology, Graduate School of Pharmaceutical Sciences, The University of Tokyo, Tokyo 113-0033, Japan

⁴Department of Physiology, Hyogo College of Medicine, Hyogo 663-8501, Japan

⁵Department of Neuropsychiatry, School of Medicine, Keio University, Tokyo 160-8582, Japan

Key points

- Autonomic respiratory rhythm is essential to maintain lives and is generated in the lower brainstem. The ventrolateral medullary region, called the pre-Bötzinger complex (preBötC), is the kernel for respiratory rhythm generation. Despite previous extensive studies focusing on neurons, the mechanism of how respiratory rhythm is generated has not been fully understood.
- Here we show that non-neuronal glial cells (a subset of putative astrocytes) in the preBötC are periodically activated preceding inspiratory neuronal activity, periodic activity of putative astrocytes persists during blockade of neuronal activity, and stimulation of astrocytes in the preBötC induces inspiratory neuronal firings.
- These findings together with the previous report that blockade of astrocytic metabolism abolishes inspiratory neural output suggest that astrocytes are functionally involved in respiratory rhythm generation.
- These results will help us better understand how respiratory rhythm is generated and how respiratory output is disturbed in various pathological conditions.

Abstract The neural inspiratory activity originates from a ventrolateral medullary region called the pre-Bötzinger complex (preBötC), yet the mechanism underlying respiratory rhythmogenesis is not completely understood. Recently, the role of not only neurons but astrocytes in the central respiratory control has attracted considerable attention. Here we report our discovery that an intracellular calcium rise in a subset of putative astrocytes precedes inspiratory neuronal firing in rhythmically active slices. Functional calcium imaging from hundreds of preBötC cells revealed that a subset of putative astrocytes exhibited rhythmic calcium elevations preceding inspiratory neuronal activity with a time lag of approximately 2 s. These preinspiratory putative astrocytes maintained their rhythmic activities even during the blockade of neuronal activity with tetrodotoxin, whereas the rhythm frequency was lowered and the intercellular phases of these rhythms were decoupled. In addition, optogenetic stimulation of preBötC putative astrocytes induced firing of inspiratory neurons. These findings raise the possibility that astrocytes in the preBötC are actively involved in respiratory rhythm generation in rhythmically active slices.

(Resubmitted 29 February 2012; accepted after revision 6 July 2012; first published online 9 July 2012)

Corresponding author Y. Okada: Division of Internal Medicine and Laboratory of Electrophysiology, Murayama Medical Center, 2-37-1 Gakuen, Musashimurayama, Tokyo 208-0011, Japan. Email: yasumasaokada@1979.jukuin.keio.ac.jp

Abbreviations aCSF, artificial cerebrospinal fluid; ChR2, channelrhodopsin-2; CV, coefficient of variation; F , fluorescence intensity; I_{CAN} , non-specific cationic current; I_{NaP} , persistent sodium current; MLC, megalencephalic leukoencephalopathy with subcortical cysts; OGB1, Oregon Green 488 BAPTA-1 AM; PC, principal component; preBötC, pre-Bötzinger complex; ROI, region of interest; tTA, tetracycline transactivator; YFP, yellow fluorescent protein.

Introduction

Medullary slice preparations made of neonatal rodents that contain the pre-Bötzinger complex (preBötC), termed rhythmically active slices (Del Negro *et al.* 2011), exhibit spontaneous inspiratory-related rhythmic activity *in vitro* (Smith *et al.* 1991) and have been widely used to investigate the mechanism of respiratory rhythm generation. Although various hypotheses have been proposed to explain the mechanism of rhythmogenesis in this region, many of which are based on single neuron kinetics and neuronal network dynamics (Koshiya & Smith, 1999; Feldman & Del Negro, 2006; Pace *et al.* 2007; Koizumi & Smith, 2008; Rubin *et al.* 2009; Morgado-Valle *et al.* 2010; Del Negro *et al.* 2011), no consensus exists among researchers in this area. Although inhibitory synaptic transmission is necessary in network-driven rhythmogenesis in mature animal preparations (Smith *et al.* 2007), inspiratory-related activity persists even after inhibitory synaptic transmission is blocked in rhythmically active slices (Shao & Feldman, 1997). It is also reported that rhythmogenesis requires subthreshold currents, such as persistent sodium current (I_{NaP}) or non-specific cationic current (I_{CAN}) (Del Negro *et al.* 2005). However, even if both I_{NaP} and I_{CAN} were abolished, the rhythm could be maintained by boosting neural excitability, suggesting that pacemaker neurons are not essential for rhythmogenesis (Del Negro *et al.* 2005). These experimental results suggest the involvement of a currently unknown regulatory factor.

Recent progresses of glial cell physiology has revealed that astrocytes play active roles in various brain functions by driving neuronal synchronization (Parri *et al.* 2001; Angulo *et al.* 2004; Fellin *et al.* 2004; Tian *et al.* 2005). Hülsmann *et al.* (2000) reported that blockade of astrocytic metabolism abolishes inspiratory-related neural output in rhythmically active slices, suggesting that astrocytic metabolic support is necessary for the maintenance of respiratory rhythm. Also, it has been indicated that astrocytes play chemosensory, modulatory and regulatory roles in the respiratory network in the medulla oblongata (Neusch *et al.* 2006; Szoke *et al.* 2006; Ballanyi *et al.* 2010; Erlichman *et al.* 2010; Gourine *et al.* 2010; Huxtable *et al.* 2010; Wenker *et al.* 2010; Gourine & Kasparov, 2011; Mulkey & Wenker, 2011). Härtel *et al.* (2009) showed that astrocytes in the ventral respiratory group are activated by neuromodulators that also stimulate respiratory neurons, suggesting that thus modulated astrocytes influence respiratory neurons in the ventral respiratory group. In humans, abnormalities of

brainstem astrocytes correlate with disturbances in central respiratory control, e.g. abnormal proliferation of brainstem astrocytes with central neurogenic hyperventilation (Rodriguez *et al.* 1982) and immaturity of brainstem astrocytes with sudden infant death syndrome (Zhang *et al.* 1992).

These reports led us to further investigate the role of astrocytes in respiratory control, and we conducted large-scale simultaneous analyses of inspiratory-related activities of putative neurons and astrocytes in rhythmically active slices. Here we show that a subset of putative astrocytes in the preBötC is activated preceding inspiratory neuronal firing.

Methods

Animal experiment ethics

All experiments were performed with the approval of the Animal Experiment Ethics Committee at the University of Tokyo (approval numbers: 19-35 and 19-43) and were in accordance with the University of Tokyo guidelines for the care and use of laboratory animals.

Preparations

Rhythmically active slices were prepared from rats for calcium imaging ($n = 16$) or from mice for optogenetic experiments ($n = 30$) between postnatal days 1 and 3. Briefly, animals were deeply anaesthetized with diethyl ether, and the brainstem was dissected and mounted with its rostral side down on an agar block, and transferred to a vibratome (VT1200S, Leica, Heidelberg, Germany). The brainstem was serially sectioned until the most caudal portion of the inferior olive, which approximately corresponded to the third or fourth hypoglossal nerve rootlet. Then a transverse slice containing the preBötC was prepared with a thickness of 550–650 μm . The rostral surface of the slice should coincide with the rostral end of the preBötC or be within approximately 50 μm rostral to the preBötC (see Fig. 1A in Ruangkittisakul *et al.* (2008)). The slice was placed in a submerged-type recording chamber with its rostral side up, and continuously superfused with oxygenated (95% O_2 and 5% CO_2) artificial cerebrospinal fluid (aCSF) containing (in mM): 118 NaCl, 3 KCl, 1.2 CaCl_2 , 1 MgCl_2 , 1 NaH_2PO_4 , 25 NaHCO_3 and 30 D-glucose (~ 330 mosmol l^{-1} , pH 7.4). The temperature was maintained at 28°C. The concentration of potassium

ions was then elevated to 8 mM to activate and maintain rhythmic activity (Smith *et al.* 1991).

Calcium imaging

A glass pipette (1–3 M Ω) for dye loading was filled with aCSF containing of 200 μ M Oregon Green 488 BAPTA-1 AM (OGB1, Invitrogen, Carlsbad, CA, USA), 2% Pluronic F-127 (Invitrogen) and 20% DMSO. The tip of the pipette was inserted into the centre of the preBötC region of rhythmically active slices. The point of pipette insertion was carefully determined making reference to Fig. 2C of Ruangkittisakul *et al.* (2009). Then, the dye solution was pressure-ejected over 3–5 min by manually controlling a 10 ml syringe press (30–70 kPa), following the calcium dye injection technique originally developed for the use *in vivo* (Stosiek *et al.* 2003) and later applied to the preBötC *in vitro* by the group of Ballanyi (Ruangkittisakul *et al.* 2006, 2008). The fluorophore was excited at 488 nm using a laser diode (2–8 mW, HPU50101, FITELE, Tokyo, Japan) and was recorded through a 507 nm long-pass emission filter. Images (512 \times 512 pixels = 512 \times 512 μ m, 14-bit intensity) were captured at 5–50 frames s⁻¹ using a Nipkow-disk confocal scanner unit (CSU-X1; Yokogawa Electric, Tokyo, Japan), a back-illuminated cooled CCD camera (iXon DU897; Andor, Belfast, Northern Ireland, UK), an upright microscope (ECLIPSE FN1; Nikon, Tokyo, Japan) and a water-immersion objective lens (16 \times , 0.8 NA or 40 \times , 0.8 NA, Nikon).

Analysis of calcium dynamics

First, a region covering the whole preBötC was set in order to show average calcium rises in the entire preBötC. Second, a small region of interest (ROI) was carefully placed on (within) each recorded cell body, and the fluorescence intensity (F) was measured. In some cases, non-cellular structures were labelled with calcium dyes, and thus we carefully avoided placing ROIs on these structures based on the following two features. First, because the somata of cells are separated at a cell-to-cell distance at least 20 μ m, the ROIs were not placed within 20 μ m from the nearest ROI. Second, each ROI was set to be 50–75 μ m² in size to avoid placing ROIs on the small non-cellular structures. Some astrocytes contact vessels through their endfeet, and these endfeet were easily identified by eye and excluded from the analysis. In each ROI, the fluorescence intensity was spatially averaged. The change in fluorescence ($\Delta F/F$) in a ROI was calculated as $(F_t - F_0)/F_0$, where F_t was the fluorescence intensity at a focused time point and F_0 was the average baseline fluorescence intensity across the 10 s periods before and after the focused time point. No image

compensation was made to cancel background inspiratory activity.

Functional classification of neurons and astrocytes

In calcium imaging experiments, recorded cells were often classified into neurons and astrocytes on the basis of distinct calcium rise patterns, i.e. faster and slower calcium rises in neurons and astrocytes, respectively (e.g. Pasti *et al.* 1997; Fellin *et al.* 2004, 2006; Bardoni *et al.* 2010). In the present study, however, we intended to classify the recorded cells into putative neurons and putative astrocytes more reliably, and conducted principal component analysis of the waveforms derived from the cellular calcium dynamics. The $\Delta F/F$ waveform was quantified with the following multivariate measures: the rise time constant, maximum rise amplitude and decay time constant of fluorescent calcium signals, based on the universally observed characteristics that the astrocytic calcium changes exhibit slower kinetics and higher amplitude rises compared with those of neurons (Pasti *et al.* 1997; Svoboda *et al.* 1997; Garaschuk *et al.* 2000; Parri *et al.* 2001; Stosiek *et al.* 2003; Fellin *et al.* 2004, 2006; Hirase *et al.* 2004; Nimmerjahn *et al.* 2004; Sasaki *et al.* 2007, 2011a; Bardoni *et al.* 2010; Takahashi *et al.* 2010). The dataset was decomposed into a lower dimensional space of the first two components of the principal component analysis, a linear data-reduction technique (Sasaki *et al.* 2008). The first two principal components were plotted in a figure. The calcium dynamics were automatically classified using a linear support-vector machine. The support-vector machine is one of the most widely used supervised learning methods for classification, initially conceived by Cortes & Vapnik (1995). Further, cell sizes and an identity factor were incorporated. The identity factor scores +1 or -1, according to subjective judgement of neurons or astrocytes based on their calcium waveforms and soma sizes (assuming soma diameter <10 μ m as astrocyte and soma diameter >10 μ m as neuron) (Ruangkittisakul *et al.* 2009; Huxtable *et al.* 2010) were used for *post hoc* confirmation of the cell types after automatic classification. Cells were thus classified into putative neurons and putative astrocytes, and this classification was validated in a selected sample by S100 β immunoreactivity and by their responses to tetrodotoxin (see Results).

Classification of astrocyte types

Based on the relative timings of astrocytic activity to inspiratory neuronal activity, recorded putative astrocytes were classified into four types: preinspiratory cells (type 1), inspiratory cells (type 2), non-respiratory cells (type 3) and silent cells (type 4). Preinspiratory cells were defined

as cells that were activated 0.5–2 s earlier than inspiratory neuronal activity. Inspiratory cells were defined as cells that were activated corresponding with inspiratory timings. Non-inspiratory cells were defined as cells that were activated irrespective of inspiratory timings. Silent cells were defined as cells that were not activated at least during our recording period of 5 min.

Electrophysiology

All electrophysiological recordings were carried out with a MultiClamp 700B amplifier and a Digidata 1440A digitizer controlled by pCLAMP 10 software (Molecular Devices, Union City, CA, USA). Spontaneous multi-unit activity was recorded from the preBötC using borosilicate glass pipettes ($\sim 1\text{ M}\Omega$) and integrated along a time window of 300 ms (Funke *et al.* 2007). This integrated signal of spontaneous multi-unit activity has been widely used as neural output of inspiratory activity (Funke *et al.* 2007; Ruangkittisakul *et al.* 2009; Schnell *et al.* 2011). The onset of integrated signal faithfully coincided with that of average fluorescence signal recorded from the entire preBötC region, and thus the average fluorescence signal was used as an alternative of electrophysiologically recorded inspiratory neural output. For cell-attached recordings, borosilicate glass pipettes (4–7 M Ω) were filled with aCSF. For current-clamp recordings, the same pipettes were filled with an internal solution consisting of (in mM): 135 potassium gluconate, 4 KCl, 10 HEPES, 10 phosphocreatine- Na_2 , 0.3 Na_2 -GTP and 4 Mg-ATP (pH 7.2). Signals were low-pass filtered at 1–2 kHz, digitized at 10 kHz and analysed with pCLAMP 10 software.

Optogenetics

Megalencephalic leukoencephalopathy with subcortical cysts (MLC) is a neurological disorder caused by mutations in the MLC1 gene in human. MLC1 and its mouse homologue, Mlc1, are specifically expressed in astrocytes (Schmitt *et al.* 2003). To selectively excite astrocytes, we developed transgenic mice that expressed the light-activated ion channel, channelrhodopsin-2 (ChR2), fused to enhanced yellow fluorescent protein (YFP), under the control of the astrocyte-specific Mlc1 gene promoter. The ChR2(C128S mutant)–YFP fusion molecule, a gift from Karl Deisseroth (Berndt *et al.* 2009), was selectively expressed in astrocytes using the tetracycline transactivator (tTA)–tetO promoter strategy (Gossen *et al.* 1995), whereby two lines of mice were bred, one expressing an astrocyte-specific promoter driving tTA expression and the other expressing the tetO promoter driving the ChR2–YFP transgene. The transgene was induced by binding of tTA to tetO. Mlc1-tTA mice (Tanaka *et al.* 2010) expressing tTA under the control of the

astrocyte specific gene (Mlc1) promoter were bred with the newly developed tetO-ChR2(C128S)–YFP mice (Tanaka *et al.* in press). ChR2 is a cation-selective membrane channel, and by photo-activation it rapidly opens to form a pore permeable to monovalent (Na^+ , K^+ and H^+) and divalent (Ca^{2+}) cations (Figueiredo *et al.* 2011). To show the rapid channel opening in photo-stimulated astrocytes, photo-induced inward currents were recorded in patch-clamped astrocytes. As light activation of ChR2 could not be combined with fluorescent OGB1 imaging due to optical interference, inspiratory-related neural activity was electrophysiologically monitored from the preBötC. Since our experiments with photo-activation of astrocytes aimed to examine whether photo-activation induces neuronal firing and also preliminary experiments indicated that suppression of spontaneous inspiratory-related activity was more advantageous to obtain clearer responses, after we observed spontaneous firings of inspiratory neurons we lowered the superfusate potassium concentration from 8 mM to 4 mM that induced ‘*in vitro* apnoea’ (Smith *et al.* 1991). While a preBötC neuron from a rhythmically active slice was current-clamped, the centre part of the preBötC region, which was determined by referring to Ruangkittisakul *et al.* (2011), was illuminated with a circular spot (diameter $\sim 350\ \mu\text{m}$) of blue (488 nm; 6.6 mW) or green (540 nm; 19.5 mW) light to selectively stimulate and suppress the preBötC astrocytes for 3–4 s, respectively.

Immunostaining

Functional classification of cells recorded by calcium imaging into neurons and astrocytes was validated in a representative rhythmically active slice of rat by *post hoc* immunoreactivity analysis of S100 β , which is an established astrocyte-specific marker protein in rodents (Huxtable *et al.* 2010; Aoyama *et al.* 2011; Sasaki *et al.* 2011a), though not necessarily astrocyte-specific in humans (Steiner *et al.* 2007). Also, ChR2–YFP-positive cells in rhythmically active slices of mouse were immunohistochemically examined to determine whether they were exclusively astrocytes.

A rhythmically active slice was fixed as a whole in 4% paraformaldehyde in 0.1 M phosphate buffer solution overnight at 4°C and permeabilized and blocked with 0.1% Triton X-100 and 5% goat serum at room temperature (Sasaki *et al.* 2011a). It was then incubated overnight with primary mouse monoclonal anti-S100 β antibody (S2532, Sigma-Aldrich, 1:1000) and 1 $\mu\text{g ml}^{-1}$ affinity-purified rabbit antibody to green fluorescent protein (gift from Takeshi Kaneko; Tamamaki *et al.* 2000) and labelled with secondary anti-rabbit IgG Alexa-488 (A11008, Invitrogen, 1:500), anti-mouse IgG Alexa-594 (A11032, Invitrogen, 1:500) and Neuro Trace 435/455

blue fluorescent Nissl (N21479, Invitrogen, 1:50), which specifically stains neuronal somata (Schmued *et al.* 1982), for 5 h at room temperature. The region where calcium imaging was conducted was identified based on the structural characteristics of the slice and the distribution of S100 β -positive cells in the region that was presumed to be at the same depth as calcium imaging was observed under a confocal microscope (MRC-1024, Bio-Rad, CA, USA).

Results

Inspiratory-related activity patterns of astrocytes and neurons

The medullary region into which OGB1 was bolus-injected emitted green fluorescence, and coincided well with the preBötC in rhythmically active slices of rats (Fig. 1A). All tested slices spontaneously exhibited intermittent elevations of total fluorescence intensity in the dye-loaded region, the onset of which faithfully

coincided with that of extracellularly recorded inspiratory neuron bursting, as previously shown (Koshiya & Smith, 1999; Ruangkittisakul *et al.* 2009) (Fig. 1B). These inspiratory-related rhythmic signals were monitored at 5–50 frames s^{-1} at a cellular resolution using a Nipkow-disk confocal microscope. Single videos were 243 ± 14 s in length, and included 161 ± 23 cells (120 ± 17 putative neurons; 41 ± 5 putative astrocytes) ($n = 25$ videos from 14 slices). The recorded cells were classified into putative astrocytes and putative neurons by principal component analysis of the waveforms derived from the cellular calcium dynamics and of the cell sizes (Fig. 1C). This functional cell classification was validated by *post hoc* immunohistochemical staining for the astrocytic marker S100 β (Fig. 1D). Cell-attached patch-clamp recordings from these identified cells confirmed that the calcium elevations in putative neurons, but not in putative astrocytes, were associated with action potentials (9.5 ± 5.8 spikes per neuron in one inspiratory phase), thereby validating our classification methodology (Fig. 1E;




Figure 1. Large-scale multi-cell calcium imaging from the preBötC

A, bolus injection of OGB1 (green) into the preBötC of a rhythmically active slice. B, spontaneous fluctuations in the mean fluorescence of the preBötC (top) correlated with the integrated inspiratory unit activity (bottom). C, Functional classification of neurons and astrocytes by a principal component (PC) analysis of the calcium waveforms. Putative neurons and putative astrocytes are indicated in black and blue, respectively. D, Immunohistochemical classification of neurons and astrocytes (right) based on S100 β immunoreactivity (middle) after OGB1 imaging (left). E, simultaneous calcium imaging ($\Delta F/F$) and cell-attached recording (unit) from a neuron (top) and an astrocyte (bottom). F, locations of 236 putative neurons and 90 putative astrocytes identified in a single video. G, raster plots of the calcium activity of the putative astrocytes and putative neurons that are represented in F. The bottom histogram indicates the percentage of coactivated putative neurons at a given time (bin = 0.2 s). Putative astrocytes and putative neurons are plotted as blue and black dots, respectively.

putative neuron, $n = 8$ cells; putative astrocyte, $n = 8$ cells). Also, any cells classified as putative neurons did not show spontaneous calcium rises in the presence of tetrodotoxin, which further validated our cell classification (data not shown).

The spatial distribution and spatiotemporal activity patterns of 236 putative neurons and 90 putative astrocytes from one experiment are illustrated in Fig. 1F and G, respectively. The inspiratory-related modulation of calcium activity was evident in the putative neuronal populations, with a rhythm frequency of 0.16 ± 0.05 Hz and with $55 \pm 19\%$ of putative neurons participating in a single respiratory cycle ($n = 14$ videos).

Activity of putative astrocytes precedes inspiratory neuronal firing

A fraction of putative astrocytes also exhibited inspiratory-related modulation of their calcium activity in slices of rats (Fig. 1G). Thus, we analysed the temporal profile of the individual putative astrocytic responses. Trace plots of the inspiratory-related activity revealed

that a subset of putative astrocytes were activated early relative to the inspiratory neuronal activity with a time lag of approximately 2 s (Fig. 2A and B and Supplementary video). The location of these preinspiratory putative astrocytes was unlikely to indicate spatial organization (Fig. 2C). In the present study, 10.5% of putative astrocytes were found to be preinspiratory, 3.2% of cells exhibited spontaneous calcium activity irrespective of inspiratory-related activity and 86.3% were either inspiratory or silent during the observation period. We avoided strict separation of inspiratory and silence cells when counting their numbers, because recorded cell activity could reflect a small contamination with background inspiratory fluorescence of neuropiles (Fig. 2D).

Intrinsic rhythmic activity of preinspiratory putative astrocytes

To analyse activity patterns of putative astrocytes without the influence of neuronal firings, we blocked neuronal activity by bath application of $1 \mu\text{M}$ tetrodotoxin, a voltage-sensitive Na^+ channel inhibitor in slices of




Figure 2. Preinspiratory activity of putative astrocytes

A, representative calcium $\Delta F/F$ traces of preinspiratory (type 1), inspiratory (type 2), non-respiratory (type 3) and silent putative astrocytes (type 4). Top: bulk $\Delta F/F$ fluctuations of the preBötC. B, peri-inspiratory $\Delta F/F$ responses of preinspiratory, inspiratory, non-respiratory and silent putative astrocytes. The traces were averaged from 20 successive events by aligning the onset of individual inspiratory events at 0 s. C, representative cell map of putative neurons and four types of putative astrocytes. We avoided strict separation of inspiratory and silence cells, and they are shown together as type 2+4. D, distribution of four types of putative astrocytes ($n = 541$ putative astrocytes from 14 movies).

rats. Preinspiratory and inspiratory putative astrocytes maintained their periodic calcium rises even during the blockade of neuronal activity with tetrodotoxin, whereas the intercellular phases of these rhythmic activities were decoupled (Fig. 3A). Tetrodotoxin also reduced the frequency of calcium rises in preinspiratory, inspiratory and non-respiratory putative astrocytes (Fig. 3B). The activity patterns of the preinspiratory and inspiratory putative astrocytes tended to be more periodic than that of non-respiratory putative astrocytes both before and after application of tetrodotoxin, although their differences were not statistically significant when their rhythm regularities were assessed by the coefficient of variation (CV) in inter-activity intervals (Fig. 3C). The regularity of rhythmic activity of preinspiratory putative astrocytes was similar to that of inspiratory neuronal activity (Fig. 3C). It was interesting to find that tetrodotoxin induced rhythmic calcium activity with unknown mechanisms in a small number of putative astrocytes that were silent before application of tetrodotoxin (bottom trace in Fig. 3A).


Selective photo-stimulation of preBötC astrocytes

To gain an insight into the functional role of preBötC astrocytes, we examined whether astrocytes in the

preBötC were capable of activating inspiratory neurons, by selectively photo-stimulating astrocytes in the centre part of the preBötC in slices of ChR2(C128S)-YFP transgenic mice (Fig. 4A). As ChR2 is rapidly inactivated after cessation of light application, we employed the C128S point mutation of ChR2 (Berndt *et al.* 2009), which alters the photocycle, enabling bistable, step-like control of cell membrane potentials by two different wavelengths. This optical property is effective to mimic the slow kinetics (2–4 s) of astrocytic calcium dynamics. To confirm the specificity of transgene expression in astrocytes, we performed simultaneous Nissl staining that specifically stains neuronal somata and immunohistochemistry for YFP and for an astrocyte-specific marker S100 β (Fig. 4B). In the preBötC, 45% of S100 β -positive cells expressed YFP signals ($n = 278$ cells from 16 slices). On the other hand, there was no overlap between Nissl-positive cells and YFP signals ($n = 895$ cells from 16 slices). To further validate the specificity of transgene expression, whole-cell patch-clamp recordings were performed randomly from astrocytes and neurons in the presence of 1 μM tetrodotoxin, 100 μM Cd²⁺ and 50 μM 6-cyano-7-nitroquinoxaline-2,3-dione (CNQX) to avoid generation of action potentials, Ca²⁺-dependent vesicular releases and non-NMDA receptor-mediated transmission,

Figure 3. Intrinsic rhythmic activities of preinspiratory putative astrocytes

A, representative overall inspiratory $\Delta F/F$ traces (top) and calcium $\Delta F/F$ traces of preinspiratory, inspiratory, non-respiratory and silent putative astrocytes before and after application of tetrodotoxin. Note that the bottom silent trace exhibits activity after application of tetrodotoxin. B, the mean frequency of calcium events in preinspiratory (blue, $n = 20$ cells), inspiratory (red, $n = 175$ cells) and non-respiratory putative astrocytes in the presence (filled bar) or absence (open bar) of tetrodotoxin. C, coefficient of variation (CV) of the time-intervals between two successive calcium events in preinspiratory, inspiratory and non-respiratory putative astrocytes in the presence or absence of tetrodotoxin. The dashed line indicates the CV of the inspiratory-related neuronal activity in the presence of tetrodotoxin, i.e. the irregularity of the inspiratory rhythm. * $P < 0.05$, ** $P < 0.01$, paired t test.



respectively, if any. In astrocytes, 5 out of 9 cells exhibited inward currents in response to brief illumination with blue light, and the currents persisted until flashed with green light (Fig. 4C, top). The average amplitude of the photocurrents was 68 ± 37 pA ($n = 5$ astrocytes). In contrast, no current responses to photo-stimulation were observed in all neurons tested ($n = 25$ cells) (Fig. 4C, bottom). These histological and electrophysiological findings clearly demonstrate that ChR2(C128S)-YFP was expressed exclusively in a subset of astrocytes. Using this transgenic mouse line, we then investigated the effect of selective activation of preBötC astrocytes on excitability of neighbouring inspiratory neurons. Photo-activation of astrocytes elicited single or burst firing of action potentials in 8 of 16 preBötC inspiratory neurons in the ChR2(C128S)-YFP transgenic mice ($n = 16$ mice), but not in the wild-type mice ($n = 9$ cells from 9 mice) (Fig. 4D and E). In the eight responsive neurons, $75 \pm 11\%$ of trials with light stimulation elicited action potentials (Fig. 4F) and the latency from the stimulus onset to the first spike was 5.5 ± 0.9 s (Fig. 4G).

Discussion

In the present study we have shown that a subset of putative astrocytes exhibited calcium elevations preceding inspiratory neuronal activity with a time lag of approximately 2 s. These preinspiratory putative astrocytes maintained their rhythmic activity even during the blockade of neuronal activity with tetrodotoxin, whereas their rhythm frequency was lowered and the intercellular phases of these rhythms were decoupled. Optogenetic stimulation of preBötC astrocytes induced firing of inspiratory neurons.

Recently, a number of studies by calcium imaging in the preBötC have been reported. However, none showed respiratory-related rhythmic calcium activity in astrocytes (e.g. Funke *et al.* 2007; Härtel *et al.* 2009; Ruangkittisakul *et al.* 2009), and the present study is the first to show respiratory-related stable rhythmic calcium activities in putative astrocytes. Therefore, the preinspiratory activity in putative astrocytes is also first reported here. Although it is not clear why respiratory-related rhythmic calcium activities in astrocytes have not previously been reported; it may be partly attributed to the experimental technique




Figure 4. Astrocyte-driven action potentials in inspiratory neurons

A, YFP fluorescence of a rhythmically active slice prepared from a Mlc1-tTA::tetO-ChR2(C128S)-YFP transgenic mouse. **B**, triple labelling of preBötC cells with an antibody against green fluorescent protein (i.e. YFP appearing in green) to indicate ChR2-expressing cells, with fluorescent Nissl to show neuronal somata (blue), and with an antibody against S100 β to identify astrocytes (red). ChR2-YFP positivity was found exclusively in S100 β -positive cells. Thus, ChR2 was expressed only in astrocytes. **C**, switch-on (blue light) and switch-off (green light) of a persistent inward current in a voltage-clamped ChR2-expressing astrocyte (top) and neuron (bottom). Voltages were clamped at -70 mV. The black bar indicates the activated period of preBötC astrocytes. **D**, photo-activation of astrocytes triggers action potentials in an inspiratory neuron from the preBötC in a ChR2(C128S)YFP transgenic (but not in a wild-type) mouse. Black bars indicate the activation periods of preBötC astrocytes. **E**, ratios of neurons that fired in response to photo-activation of astrocytes in wild-type ($n = 9$ neurons) and in ChR2(C128S)YFP transgenic mice ($n = 16$ neurons from 5 animals). **F**, ratios of the trials in which light stimulation elicited action potentials ($n = 8$ responsive neurons). **G**, latency from the light-stimulus onset to the first spike ($n = 8$ neurons).

in previous studies, i.e. calcium indicator dyes were injected into the midline of the slice to selectively label neurons through axons of passage (Koizumi *et al.* 2008; Koshiya & Smith, 1999), imaging areas were rather small (Koizumi *et al.* 2008) or injected dyes preferentially stained neurons. On the contrary, we directly stained the preBötC region and conducted large-scale imaging, and the dye we used, OGB1, appears to preferentially stain astrocytes (Funke *et al.* 2007; Sasaki *et al.* 2011a). Recently, Schnell *et al.* (2011) analysed functional coupling between astrocytes and neurons, detected a rhythmic inwardly directed current that was entrained to neuron population discharges in 10% of preBötC astrocytes by whole-cell patch recording, and attributed the rhythmic current fluctuation to periodic elevations of extracellular potassium that were produced by efflux from rhythmically oscillating neurons and to consequent changes in potassium equilibrium potentials of astrocytes (Brockhaus *et al.* 1993; Okada *et al.* 2005). However, Schnell *et al.* (2011) did not detect periodic calcium rises in astrocytes coupled with neuronal rhythmic activities. Although they observed spontaneous calcium fluctuation in astrocytes, it did not propagate to or from other cells. To explain the discrepancy between their and our observations, further studies are needed.

Optogenetic stimulation is a powerful method to study the function of astrocytes (Figueiredo *et al.* 2011). It has been reported that optogenetic stimulation of chemosensory astrocytes near the ventral surface of the medulla oblongata induces respiratory augmentation (Gourine *et al.* 2010). In the present study we applied an optogenetic stimulation technique, and carefully focused the photo-stimulating region within the preBötC. Thus, chemosensory astrocytes near the ventral surface should not have been excited. Although we could induce inspiratory neuronal firings by optogenetic stimulation of astrocytes in the preBötC, not only preinspiratory but all other types of astrocytes expressing ChR2 should have been excited with this technique. Also, the latency approximately 5 s after photo-stimulation to inspiratory neuronal firings, as well as the latency approximately 2 s after activation of preinspiratory putative astrocytes to inspiratory neuronal firings, appear too long, if assuming a pacemaking role in preinspiratory astrocytes to trigger inspiratory neurons especially in *in vivo* conditions where breathing rates are notably faster than 1 Hz in rodents. Further studies are necessary to examine the response of inspiratory neurons to selective activation of preinspiratory astrocytes and to clarify the role of astrocytes in respiratory control in *in vitro* and *in vivo* conditions.

Even if preinspiratory astrocytes play an active role in initiating inspiratory neuron firings, the notion does

not necessarily contradict any previously proposed neuronal pacemaker or neuronal group pacemaker hypotheses (Koshiya & Smith, 1999; Feldman & Del Negro, 2006; Pace *et al.* 2007; Mironov, 2008; Del Negro *et al.* 2011). Although calcium influxes in the somata of inspiratory neurons have been shown to be a consequence of membrane depolarization and through a voltage-dependent calcium channels (Morgado-Valle *et al.* 2008), those in the dendrites precede somatic inspiratory activity with a time lag of a few hundred milliseconds (Mironov, 2008; Del Negro *et al.* 2011). According to the group pacemaker hypothesis, preBötC excitatory interneurons mutually interact to elevate dendritic calcium, activate I_{CAN} and periodically trigger inspiration (Mironov, 2008; Rubin *et al.* 2009). Our findings raise an alternative possibility that the preinspiratory dendritic calcium elevation and depolarization are evoked, at least partly, by the preceding astrocytic activity (Sasaki *et al.* 2011b). Our observation that blockade of neuronal activity with tetrodotoxin slowed and decoupled the periodic calcium rises of preinspiratory putative astrocytes suggests that the neuronal network facilitates the occurrence of calcium transient of preinspiratory astrocytes and synchronizes them. Inspiratory neurons appear to have an influence on the rhythmic glial activities and thus preinspiratory astrocytes and inspiratory neurons seem to have mutual interactions in the preBötC.

These findings, together with previous reports that inhibition of astrocytic metabolism abolishes inspiratory-related neural output in rhythmically active slices (Hülsmann *et al.* 2000) and decreases respiratory frequency in *in vivo* rat pups (Young *et al.* 2005), raise the possibility that astrocytes are actively involved in respiratory rhythm generation in the preBötC. As astrocyte-driven neural synchronization has been reported in many other brain regions (Parri *et al.* 2001; Angulo *et al.* 2004; Fellin *et al.* 2004; Tian *et al.* 2005), we propose a hypothesis that preinspiratory astrocytes are actively involved in respiratory rhythm generation at least in rhythmically active slices. This issue must be thoroughly investigated in the future, in a precisely 'calibrated' rhythmically active slice that can generate an eupnoea-like inspiratory burst pattern in a more physiological superfusate potassium concentration (Ballanyi & Ruangkittisakul, 2009) and by experiments including long-term calcium imaging and patch-clamp recording from astrocytes that are identified, e.g. genetically (e.g. Härtel *et al.* 2009; Ruangkittisakul *et al.* 2009). Also, pharmacological modulation of the bursting rhythm by glial toxins and agonist/antagonist of possible gliotransmitter receptors must be thoroughly examined to clarify whether preinspiratory calcium rise in astrocytes leads inspiratory neuronal firings.

References

- Angulo MC, Kozlov AS, Charpak S & Audinat E (2004). Glutamate released from glial cells synchronizes neuronal activity in the hippocampus. *J Neurosci* **24**, 6920–6927.
- Aoyama R, Okada Y, Yokota S, Yasui Y, Fukuda K, Shinozaki Y, Yoshida H, Nakamura M, Chiba K, Yasui Y, Kato F & Toyama Y (2011). Spatiotemporal and anatomical analyses of P2X receptor-mediated neuronal and glial processing of sensory signals in the rat dorsal horn. *Pain* **152**, 2085–2097.
- Ballanyi K & Ruangkittisakul A (2009). Structure-function analysis of rhythmic inspiratory pre-Bötzinger complex networks in ‘calibrated’ newborn rat brainstem slices. *Respir Physiol Neurobiol* **168**, 158–178.
- Ballanyi K, Panaitescu B & Ruangkittisakul A (2010). Control of breathing by ‘nerve glue’. *Sci Signal* **3**, pe41.
- Bardoni R, Ghirri A, Zonta M, Betelli C, Vitale G, Ruggieri V, Sandrini M & Carmignoto G (2010). Glutamate-mediated astrocyte-to-neuron signalling in the rat dorsal horn. *J Physiol* **588**, 831–846.
- Berndt A, Yizhar O, Gunaydin LA, Hegemann P & Deisseroth K (2009). Bi-stable neural state switches. *Nat Neurosci* **12**, 229–234.
- Brockhaus J, Ballanyi K, Smith JC & Richter DW (1993). Microenvironment of respiratory neurons in the *in vitro* brainstem–spinal cord of neonatal rats. *J Physiol* **462**, 421–445.
- Cortes C & Vapnik V (1995). Support-vector networks. *Mach Learn* **20**, 273–297.
- Del Negro CA, Hayes JA & Rekling JC (2011). Dendritic calcium activity precedes inspiratory bursts in preBötzinger complex neurons. *J Neurosci* **31**, 1017–1022.
- Del Negro CA, Morgado-Valle C, Hayes JA, Mackay DD, Pace RW, Crowder EA & Feldman JL (2005). Sodium and calcium current-mediated pacemaker neurons and respiratory rhythm generation. *J Neurosci* **25**, 446–453.
- Erlichman JS, Leiter JC & Gourine AV (2010). ATP, glia and central respiratory control. *Respir Physiol Neurobiol* **173**, 305–311.
- Feldman JL & Del Negro CA (2006). Looking for inspiration: new perspectives on respiratory rhythm. *Nat Rev Neurosci* **7**, 232–242.
- Fellin T, Gomez-Gonzalo M, Gobbo S, Carmignoto G & Haydon PG (2006). Astrocytic glutamate is not necessary for the generation of epileptiform neuronal activity in hippocampal slices. *J Neurosci* **26**, 9312–9322.
- Fellin T, Pascual O, Gobbo S, Pozzan T, Haydon PG & Carmignoto G (2004). Neuronal synchrony mediated by astrocytic glutamate through activation of extrasynaptic NMDA receptors. *Neuron* **43**, 729–743.
- Figueiredo M, Lane S, Tang F, Liu BH, Hewinson J, Marina N, Kasymov V, Souslova EA, Chudakov DM, Gourine AV, Teschemacher AG & Kasparov S (2011). Optogenetic experimentation on astrocytes. *Exp Physiol* **96**, 40–50.
- Funke F, Dutschmann M & Müller M (2007). Imaging of respiratory-related population activity with single-cell resolution. *Am J Physiol Cell Physiol* **292**, C508–C516.
- Garaschuk O, Linn J, Eilers J & Konnerth A (2000). Large-scale oscillatory calcium waves in the immature cortex. *Nat Neurosci* **3**, 452–459.
- Gossen M, Freundlieb S, Bender G, Müller G, Hillen W & Bujard H (1995). Transcriptional activation by tetracyclines in mammalian cells. *Science* **268**, 1766–1769.
- Gourine AV & Kasparov S (2011). Astrocytes as brain interoceptors. *Exp Physiol* **96**, 411–416.
- Gourine AV, Kasymov V, Marina N, Tang F, Figueiredo MF, Lane S, Teschemacher AG, Spyer KM, Deisseroth K & Kasparov S (2010). Astrocytes control breathing through pH-dependent release of ATP. *Science* **329**, 571–575.
- Härtel K, Schnell C & Hülsmann S (2009). Astrocytic calcium signals induced by neuromodulators via functional metabotropic receptors in the ventral respiratory group of neonatal mice. *Glia* **57**, 815–827.
- Hirase H, Qian L, Barthó P & Buzsáki G (2004). Calcium dynamics of cortical astrocytic networks *in vivo*. *PLoS Biol* **2**, E96.
- Hülsmann S, Oku Y, Zhang W & Richter DW (2000). Metabolic coupling between glia and neurons is necessary for maintaining respiratory activity in transverse medullary slices of neonatal mouse. *Eur J Neurosci* **12**, 856–862.
- Huxtable AG, Zwicker JD, Alvares TS, Ruangkittisakul A, Fang X, Hahn LB, Posse de Chaves E, Baker GB, Ballanyi K & Funk GD (2010). Glia contribute to the purinergic modulation of inspiratory rhythm-generating networks. *J Neurosci* **30**, 3947–3958.
- Koizumi H & Smith JC (2008). Persistent Na⁺ and K⁺-dominated leak currents contribute to respiratory rhythm generation in the pre-Bötzinger complex *in vitro*. *J Neurosci* **28**, 1773–1785.
- Koizumi H, Wilson CG, Wong S, Yamanishi T, Koshiya N & Smith JC (2008). Functional imaging, spatial reconstruction, and biophysical analysis of a respiratory motor circuit isolated *in vitro*. *J Neurosci* **28**, 2353–2365.
- Koshiya N & Smith JC (1999). Neuronal pacemaker for breathing visualized *in vitro*. *Nature* **400**, 360–363.
- Mironov SL (2008). Metabotropic glutamate receptors activate dendritic calcium waves and TRPM channels which drive rhythmic respiratory patterns in mice. *J Physiol* **586**, 2277–2291.
- Morgado-Valle C, Baca SM & Feldman JL (2010). Glycinergic pacemaker neurons in preBötzinger complex of neonatal mouse. *J Neurosci* **30**, 3634–3639.
- Morgado-Valle C, Beltran-Parral L, DiFranco M, Vergara JL & Feldman JL (2008). Somatic Ca²⁺ transients do not contribute to inspiratory drive in preBötzinger complex neurons. *J Physiol* **586**, 4531–4540.
- Mulkey DK & Wenker IC (2011). Astrocyte chemoreceptors: mechanisms of H⁺ sensing by astrocytes in the retrotrapezoid nucleus and their possible contribution to respiratory drive. *Exp Physiol* **96**, 400–406.
- Neusch C, Papadopoulos N, Müller M, Maletzki I, Winter SM, Hirrlinger J, Handschuh M, Bähr M, Richter DW, Kirchhoff F & Hülsmann S (2006). Lack of the Kir4.1 channel subunit abolishes K⁺ buffering properties of astrocytes in the ventral respiratory group: impact on extracellular K⁺ regulation. *J Neurophysiol* **95**, 1843–1852.
- Nimmerjahn A, Kirchhoff F, Kerr JN & Helmchen F (2004). Sulforhodamine 101 as a specific marker of astroglia in the neocortex *in vivo*. *Nat Methods* **1**, 31–37.

- Okada Y, Kuwana S, Kawai A, Mückenhoff K & Scheid P (2005). Significance of extracellular potassium in central respiratory control studied in the isolated brainstem-spinal cord preparation of the neonatal rat. *Respir Physiol Neurobiol* **146**, 21–32.
- Pace RW, Mackay DD, Feldman JL & Del Negro CA (2007). Inspiratory bursts in the preBötzinger complex depend on a calcium-activated non-specific cation current linked to glutamate receptors in neonatal mice. *J Physiol* **582**, 113–125.
- Parri HR, Gould TM & Crunelli V (2001). Spontaneous astrocytic Ca^{2+} oscillations *in situ* drive NMDAR-mediated neuronal excitation. *Nat Neurosci* **4**, 803–812.
- Pasti L, Volterra A, Pozzan T & Carmignoto G (1997). Intracellular calcium oscillations in astrocytes: a highly plastic, bidirectional form of communication between neurons and astrocytes *in situ*. *J Neurosci* **17**, 7817–7830.
- Rodriguez M, Baele PL, Marsh HM & Okazaki H (1982). Central neurogenic hyperventilation in an awake patient with brainstem astrocytoma. *Ann Neurol* **11**, 625–628.
- Ruangkittisakul A, Okada Y, Oku Y, Koshiya N & Ballanyi K (2009). Fluorescence imaging of active respiratory networks. *Respir Physiol Neurobiol* **168**, 26–38.
- Ruangkittisakul A, Panaitecu B & Ballanyi K (2011). K^+ and Ca^{2+} dependence of inspiratory-related rhythm in novel ‘calibrated’ mouse brainstem slices. *Respir Physiol Neurobiol* **175**, 37–48.
- Ruangkittisakul A, Schwarzacher SW, Secchia L, Ma Y, Bobocea N, Poon BY, Funk GD & Ballanyi K (2008). Generation of eupnea and sighs by a spatiochemically organized inspiratory network. *J Neurosci* **28**, 2447–2458.
- Ruangkittisakul A, Schwarzacher SW, Secchia L, Poon BY, Ma Y, Funk GD & Ballanyi K (2006). High sensitivity to neuromodulator-activated signaling pathways at physiological $[K^+]$ of confocally imaged respiratory center neurons in on-line-calibrated newborn rat brainstem slices. *J Neurosci* **26**, 11870–11880.
- Rubin JE, Hayes JA, Mendenhall JL & Del Negro CA (2009). Calcium-activated nonspecific cation current and synaptic depression promote network-dependent burst oscillations. *Proc Natl Acad Sci U S A* **106**, 2939–2944.
- Sasaki T, Kuga N, Namiki S, Matsuki N & Ikegaya Y (2011a). Locally synchronized astrocytes. *Cereb Cortex* **21**, 1889–1900.
- Sasaki T, Matsuki N & Ikegaya Y (2007). Metastability of active CA3 networks. *J Neurosci* **27**, 517–528.
- Sasaki T, Matsuki N & Ikegaya Y (2011b). Action-potential modulation during axonal conduction. *Science* **331**, 599–601.
- Sasaki T, Takahashi N, Matsuki N & Ikegaya Y (2008). Fast and accurate detection of action potentials from somatic calcium fluctuations. *J Neurophysiol* **100**, 1668–1676.
- Schmitt A, Gofferje V, Weber M, Meyer J, Mössner R & Lesch KP (2003). The brain-specific protein MLC1 implicated in megalencephalic leukoencephalopathy with subcortical cysts is expressed in glial cells in the murine brain. *Glia* **44**, 283–295.
- Schmued LC, Swanson LW & Sawchenko PE (1982). Some fluorescent counterstains for neuroanatomical studies. *J Histochem Cytochem* **30**, 123–128.
- Schnell C, Fresemann J & Hülsmann S (2011). Determinants of functional coupling between astrocytes and respiratory neurons in the pre-Bötzinger complex. *PLoS One* **6**, e26309.
- Shao XM & Feldman JL (1997). Respiratory rhythm generation and synaptic inhibition of expiratory neurons in pre-Bötzinger complex: differential roles of glycinergic and GABAergic neural transmission. *J Neurophysiol* **77**, 1853–1860.
- Smith JC, Abdala AP, Koizumi H, Rybak IA & Paton JF (2007). Spatial and functional architecture of the mammalian brain stem respiratory network: a hierarchy of three oscillatory mechanisms. *J Neurophysiol* **98**, 3370–3387.
- Smith JC, Ellenberger HH, Ballanyi K, Richter DW & Feldman JL (1991). Pre-Bötzinger complex: a brainstem region that may generate respiratory rhythm in mammals. *Science* **254**, 726–729.
- Steiner J, Bernstein HG, Bielau H, Berndt A, Brisch R, Mawrin C, Keilhoff G & Bogerts B (2007). Evidence for a wide extra-astrocytic distribution of S100B in human brain. *BMC Neurosci* **8**, 2.
- Stosiek C, Garaschuk O, Holthoff K & Konnerth A (2003). *In vivo* two-photon calcium imaging of neuronal networks. *Proc Natl Acad Sci U S A* **100**, 7319–7324.
- Svoboda K, Denk W, Kleinfeld D & Tank DW (1997). *In vivo* dendritic calcium dynamics in neocortical pyramidal neurons. *Nature* **385**, 161–165.
- Szoke K, Härtel K, Grass D, Hirrlinger PG, Hirrlinger J & Hülsmann S (2006). Glycine transporter 1 expression in the ventral respiratory group is restricted to protoplasmic astrocytes. *Brain Res* **1119**, 182–189.
- Takahashi N, Sasaki T, Matsumoto W, Matsuki N & Ikegaya Y (2010). Circuit topology for synchronizing neurons in spontaneously active networks. *Proc Natl Acad Sci U S A* **107**, 10244–10249.
- Tamamaki N, Nakamura K, Furuta T, Asamoto K & Kaneko T (2000). Neurons in Golgi-stain-like images revealed by GFP-adenovirus infection *in vivo*. *Neurosci Res* **38**, 231–236.
- Tanaka KF, Ahmari SE, Leonardo ED, Richardson-Jones JW, Budreck EC, Scheiffle P, Sugio S, Inamura N, Ikenaka K & Hen R (2010). Flexible Accelerated STOP Tetracycline Operator-knockin (FAST): a versatile and efficient new gene modulating system. *Biol Psychiatry* **67**, 770–773.
- Tanaka KF, Matsui K, Sasaki T, Sano H, Sugio S, Fan K, Hen R, Nakai J, Yanagawa Y, Hasuwa H, Okabe M, Deisseroth K, Ikenaka K & Yamanaka A. Expanding the repertoire of optogenetically targeted cells with an enhanced gene expression system. *Cell Reports*, in press.
- Tian GF, Azmi H, Takano T, Xu Q, Peng W, Lin J, Oberheim N, Lou N, Wang X, Zielke HR, Kang J & Nedergaard M (2005). An astrocytic basis of epilepsy. *Nat Med* **11**, 973–981.
- Wenker IC, Kréneisz O, Nishiyama A & Mulkey DK (2010). Astrocytes in the retrotrapezoid nucleus sense H^+ by inhibition of a Kir4.1-Kir5.1-like current and may contribute to chemoreception by a purinergic mechanism. *J Neurophysiol* **104**, 3042–3052.
- Young JK, Dreshaj IA, Wilson CG, Martin RJ, Zaidi SI & Haxhiu MA (2005). An astrocyte toxin influences the pattern of breathing and the ventilatory response to hypercapnia in neonatal rats. *Respir Physiol Neurobiol* **147**, 19–30.

Zhang WD, Tsukamoto T, Yamada Y, Itakura Y, Oono T, Nagao M & Takatori T (1992). Immunohistochemical classification of astrocytes in infants by glial fibrillary acidic protein staining and application to forensic practice. *Nihon Hoigaku Zasshi* **46**, 189–197.

Author contributions

The experiments were performed at the laboratory of Y.I. Y.O.: Conception of the experiments and drafting the article. T.S. and Y.O.: Collection, analysis and interpretation of data and drafting the article. N.T., M.S. and S.U.: Collection of data. K.F.T.: Development of transgenic mice. N.M.: Supervision of experiments and critical revision of the article. Y.I.: Design of

the experiments, analysis and interpretation of data, and drafting the article. All authors approved this manuscript.

Acknowledgements

This work was supported in part by Grants-in-Aid for Science Research (nos. 20590218, 22115003, 22650080, 22680025, 23650218, 24300108 and 24500365) from the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Suzuken Memorial Foundation; the Kanae Foundation for the Promotion of Medical Science; the Daiichi-Sankyo Foundation of Life Science; Health and Labour Sciences Research Grants; and the Funding Program for Next Generation World-Leading Researchers (no. LS023). Authors declare no conflict of interest.