

Clinical Article

Bone Cement-Augmented Short Segment Fixation with Percutaneous Screws for Thoracolumbar Burst Fractures Accompanied by Severe Osteoporosis

Hyun Jin Jung, M.D.,¹ Seok Won Kim, M.D.,¹ Chang Il Ju, M.D.,¹ Sung Hoon Kim, M.D.,² Hyen Sung Kim, M.D.³

Department of Neurosurgery,¹ School of Medicine, Chosun University, Gwangju, Korea

Department of Rehabilitation,² Wonju College of Medicine, Yonsei University, Wonju, Korea

Department of Neurosurgery,³ Heori Sarang Hospital, Daejeon, Korea

Objective : The purpose of this study was to determine the efficacy of bone cement-augmented short segment fixation using percutaneous screws for thoracolumbar burst fractures in a background of severe osteoporosis.

Methods : Sixteen patients with a single-level thoracolumbar burst fracture (T11-L2) accompanying severe osteoporosis treated from January 2008 to November 2009 were prospectively analyzed. Surgical procedures included postural reduction for 3 days and bone cement augmented percutaneous screw fixation at the fracture level and at adjacent levels without bone fusion. Due to the possibility of implant failure, patients underwent implant removal 12 months after screw fixation. Imaging and clinical findings, including involved vertebral levels, local kyphosis, canal encroachment, and complications were analyzed.

Results : Prior to surgery, mean pain score (visual analogue scale) was 8.2 and this decreased to a mean of 2.2 at 12 months after screw fixation. None of the patients complained of pain worsening during the 6 months following implant removal. The percentage of canal compromise at the fractured level improved from a mean of 41.0% to 18.4% at 12 months after surgery. Mean kyphotic angle was improved significantly from 19.8° before surgery to 7.8 at 12 months after screw fixation. Canal compromise and kyphotic angle improvements were maintained at 6 months after implant removal. No significant neurological deterioration or complications occurred after screw removal in any patient.

Conclusion : Bone cement augmented short segment fixation using a percutaneous system can be an alternative to the traditional open technique for the management of selected thoracolumbar burst fractures accompanied by severe osteoporosis.

Key Words : Burst fracture · Fusion · Percutaneous.

INTRODUCTION

Thoracolumbar burst fractures are common spinal injuries caused by vertical loading, and affect at least two columns of the spine predominantly in younger patients, whereas thoracolumbar burst fractures accompanied by osteoporosis increase with age⁵. The goals of surgical treatment for thoracolumbar burst fractures are to restore stability of the vertebral column and to decompress the spinal canal, which when achieved, lead to early mobilization. Despite general agreement regarding the goals of surgical treatment, the management of thoracolumbar burst fractures accompanying osteoporosis without neurological def-

icit remains controversial. The traditional procedures for treating thoracolumbar burst fractures in patients with osteoporosis entail bone fusion with supplemental long-level instrumentation. However, these surgeries have distinct disadvantages. Extensive dissection of paraspinal muscles may be associated with massive blood loss and lead to persisting sequelae, such as, muscular denervation, atrophy, and pain. On the other hand, recent reports on burst fractures have demonstrated the efficacy of bone cement augmented short segment fixation in terms of preventing implant failure^{1,3}. Furthermore, percutaneous screw fixation has been proposed and applied to minimize the negative consequences of traditional open surgery, which include

• Received : March 29, 2012 • Revised : August 28, 2012 • Accepted : October 4, 2012

• Address for reprints : Seok Won Kim, M.D.

Department of Neurosurgery, School of Medicine, Chosun University, 365 Pilmun-daero, Dong-gu, Gwangju 501-717, Korea

Tel : +82-62-220-3126, Fax : +82-62-227-4575, E-mail : chosunns@chosun.ac.kr

• This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

muscular denervation and atrophy^{6,8}). However, no prospective study has been conducted on the clinical outcomes of short segment fixation by percutaneous screw insertion without bone fusion for burst fractures in a background of osteoporosis. Accordingly, we designed this prospective study to evaluate the clinical outcomes of short segment fixation without bone fusion of thoracolumbar burst fractures in osteoporotic patients. We also examined whether implant removal aggravates kyphotic deformities and adversely affects clinical outcomes. The hypothesis of this study was that screw removal would not affect spinal stability.

MATERIALS AND METHODS

This study included 16 patients (5 males, 11 females) with a thoracolumbar burst fracture. The inclusion criteria were limited to : 1) neurologically intact patients with a single level thoracolumbar burst fracture (canal compromise of >30%); 2) kyphotic angle of >20° and anterior vertebral height loss of >40%; 3) intact bilateral pedicles, enabling screw insertion at the fractured vertebra; and 4) an osteoporotic spine (T-score by bone mineral densitometry of <-3.0). Patients with conditions requiring anterior decompression for neurologic deficits, or conditions preventing screw insertion at the fracture level were excluded. However, patients with a posterior column fracture were not excluded (Table 1). All patients underwent implant removal at 12 months after screw fixation due to possibility of implant failure (the need for this procedure was fully explained before surgery). After postural reduction for 3 days using a soft roll under the collapsed vertebra in the supine position, bone cement augmented, percutaneous screw fixation confined to adjacent levels

(one above and one below the injured vertebra and including the fractured vertebra) was performed. Bone cement augmentation was achieved using polymethylmethacrylate (PMMA), which was injected under C-arm guidance through a vertebroplasty needle and allowed to localize in the vertebral body area. Approximately 2 cc of PMMA was injected through the vertebroplasty needle - a total of 4 cc per vertebra. A transpedicular screw system with a detachable screw extender, Apollon® Solco Medical Korea (Fig. 1) was used to insert screws. Intraoperatively, following the induction of general or spinal anesthesia, patients were placed in the prone position with cushions under the iliac crests and thorax to maintain a hyperextended spine.

Lateral plain radiography, computed tomography (CT), and magnetic resonance imaging were performed at three time points, that is, 1) preoperatively, 2) at 12 months after surgery, just before implant removal, 3) at six months after implant removal. The axial CT images showing greatest canal encroachment by retropulsed bone fragments were selected to measure canal compromise. Kyphotic angle was measured by making a straight line from the superior endplate of the vertebra one level above the fractured body and a straight line from the inferior endplate of the vertebra one level below the fractured body. Imaging and clinical findings, including involved vertebral level, vertebral height restoration, local kyphosis, clinical outcomes, and complications were analyzed.

Safety and outcome evaluation

Visual analogue scale pain scores were evaluated at the above-mentioned time points. At six months after implant removal, the patients were evaluated according to a modified version of MacNab’s criteria for characterizing clinical outcomes after spi-

Table 1. Clinical and radiological data of the patients included in this series

Age/Sex	Level	Combined injury	Anesthesia	Canal encroachment (%)			Kyphotic angle (°)		
				Preoperative	12 months after surgery	6 months after implant removal	Preoperative	12 months after surgery	6 months after implant removal
51/F	T12	-	General	40	30	30	30	8	8
48/F	L1	-	General	40	15	15	15	8	12
58/M	L2	Lamina fracture	General	45	20	20	10	5	7
50/M	L2	-	Spinal	60	25	25	24	12	12
49/F	L1	-	Spinal	55	15	15	34	10	11
50/F	L1	-	General	50	25	25	20	7	8
51/F	L2	-	General	55	30	30	20	6	10
45/M	L1	-	Spinal	40	15	15	22	6	10
68/F	L2	-	General	30	15	15	12	6	10
68/F	T12	Lamina fracture	Spinal	30	10	10	15	6	8
67/F	L1	-	General	40	10	10	12	10	10
69/F	L1	-	General	45	25	25	14	10	12
58/F	L2	-	Spinal	30	15	15	30	6	8
55/F	L2	-	General	30	15	15	13	8	10
61/F	L1	-	General	30	20	20	23	8	10
72/F	L1	-	Spinal	35	10	10	23	8	10

nal surgery.

Statistics

Statistical analysis was performed using SAS software version 6.12 (SAS Institute, Inc., Cary, NC, USA). Comparisons at different time points were conducted using the paired Student's t-test. Results are presented as mean values and standard deviations, and differences were considered statistically significant for *p* values of <0.05.

RESULTS

With respect to the anatomic distribution of fractures, T12 (n=2), L1 (n=8), and L2 (n=6) were involved in the present study. Mean operation time was 66.5±30.5 minutes and mean blood loss was 63.1±45.0 mL.

Overall reduction of kyphotic deformity

Intraoperative correction of kyphotic deformities by longitudinal distraction was possible. Mean kyphotic angle improved significantly from 19.8±8.2° (range : 10-34°) before surgery to 7.8° (range : 6-12°) at 12 months after surgery.

At six months after implant removal, the average amount of correction loss of kyphotic deformities was 2.0°, which represented a 10° change in angulation from the original position (*p*<0.01) (Fig. 1).

Proportion of canal compromise

Mean preoperative canal compromise, due to retropulsed bone fragments, was 41.0%. CT at 1 year after surgery and just before implant removal, revealed an improvement in canal compromise to 18.4%. Furthermore, this improvement was maintained on CT scans at 6 months after implant removal (Fig. 2), and was statistically significant (*p*<0.001). No patient experienced neurological deterioration due to a retropulsed bony fragment after implant removal.

Clinical outcomes

No patient needed a drainage catheter and all 16 patients were able to ambulate at 24 hours after surgery wearing a thoracolumbo-sacral-orthosis. At 6 months after implant removal, all patients achieved an excellent or good outcome (excellent in 14 patients and good in 2). Prior to surgery, mean pain score was 8.2 and this decreased to 2.2 at 12 months after surgery and maintained at 6 months after implant removal (Fig. 3). No significant complications or neurological deterioration occurred after screw removal in any patient (Fig. 4, 5).

DISCUSSION

The goals of the surgical treatment of thoracolumbar burst fractures are; to restore vertebral column stability, to decompress the spinal canal, and to facilitate early ambulation. Stabili-

zation by open surgery continues to be widely used to treat thoracolumbar burst fractures, but traditional open surgery is highly invasive and requires a long skin incision. Furthermore, when performed anteriorly, it requires significant invasion of the retroperitoneal space or thoracic cavity and a wide dissection margin^{2,9)}. Therefore, when there is no need to decompress neural elements, multi-segment fixation using a posterior ap-




Fig. 1. Changes in kyphotic angles.




Fig. 2. Fraction of canal compromise.




Fig. 3. Pain score improvement. VAS : visual analogue scale.




Fig. 4. The case of a neurologically intact 58-year-old woman with an L2 bursting fracture. A, B and C : Preoperative simple radiograph and computed tomography scans show about 45% height loss and 45% canal compromise. Posterior element fracture is also seen. D and E : Simple radiograph and a computed tomography scan taken at 12 months after screw fixation show restored vertebral height and improved kyphotic deformity. F and G : Simple radiograph and a magnetic resonance image taken at 6 months after implant removal reveal a well maintained kyphotic angle and much reduced canal compromise.




Fig. 5. The case of neurologically intact 45-year-old man with an L1 bursting fracture. A, B and C : Preoperative simple radiograph, computed tomography scan and magnetic resonance image show about 50% height loss and 40% canal compromise. D and E : Simple radiograph and a computed tomography scan taken at 12 months after screw fixation show a restored vertebral height and improved kyphotic deformity. F and G : Simple radiograph and a magnetic resonance image taken at 6 months after implant removal reveal a well maintained kyphotic angle and much reduced canal compromise.

proach is usually performed. However, this also requires extensive dissection of the paraspinal muscles of the back and resection of the posterior elements of the spine, which makes it difficult to preserve motion segments¹⁵. Due to several problems associated with extensive long level fusion, posterior short-segment fusion is now preferred for the treatment of thoracolumbar burst fractures¹³. The main advantage of short-segment fixation is that it preserves motion segments more so than long-level fixation. However, failure to restore anterior column support can lead to secondary kyphosis, which primarily depends on the residual load transfer capacity of the fractured vertebral body. To date, the results of short-segment fixation have not been good, and implant failure rates of 10 to 30% have been reported^{7,10,16}. Moreover, in severe osteoporotic patients, the anchoring effect that holds screws in place is diminished,

and the probability of hardware failure is high, and thus, the risk of non-union is increased¹⁷. In 2006, Wang et al.¹⁸ reported satisfactory results for short-segment fixation without fusion in surgically-treated burst fractures of the thoracolumbar spine, and found that for surgically treated burst fractures, clinical results were similar for those treated with or without bone fusion. The advantages of instrumentation without fusion are the elimination of donor site complications, the preservation of motion segments, reduced blood loss, and a shorter operation time. More recently, some investigators have reported successful surgical outcomes for percutaneous non-fusion short-segment fixation^{6,8}. The preservation of the segmental stability obtained by posterior screw fixation is not easily achieved before the fusion of fractured vertebra, and screw loosening or vertebral collapse may occur more frequently after percutaneous screwing proce-

dures. In addition, these problems may be more significant in patients with severe osteoporosis. To overcome these limitations, we attempted bone cement augmentation at adjacent vertebrae. In addition, the insertion of bone cement-augmented pedicle screws at fracture levels resulting in segmental constructs with improved biomechanical stability due to the protection afforded fractured vertebral bodies and indirect support of the anterior column^{11,12}. Furthermore, the insertion of bone cement-augmented pedicle screws can significantly reduce pedicle screw bending and increase initial stiffness in the flexion-extension plane. In a previous study, it was found that PMMA can increase pull-out strength from 96 to 262% and transverse bending stiffness by up to 153%¹⁹. In addition, spontaneous remodeling of the spinal canal was shown to occur after a burst fracture, regardless of the treatment modality used. Scapinelli and Candioto¹⁴ concluded that rhythmic respiratory oscillations in cerebrospinal fluid pressure and loss of mechanical loading are important components of bone remodeling. de Klerk et al.⁴ reported that the process of remodeling mainly takes place during the first year after injury, and that subsequently, little remodeling takes place, which is why, we removed implants at 1 year after screw fixation. Furthermore, after bone cement-augmented short segment fixation without bone fusion, early ambulation was possible and motion segments were preserved by screw removal, which contrasts to that observed after long level instrumentation and fusion. The advantages of short segment fixation without bone fusion also include immediate pain relief due to the elimination of donor site pain, reduced blood loss, and a short operative time. The present study demonstrates the efficacy of short segment fixation without fusion in patients with a thoracolumbar burst fracture but without neurologic deficit, but this method is by no means indicated in all patients with a thoracolumbar burst fracture, and in particular, may be contraindicated in patients with a highly unstable fracture and severe neurologic injury. In the present study, we addressed the question as to whether simultaneous fusion is necessary when treating a thoracolumbar burst fracture by percutaneous short-segment fixation rather than addressing controversial issues related to the indications for operative or nonoperative treatment. Furthermore, we acknowledge that some of the patients in the present series might also have been appropriate candidates for balloon kyphoplasty. In fact, we know that nonoperative treatment is also safe and effective for selected patients with an osteoporotic thoracolumbar burst fracture, as demonstrated by improvements in pain and work status. Randomized, comparative, clinical trials in larger populations are required in the near future.

CONCLUSION

Bone cement augmented, percutaneous short segment fixation following postural reduction without fusion might be an alternative to traditional long level fusion for the management

of patients with a thoracolumbar burst fracture accompanied by severe osteoporosis but without neurologic deficit. In the present study, this technique was found to reduce the number of screw fixation levels, preserve motion segments, and correct kyphotic deformities. Furthermore, improved canal compromise and kyphotic deformity were maintained during follow-up after screw removal.

References

- Acosta FL Jr, Aryan HE, Taylor WR, Ames CP : Kyphoplasty-augmented short-segment pedicle screw fixation of traumatic lumbar burst fractures : initial clinical experience and literature review. *Neurosurg Focus* 18 : e9, 2005
- Arrington ED, Smith WJ, Chambers HG, Bucknell AL, Davino NA : Complications of iliac crest bone graft harvesting. *Clin Orthop Relat Res* : 300-309, 1996
- Cho DY, Lee WY, Sheu PC : Treatment of thoracolumbar burst fractures with polymethyl methacrylate vertebroplasty and short-segment pedicle screw fixation. *Neurosurgery* 53 : 1354-1360; discussion 1360-1361, 2003
- de Klerk LW, Fontijne WP, Stijnen T, Braakman R, Tanghe HL, van Linge B : Spontaneous remodeling of the spinal canal after conservative management of thoracolumbar burst fractures. *Spine (Phila Pa 1976)* 23 : 1057-1060, 1998
- Denis F : The three column spine and its significance in the classification of acute thoracolumbar spinal injuries. *Spine (Phila Pa 1976)* 8 : 817-831, 1983
- Gahr RH, Strasser S, Strasser E, Schmidt OI : Percutaneous internal fixation of thoracolumbar spine fractures. *Top Spinal Cord Inj Rehabil* 12 : 45-54, 2006
- Hakalo J, Wroński J : [Complications of a transpedicular stabilization of thoraco-lumbar burst fractures]. *Neurol Neurochir Pol* 40 : 134-139, 2006
- Huang QS, Chi YL, Wang XY, Mao FM, Lin Y, Ni WF, et al. : [Comparative percutaneous with open pedicle screw fixation in the treatment of thoracolumbar burst fractures without neurological deficit]. *Zhonghua Wai Ke Za Zhi* 46 : 112-114, 2008
- Kaneda K, Taneichi H, Abumi K, Hashimoto T, Satoh S, Fujiya M : Anterior decompression and stabilization with the Kaneda device for thoracolumbar burst fractures associated with neurological deficits. *J Bone Joint Surg Am* 79 : 69-83, 1997
- Lee YS, Sung JK : Long-term follow-up results of short-segment posterior screw fixation for thoracolumbar burst fractures. *J Korean Neurosurg Soc* 37 : 416-421, 2005
- Lü FX, Huang Y, Zhang Q, Shi FL, Zhao DS, Hu Q : Pedicle screw fixation against burst fracture of thoracolumbar vertebrae. *Chin J Traumatol* 10 : 349-352, 2007
- Mahar A, Kim C, Wedemeyer M, Mitsunaga L, Odell T, Johnson B, et al. : Short-segment fixation of lumbar burst fractures using pedicle fixation at the level of the fracture. *Spine (Phila Pa 1976)* 32 : 1503-1507, 2007
- Parker JW, Lane JR, Karaikovic EE, Gaines RW : Successful short-segment instrumentation and fusion for thoracolumbar spine fractures : a consecutive 41/2-year series. *Spine (Phila Pa 1976)* 25 : 1157-1170, 2000
- Scapinelli R, Candioto S : Spontaneous remodeling of the spinal canal after burst fractures of the low thoracic and lumbar region. *J Spinal Disord* 8 : 486-493, 1995
- Sihvonen T, Herno A, Paljärvi L, Airaksinen O, Partanen J, Tapaninaho A : Local denervation atrophy of paraspinal muscles in postoperative failed back syndrome. *Spine (Phila Pa 1976)* 18 : 575-581, 1993
- Son BG, Bae KJ, Jung ES, Choi ES, Shin JH, Kim MJ, et al. : Results of

- posterior short segment fixation cases which are more than 65% canal compromise of thoracolumbar burst fractures (Minimum 3 years follow-up). *Korean J Spine* 4 : 199-204, 2007
17. Soshi S, Shiba R, Kondo H, Murota K : An experimental study on transpedicular screw fixation in relation to osteoporosis of the lumbar spine. *Spine (Phila Pa 1976)* 16 : 1335-1341, 1991
18. Wang ST, Ma HL, Liu CL, Yu WK, Chang MC, Chen TH : Is fusion necessary for surgically treated burst fractures of the thoracolumbar and lumbar spine? : a prospective, randomized study. *Spine (Phila Pa 1976)* 31 : 2646-2652; discussion 2653, 2006
19. Wittenberg RH, Lee KS, Shea M, White AA 3rd, Hayes WC : Effect of screw diameter, insertion technique, and bone cement augmentation of pedicular screw fixation strength. *Clin Orthop Relat Res* : 278-287, 1993