

Are socioeconomic disparities in tobacco consumption increasing in India? A repeated cross-sectional multilevel analysis

Nandita Bhan,¹ Swati Srivastava,² Sutapa Agrawal,³ Malavika Subramanyam,⁴ Christopher Millett,⁵ Sakhivel Selvaraj,² S V Subramanian¹

To cite: Bhan N, Srivastava S, Agrawal S, *et al*. Are socioeconomic disparities in tobacco consumption increasing in India? A repeated cross-sectional multilevel analysis. *BMJ Open* 2012;**2**:e001348. doi:10.1136/bmjopen-2012-001348

► Prepublication history and additional material for this paper are available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2012-001348>).

Accepted 28 August 2012

This final article is available for use under the terms of the Creative Commons Attribution Non-Commercial 2.0 Licence; see <http://bmjopen.bmj.com>

For numbered affiliations see end of article

Correspondence to

Dr S V Subramanian; svsubram@hsph.harvard.edu

ABSTRACT

Objectives: India bears a significant portion of the global tobacco burden with high prevalence of tobacco use. This study examines the socioeconomic patterning of tobacco use and identifies the changing gender and socioeconomic dynamics in light of the *Cigarette Epidemic Model*.

Design: Secondary analyses of second and third National Family Health Survey (NFHS) data.

Setting and participants: Data were analysed from 201 219 men and 255 028 women over two survey rounds.

Outcomes and methods: Outcomes included *smoking* (cigarettes, *bidis* and pipes/cigar), *chewed tobacco* (*paan masala*, *gutkha* and others) and *dual use*, examined by education, wealth, living environment and caste. Standardised prevalence and percentage change were estimated. Pooled multilevel models estimated the effect of socioeconomic covariates on the log odds of tobacco use by gender, along with fixed and random parameters.

Findings: Among men (2005–2006), gradients in smoking by education (illiterates: 44% vs postgraduates: 15%) and chewing (illiterates: 47% vs postgraduates: 19%) were observed. Inverse gradients were also observed by wealth, living environment and caste. Chewed tobacco use by women showed inverse socioeconomic status (SES) gradients comparing the illiterates (7.4%) versus postgraduates (0.33%), and poorest (17%) versus richest (2%) quintiles. However, proportional increases in smoking were higher among more educated (postgraduates (98%) vs high schooling only (17%)) and chewing among richer (richest quintile (49%) vs poorest quintile (35%)). Among women, higher educated showed larger declines for smoking—90% (postgraduates) versus 12% (illiterates). Younger men (15–24 years) showed increasing tobacco use (smoking: 123% and chewing: 112%). Older women (35–49 years) show higher prevalence of smoking (3.2%) compared to younger women (0.3%).

Conclusions: Indian tobacco use patterns show significant diversions from the *Cigarette Epidemic Model*—from gender and socioeconomic perspectives. Separate analysis by type is needed to further understand social determinants of tobacco use in India.

ARTICLE SUMMARY

Article focus

- India bears a significant burden of tobacco consumption, with high prevalence of smoking and chewing among men and women.
- Previous studies have established a unique social and spatial gradient in tobacco use. However, no studies have yet reported estimates for changing patterns in tobacco use prevalence or relative risk over time.
- This study estimates socioeconomic patterns and examines the changing gender and socioeconomic dynamics of tobacco use in light of the *Cigarette Epidemic Model*.

Key messages

- Among men, higher prevalence of smoking and chewing for less educated, poorer, rural and lower caste. Sharp and rising inequalities by survey year, but percentage change shows increases are greater among higher socioeconomic status (SES) groups—higher education, urban, richer populations, previously unreported.
- Low and declining risks of smoking and chewing among women. Higher rates of chewing compared to smoking. Increase in smoking with urbanisation for women. Greater declines over time for higher-educated women.
- Significant changing trend by wealth, education and living environment in smoking among men and in chewing among women. Increases in smoking prevalence among younger men (15–24 years) and chewing among younger women (15–24 years).

Strengths and limitations of this study

- First systematic examination of socioeconomic patterns in tobacco use in India, highlighting SES gradients in use and risks among vulnerable populations.
- Large sample, representative and generalisable surveys providing repeated and comparable estimates over time.
- Limitations: (1) cross-sectional data, limiting scope for causal inference, (2) lack of data by tobacco type or volume of use and (3) data from a reproductive health survey may suffer from social desirability bias.

INTRODUCTION

Global estimates indicate that 1 in 10 adult deaths can be attributed to tobacco consumption, leading to approximately five million global deaths per year.^{1–4} Of these, 2.4 million deaths occur in developing countries. India bears a significant portion of this global tobacco burden.^{3–4} Consumption of both smoked and smoke-less (chewed and inhaled) forms of tobacco is highly prevalent among men (47%) and women (14%).⁵ However, previous studies have indicated that tobacco use, like other non-communicable disease risk factors, is unequally distributed across different social determinants in India—education, caste and wealth—among both men and women,^{1–7} indicating a distinct ‘economic and spatial distribution’ in tobacco use.⁶ No studies have yet systematically examined patterns and changes in the prevalence of tobacco consumption in India by socioeconomic factors over time.

In this study, we aim to provide estimates and inferences on the changing gradient of tobacco consumption in India, analysing prevalence and OR patterns from the National Family Health Surveys (NFHS).^{8–9} We discuss our findings in light of the Cigarette Epidemic Model^{10–11} and examine what populations show higher and lower prevalence of tobacco consumption over time.

METHODS

Data were analysed from two rounds of the Indian NFHS (2 and 3) conducted during 1998–1999 and 2005–2006, respectively. NFHS is a national representative cross-sectional survey that is collected and managed by the Indian Institute of Population Sciences (IIPS) in Mumbai, India. These surveys provide vital sources of information on demographic, health and socioeconomic behaviour of Indian households. Data from men and women in the age group of 15–49 years were used from both survey rounds to ensure comparability. Data are representative of all Indian states (except for the small Union Territories), hence covering almost 99% of the country’s population. The surveys were collected using multistage cluster random sampling techniques. Rural and urban areas were sampled separately and a uniform sample design was followed in each state; states and PSUs are considered as levels. Individual questionnaires for men and women were used to interview usual residents of the household or visitors who stayed in the house the night before. Further details on sample design, including sampling framework and sample implementation, are provided in the basic survey reports by IIPS.^{8–9}

Outcomes of interest included *smoking* (cigarettes, *bidis*ⁱ and pipes/cigar), *chewing tobacco* (*paan masala*,ⁱⁱ *gutkha*ⁱⁱⁱ)

ⁱ*Bidis* are local inexpensive cigarettes, which are thinner and contain tobacco flakes rolled inside tendu leaves. Bidis are often smoked by poorer populations.

ⁱⁱ*Paan masala* is a powdered preparation of betel leaves combined with cured tobacco and/or areca nut, which has stimulating properties.

ⁱⁱⁱ*Gutkha* is a savoury or sweet preparation containing areca nut, tobacco, catechu, paraffin and slaked lime.

or other chewed forms of tobacco) and *dual use* (consuming both smoked and chewed forms) of tobacco. NFHS-3 provides details on the different types of smoked and chewed tobacco products, but this information was unavailable in NFHS-2. The main covariates of interest were age, marital status and education at the individual level, and household wealth, area of residence (urban/rural), religion and caste/tribe status at the household level (*variable definitions are provided in table 1*). Survey-weighted age-standardised prevalence estimates of smoking, chewing and dual use of tobacco along with percentage change were calculated. Pooled multilevel models with state, local area and individual as analytical levels were used to estimate the effect of wealth, education, living environment and caste on the log odds of smoking and chewing among men and women. Regression models were adjusted for age, religion and marital status. Survey year was used in the interaction terms to estimate a time trend in socioeconomic determinants of tobacco. Tests for trend included joint tests for significance of fixed parameters and significance tests for random parameters. Model estimates were maximum likelihood based using the Iterative Generalised Least-Squares (IGLS) algorithm as implemented within the MLwin software programme (V.2.23).

RESULTS

Data used in this analysis covered 131 464 men and 130 886 women residing in 92 486 households in NFHS-2 and 69 755 men and 124 142 women residing in 109 041 households in NFHS-3, with an overall response rate of 96% for NFHS-2 and 98% for NFHS-3. Prevalence (%) of smoking, chewing and dual use of tobacco over two survey rounds are presented by the three primary markers of socioeconomic status (SES)—education, wealth and caste (*table 1*) along with percentage change estimates. Estimates by living environment, marital status, age and religion are presented in the appendix (see web *table 1*). Among men, the prevalence of tobacco use (smoking, chewing and dual use) was seen to increase across all socioeconomic groups. For instance, smoking has risen from 35.5% to 40.6% in the fifth (poorest) quintile, 30.6% to 36.5% in the fourth quintile, 25.6% to 31.4% in the middle quintile, 19.3% to 25.8% in the second quintile and 11.9% to 19.9% in the first (richest) quintile (*table 1*). Chewed tobacco use increased from 34.4% to 47.1% among the illiterate populations, 30.2–41.9% among those with primary schooling only, 23.3–33.1% among those with high-school education, 14.9–23.9% among those with college education and 12.4–18.5% among those with postgraduate degree. Higher prevalence of tobacco use among men in each survey round was seen for socioeconomically disadvantaged groups—with less educated, lower wealth, rural residence or lower caste showing an inverse SES gradient; however, greater proportional increases in

Table 1 Prevalence (%) of tobacco smoking, chewing and dual use (smoking and chewing) by wealth, education and caste/tribe status among men and women in the National Family Health Surveys (NFHSs) 1998–1999 and 2005–2006

Sample population	Men									Women									
	Smoke		Chew		Percentage of Δ^{\ddagger}	Dual		Smoke		Chew		Percentage of Δ^{\ddagger}	Dual						
	1998–1999	2005–2006	1998–1999	2005–2006		1998–1999	2005–2006	1998–1999	2005–2006	1998–1999	2005–2006		1998–1999	2005–2006					
<i>Caste/tribe status*</i>																			
General	(M) 50 939	(M) 21 850	22.2	28.8	30	22.2	33.2	50	7.4	10.8	1.0	0.8	-20	6.6	7.24	10	0.24	0.2	
	(F) 50 526	(F) 41 844	(21.4, 22.9)	(27.7, 29.9)		(21.4, 23.02)	(31.8, 34.5)		(6.9, 7.8)	(10.0, 11.5)	(0.9, 1.1)	(0.55, 1.0)		(6.1, 7.1)	(6.6, 7.85)		(0.17, 0.3)	(0.11, 0.27)	
SC	(M) 21 491	(M) 11 953	31.5	39.3	25	27.8	40.5	46	12.1	15.8	2.3	2.4	4	10.9	12.1	11	0.5	0.4	
	(F) 21 045	(F) 20 566	(30.5, 32.5)	(37.8, 40.9)		(26.6, 29.1)	(38.8, 42.1)		(11.4, 12.8)	(14.7, 16.9)	(1.9, 2.8)	(2.05, 2.8)		(10.1, 11.8)	(11.2, 12.9)		(0.4, 0.7)	(0.3, 0.6)	
ST	(M) 16 187	(M) 8453	30.6	36.6	20	38.6	52.6	36	14.5	18.6	3.0	2.7	-10	18.5	25.08	36	1.0	0.9	
	(F) 16 520	(F) 16 518	(28.8, 32.3)	(34.1, 39.1)		(36.9, 40.4)	(49.9, 55.3)		(13.3, 15.6)	(16.7, 20.4)	(2.5, 3.6)	(2.04, 3.4)		(17.0, 20.1)	(22.8, 27.4)		(0.7, 1.2)	(0.6, 1.3)	
OBC	(M) 36 381	(M) 25 144	24.7	31.2	26	25.4	36.2	43	9.9	13.1	1.5	1.4	-7	7.3	7.08	-3	0.3	0.14	
	(F) 36 290	(F) 29 561	(23.9, 25.5)	(30.2, 32.2)		(24.5, 26.3)	(35.0, 37.5)		(9.4, 10.5)	(12.3, 13.8)	(1.3, 1.7)	(1.2, 1.7)		(6.8, 7.7)	(6.6, 7.6)		(0.26, 0.4)	(0.09, 0.21)	
No caste	(M) 6466	(M) 2355	31.8	37.9	19	23.3	35.5	52	10.7	13.0	1.4	0.9	-36	10.3	12.8	24	0.4	0.31	
	(F) 6505	(F) 5653	(29.4, 34.3)	(34.2, 41.6)		(20.9, 25.6)	(31.6, 39.3)		(9.3, 12.1)	(10.9, 15.2)	(0.8, 1.9)	(0.36, 1.48)		(9.1, 11.6)	(10.8, 14.8)		(0.2, 0.6)	(0.03, 0.59)	
<i>Education level†</i>																			
Postgraduate	(M) 3432	(M) 2920	7.6	15.05	98	12.4	18.5	49	1.7	4.04	0.2	0.02	-90	1.2	0.33	-73	0.2	0.004	
	(F) 1963	(F) 3526	(6.5, 8.7)	(13.01, 17.1)		(10.8, 14.1)	(16.0, 21.1)		(1.2, 2.2)	(3.0, 5.1)	(-0.2, 0.6)	(-0.007, 0.05)		(0.45, 1.9)	(0.14, 0.5)		(-0.17, 0.5)	(-0.002, 0.012)	
College	(M) 11 340	(M) 7811	11.1	20.7	86	14.9	23.9	60	3.5	6.7	0.1	0.11	10	1.3	1.8	39	0.05	0.04	
	(F) 6586	(F) 9424	(10.2, 11.9)	(19.2, 22.1)		(13.9, 15.8)	(22.3, 25.6)		(3.1, 3.9)	(5.8, 7.6)	(-0.01, 0.2)	(0.03, 0.19)		(0.9, 1.6)	(1.4, 2.2)		(-0.04, 0.14)	(-0.017, 0.1)	
High school	(M) 69 996	(M) 26 100	21.2	24.7	17	23.3	33.1	42	7.8	9.9	0.2	0.07	-65	4.2	3.4	-19	0.1	0.04	
	(F) 46 629	(F) 34 338	(20.7, 21.8)	(23.8, 25.5)		(22.7, 23.9)	(32.0, 34.1)		(7.5, 8.2)	(9.3, 10.6)	(0.17, 0.3)	(0.04, 0.1)		(3.9, 4.6)	(3.04, 3.7)		(0.06, 0.13)	(0.02, 0.06)	
Primary school	(M) 21 730	(M) 12 622	32.7	35.6	1	30.2	41.9	39	12.1	14.8	0.7	0.2	-71	9.0	7.4	-18	0.18	0.07	
	(F) 20 604	(F) 19 451	(31.7, 33.6)	(34.3, 36.9)		(29.2, 31.2)	(40.5, 43.4)		(11.4, 12.7)	(13.8, 15.7)	(0.5, 0.8)	(0.13, 0.3)		(8.4, 9.6)	(6.8, 8.02)		(0.13, 0.24)	(0.03, 0.11)	
Illiterate	(M) 24 966	(M) 20 302	38.9	43.9	13	34.4	47.1	37	15.9	18.9	2.6	2.3	-12	11.9	13.3	12	0.6	0.42	
	(F) 55 104	(F) 57 403	(37.9, 39.9)	(42.8, 45.2)		(33.2, 35.5)	(45.7, 48.5)		(15.1, 16.6)	(17.9, 19.9)	(2.4, 2.9)	(2.02, 2.6)		(11.4, 12.6)	(12.6, 13.0)		(0.53, 0.72)	(0.33, 0.51)	
<i>Wealth quintiles</i>																			
Richest	(M) 26 291	(M) 13 706	11.9	19.9	63	13.7	20.4	49	3.3	5.8	0.2	0.14	-30	2.2	2.02	-8	0.05	0.042	
	(F) 26 177	(F) 24 837	(11.2, 12.5)	(18.8, 21.1)		(12.8, 14.6)	(19.0, 21.8)		(2.97, 3.6)	(5.1, 6.4)	(0.1, 0.3)	(0.08, 0.2)		(1.9, 2.5)	(1.7, 2.3)		(0.01, 0.07)	(0.009, 0.07)	
Richer	(M) 26 293	(M) 13 946	19.3	25.8	37	19.0	30.2	59	5.6	9.3	0.46	0.37	-20	5.4	4.9	-9	0.1	0.06	
	(F) 26 177	(F) 24 837	(18.5, 20.04)	(24.6, 27.1)		(18.1, 19.9)	(28.7, 31.7)		(5.2, 6.02)	(8.4, 10.2)	(0.37, 0.54)	(0.27, 0.47)		(4.9, 5.9)	(4.4, 5.4)		(0.07, 0.18)	(0.02, 0.09)	
Middle	(M) 26 294	(M) 14 075	25.6	31.4	25	22.9	34.9	52	7.6	11.5	1.1	0.7	-36	7.8	6.9	-11	0.23	0.07	
	(F) 26 174	(F) 24 826	(24.7, 26.4)	(30.1, 32.7)		(22.0, 23.9)	(33.4, 36.3)		(7.2, 8.1)	(10.6, 12.3)	(0.9, 1.3)	(0.6, 0.9)		(7.2, 8.3)	(6.4, 7.4)		(0.16, 0.3)	(0.04, 0.11)	
Poorer	(M) 26 293	(M) 14 007	30.6	36.5	21	28.9	39.5	37	11.6	14.5	1.7	1.7	0	10.9	10.5	-4	0.4	0.24	
	(F) 26 179	(F) 24 814	(29.7, 31.5)	(35.2, 37.8)		(28.0, 29.9)	(38.03, 40.9)		(10.9, 12.2)	(13.6, 15.5)	(1.5, 1.9)	(1.4, 1.9)		(10.3, 11.6)	(9.8, 12.2)		(0.3, 0.5)	(0.17, 0.31)	
Poorest	(M) 26 293	(M) 14 021	35.5	40.6	13	36.8	49.7	35	16.7	19.4	3.5	3.2	-9	14.1	17.1	21	0.9	0.7	
	(F) 26 179	(F) 24 828	(34.4, 36.5)	(39.3, 41.9)		(35.7, 37.9)	(48.2, 51.2)		(15.9, 17.5)	(18.4, 20.5)	(3.1, 3.9)	(2.8, 3.7)		(13.3, 15.0)	(16.03, 18.1)		(0.7, 1.0)	(0.52, 0.86)	
Total	(M) 131 464	(M) 69 755																	
	(F) 130 886	(F) 124 142																	

*Scheduled castes (SC) and scheduled tribes (ST) are identified by the Government of India as socially and economically backward and needing protection from social injustice and exploitation. Other backward class is a diverse collection of intermediate castes that were considered low in the traditional caste hierarchy but are clearly above SC. General is thus a default residual group that enjoys higher status in the caste hierarchy.

†Postgraduate: 15 or more years of education; college: 13–15 years of education; high school: 8–12 years of education; secondary: 5–8 years of education; primary: 0–5 years of education; illiterate: 0 years of education.

‡Percentage change (% Δ) numbers have been rounded to the nearest integer.

All results for prevalence are survey adjusted and age standardised. Prevalence results are all in percentages.

prevalence over time were seen among higher SES groups. For instance, higher absolute smoking prevalence in NFHS-3 was seen among men in lower wealth quintiles compared to those in higher wealth quintiles (41% for fifth (poorest) quintile and 37% for fourth quintile, compared to 20% in first (richest) quintile and 26% in second quintile); higher percentage increases in smoking were recorded among first (richest) quintile (63%) and second quintile (37%) compared to fourth quintile (21%) and fifth (poorest) quintile (13%). Similar trend was seen for education with a 98% increase in prevalence among those with postgraduate education and 13% increase in prevalence among those with no education over the two survey rounds. Sharper inequalities with higher inter-group differences were seen for smoking compared to chewing. Prevalence of chewing among men in the richest quintile and with postgraduate education each increased by 49%, while that for the poorest increased by 35% and for those with no education increased by 37%.

Socioeconomic patterns for tobacco use among women differed distinctly. Overall prevalence rates of tobacco use among women were significantly lower than men. In 2005–2006, prevalence of smoking and chewing among women with no education was 2.3% and 13.3%, respectively, while the same for men was 43.9% and 47.1%, respectively (table 1). Women in most SES categories showed a declining trend for tobacco use, and only scheduled caste women and those with college education showed small increases in smoking and chewing. Higher and more consistent declines in prevalence were seen for education, compared to wealth and caste. For instance, women with postgraduate education noted a 90% decrease in smoking and a 73% decrease in chewing (table 1). Women in the first (richest) quintile showed a decline of 30% for smoking and 8% for chewing (table 1). Results by area of residence (see web table 1) showed an increase in risks for tobacco use with urbanisation among both men and women, except in the prevalence of chewed tobacco among women. Figures 1 and 2 show the percentage change in smoking and chewing by education and wealth for men and women reflecting findings from table 1.

Tables 2 and 3 present results from pooled multilevel models showing ORs for smoking and chewing by education, wealth, living environment and caste, along with interactions with survey year. Gradients in ORs (95% CI) are seen by all four markers of SES among men and women for smoking and chewing with sharper inequalities seen for education and wealth, compared to other markers. Controlling for wealth, caste and living environment and compared to those with postgraduate education, the OR of smoking for men with no education: 3.18 (95% CI 2.96 to 3.43), with primary education: 2.73 (95% CI 2.54 to 2.94), with high school education: 1.81 (95% CI 1.69 to 1.85) and with college education: 1.38 (95% CI 1.28 to 1.49). Controlling for education, caste and living environment and compared to the first (richest) quintile,

OR for chewing in the second quintile: 1.43 (95% CI 1.38 to 1.48), middle quintile: 1.75 (95% CI 1.68 to 1.82), fourth quintile: 1.92 (95% CI 1.84 to 2.01) and fifth (poorest) quintile: 2.1 (95% CI 1.99 to 2.2). Interaction terms in the two tables provide effect estimates for change over the two survey rounds. Among men, significant parameters for interaction terms for smoking are seen by wealth, education (except college educated) and living environment (except for small city); and for chewing by wealth, higher education and by residence in towns. The joint test for interaction of fixed terms is significant for smoking by wealth (joint test: 174.31, $p < 0.001$), education (joint test: 13.31, $p = 0.009$) and living environment (joint test: 13.44, $p = 0.003$) and for chewing by wealth (joint test: 15.63, $p = 0.003$), representing robust change over time. χ^2 values for random parameters are significant both at state (smoking χ^2 : 12.82, $p = 0.0003$ and chewing χ^2 : 12.89, $p = 0.0003$) and local area (smoking χ^2 : 650.41, $p < 0.0001$ and chewing χ^2 : 801.4, $p < 0.0001$) level showing variation at both levels.

Among women, controlling for education, caste and living environment and compared to those in the first (richest) quintile, the OR of smoking in second quintile: 1.48 (95% CI 1.28 to 1.71), middle quintile: 1.9 (95% CI 1.64 to 2.2), fourth quintile: 2.75 (95% CI 2.37 to 3.19) and fifth (poorest) quintile: 3.95 (95% CI 3.39 to 4.6). Controlling for wealth, caste and living environment and compared to those with postgraduate education, OR of chewing among women with college education: 1.84 (95% CI 1.55 to 2.19), high-school education: 2.19 (95% CI 1.86 to 2.57), primary schooling: 2.87 (95% CI 2.44 to 3.4) and no education: 3.85 (95% CI 3.27 to 4.53). Significant ORs for interaction terms are seen for smoking by wealth (joint test: 19.128, $p < 0.0001$) and for chewing by wealth (joint test: 31.96, $p < 0.0001$), education (joint test: 17.42, $p < 0.0001$) and living environment (joint test: 157.008, $p < 0.0001$). χ^2 values for random parameters are significant for both state (smoking χ^2 : 12.91, $p = 0.0004$ and chewing χ^2 : 12.94, $p = 0.011$) and local area (smoking χ^2 : 264, $p < 0.0001$ and chewing χ^2 : 839, $p < 0.0001$), showing variation at both levels. Figure 3 presents adjusted probability estimates for smoking and chewing among men and women by wealth and education from multilevel models, which show findings similar to prevalence estimates.

DISCUSSION

In 1994, Lopez *et al*¹⁰ proposed the four-stage *Cigarette Epidemic Model* discussing transitions in smoking prevalence, consumption amount and mortality in developed countries. As per the model in stage I, male smoking prevalence is comparatively low and rising (<20%) and female smoking prevalence does not exceed 5% due to sociocultural factors. In stage II, tobacco prevalence among men starts to rise rapidly and peaks around 50–80% with female smoking increasing at a lagged pace behind men. In stage III, prevalence rates for smoking




Figure 1 Percentage change in smoking among men and women by education level and wealth quintiles.

among men start to fall, with both male and female smoking converging. Increases are seen for smoking rates among younger compared to older populations. In stage IV, prevalence of smoking begins to decline for both men and women. Mortality attributable to smoking rises to about one-third for all men, with much lower mortality rates seen among women. This model was developed based on empirical data from developed countries and has not been tested in developing countries. However, in 2011, Thun *et al*¹¹ proposed




Figure 2 Percentage change in chewing among men and women by education level and wealth quintiles.

modifications to the model potentially relevant for developing countries and were the first to note that smoking patterns by gender in developing countries distinctly differ from patterns noted in developed countries. India shows a high and complex burden of tobacco consumption, as also reported in tobacco surveillance studies.^{2 12 13} This paper uses empirical evidence to show that India is currently between stages II and III of the Cigarette Epidemic model on the basis of estimates of smoking for men, but distinctly differs from the model on the patterns seen for women.

Overall, several dissimilarities are noted in the Indian experience from this model. *First*, India's unique tobacco experience comprises a 'double burden' of smoked *and* chewed tobacco consumption. Patterns for smoking and chewing seem to follow trajectories that differ by education, living environment and wealth. Further, within smoking, differences potentially exist by SES in the consumption of cigarettes from *bidis*, which most data are unable to distinguish. The quantity and quality of these products may differentially determine the mortality burden attributable to tobacco use in India.¹⁴⁻¹⁶ Assessment of the disease, mortality and cost burden of the tobacco epidemic needs to account for this complexity.^{14 15}

Second, socioeconomic and sociocultural dynamics play profound roles in impacting tobacco use in India. Differences in tobacco consumption are seen by major SES markers such as wealth, education, living environment and caste. Findings from this analysis indicate a dichotomy between *higher absolute prevalence* by lower caste, wealth and education levels; but *higher relative change in prevalence* over time by higher caste, wealth and education levels. *No previous study has reported this finding for India.* Urbanisation seems to be playing an increasing role in impacting tobacco use for men and women. Further analyses by type and amount of tobacco consumed are needed to systematically understand these patterns.

Third, social gradients in tobacco use (overall and by type of tobacco) in India distinctly differ by gender. Despite women's empowerment, large-scale increases in women's smoking as predicted by the *Cigarette Epidemic Model* are yet to be seen in India.^{10 11 17 18} Aggregate estimates show that women are far behind men in prevalence rates for smoking; and smoking and chewing rates among women, barring a few groups, seem to be declining. The reasons for this could be several. *First*, that Indian sociocultural realities and lower acceptability of smoking among women lead to delays in age of initiation of smoking and higher rates among older compared to younger women (see web table 1). Women's smoking has been linked to their empowerment, but this may be confined to urban areas and it is possible that on average, smoking continues to remain a social taboo among women. Representation of smoking in the media may also explain the gender patterns in the use of tobacco; smoking has been projected as an expression

Table 2 Pooled regression models showing ORs (95% CIs) for smoking and chewing among men and interactions for wealth, education, residence and caste over time

Covariates	Smoking					Chewing				
	Model 1: full model	Model 2: interaction with wealth	Model 3: interaction with education	Model 4: interaction with living environment	Model 5: interaction with caste	Model 1: full model	Model 2: interaction with wealth	Model 3: interaction with education	Model 4: interaction with living environment	Model 5: interaction with caste
<i>Wealth quintiles</i> (richest)	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Richer	1.37 (1.32 to 1.43)	1.51 (1.44 to 1.59)	1.37 (1.31 to 1.42)	1.36 (1.3 to 1.41)	1.37 (1.31 to 1.42)	1.43 (1.38 to 1.48)	1.33 (1.27 to 1.4)	1.42 (1.36 to 1.47)	1.42 (1.36 to 1.47)	1.41 (1.36 to 1.47)
Middle	1.71 (1.64 to 1.78)	1.99 (1.89 to 2.1)	1.71 (1.64 to 1.78)	1.68 (1.61 to 1.75)	1.7 (1.63 to 1.77)	1.75 (1.68 to 1.82)	1.63 (1.55 to 1.72)	1.73 (1.66 to 1.8)	1.72 (1.65 to 1.8)	1.72 (1.65 to 1.79)
Poorer	2.06 (1.97 to 2.16)	2.51 (2.37 to 2.65)	2.05 (1.96 to 2.14)	2.02 (1.93 to 2.11)	2.04 (1.95 to 2.14)	1.92 (1.84 to 2.01)	1.77 (1.67 to 1.87)	1.89 (1.8 to 1.97)	1.86 (1.78 to 1.95)	1.87 (1.79 to 1.96)
Poorest	2.33 (2.22 to 2.46)	2.88 (2.72 to 3.06)	2.31 (2.19 to 2.43)	2.29 (2.18 to 2.41)	2.3 (2.19 to 2.42)	2.1 (1.99 to 2.2)	1.93 (1.82 to 2.05)	2.03 (1.92 to 2.13)	2.03 (1.932.13)	2.02 (1.93 to 2.12)
<i>Education</i> (postgraduate)	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
College	1.38 (1.28 to 1.49)	1.41 (1.31 to 1.53)	1.34 (1.19 to 1.5)	1.4 (1.29 to 1.51)	1.4 (1.3 to 1.51)	1.26 (1.17 to 1.36)	1.28 (1.19 to 1.39)	1.14 (1.02 to 1.27)	1.29 (1.19 to 1.39)	1.29 (1.2 to 1.39)
High school	1.81 (1.69 to 1.95)	1.91 (1.78 to 2.05)	2.16 (1.95 to 2.4)	1.87 (1.74 to 2.01)	1.87 (1.74 to 2.01)	1.53 (1.43 to 1.65)	1.64 (1.53 to 1.76)	1.46 (1.33 to 1.61)	1.65 (1.54 to 1.77)	1.66 (1.54 to 1.78)
Primary	2.73 (2.54 to 2.94)	2.81 (2.61 to 3.04)	3.4 (3.05 to 3.78)	2.77 (2.57 to 2.99)	2.77 (2.57 to 2.99)	1.98 (1.84 to 2.13)	2.04 (1.9 to 2.2)	1.84 (1.66 to 2.04)	2.05 (1.9 to 2.21)	2.06 (1.91 to 2.21)
No education	3.18 (2.96 to 3.43)	3.27 (3.03 to 3.53)	4.07 (3.66 to 4.53)	3.19 (2.96 to 4.34)	3.18 (2.95 to 3.42)	2.09 (1.94 to 2.26)	2.06 (1.91 to 2.22)	1.93 (1.74 to 2.14)	2.07 (1.92 to 2.23)	2.08 (1.93 to 2.24)
<i>Living environment</i> (large city)	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Small city	0.77 (0.72 to 0.82)	0.81 (0.76 to 0.87)	0.8 (0.74 to 0.85)	0.82 (0.75 to 0.9)	0.79 (0.74 to 0.84)	0.96 (0.89 to 1.02)	1.02 (0.96 to 1.09)	1.02 (0.96 to 1.09)	1.06 (0.97 to 1.16)	1.03 (0.97 to 1.1)
Town		0.83 (0.79 to 0.88)	0.81 (0.77 to 0.85)	0.88 (0.82 to 0.95)	0.81 (0.77 to 0.85)		1.01 (0.95 to 1.06)	1.01 (0.95 to 1.06)	1.06 (0.98 to 1.15)	1.01 (0.96 to 1.07)

Continued

Table 2 Continued

Covariates	Smoking					Chewing					
	Model 1: full model	Model 2: interaction with wealth	Model 3: interaction with education	Model 4: interaction with living environment	Model 5: interaction with caste	Model 1: full model	Model 2: interaction with wealth	Model 3: interaction with education	Model 4: interaction with living environment	Model 5: interaction with caste	
Village	0.78 (0.74 to 0.82)	0.75 (0.71 to 0.78)	0.73 (0.7 to 0.77)	0.87 (0.82 to 0.92)	0.73 (0.7 to 0.76)	0.92 (0.87 to 0.97)	0.76 (0.73 to 0.79)	0.92 (0.88 to 0.96)	0.92 (0.88 to 0.96)	0.93 (0.88 to 0.99)	0.93 (0.88 to 0.97)
Caste/tribe (Other)	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
SC	1.16 (1.12 to 1.2)	1.15 (1.11 to 1.19)	1.15 (1.11 to 1.18)	1.15 (1.11 to 1.19)	1.16 (1.11 to 1.21)	1.12 (1.08 to 1.15)	1.1 (1.06 to 1.13)	1.1 (1.06 to 1.13)	1.1 (1.06 to 1.13)	1.1 (1.06 to 1.13)	1.1 (1.06 to 1.14)
ST	1.14 (1.09 to 1.2)	1.13 (1.08 to 1.2)	1.14 (1.08 to 1.18)	1.14 (1.09 to 1.2)	1.15 (1.09 to 1.21)	3.02 (2.88 to 3.15)	1.1 (1.06 to 1.15)	1.1 (1.05 to 1.15)	1.1 (1.05 to 1.15)	1.1 (1.05 to 1.15)	1.17 (1.11 to 1.24)
OBC	1.00 (0.98 to 1.04)	0.99 (0.97 to 1.02)	0.99 (0.96 to 1.02)	0.99 (0.97 to 1.02)	1.03 (0.99 to 1.07)	1.05 (1.02 to 1.08)	1.01 (0.97 to 1.03)	1.002 (0.97 to 1.03)	1.003 (0.97 to 1.03)	1.003 (0.97 to 1.03)	1.04 (0.99 to 1.07)
No caste or missing	1.06 (1.01 to 1.13)	1.07 (1.01 to 1.13)	1.07 (1.01 to 1.13)	1.08 (1.02 to 1.14)	1.05 (0.98 to 1.13)	0.97 (0.92 to 1.03)	1.02 (0.96 to 1.08)	1.01 (0.96 to 1.08)	1.02 (0.96 to 1.08)	1.02 (0.96 to 1.08)	0.97 (0.9 to 1.04)
Survey year		1.74 (1.7 to 1.88)	1.67 (1.46 to 1.91)	1.49 (1.4 to 1.58)	1.24 (1.19 to 1.3)		1.47 (1.38 to 1.56)	1.34 (1.17 to 1.54)	1.69 (1.58 to 1.8)	1.72 (1.65 to 1.79)	
Year* Richer		0.78 (0.73 to 0.84)					1.15 (1.07 to 1.24)				
Year * Middle		0.68 (0.63 to 0.73)					1.13 (1.05 to 1.22)				
Year* Poorer		0.60 (0.56 to 0.65)					1.15 (1.07 to 1.24)				
Year* Poorest		0.58 (0.53 to 0.62)					1.11 (1.03 to 1.2)				
Year* College			1.13 (0.97 to 1.32)					1.27 (1.1 to 1.48)			
Year* High school			0.78 (0.68 to 0.89)					1.28 (1.11 to 1.47)			
Year* Primary			0.66 (0.57 to 0.77)					1.02 (0.88 to 1.18)			

Continued

Table 2 Continued

Covariates	Smoking					Chewing				
	Model 1: full model	Model 2: interaction with wealth	Model 3: interaction with education	Model 4: interaction with living environment	Model 5: interaction with caste	Model 1: full model	Model 2: interaction with wealth	Model 3: interaction with education	Model 4: interaction with living environment	Model 5: interaction with caste
Year* No education			0.61 (0.53 to 0.70)					1.01 (0.87 to 1.17)		
Year* Small city				0.99 (0.87 to 1.12)					0.94 (0.82 to 1.06)	
Year* Town				0.89 (0.81 to 0.98)					0.9 (0.82 to 0.99)	
Year* Village				0.73 (0.68 to 0.79)					0.98 (0.91 to 1.05)	
Year* SC					0.99 (0.92 to 1.05)					0.99 (0.93 to 1.06)
Year* ST					0.97 (0.9 to 1.05)					0.84 (0.78 to 0.91)
Year* OBC					0.92 (0.87 to 0.97)					0.92 (0.87 to 0.97)
Year* No caste					1.07 (0.94 to 1.21)					1.15 (1.02 to 1.29)
<i>Fixed part of the model</i>										
Joint χ^2 test for interaction (p value)	174.31 (p<0.001)	13.31 (p=0.009)	13.44 (p=0.003)	0.318 (p=0.98)		15.63 (p=0.003)	8.52 (p=0.074)	2.6 (p=0.46)	1.387 (p=0.85)	
<i>Random part of the model</i>										
χ^2 for level 3: state (p value)	12.82 (p=0.003)	12.82 (p=0.003)	12.82 (p=0.003)	12.82 (p=0.003)		12.89 (p=0.0003)	12.89 (p=0.0003)	12.89 (p=0.0003)	12.89 (p=0.0003)	12.89 (p=0.0003)
χ^2 for level 2: local area (p value)	650.41 (p<0.001)	655.7 (p<0.001)	654.2 (p<0.001)	660.7 (p<0.001)		801.6 (p<0.001)	802.4 (p<0.001)	802.4 (p<0.001)	801.7 (p<0.001)	801.7 (p<0.001)

*Models are controlled for age (centred at 29 years), marital status and religion.

Table 3 Pooled regression models showing ORs (95% CIs) for smoking and chewing among women and interactions for wealth, education, residence and caste over time.

Covariates	Smoking					Chewing				
	Model 1: full model	Model 2: interaction with wealth	Model 3: interaction with education	Model 4: interaction with living environment	Model 5: interaction with caste	Model 1: full model	Model 2: interaction with wealth	Model 3: interaction with education	Model 4: interaction with living environment	Model 5: interaction with caste
<i>Wealth quintiles</i> (richest)	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Richer	1.48 (1.28 to 1.71)	1.82 (1.49 to 2.22)	1.48 (1.28 to 1.71)	1.48 (1.28 to 1.71)	1.49 (1.29 to 1.72)	1.48 (1.41 to 1.56)	1.41 (1.32 to 1.51)	1.48 (1.4 to 1.55)	1.51 (1.43 to 1.59)	1.48 (1.41 to 1.56)
Middle	1.9 (1.64 to 2.19)	2.63 (2.17 to 3.19)	1.87 (1.62 to 2.16)	1.88 (1.62 to 2.18)	1.9 (1.64 to 2.19)	1.75 (1.66 to 1.85)	1.58 (1.47 to 1.7)	1.75 (1.65 to 1.84)	1.81 (1.72 to 1.92)	1.76 (1.66 to 1.85)
Poorer	2.75 (2.37 to 3.19)	3.72 (3.07 to 5.52)	2.8 (2.41 to 3.25)	2.81 (2.42 to 3.26)	2.85 (2.45 to 3.30)	2.14 (2.02 to 2.27)	1.86 (1.72 to 2.002)	2.13 (2.01 to 2.25)	2.2 (2.07 to 2.32)	2.14 (2.02 to 2.27)
Poorest	3.95 (3.39 to 4.6)	4.83 (3.97 to 5.88)	4.03 (3.46 to 4.69)	4.05 (3.47 to 4.72)	4.08 (3.5 to 4.75)	2.67 (2.5 to 2.84)	2.14 (1.99 to 2.32)	2.65 (2.49 to 2.82)	2.7 (2.54 to 2.88)	2.7 (2.5 to 2.84)
<i>Education</i> (postgraduate)	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
College	1.59 (0.91 to 2.8)	1.61 (0.92 to 2.82)	0.98 (0.37 to 2.58)	1.58 (0.9 to 2.78)	1.57 (0.9 to 2.75)	1.84 (1.55 to 2.19)	1.83 (1.54 to 2.17)	1.1 (0.85 to 1.42)	1.82 (1.53 to 2.17)	1.84 (1.55 to 2.19)
High school	1.78 (1.06 to 2.99)	1.67 (1.01 to 2.82)	1.95 (0.83 to 4.56)	1.62 (0.96 to 2.72)	1.61 (0.99 to 2.7)	2.19 (1.86 to 2.57)	2.13 (1.82 to 2.51)	1.47 (1.17 to 1.85)	2.15 (1.82 to 2.54)	2.21 (1.88 to 2.6)
Primary	2.78 (1.66 to 4.68)	2.62 (1.56 to 4.41)	3.32 (1.42 to 7.76)	2.54 (1.51 to 4.29)	2.52 (1.5 to 4.24)	2.87 (2.44 to 3.4)	2.83 (2.4 to 3.33)	2.03 (1.61 to 2.56)	2.86 (2.42 to 3.37)	2.89 (2.45 to 3.41)
No education	4.78 (2.84 to 8.04)	4.91 (2.93 to 8.23)	6.89 (2.53 to 13.73)	4.72 (2.81 to 7.93)	4.66 (2.77 to 7.81)	3.85 (3.27 to 4.53)	3.75 (3.19 to 4.42)	2.58 (2.04 to 3.24)	3.8 (3.22 to 4.48)	3.85 (3.27 to 4.53)
<i>Living environment</i> (large city)	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Small city	0.91 (0.75 to 1.09)	0.98 (0.82 to 1.17)	1.01 (0.84 to 1.21)	0.78 (0.58 to 1.04)	1.004 (0.84 to 1.2)	1.23 (1.13 to 1.32)	1.25 (1.15 to 1.35)	1.22 (1.13 to 1.32)	0.97 (0.86 to 1.1)	1.2 (1.11 to 1.29)
Town										

Continued

Table 3 Continued

Covariates	Smoking					Chewing				
	Model 1: full model	Model 2: interaction with wealth	Model 3: interaction with education	Model 4: interaction with living environment	Model 5: interaction with caste	Model 1: full model	Model 2: interaction with wealth	Model 3: interaction with education	Model 4: interaction with living environment	Model 5: interaction with caste
Village	1.19 (1.03 to 1.37)	1.2 (1.04 to 1.38)	1.22 (1.05 to 1.4)	1.31 (1.06 to 1.62)	1.23 (1.1 to 1.41)	1.36 (1.27 to 1.45)	1.37 (1.29 to 1.47)	1.36 (1.27 to 1.45)	0.95 (0.86 to 1.04)	1.34 (1.25 to 1.43)
Caste/tribe (other)	1.4 (1.24 to 1.59)	1.28 (1.13 to 1.46)	1.31 (1.16 to 1.49)	1.37 (1.14 to 1.66)	1.32 (1.16 to 1.49)	1.07 (1.01 to 1.14)	1.09 (1.02 to 1.16)	1.08 (1.02 to 1.15)	0.69 (0.64 to 0.75)	1.07 (1.005 to 1.13)
SC	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
ST	1.5 (1.39 to 1.62)	1.55 (1.43 to 1.67)	1.54 (1.43 to 1.67)	1.54 (1.43 to 1.67)	1.4 (1.27 to 1.55)	1.28 (1.23 to 1.33)	1.28 (1.23 to 1.36)	1.28 (1.23 to 1.33)	1.29 (1.24 to 1.34)	1.28 (1.22 to 1.35)
OBC	2.04 (1.86 to 2.24)	2.11 (1.92 to 2.3)	2.11 (1.93 to 2.31)	2.11 (1.93 to 2.31)	1.99 (1.79 to 2.23)	1.53 (1.46 to 1.6)	1.53 (1.46 to 1.61)	1.53 (1.46 to 1.61)	1.52 (1.45 to 1.6)	1.48 (1.39 to 1.58)
No caste or missing	1.11 (1.03 to 1.2)	1.17 (1.08 to 1.26)	1.16 (1.08 to 1.26)	1.16 (1.08 to 1.25)	1.13 (1.03 to 1.24)	1.03 (0.99 to 1.07)	1.03 (0.99 to 1.07)	1.03 (0.99 to 1.07)	1.03 (0.99 to 1.07)	1.06 (1.02 to 1.12)
Survey year	0.7 (0.6 to 0.81)	0.74 (0.64 to 0.86)	0.72 (0.62 to 0.85)	0.73 (0.62 to 0.85)	1.01 (0.85 to 1.2)	1.02 (0.95 to 1.09)	1.02 (0.96 to 1.09)	1.02 (0.95 to 1.09)	1.01 (0.95 to 1.08)	0.92 (0.85 to 1.01)
Year* Richer		1.004 (0.98 to 1.03)	0.89 (0.31 to 2.56)	0.67 (0.55 to 0.83)	0.57 (0.51 to 0.63)		0.81 (0.75 to 0.87)	0.54 (0.4 to 0.74)	0.52 (0.47 to 0.56)	1.02 (0.97 to 1.07)
Year* Middle		0.66 (0.5 to 0.87)					1.09 (0.99 to 1.21)			
Year* Poorer		0.47 (0.36 to 0.61)					1.21 (1.1 to 1.34)			
Year* Poorest		0.56 (0.44 to 0.72)					1.3 (1.18 to 1.43)			
Year* College		0.71 (0.56 to 0.91)					1.5 (1.37 to 1.64)			
Year* High school			1.98 (0.6 to 6.52)					2.26 (1.6 to 3.21)		
Year* Primary			0.77 (0.26 to 2.22)					1.94 (1.41 to 2.67)		
			0.61 (0.21 to 1.75)					1.74 (1.27 to 2.4)		

Continued

Table 3 Continued

Covariates	Smoking					Chewing				
	Model 1: full model	Model 2: interaction with wealth	Model 3: interaction with education	Model 4: interaction with living environment	Model 5: interaction with caste	Model 1: full model	Model 2: interaction with wealth	Model 3: interaction with education	Model 4: interaction with living environment	Model 5: interaction with caste
Year* No education			0.7 (0.24 to 2.0)					1.94 (1.41 to 2.66)		
Year* Small city				1.44 (0.99 to 2.07)					1.61 (1.38 to 1.88)	
Year* Town				0.89 (0.68 to 1.16)					1.94 (1.72 to 2.19)	
Year* Village				0.92 (0.74 to 1.15)					2.26 (2.06 to 2.48)	
Year* SC					1.29 (1.11 to 1.51)					0.99 (0.93 to 1.07)
Year* ST					1.16 (0.99 to 1.36)					1.07 (0.98 to 1.16)
Year* OBC					1.1 (0.96 to 1.27)					0.95 (0.89 to 1.01)
Year* No caste					0.45 (0.32 to 0.63)					1.22 (1.07,1.38)
<i>Fixed part of the model</i>										
Overall χ^2 for interaction (p value)	19.128 (p<0.001)	0.041 (p=0.99)	50.195 (p<0.001)	0.992 (p=0.91)		31.96 (p<0.0001)	17.42 (p=0.001)	157.008 (p<0.0001)	2.665 (p=0.615)	
<i>Random part of the model</i>										
χ^2 for level 3: state (p value)	12.91 (p=0.011)	12.91 (p=0.011)	12.91 (p=0.004)	12.91 (p=0.011)		12.94 (p=0.011)	12.94 (p=0.011)	12.94 (p=0.004)	12.94 (p=0.011)	
χ^2 for level 2: local area (p value)	260.98 (p<0.001)	266.3 (p<0.001)	264.9 (p<0.001)	263.5 (p<0.001)		844.91 (p<0.0001)	839.3 (p<0.0001)	824.92 (p<0.0001)	837.64 (p<0.0001)	

*Models are controlled for age (centred at 29 years), marital status and religion.




Figure 3 Probability of smoking and chewing among men and women by education and wealth. NFHS, National Family Health Survey.

of masculinity among men and has moralistic connotations for women.^{19–21} *Second*, an economic perspective explaining the lower smoking rates among women in India may attribute this statistic to women’s unequal participation in the labour market and limited access to personal disposable income. Higher smoking among women in cities may partly indicate greater uptake of smoking by employed women. *Third*, given that data for this analysis come from a reproductive health survey, it is possible that results for women are an underestimate. Web table 1 indicates that older (above 35 years of age) women are more likely to use tobacco. However, patterns in this analysis match findings from other tobacco studies such as the GATS (global adult tobacco survey) in India (IIPS),^{5 6} providing a counter to this argument.

Finally, evidence on the socioeconomic gradient in tobacco use in India needs to be linked to its implications for tobacco-related mortality and morbidity.^{14 15} Dikshit *et al*¹⁵ provide the first estimates of cancer mortality in India, attributing a major component of age-standardised cancer mortality from lung and oral cancers to high rates of tobacco use in India. Analyses stratifying cancer morbidity and mortality in India by SES have not yet been attempted, and the lack of reliable surveillance data for chronic diseases prevent exhaustive assessments of the impact of tobacco use on Indian current and future chronic disease burden.²²

This study provides a systematic examination of the socioeconomic patterns in tobacco use in India over time. Data in this analysis come from the NFHS, which is a large, representative and generalisable survey, providing a comparative picture of tobacco patterns over time. The

limitations of this analysis are the following. First, the surveys are cross-sectional, hence limiting scope for causal inference. NFHS does not provide detailed data by type or volume of tobacco. Finally, NFHS is a reproductive health survey where women in the ages of 15–49 years are sampled. Men are sampled in the households of the female sample. This introduces the potential for two downward biases. The first pertains to *social desirability bias* particularly related to underestimation of smoking patterns in women’s childbearing years. Second, since the sample of men is conditional on the households from which women were sampled, the pool of men sampled may not be representative. Despite these caveats, NFHS (and in general the demographic and health surveys (DHSs)) has proven to be representative and generalisable, and continues to be used in a number of studies related to tobacco.^{5 6} In addition, our findings are consistent with estimates from studies using other surveys assessing the burden of tobacco and its drivers in India.^{2 23} Tobacco burden in the ‘productive’ populations (14–50 years) not only represents the current burden of tobacco but may predict future morbidity.

We present empirical evidence that India is experiencing a unique economic and social transition in tobacco consumption, quite distinct from the experience of developed countries that is likely to manifest in a number of morbidities.^{2 14 15} In order to ensure policy effectiveness to prevent and reduce the exposure to tobacco, there is a need to systematically monitor and examine the social inequities in tobacco use over time and channel interventions to the social groups that are most vulnerable to these inequalities.

Author affiliations

¹Department of Society, Human Development and Health, Harvard School of Public Health, Harvard University, Boston, Massachusetts, USA

²Public Health Foundation of India (PHFI), New Delhi, India

³South Asia Network for Chronic Diseases (SANCD), New Delhi, India

⁴Center for Integrative Approaches to Health Disparities, School of Public Health, University of Michigan, Michigan, USA

⁵Department of Primary Care and Social Medicine, School of Public Health, Imperial College London, London, UK

Acknowledgements We acknowledge the support of Macro International Inc (Washington, DC) for providing access to the surveys.

Contributors SVS and NB conceptualised and planned the study. NB led the data analysis, interpretation and writing of the manuscript. SS, MAS and SA contributed to data analysis and writing of the manuscript. CJM and SS contributed to the interpretation and writing of the manuscript. SVS contributed to the interpretation and writing of the manuscript, and provided the overall supervision.

Funding NB is supported by the Presidential Scholarship at Harvard University. SA is supported by a Wellcome Trust Strategic Award Grant No. Z/041825. CM conducted this work during a Study Abroad Fellowship funded by the Leverhulme Trust. CM is also funded by the Higher Education Funding Council for England and the National Institute for Health Research.

Competing interests None.

Ethics approval Secondary analysis of open-source datasets.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement Data for the analysis comes from the India chapter of the Demographic and Health Surveys (DHS) and is freely available online at the Measure DHS website. It is also available from the International Institute for Population Sciences (IIPS), Mumbai, India.

REFERENCES

- Reddy KS, Prabhakaran D, Jeemon P, *et al.* Educational status and cardiovascular risk profile in Indians. *Proc Natl Acad Sci* 2007;104.41:16263–8.
- WHO. *WHO report on the global tobacco epidemic, 2008: the MPOWER package*. Geneva: World Health Organization, 2008.
- Murray CJL, Lopez AD. Global mortality, disability and the contribution of risk factors: global burden of disease study. *Lancet* 1997a;349:1436–42.
- Murray CJL, Lopez AD. Alternative projections of mortality and disability by cause 1990–2020: global burden of disease study. *Lancet* 1997b;349:1498–504.
- Rani M, Bonu S, Jha P, *et al.* Tobacco use in India: prevalence and predictors of smoking and chewing in a national cross sectional household survey. *Tobacco Control* 2003;12.4:e4.
- Subramanian SV, Nandy S, Kelly M, *et al.* Patterns and distribution of tobacco consumption in India: cross sectional multilevel evidence from the 1998–9 National Family Health Survey. *BMJ* 2004;328.7443:801–6.
- John RM, Rao RK, Rao MG, *et al.* *The economics of tobacco and tobacco taxation in India*. Paris: International Union against Tuberculosis and Lung Disease, 2010.
- IIPS, & ORC Macro. *National Family Health Survey (NFHS-2), 1998–99. Data*. Mumbai, India: IIPS, 2000.
- IIPS, & ORC Macro. *National Family Health Survey (NFHS-3), 2005–06: India: Vol I. Data*. Mumbai, India: IIPS, 2007.
- Lopez AD, Collishaw NE, Piha T. A descriptive model of the cigarette epidemic in developed countries. *Tobacco Control* 1994;3:242–7.
- Thun M, Peto R, Boreham J, *et al.* Stages of the cigarette epidemic on entering its second century. *Tobacco Control* 2011;21:96–101.
- Sinha DN, Palipudi KM, Rolle I, *et al.* Tobacco use among youth and adults in member countries of South-East Asia region: review of findings from surveys under the Global Tobacco Surveillance System. *Indian J Public Health* 2011;55:169–76.
- GOI. *Global Adult Tobacco Survey. GATS India 2009–10. Report*. Ministry of Health & Family Welfare, Government of India, 2010.
- Jha P, Jacob B, Gajalakshmi V, *et al.* A nationally representative case-control study of smoking and death in India. *N Engl J Med* 2008;358:1137–47.
- Dikshit R, Gupta PC, Ramasundarahettige C, *et al.* Cancer mortality in India: a nationally representative survey. *Lancet* 2012;379:1807–16.
- Critchley JA, Unal B. 2003. Health effects associated with smokeless tobacco: a systematic review. *Thorax* 2003;58:435–43.
- WHO. *WHO Report on the Global Tobacco Epidemic, 2008: The MPOWER package. Gender, Women and the Tobacco Epidemic. Summary and Overview*. Geneva: World Health Organization, 2008.
- Fernandez E, Garcia M, Schiaffino A, *et al.* Smoking initiation and cessation by gender and educational level in Catalonia, Spain. *Prev Med* 2011;32:218–23.
- Viswanath K, Ackerson LK, Sorensen G, *et al.* 2010. Movies and TV influence tobacco use in India: findings from a national survey. *PLoS One* 2010;5:e11365.
- Stigler M, Dhavan P, Van Dusen D, *et al.* Westernization and tobacco use among young people in Delhi, India. *Soc Sci Med* 2010;71:891–7.
- Shah PB, Pednekar MS, Gupta PC, *et al.* The relationship between tobacco advertisements and smoking status of youth in India. *Asian Pac J Cancer Prev* 2008;9:637–42.
- Ebrahim S. Surveillance and monitoring for chronic diseases: a vital investment. *Natl Med J India Editorial* 2011;22.3:129–32.
- Palipudi KM, Gupta PC, Sinha DN, *et al.* Social determinants of health and tobacco use in thirteen low and middle income countries: evidence from Global Adult tobacco Survey. *PLoS One* 2012; 7:e33466.