

# Two Outer Membrane Lipoproteins from *Histophilus somni* Are Immunogenic in Rabbits and Sheep and Induce Protection against Bacterial Challenge in Mice

Carolina Guzmán-Brambila,<sup>a</sup> Argelia E. Rojas-Mayorquín,<sup>b,c</sup> Beatriz Flores-Samaniego,<sup>d</sup> and Daniel Ortuño-Sahagún<sup>a</sup>

Laboratorio de Desarrollo y Regeneración Neural, Instituto de Neurobiología, Departamento de Biología Celular y Molecular, CUCBA, Universidad de Guadalajara, Las Aguas, Zapopan, Jalisco, Mexico<sup>a</sup>; Departamento de Ciencias Ambientales, Instituto de Neurociencias, CUCBA, Universidad de Guadalajara, Guadalajara, Mexico<sup>b</sup>; Departamento de Investigación Básica, Instituto Nacional de Geriátrica, Mexico City, Mexico<sup>c</sup>; and Laboratorio de Investigación y Desarrollo Bio-Zoo, S.A. de C.V., Guadalajara, Mexico<sup>d</sup>

***Histophilus somni* is an economically important pathogen of cattle and other ruminants and is considered one of the key components of the bovine respiratory disease (BRD) complex, the leading cause of economic loss in the livestock industry. BRD is a multifactorial syndrome, in which a triad of agents, including bacteria, viruses, and predisposing factors or “stressors,” combines to induce disease. Although vaccines against *H. somni* have been used for many decades, traditional bacterins have failed to demonstrate effective protection in vaccinated animals. Hence, the BRD complex continues to produce strong adverse effects on the health and well-being of stock and feeder cattle. The generation of recombinant proteins may facilitate the development of more effective vaccines against *H. somni*, which could confer better protection against BRD. In the present study, primers were designed to amplify, clone, express, and purify two recombinant lipoproteins from *H. somni*, p31 (Plp4) and p40 (LppB), which are structural proteins of the outer bacterial membrane. The results presented here demonstrate, to our knowledge for the first time, that when formulated, an experimental vaccine enriched with these two recombinant lipoproteins generates high antibody titers in rabbits and sheep and exerts a protective effect in mice against septicemia induced by *H. somni* bacterial challenge.**

Bovine respiratory disease (BRD) is a major cause of economic losses in the livestock industry (15, 47). Technological, biological, and pharmacological advances have facilitated the development of several products to combat BRD, including vaccines. Nevertheless, the bacterial component of the BRD complex continues to provoke important adverse effects on the health and well-being of stock and feeder cattle (14, 20, 33).

BRD is a multifactorial syndrome, in which a triad of agents, including bacteria, viruses, and predisposing factors or “stressors,” combines to induce disease (reviewed in reference 49). BRD involves complex interactions among viral and bacterial pathogens that can lead to intense pulmonary inflammation (fibrinous pleuropneumonia) (10). *Histophilus somni* is among the bacterial pathogens associated with this disease (formerly *Haemophilus somnus*) (1), an unencapsulated Gram-negative coccobacillus and member of the *Pasteurellaceae* family that exhibits strict ruminant host specificity (9, 23, 39). *H. somni* is considered one of the key components of the BRD complex (11), and it is an economically important pathogen that causes respiratory disease, septicemia, thrombotic meningoencephalitis, myocarditis, arthritis, and abortion (9, 37, 44). *H. somni* is an opportunistic normal habitant of the lower reproductive tract and upper respiratory tract of cattle and other ruminants (9, 39), and it acts as a pathogen under certain circumstances via mechanisms that remain poorly understood (55).

Pathogens involved in BRD have developed intricate mechanisms to thwart both the innate and adaptive immune responses of their hosts. These immune evasive strategies probably contribute to the failure of currently available vaccines to provide complete protection against these pathogens (24, 29, 48). Consequently, the combination of multiple antigens in an enriched vaccine may provide more effective protection.

Lipooligosaccharide and various outer membrane proteins (OMPs) have been proposed as potential virulence factors of *H. somni*, although experimental evidence is lacking (7). A comparative genomic analysis of virulence factors from two *H. somni* strains (2336 and 129Pt) identified genes and gene products that are putatively involved in *H. somni* virulence, placing particular emphasis on those that were common to both strains (44). Nonetheless, a recent detailed sequence analysis of the *H. somni* transcriptional map (28) suggests that other immune-active proteins may also be involved.

To identify potentially protective antigens to formulate an experimental vaccine against *H. somni* infection during BRD, we selected two OMPs and evaluated their antigenicity/immunogenicity as immunogenic proteins. Several immunodominant surface antigens of *H. somni* were identified previously (8), including a 40-kDa protein (p40) that was subsequently identified and cloned (50), and designated LppB for lipoprotein B (GenBank accession no. AAA72348.1). According to domain analyses performed using the Motif Scan function of MyHits (38), this protein contains a prokaryotic membrane lipoprotein lipid attachment site profile from amino acids 1 to 17, and a LysM domain from amino acids 120 to 163 that is also found in a variety of enzymes involved in bacterial cell wall degradation (25). Indeed, the struc-

Received 23 July 2012 Returned for modification 28 August 2012


Accepted 6 September 2012

Published ahead of print 12 September 2012

Address correspondence to Daniel Ortuño-Sahagún, dortuno@cucba.udg.mx.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/CVI.00451-12



**FIG 1** (A) p40 protein domain structure. The prokaryotic membrane lipoprotein lipid attachment site (a), LysM domain (b), and peptidase family M23 domain (c) are indicated. (B) p31 protein domain structure. The prokaryotic membrane lipoprotein lipid attachment site (d), SmpA/OmlA family signature domain (e), and OmpA-like (outer membrane protein) domain signature (f) are indicated. The sequences were obtained from GenBank (accession numbers [AAA72348.1](#) and [AAA24941.1](#), respectively). Analyses were performed by Motif Scan using MyHits (35).

ture of this domain was defined (3), and it may have a general peptidoglycan binding function. Finally, LppB contains a peptidase family M23 domain from amino acids 243 to 263 (Fig. 1A).

Another relevant immunogenic protein, p31, has also been identified (52) (GenBank accession no. [AAA24941.1](#)), which shares sequence homology with the *plpD* gene that encodes a 31-kDa lipoprotein (Plp4) present in *Pasteurella haemolytica* A1 (34), and with a 19.2-kDa antigen from *Neisseria meningitidis*. A domain analyses performed using Motif Scan from MyHits (38) demonstrated that p31 possesses 273 amino acids and that it contains a prokaryotic membrane lipoprotein lipid attachment site from amino acids 1 to 19 and a SmpA/OmlA family signature domain from amino acids 19 to 122. This latter domain is found in some bacterial outer membrane lipoproteins and may be involved in maintaining the structural integrity of the cell envelope (36). In addition, this protein possess an OMPA-like domain signature from amino acids 129 to 251, which is thought to be responsible for noncovalent interactions with peptidoglycan (21) (Fig. 1B).

While vaccines against *H. somni* have been available for several decades, traditional bacterins do not work properly and have failed to demonstrate effective protection in vaccinated animals. A multivalent vaccine for bovine bacterial respiratory disease was developed to simplify the vaccination schedule and increase the range of protection, composed of two immunogens and five bacterins (including attenuated *H. somni*) (5). However, similarly developed commercial vaccines produce diverse secondary effects due to the high lipooligosaccharide content of the cell walls of Gram-negative bacteria. Consequently, the generation of recombinant proteins may facilitate the formulation of recombinant vaccines against *H. somni* that could confer better protection against BRD.

We previously formulated an experimental vaccine using two fragments from *Mannheimia haemolytica* OMPs that elicited high antibody titers in rabbits and sheep and that protected mice against bacterial challenge (22). However, since BRD is a complex pathology a multivalent vaccine is more desirable, both to increase the protective range and to simplify the vaccination schedule. Hence, in the present study we selected two lipoproteins from *H. somni*, p31 (homologous to Plp4) and p40 (LppB), to induce antibodies with which to formulate an experimental vaccine to protect against *H. somni* challenge. As a vaccine adjuvant we used

aluminum hydroxide [Al(OH)<sub>3</sub>], which stimulates immunity (13) by potentiating the immune response (46) and serves as a useful alternative to cholera toxin or Freund incomplete adjuvant. Our results clearly show that the combination of the p31 and p40 fusion proteins in this experimental vaccine formulation yielded high antibody titers in rabbits and sheep and protected mice from *H. somni* infection.

(The results presented here are included in a masters and doctoral thesis prepared by Carolina Guzmán-Brambila at the Universidad de Guadalajara, Guadalajara, Mexico.)

## MATERIALS AND METHODS

**Bacterial strains and culture conditions.** *Escherichia coli* (TOP10 or M15; Invitrogen) was used as the host to clone and propagate plasmids, which were cultured in Luria-Bertani (LB) broth supplemented with thymine (50 mg/ml) and ampicillin (100 mg/ml), chloramphenicol (25 mg/ml), or kanamycin (50 mg/ml), as necessary. *H. somni*, obtained from the American Type Culture Collection (strain ATCC 2336), was used to obtain bacterial DNA for PCR.

**Extraction and quantification of genomic DNA and recombinant methods.** Bacterial genomic DNA was obtained using the Illustra bacteria genomic Prep Mini spin kit (GE Healthcare, United Kingdom). DNA integrity was routinely evaluated by electrophoresis in agarose gels stained with Safe DNA gel stain (Invitrogen), and its quality was evaluated by determining the absorbance ratio at A<sub>260</sub>/A<sub>280</sub>. All DNA cloning and ligation was carried out according to standard recombinant DNA techniques (2, 43).

**Oligonucleotide design and PCR.** Oligonucleotide sequences of the primers used for PCR were as follows. The 5'-GGTAGGCCTATGAACTATCACGTTTTGTAT-3' sense primer and the 5'-TAATCTCTCTTGA TATAGGTAAGCTTATCA-3' antisense primer were used to amplify an 852-bp fragment encoding p31. The 5'-TAAAGTAACGGAGAATTAC ATGAA-3' sense primer and 5'-TTAATAAAGCTTAAATTACCATATC CACG-3' antisense primer were used to amplify an 870-bp fragment that encoded p40. Primers were designed with the *Stu*I restriction enzyme site in both forward primers and *Hind*III in both reverse primers to achieve directional cloning in the expression vector.

PCR was performed using a Perkin-Elmer GeneAmp PCR System 2400 thermocycler (Perkin-Elmer, Foster City, CA). For all PCR experiments, Platinum PCR SuperMix High Fidelity (Invitrogen) was used. PCR products were visualized on 1.5% agarose gels stained with Safe DNA gel stain (Invitrogen). Optimal annealing temperatures were initially established by testing a temperature gradient. All PCRs were performed after a single denaturation step at 94°C for 5 min and involved 30 denaturation cycles at 94°C for 1 min, annealing for 1 min at 62°C for the p31 primers and at 58°C for the p40 primers, and extension at 72°C for 1 min, followed by a final extension at 72°C for 5 min.

**Construction of p31-pQE 30 Xa and p40-pQE 30 Xa and molecular cloning.** To clone the PCR products, fragments were purified from agarose gels using the GFX PCR DNA and a gel band purification kit (GE Healthcare) and cloned directly using the pCR TOPO 2.1 vector (Invitrogen). TOP10 bacteria were transformed by the CaCl<sub>2</sub> method, and positive clones were selected using the β-galactosidase reaction. Plasmid DNA was obtained from *E. coli* using the plasmid Prep Mini spin kit (GE Healthcare) and, for each insert, at least three clones were sequenced to verify the sequence of both genes. Only clones that precisely matched the reported sequence were used for subcloning and to express the fusion protein. Sequences were obtained in a capillary ABI Prism 310 sequencer (Applied Biosystems) and were compiled with the Chromas v2.31 and DNASTAR, Inc., software packages before they were compared to the sequences in databases using the BLAST program.

The pQE 30 Xa vector (Qiagen) was used to generate the fusion protein, which introduces a six-histidine tag and a site for factor Xa cleavage. Previously cloned plasmid DNA with the PCR inserts was digested and religated at the *Stu*I and *Hind*III sites located at the ends of the p31 and

TABLE 1 Rabbit and sheep groups

Group	Rabbit immunogen	Sheep immunogen	Vaccination times (days)	Sampling times (days) for blood (sera)
A	Biobac 11 vías	Biobac 11 vías	0, 14	0, 14, 21, 28, 35, 42
B	Recombinant (30 µg)	Recombinant (50 µg)	0, 14	0, 14, 21, 28, 35, 42
C	Recombinant (60 µg)	Recombinant (100 µg)	0, 14	0, 14, 21, 28, 35, 42
D	Biobac 7 vías	Biobac 7 vías	0, 14	0, 14, 21, 28, 35, 42
E	None [PBS + Al(OH) <sub>3</sub> ]	None [PBS + Al(OH) <sub>3</sub> ]	0, 14	0, 14, 21, 28, 35, 42

p40 sequences, thereby generating two constructs expressing fusion proteins, designated p31-pQE 30 Xa and p40-pQE 30 Xa, and theoretically expressing 32.4- and 32-kDa derivatives, respectively. These constructs were transformed into *E. coli* M15 (Qiagen) to express the fusion proteins.

**Fusion protein expression by IPTG and purification.** IPTG (isopropyl-β-D-thiogalactopyranoside) was used to express the fusion proteins. Bacterial cultures were grown overnight, of which 1 ml was used to inoculate 50 ml of LB medium and allowed to grow for 2 h. Next, 1 mM IPTG was added, and the culture was grown for a further 3 h. Finally, the crude total extracts were obtained and examined by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE).

Fusion proteins were purified using the QIAexpress system (Qiagen), based on the affinity of the histidine tag included in the fusion protein for nickel-nitrilotriacetic acid (Ni-NTA) resin. Acid or basic buffers were then used to elute fusion protein from the column in 500-µl fractions and factor Xa was used to isolate the protein from the total extracts. The total protein was quantified by the Lowry method (31), and a small aliquot of the extract (5 µg of total protein) was examined by SDS-PAGE to confirm the presence of proteins corresponding to the p31 and p40 fusion proteins.

**Production of anti-p31 and anti-p40 antibodies in rabbits.** To prepare the immunogen, the purified proteins in the elution buffer (8 M urea, 0.01 M Tris-HCl, 0.1 M NaH<sub>2</sub>PO<sub>4</sub>) were mixed (1/8 [vol/vol]) with the adjuvant [0.25% Al(OH)<sub>3</sub>] in sterile phosphate-buffered saline (PBS; pH 7.2). Once prepared, the immunogen was subcutaneously inoculated into New Zealand White rabbits weighing approximately 2 kg. Two rabbits were subcutaneously inoculated with each fusion protein (90 µg) on days 0, 14, and 21, and serum was collected prior to the first inoculation on day 0 (preimmune serum control) and after each inoculation (days 14 and 21). Finally, total serum was obtained by complete exsanguination of the rabbit on day 31. The crude serum was heat inactivated and stored at -20°C until it was analyzed in Western blots.

**Protein electrophoresis and Western blotting.** Western blots were used to detect the production of antibodies against p31 and p40 in rabbits immunized with the purified fusion proteins. Crude extracts of total protein were obtained in lysis buffer (PBS [pH 7.2], 4% SDS), and the homogenates were then centrifuged at 14,000 × g for 15 min at 4°C. The supernatants were collected, and the protein content was determined using the Lowry method. Samples were diluted in PBS (pH 7.2) containing 4% SDS, denatured by boiling, and separated by electrophoresis on 10% acrylamide gels. The proteins were then transferred onto nitrocellulose membranes (Millipore, Bedford, MA), and the membranes were blocked overnight at 4°C with 80 g/liter of nonfat milk in 0.1% PBS-Tween-Tris-buffered saline (TBST). After the membranes were incubated with crude rabbit serum (1:1,000) for 1 h at room temperature, the blots were washed thoroughly in TBST and incubated for 1 h with horseradish peroxidase (HRP)-conjugated anti-rabbit IgG (1:5,000; Millipore). Immunoreactive proteins were detected by ECL (ECL Western blot analysis system; GE Healthcare) and analyzed using ChemiDoc (Bio-Rad Laboratories, Hercules, CA).

**Vaccination and production of polyclonal antibodies in rabbits and sheep.** Twenty-five New Zealand White rabbits weighing ~2 kg were divided into five groups (Table 1). The experimental vaccines (500 µl of

each) were subcutaneously inoculated on days 0 and 14. Group A (positive control) was inoculated with a commercial bacterin (Biobac 11 vías) composed of *Clostridium chauvoei*, *Clostridium septicum*, *Clostridium novyi*, *Clostridium sordellii*, *Clostridium perfringens* type C, *Clostridium perfringens* type D, *Pasteurella multocida* type A, *Pasteurella multocida* type D, *Mannheimia haemolytica* serotype A1, *Histophilus somni*, and adjuvant [Al(OH)<sub>3</sub>]. Group B was vaccinated with the recombinant preparation containing 30 µg of recombinant p31 (rp31) and 30 µg of recombinant p40 (rp40) and complemented with the commercial vaccine Biobac 7 vías (*Clostridium chauvoei*, *Clostridium septicum*, *Clostridium novyi*, *Clostridium sordellii*, *Clostridium perfringens* type C, *Clostridium perfringens* type D, and Al(OH)<sub>3</sub>). Rabbits of group C were inoculated with a recombinant preparation composed of 60 µg of rp31 plus 60 µg of rp40 plus Biobac 7 vías. Group D (negative control) was inoculated with Biobac 7 vías, and group E (negative control) was inoculated with adjuvant alone.

For the second experiment, 10 healthy female sheep (hybrids of Pelibuey, Katahdin, and Blackbelly) weighing ~30 kg, were housed in corrals and dewormed orally with 5% closantel. The animals were divided into five groups and vaccinated in the lateral neck region on days 0 and 14 with 2.5 ml of each formulation. The groups were as follows: group A (positive control) was immunized with the recombinant preparation alone; group B was immunized with recombinant preparation containing 50 µg of rp31, 50 µg of rp40, and Biobac 7 vías; group C was immunized with a recombinant preparation containing 100 µg of rp31, 100 µg of rp40, and Biobac 7 vías; group D (negative control) was immunized with Biobac 7 vías; and group E (negative control) with PBS and Al(OH)<sub>3</sub>.

For antibody determination, serum samples were collected on days 0, 14, 21, 28, 35, and 42 and assayed for rp31, rp40, and *H. somni* whole bacterial cell crude protein extract (WC) using an enzyme-linked immunosorbent assay (ELISA). Protein extracts (WC) were prepared as described previously, and 2 µg of total protein was assayed for each group.

**ELISA.** Polystyrene microplates (96-well) were coated separately with *H. somni* WC in lysis buffer and recombinant proteins. The plates were coated with 100 ng of each protein/well diluted in carbonate buffer (0.1 M sodium carbonate, 0.1 M bicarbonate sodium [pH 9.6]), followed by incubation for 16 to 18 h at 4°C. The supernatant was removed, and the plates were washed four times in buffer (1.25 M NaCl, 250 mM Tris-HCl [pH 7.9], 1% Tween 20), blocked with 2% nonfat milk in 0.1% TBST, and incubated at 37°C for 1 h. Subsequently, the plates were washed, and the WC, rp31, and rp40 samples were applied at a 1:800 dilution. After incubation for 1 h at 37°C, the microplates were washed four times, followed by incubation with the secondary antibody (HRP-conjugated anti-rabbit IgG [1:1,000]; Millipore) for 1 h at 37°C. After *o*-phenylenediamine (OPD; Sigma) was added as a peroxidase substrate, the plates were incubated for 5 min at room temperature, and the absorbance was then read at 415 nm. The results were reported as the median values from independent determinations each performed in triplicate and corrected versus the background control. To determine the levels of anti-rp31 and anti-rp40 for the sheep IgG antibodies, the previous steps were repeated diluting the serum 1:800 and using HRP-conjugated rabbit anti-sheep IgG (H+L, 1:1,000; Kirkegaard & Perry Laboratories, Inc., Gaithersburg, MD) as the secondary antibody.






FIG 2 Expression and purification of p31-pQE 30 Xa. (A) Induction of fusion protein expression; (B) protein purification after elution from the column.

**Bacterial challenge in BALB/c mice.** As previously demonstrated (17, 18, 45), *H. somni* causes septicemia when inoculated intraperitoneally in mice. Therefore, we selected this model to test for the efficacy of our experimental vaccine.

Seventy-five male BALB/c mice weighing 26 to 32 g were housed under controlled conditions of temperature, humidity, and lighting. The mice were divided into five groups and injected intraperitoneally on days 0 and 14 with 250  $\mu$ l of each formulation. Group A (positive control) was immunized with Biobac 11 vias. Group B was immunized with a recombinant preparation containing 10  $\mu$ g of rp31, 10  $\mu$ g of rp40, and Biobac 7 vias. Group C was immunized with a recombinant preparation containing 20  $\mu$ g of rp31 plus 20  $\mu$ g of rp40 plus Biobac 7 vias, and the negative control groups D and E were immunized with Biobac 7 vias and PBS-Al(OH)<sub>3</sub>, respectively. Since the final objective was to formulate and test a combined experimental vaccine, we did not include groups to test each recombinant protein individually.

All mice were challenged 28 days after the initial immunization with approximately  $3.3 \times 10^8$  CFU of virulent *H. somni* (ATCC 2336), injected intraperitoneally, and their survival was monitored for 10 days. Necropsies were performed to verify the effect of the bacterial challenge.

**Animal care and use.** The present study complied with all Institutional Guidelines and the Official Mexican Regulations (35) for the production, care, and use of laboratory animals. The protocol was approved by the local Animal Ethics Committee. Every effort was made to minimize animal suffering and the number of animals used.

**Statistical analyses.** The Fisher exact test was used to evaluate the protection against *H. somni*, and the results are presented as a Kaplan-Meier curve. ELISA results were compared by analysis of variance.

**RESULTS**

**Induction of rp31-pQE 30 Xa and rp40-pQE 30 Xa expression.** PCR fragments from of each gene were initially cloned into pCR 2.1 TOPO (Invitrogen), digested and the gel-purified StuI-HindIII restriction fragments were subcloned into the pQE-30 Xa expression vector (Qiagen). The fusion proteins were expressed in M-15 *E. coli* (Invitrogen) transformed by heat shock.

In the case of p31, the whole protein was cloned (Fig. 1A) as a 283-amino-acid polypeptide to generate a fusion protein with a theoretical molecular mass of 32 kDa that included the histidine tag and the Xa factor recognition site. The fusion protein was clearly induced by IPTG, and its electrophoretic migration corresponded well with the theoretical molecular mass (Fig. 2A). The




FIG 3 Expression and purification of p40-pQE 30 Xa. (A) Induction of fusion protein expression; (B) protein purification after elution from the column.

construct used for p40 (LppB) contained 289 amino acids and included the tag and protease recognition site. The resulting fusion protein had a theoretical molecular mass of 32 kDa. Although this protein was also clearly induced by IPTG, it had a slightly higher electrophoretic mobility than expected (Fig. 3A).

**Fusion protein purification and antibody production.** Fusion proteins were purified from the crude extracts using the QIA-express assay system (Qiagen), and they were analyzed by SDS-PAGE (Fig. 2B and 3B). Three New Zealand White rabbits were immunized with 30  $\mu$ g of each protein (a total of 90  $\mu$ g) on days 0, 14, and 21 after collecting preimmune serum prior to the first immunization as an internal control. Ten days after the last immunization the rabbits were sacrificed and exsanguinated to obtain the final serum sample. All samples were analyzed in Western blots of total protein extracts, with or without the induced proteins (Fig. 4A [p31] and B [p40]).

**ELISA analysis of the antibody response.** Antibodies against *H. somni*, rp31 and rp40, were evaluated by ELISA in two experimental models (rabbits and sheep). Vaccination of rabbits with commercial bacterin (Biobac 11 vias) and recombinant vaccine formulations stimulated detectable levels of anti-p31 and anti-p40 antibodies (Fig. 5A), with significant increases in antibody pro-





FIG 4 (A and B) Western blot analysis of rabbit serum. Serum containing antibodies against p31 (A) and p40 (B). M, molecular mass marker; +IPTG, total bacterial extract in which expression was induced; P5, 5  $\mu$ g of the purified fusion protein; P2, 2  $\mu$ g of purified fusion protein.



**FIG 5** Results from ELISA analysis. (A) Rabbits. Group A (positive control) was inoculated with a commercial bacterin (Biobac 11 vías [see Materials and Methods]). Group B was vaccinated with a recombinant preparation containing 30  $\mu\text{g}$  of recombinant p31 (rp31) and 30  $\mu\text{g}$  of recombinant p40 (rp40) and complemented with the commercial vaccine Biobac 7 vías. Group C was inoculated with a recombinant preparation containing 60  $\mu\text{g}$  of rp31, 60  $\mu\text{g}$  of rp40, and Biobac 7 vías. Negative control groups D and E were inoculated with Biobac 7 vías and PBS-adjuvant  $[\text{Al}(\text{OH})_3]$ , respectively. (B) Sheep. Group A (positive control) was immunized with the recombinant preparation alone. Group B was immunized with a recombinant preparation containing 50  $\mu\text{g}$  of rp31, 50  $\mu\text{g}$  of rp40, and Biobac 7 vías. Group C was immunized with a recombinant preparation containing 100  $\mu\text{g}$  of rp31, 100  $\mu\text{g}$  of rp40, and Biobac 7 vías. Negative control groups D and E were inoculated with Biobac 7 vías and PBS-adjuvant  $[\text{Al}(\text{OH})_3]$ , respectively.

duction observed from day 21. For rp31, the antibody response was significantly greater in group C (rp31, 60  $\mu\text{g}$ ) than in groups A and B, whereas no antibody production was observed in groups D or E (negative controls). For rp40, the antibody response was significantly greater in group B (rp40, 30  $\mu\text{g}$ ) than in groups A and C, also was higher in group C (rp40, 60  $\mu\text{g}$ ) than in group A (commercial vaccine).

In the second experiment (Fig. 5B), serum samples from sheep were analyzed for rp31 and rp40 and for WC *H. somni*. Increases in anti-p31 and anti-rp40 were clearly observed from day 21 in groups B (rp31 + rp40, 50  $\mu\text{g}$ ) and C (rp31 + rp40, 100  $\mu\text{g}$ ) (recombinant vaccine formulations), with greater responses in the latter group for p31, when comparing with controls but also with commercial vaccine (group A). Increased levels of anti-WC *H. somni* were detected from day 14 in all groups treated with the commercial and recombinant vaccine formulations, at a similar level until day 42.


Immunization of rabbits and sheep with experimental vaccines complemented with recombinant proteins strongly stimulated the dose-dependent production of antibodies against p31 and p40 (Fig. 5). Hence, the recombinant proteins used in these experiments are clearly immunogenic and can be used to formulate vaccines against BRD.

**Bacterial challenge.** Protection against bacterial challenge with pathogenic *H. somni* was analyzed by the Fisher exact test, and the results are summarized on a Kaplan-Meier curve (Fig. 6). Mice were monitored for 10 days after challenge with  $3.3 \times 10^8$  CFU of virulent *H. somni*, and protection was observed in groups A, B, and C as follows: group A (Biobac 11 vías), 93% survival;

group B (immunized with a low dose of recombinant vaccine (rp31 + rp40, 10  $\mu\text{g}$  of each), 87% survival; and group C (immunized with a high dose of recombinant vaccine (20  $\mu\text{g}$  of rp31 + rp40), 100% survival). No protective effects were observed in the negative control groups (D and E), in which no survival was observed following the challenge. The protective effect of the commercial (Biobac 11 vías) and recombinant vaccines (containing rp31 and rp40) was significantly greater ( $P < 0.0001$ ) than that of the negative controls (Biobac 7 vías and adjuvant).

## DISCUSSION

BRD is generally detected in cattle raised on farms with poor or nonexistent cattle health management plans (20), but many feed-



**FIG 6** Result of challenge test in BALB/c mice. A Kaplan-Meier curve shows the percent survival over time.

lots, calf raising facilities, and other facilities with excellent management also still have a considerable BRD problem (14, 15, 20, 33, 47). Therefore, the use of vaccines as a preventative measure is important to prevent the spread of this disease (40). Although several commercial vaccines were developed prior to the 1990s (reviewed in reference 41), most caused secondary adverse reactions (12), emphasizing the need to develop recombinant vaccines in which the presence of more than one antigen may increase efficacy. The results presented here demonstrate that a recombinant experimental vaccine containing protein fragments of two OMP *H. somni* antigens, p31 (Plp4) and p40 (LppB), induced high antibody titers in mice and sheep. Furthermore, this vaccine formulation protected mice from septicemia after bacterial challenge. In addition, the use of Al(OH)<sub>3</sub> as an adjuvant potentiated the immune response, thereby maximizing the potency and efficacy of the antigens, which are generated in limited amounts (13), and enhancing the immune response (46).

**Protective effects of antibodies against the *H. somni* OMPs p31 (Plp4) and p40 (LppB).** The immunogenic potential of some *H. somni* proteins has been previously assessed, demonstrating that they can produce partial protection against bacterial infection. Among antigens most probably involved in stimulating host defense as well as immunopathology, OMPs are relevant as virulence factors (6, 9).

Proteins on the cell surface undoubtedly play an important role in *H. somni* virulence and host immunity (reviewed in reference 44). Because Gram-negative bacteria exhibit a high degree of genomic variability in some of its proteins, conserved OMPs became relevant immunogenic targets that are able to elicit cellular mechanisms of host defense involving the antigen-induced release of cytokines from lymphocytes and the resulting activation of macrophages with the ability to kill the pathogen.

For example, a 40-kDa OMP from *H. somni* was proposed as a candidate protective protein against pneumonia in calves following active immunization (19). However, according to a recent analysis of genes and gene products putatively involved in *H. somni* strain 2336 virulence, this protein corresponds to a different OMP than the p40 (LppB) used here (see Table 2 in reference 44). A 78-kDa OMP antigen that was also shown to be consistently and intensely immunoreactive in Western blots of *H. somni* WC reacted with convalescent-phase serum from cattle with experimental *H. somni* pneumonia (26). However, this antigen failed to elicit protective effects (19).

An immunoglobulin binding protein A (IbpA) containing a Fic motif involved in the virulence of several pathogens (42, 53) was recently described (30, 55) as a viable vaccine candidate in the bovine host. Immunization with the IbpA DR2 subunit from *H. somni* was demonstrated to partially protect against bacterial infection in a natural host (16, 30). However, further studies will be necessary to determine the immunogenic properties of the different virulence factors involved in *H. somni* infection, since the evidence for protection against pneumonia by current vaccines remains controversial (29, 48).

Although further immunogenic studies in beef cattle are required, the results presented here demonstrate for the first time that the antibody response in rabbits and in sheep against two *H. somni* OMPs, p31 (Plp4) and p40 (LppB), is relevant. In addition, when combined with a commercial vaccine for other bacterial diseases (Biovac 7 vías), recombinant fragments of p31 (Plp4) and p40 (LppB), which appear to be conserved structural proteins of

the outer bacterial membrane, exert a protective effect against *H. somni* bacterial challenge in mice.

**p31 and p40 structure and possible functional implications.** The electrophoretic mobility observed for the p40 fusion protein corresponded to that originally reported for this protein (8, 50), although it was slightly higher than theoretically expected. This protein is rich in proline (8%), asparagine (8.0%), and isoleucine (9.7%) compared to the average amino acid composition of vertebrate proteins (54), a profile that may influence its electrophoretic mobility. The fusion protein also contains some putative peptidoglycan binding sites (32), which may modify its relative molecular mobility. In addition, the increased electrophoretic mobility described here may be partially due to the addition of a histidine tag and the factor Xa recognition site. Nonetheless, when the clones were sequenced they precisely matched the reported sequence, and thus the induced fusion protein corresponds to p40.

It is of interest to determine whether the organization of the distinct domains in these two proteins are implicated in the effects of *H. somni* on endothelial cells and in the aggregation of platelets to form thrombi in blood vessels (19). These phenomena induce endothelial cell proinflammatory responses and platelet internalization (27), and they trigger cytoskeletal alterations that increase the permeability of the endothelium (4), resulting in the redistribution of PECAM 1 on the surfaces of bronchial endothelial cells (51).

Further studies are thus required with antibodies that identify p31 (Plp4) and p40 (LppB), such as those described here, that can be used to study their effects at the cellular level (e.g., by analyzing the antibody neutralization of these two proteins *in vitro*).

Finally, the basis of viral/bacterial synergism and the manner in which cattle respond to the virulence strategies of bacterial pathogens remain poorly understood (10). The ability of *H. somni* to resist leukocytes while creating a proinflammatory and procoagulation environment at the endothelial cell surface and the ability of *M. haemolytica* to circumvent leukocyte antibacterial activity via its leukotoxin LKTA probably contribute to the intense inflammation that characterizes BRD (10). As such, it seems feasible to propose that a vaccine that combines antigenic surface proteins from *M. haemolytica* (e.g., different fragments of LktA and Plp) (7, 22) and *H. somni* (e.g., p31 and p40, as demonstrated here) may be useful in preventing infection and reducing the incidence of BRD. Further experiments would be needed to test this hypothesis.

## ACKNOWLEDGMENTS

This study was partially supported by CONACYT grants 2007-C01-71496 to B.F.-S., 2009-111203 and 2010-129167 to D.O.-S., and 237076 to C.G.-B.

We thank Bio-Zoo laboratories for the use of their field-testing facilities.

## REFERENCES

1. Angen Ø, Ahrens P, Kuhnert P, Christensen H, Mutters R. 2003. Proposal of *Histophilus somni* gen. nov., sp. nov. for the three species incertae sedis “*Haemophilus somnus*,” “*Haemophilus agni*,” and “*Histophilus ovis*.” Int. J. Syst. Evol. Microbiol. 53:1449–1456.
2. Ausubel F, et al. 2001. Current protocols in molecular biology. John Wiley & Sons, Inc, New York, NY.
3. Bateman A, Bycroft M. 2000. The structure of a LysM domain from *Escherichia coli* membrane-bound lytic murein transglycosylase D (MltD). J. Mol. Biol. 299:1113–1119.
4. Behling-Kelly E, McClenahan D, Kim KS, Czuprynski CJ. 2007. Viable



- "*Haemophilus somnus*" induces myosin light-chain kinase-dependent decrease in brain endothelial cell monolayer resistance. *Infect. Immun.* 75:4572–4581.
5. Cho YS, et al. 2008. Safety and efficacy testing of a novel multivalent bovine bacterial respiratory vaccine composed of five bacterins and two immunogens. *J. Vet. Med. Sci.* 70:959–964.
  6. Confer AW. 2009. Update on bacterial pathogenesis in BRD. *Anim. Health Res. Rev.* 10:145–148.
  7. Confer AW, et al. 2009. Immunity of cattle following vaccination with a *Mannheimia haemolytica* chimeric PlpE-LKT (SAC89) protein. *Vaccine* 27:1771–1776.
  8. Corbeil LB, Kania SA, Gogolewski RP. 1991. Characterization of immunodominant surface antigens of *Haemophilus somnus*. *Infect. Immun.* 59:4295–4301.
  9. Corbeil LB. 2007. *Histophilus somni* host-parasite relationships. *Anim. Health Res. Rev.* 8:151–160.
  10. Czuprynski CJ. 2009. Host response to bovine respiratory pathogens. *Anim. Health Res. Rev.* 10:141–143.
  11. Ellis JA. 2001. The immunology of the bovine respiratory disease complex. *Vet. Clin. N. Am. Food Anim. Pract.* 17:535–550.
  12. Ellis JA, Yong C. 1997. Systemic adverse reactions in young Simmental calves following administration of a combination vaccine. *Can. Vet. J.* 38:45–47.
  13. Exley C, Siesjo P, Eriksson H. 2010. The immunobiology of aluminum adjuvants: how do they really work? *Trends Immunol.* 31:103–109.
  14. Fulton RW. 2009. Bovine respiratory disease research (1983–2009). *Anim. Health Res. Rev.* 10:131–139.
  15. Gagea MI, et al. 2006. Diseases and pathogens associated with mortality in Ontario beef feedlots. *J. Vet. Diagn. Invest.* 18:18–28.
  16. Geertsema RS, et al. 2011. IbpA DR2 subunit immunization protects calves against *Histophilus somni* pneumonia. *Vaccine* 29:4805–4912.
  17. Geertsema RS, Kimball RA, Corbeil LB. 2007. Bovine plasma proteins increase virulence of *Haemophilus somnus* in mice. *Microb. Pathog.* 42:22–28.
  18. Geertsema RS, et al. 2008. Protection of mice against *H. somni* septicemia by vaccination with recombinant immunoglobulin binding protein subunits. *Vaccine* 26:4506–4512.
  19. Gogolewski RP, Leathers CW, Liggitt, HD, Corbeil LB. 1987. Experimental *Haemophilus somnus* pneumonia in calves and immunoperoxidase localization of bacteria. *Vet. Pathol.* 24:250–256.
  20. Griffin D. 2010. Bovine pasteurellosis and other bacterial infections of the respiratory tract. *Vet. Clin. N. Am. Food Anim. Pract.* 26:57–71.
  21. Grizot S, Buchanan SK. 2004. Structure of the OmpA-like domain of RmpM from *Neisseria meningitidis*. *Mol. Microbiol.* 51:1027–1037.
  22. Guzmán-Brambila C, et al. 2012. LKTA and PlpE small fragments fusion protein protect against *Mannheimia haemolytica* challenge. *Res. Vet. Sci.* 93:1293–1300.
  23. Harris FW, Janzen ED. 1989. The *Haemophilus somnus* disease complex (hemophilosis): a review. *Can. Vet. J.* 30:816–822.
  24. Howard MD, et al. 2011. Genetics and molecular specificity of sialylation of *Histophilus somni* lipooligosaccharide (LOS) and the effect of LOS sialylation on Toll-like receptor-4 signaling. *Vet. Microbiol.* 153:163–172.
  25. Joris B, et al. 1992. Modular design of the *Enterococcus hirae* muramidase-2 and *Streptococcus faecalis* autolysin. *FEMS Microbiol. Lett.* 70:257–264.
  26. Kania SA, Gogolewski RP, Corbeil LB. 1990. Characterization of a 78-kilodalton outer membrane protein of *Haemophilus somnus*. *Infect. Immun.* 58:237–244.
  27. Kuckleburg CJ, McClenahan DJ, Czuprynski CJ. 2008. Platelet activation by *Histophilus somni* and its lipooligosaccharide induces endothelial cell proinflammatory responses and platelet internalization. *Shock* 29:189–196.
  28. Kumar R, et al. 2012. RNA-seq-based transcriptional map of bovine respiratory disease pathogen "*Histophilus somni* 2336." *PLoS One* 7:e29435. doi:10.1371/journal.pone.0029435.
  29. Larson RL, Step DL. 2012. Evidence-based effectiveness of vaccination against *Mannheimia haemolytica*, *Pasteurella multocida*, and *Histophilus somni* in feedlot cattle for mitigating the incidence and effect of bovine respiratory disease complex. *Vet. Clin. N. Am. Food Anim. Pract.* 28:97–106.
  30. Lo KL, et al. 2012. Antibody responses of calves to *Histophilus somni* recombinant IbpA subunits. *Comp. Immunol. Microbiol. Infect. Dis.* 35:453–459.
  31. Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. 1951. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* 193:265–275.
  32. Marchler-Bauer A, et al. 2011. CDD: a conserved domain database for the functional annotation of proteins. *Nucleic Acids Res.* 39:225–229.
  33. McVey DS. 2009. BRD research needs in the next 10–20 years. *Anim. Health Res. Rev.* 10:165–167.
  34. Nardini PM, Mellors A, Lo RY. 1998. Characterization of a fourth lipoprotein from *Pasteurella haemolytica* A1 and its homology to the OmpA family of outer membrane proteins. *FEMS Microbiol. Lett.* 165:71–77.
  35. Norma Oficial Mexicana. 2008. Especificaciones técnicas para la producción, cuidado y uso de los animales de laboratorio. Document NOM-062-ZOO-1999. Modified 26 November 2008. Senasica, Mexico City, Mexico. [www.senasica.gob.mx/default.asp?doc=743](http://www.senasica.gob.mx/default.asp?doc=743).
  36. Ochsner UA, Vasil AL, Johnson Z, Vasil ML. 1999. *Pseudomonas aeruginosa* fur overlaps with a gene encoding a novel outer membrane lipoprotein, OmlA. *J. Bacteriol.* 181:1099–1109.
  37. O'Toole D, Allen T, Hunter R, Corbeil LB. 2009. Diagnostic exercise: myocarditis due to *Histophilus somni* in feedlot and backgrounded cattle. *Vet. Pathol.* 46:1015–1017.
  38. Pagni M, et al. 2007. MyHits: improvements to an interactive resource for analyzing protein sequences. *Nucleic Acids Res.* 35:W433–W437.
  39. Pérez-Romero N, Aguilar-Romero F, Arellano-Reynoso B, Díaz-Aparicio E, Hernández-Castro R. 2011. Isolation of *Histophilus somni* from the nasal exudates of a clinically healthy adult goat. *Trop. Anim. Health Prod.* 43:901–903.
  40. Potter A, Gerds V, Littel-van den Hurk SD. 2008. Veterinary vaccines: alternatives to antibiotics? *Anim. Health Res. Rev.* 9:187–199.
  41. Rice JA, Carrasco-Medina L, Hodgins DC, Shewen PE. 2007. *Mannheimia haemolytica* and bovine respiratory disease. *Anim. Health Res. Rev.* 8:117–128.
  42. Roy CR, Mukherjee S. 2009. Bacterial FIC proteins AMP up infection. *Sci. Signal.* 2:pe14.
  43. Sambrook J, Fritsch EF, Maniatis T. 2001. Molecular cloning: a laboratory manual, 3rd ed. Cold Spring Harbor Laboratory Press, New York, NY.
  44. Sandal I, Inzana TJ. 2010. A genomic window into the virulence of *Histophilus somni*. *Trends Microbiol.* 18:90–99.
  45. Sanders JD, Bastida-Corcuera FD, Arnold KF, Wunderlich AC, Corbeil LB. 2003. Genetic manipulation of immunoglobulin binding proteins of *Haemophilus somnus*. *Microb. Pathog.* 34:131–139.
  46. Seubert A, Monaci E, Pizza M, O'Hagan DT, Wack A. 2008. The adjuvants aluminum hydroxide and MF59 induce monocyte and granulocyte chemoattractants and enhance monocyte differentiation toward dendritic cells. *J. Immunol.* 180:5402–5412.
  47. Snowden GD, Van Vleck LD, Cundiff LV, Bennett GL. 2006. Bovine respiratory disease in feedlot cattle: environmental, genetic, and economic factors. *J. Anim. Sci.* 84:1999–2008.
  48. Srikumaran S, Kelling CL, Ambagala A. 2007. Immune evasion by pathogens of bovine respiratory disease complex. *Anim. Health Res. Rev.* 8:215–229.
  49. Taylor JD, et al. 2010. The epidemiology of bovine respiratory disease: what is the evidence for predisposing factors? *Can. Vet. J.* 51:1095–1102.
  50. Theisen M, Rioux CR, Potter AA. 1993. Molecular cloning, nucleotide sequence, and characterization of *lppB*, encoding an antigenic 40-kilodalton lipoprotein of *Haemophilus somnus*. *Infect. Immun.* 61:1793–1798.
  51. Tiwari R, Sullivan J, Czuprynski CJ. 2009. PECAM-1 is involved in neutrophil transmigration across *Histophilus somni*-treated bovine brain endothelial cells. *Microb. Pathog.* 47:164–170.
  52. Won J, Griffith RW. 1993. Cloning and sequencing of the gene encoding a 31-kilodalton antigen of *Haemophilus somnus*. *Infect. Immun.* 61:2813–2821.
  53. Worby CA, et al. 2009. The *fic* domain: regulation of cell signaling by adenylylation. *Mol. Cell* 34:93–103.
  54. Wilkins MR, et al. 1996. From proteins to proteomes: large-scale protein identification by two-dimensional electrophoresis and amino acid analysis. *Biotechnology (NY)* 14:61–65.
  55. Zekarias B, et al. 2010. *Histophilus somni* IbpA DR2/Fic in virulence and immunoprotection at the natural host alveolar epithelial barrier. *Infect. Immun.* 78:1850–1858.