



Published in final edited form as:

J Am Chem Soc. 2012 October 24; 134(42): 17643–17652. doi:10.1021/ja307102g.

Promiscuous 8-Alkoxyadenosines in the Guide Strand of an siRNA: Modulation of Silencing Efficacy and Off-Pathway Protein Binding

Uday Ghanty, Erik Fostvedt, Rachel Valenzuela, Peter A. Beal, and Cynthia J. Burrows

Department of Chemistry, University of Utah, 315 S. 1400 East, Salt Lake City, UT 84112-0850, United States, and Department of Chemistry, University of California Davis, One Shields Ave., Davis, CA 95616, United States

Cynthia J. Burrows: burrows@chem.utah.edu

Abstract

8-Alkoxyadenosines have the potential to exist in *anti* or *syn* conformations around the glycosidic bond when paired opposite to U or G in the complementary strands, thereby placing the sterically demanding 8-alkoxy groups in the major or minor groove, respectively, of duplex RNA. These modified bases were used as ‘base switches’ in the guide strands of an siRNA to prevent off-pathway protein binding during delivery via placement of the alkoxy group in the minor groove, while maintaining significant RNAi efficacy by orienting the alkoxy group in the major groove. 8-Alkoxyadenosine phosphoramidites were synthesized and incorporated into the guide strand of caspase 2 siRNA at four different positions--two in the seed region, one at the cleavage junction and another nearer to the 3'-end of the guide strand. Thermal stabilities of the corresponding siRNA duplexes showed that U is preferred over G as the base-pairing partner in the complementary strand. When compared to the unmodified positive control siRNAs, singly modified siRNAs knocked down the target mRNA efficiently and with little or no loss of efficacy. Doubly modified siRNAs were found to be less effective and lose their efficacy at low nanomolar concentrations. SiRNAs singly modified at positions 6 and 10 of the guide strand were found to be effective in blocking binding to the RNA-dependent protein kinase PKR, a cytoplasmic dsRNA-binding protein implicated in sequence independent off target effects.

Introduction

Chemical modification of small interfering RNAs (siRNAs) is necessary to achieve potent RNA interference (RNAi) therapeutics. Modifications of the ribose moiety^{1–18} and phosphodiester backbone^{1,2,7} have improved siRNA stability,^{1,9} ribonuclease resistance,^{1,3,6,8,12} potency,^{3,8,17} and specificity.^{1,19} Polymer,^{20–22} lipid,²³ cholesterol,^{24–26} carbohydrate,^{20,21} peptide,²⁷ small molecule²⁴ terminal modifications and delivery systems of siRNAs exhibit effective RNAi and polymer- and cholesterol-based conjugates facilitated intracellular delivery of siRNA both *in vitro* and *in vivo*. Although more challenging, a few successful siRNA base modifications have also been reported.^{2,28–34} SiRNA base modifications³⁵ are relatively less explored compared to sugar modifications or terminal modifications, because preservation of necessary hydrogen-bonding interactions and the

Correspondence to: Cynthia J. Burrows, burrows@chem.utah.edu.

Supporting Information

Complete details of syntheses of modified phosphoramidites, proton and carbon NMR spectra, figure of the caspase 2 inserted sequence and ESI-MS masses of the modified guide strands This material is available free of charge via the Internet at <http://pubs.acs.org>.

stable A-form duplex are crucial in effective RNAi. Disruption of hydrogen bonding (e.g. in N^3 -Me-U) or steric occlusion of siRNA-RISC interactions (5-IU) can adversely affect the RNAi efficacy and specificity.² For this reason, most of the successful base modifications have been introduced into either passenger strands or at the 3'-end of guide strands.^{28,32,34}

Base modifications have been shown to enhance thermal stability³¹ and nuclease stability;³⁰ these also helped visualize intracellular trafficking³⁴ and reduced off-target effects.^{28,32,33} Kool and coworkers, and Manoharan, Egli and coworkers reported substitution of pyrimidines (uridine) in the guide strands of siRNAs with more hydrophobic aromatic ring systems (e.g. 2,4-difluorobenzene²⁹ and 2,4-difluorotoluene^{30,36}), although here silencing efficacy varied depending on the position of the substitution. Recently, major groove substitution of modified purines in various positions of the guide strand have been reported; here also RNAi efficacy was found to be position dependent.³⁷

Along with the targeted delivery issue, off-target effects are still one of the major limitations for specific and selective action of therapeutic siRNA *in vivo*. Off-target effects leading to immunostimulation pose a significant limitation on the therapeutic use of siRNA. Specific siRNA sequences can bind to toll-like receptors and lead to immunostimulation.³⁸⁻⁴² In addition, sequence independent off-target effects involve binding of virtually any siRNA with double-stranded RNA binding motif (dsRBM) containing proteins, e.g., RNA-dependent protein kinase (PKR), adenosine deaminases (ADARs) and other intracellular proteins.^{43,44} PKR activation leads to phosphorylation of eIF2 α , inhibition of translation initiation and other antiviral signaling events.⁴⁵ Although siRNAs are shorter than ligands that trigger high PKR activity, lower level activation of PKR has been observed with siRNAs *in vitro* and when transfected into certain cell types.³⁸ ADARs also bind dsRNA and have been shown to interact with substrates within the RNAi pathway.^{46,47} The full-length isoform of ADAR1 (ADAR1p150) has been linked to decreased siRNA potency in mammalian cells, presumably due to the formation of high affinity siRNA•ADAR1p150 complexes within the cytoplasm.⁴⁴ Additionally, ADAR has been shown to reduce RNAi efficiency in *Drosophila* cell culture.⁴⁷ Therefore, prevention of siRNA binding to dsRBM proteins off the RNAi pathway is important for potency and specificity of therapeutic RNAi. Some nucleoside analogs⁴⁸ have shown promise in preventing PKR binding, but a complete study with modified sugars to address such sequence-independent off-target effect is not available. On the other hand, altering the RNA minor groove through base modifications has shown potential in modulating siRNA properties and siRNA-protein interactions. The Beal laboratory initiated a systematic study on siRNA base modifications that could block off-pathway protein binding,³⁵ and they have also shown that pendent minor groove modifications of guanines²⁸ and 2-aminopurines³² in the passenger caspase 2 siRNA can successfully prevent siRNA-dsRBM interactions, while maintaining siRNA efficacy.

Base modifications in the guide strand are more challenging, since many modifications at crucial siRNA sites (e.g., in the middle of the seed region or at the cleavage site) will drastically reduce the silencing efficacy or might even completely abolish the knockdown of the desired mRNA. In contrast, modifications at those sites might, in fact, turn out to be more exciting and lead to elucidation of crucial mechanisms in the RISC complex. For example, appropriate modifications in the minor or major groove of an siRNA can help explore the siRNA mechanisms in detail, and placement of designer modifications in those grooves can provide further novel insights into the siRNA-RISC interactions. Our laboratories have previously shown that N^2 -alkylated 8-oxo-7,8-dihydro-2'-deoxyguanosine-containing siRNAs can substantially prevent siRNA-PKR interaction by placing a 'switchable' alkyl group in the minor groove of an siRNA during delivery; in the RISC, the modified siRNAs can flip the sterically encumbering group into the major groove and maintain good RNAi efficacy.³³ However 8-oxoguanine-containing DNA oligomers are

known to be inflammatory and immunostimulatory themselves.^{49,50} Therefore we chose to explore alternative purine modifications in the guide strand. Here we report the synthesis, thermal stability and biological activity of 8-alkoxyadenosine-substituted siRNAs.

8-AlkoxyA phosphoramidites were synthesized and incorporated into the guide strand of a siRNA targeting the caspase 2 mRNA sequence. Modified siRNAs were tested for their RNAi efficacy and ability to address off-target effects due to siRNA-PKR interactions. 8-Substituted adenosines and guanosines have been shown to exist in an equilibrium mixture of *syn/anti* conformers. In this work, 8-alkoxyadenosines are postulated to flip between *anti* and *syn* conformations depending on the base-pairing partner. In the natural *anti* conformation, 8-alkoxyA will base pair with U; whereas, in the *syn* conformation, the Hoogsteen face of the nucleoside will be exposed for base pairing, and its best complement would be *anti*G. We propose that during delivery of the siRNA, 8-alkoxyA in the guide strand (opposite to G in the passenger strand) would project its steric blockade into the minor groove of siRNA, thereby preventing intracellular protein binding onto the RNA. When the siRNA is recruited into the RISC assembly, the 8-alkoxyA in the guide siRNA would encounter U in the mRNA and would flip the bulky appendage into the major groove, thereby allowing necessary guide strand-mRNA-RISC assembly to form (Figure 1).

Alkyl groups were chosen based on their size and shape: propargyl, phenethyl and cyclohexylethyl. The rationale behind the choices is that smaller groups (propargyl) might exhibit higher duplex stabilities and better mRNA knock down efficiencies whereas a larger group (cyclohexylethyl) might prevent immunostimulation to a greater extent, and a medium sized group (phenethyl) might serve both the purposes equally efficiently. Our study suggests that the mRNA knock down ability of the singly modified siRNAs is quite similar to that of the unmodified positive control siRNA. Singly modified siRNA oligomers have also exhibited reduced tendency to bind PKR. With additional alkoxy substitution, the efficiency of the siRNA is significantly reduced; therefore multiply modified siRNAs were not tested for off-pathway protein interactions.

Results and Discussion

Design and Synthesis of 8-Alkoxyadenosine Phosphoramidites

8-AlkoxyA phosphoramidites were synthesized in multiple steps from adenosine (Figure 2). Detailed step-by-step syntheses can be found in the Supporting Information. Adenosine was first brominated at C8 following a standard procedure⁵¹ and sugar hydroxyl groups were protected by silylating agents. 5'-OH and 3'-OH groups were protected using a bidentate silylating agent⁵² and the remaining 2'-OH was protected using a TBDMS protecting group. Next, the bromide ion was displaced with an alkoxy groups via an aromatic nucleophilic substitution reaction. The alkoxide anion was generated in situ by adding dropwise nBuLi into the corresponding alcohol in THF. The exocyclic amine of adenosine was protected using a benzoyl group. Then, the 5'- and 3'-hydroxyls were deprotected using HF-pyridine, a mild defluorinating reagent, leaving 2'-OH protection intact. The 5'-OH was then protected with a standard DMT group, and the 2'-OH was converted to a phosphoramidite derivative.

Propargyl, phenethyl and cyclohexylethyl groups were chosen based on the size and shape of the alkyl groups. The choice of alkyl groups was also considered in terms of prevention of siRNA-PKR interaction as well as maintenance of target mRNA knockdown. We thought that methyl or ethyl groups might be too small to have any significant impact on the siRNA-protein interactions. However, it was also noted that too large a group in the guide strand might reduce significantly or even completely abolish the RNAi efficacy. Hence, we started the series from the propargyloxy modification, which was expected to retain the RNAi

efficacy, as well as prohibit the unwanted protein binding to some extent. The phenylethoxy group was chosen as the most promising group in the guide strand, which could serve both purposes. We were optimistic that cyclohexylethyl modifications will prevent dsRBM-containing proteins from binding siRNAs, but were interested to explore whether such a large modification might also compromise RNAi efficacy.

Design of the Caspase 2 siRNA and Synthesis of Modified Guide Strands

Caspase 2 siRNA (A:U) and the corresponding negative control siRNA as a scrambled sequence were designed using Ambion's siRNA designing tool and checked for sequence similarity using nucleotide Basic Local Alignment Search Tool (BLAST). No significant similarity was found with other genes. To analyze caspase 2 expression levels quickly and reliably, a plasmid-based dual luciferase assay system (psiCHECK2 vector) (Supporting Figure S38) was employed. A fragment of the caspase 2 mRNA sequence (Supporting Figure S39) was inserted into the vector, and the resulting reporter plasmid was used to evaluate caspase 2 mRNA knockdown. The siRNAs successfully knocked down caspase 2 mRNA in cultured HeLa cells. Using standard solid phase RNA synthesis, 8-alkoxyA phosphoramidites were effectively coupled into the guide strand to afford four singly modified and three doubly modified oligonucleotides.

Thermal Analysis of dsRNAs with 8-AlkoxyAs

Unmodified and modified (8-alkoxyA-containing) guide strands were annealed to the complementary passenger strands (Figure 3). The guide and passenger strands were designed in such a way that 8-alkoxyA faces G or U in the corresponding passenger strands. The unmodified A:U duplex was used as a reference, based upon which the thermal stabilities of the singly- and doubly-modified duplexes (both 8-ROA:U and 8-ROA:G) were analyzed. In the case of the 8-ROA:U base pair, the 8-ROA nucleoside can adopt the canonical *anti* orientation around the glycosidic bond. For the 8-ROA:G pair, 8-ROA_{syn}:G_{anti} and 8-ROA_{anti}:G_{anti} combinations were considered, while other A:G combinations were deemed too distorting to exist in an A-form duplex RNA. However, the 8-ROA_{anti}:G_{anti} pair will widen the helix diameter considerably and introduce strain into the duplex; additionally, the presence of a significantly large alkoxy group at position 8 of the purine ring favors the 8-ROA_{syn}:G_{anti} combination. Thus, the existence of 8-ROA_{anti}:G_{anti} cannot be ruled out, but 8-ROA_{syn}:G_{anti} is more likely to occur in A-form duplex RNA.

All of the singly and doubly modified duplexes displayed lower melting temperatures (T_m) compared to unmodified duplexes (Figures 4 and 5). Therefore, the alkoxy group appears to introduce instability in the RNA duplexes. In all duplexes, the 8-ROA:U base pair was found to be more stable than the 8-ROA:G base combination implying that the Watson-Crick hydrogen-bonding is still preferred in these modified adenosines, provided that 8-ROA faces U as the complementary base. Although 8-ROA:G also has a greater potential to expose its Hoogsteen face to *anti*G, steric clash between the alkoxy group of 8-ROA and the exocyclic amine of G in the minor groove might undermine the strength of hydrogen bonding resulting in lower T_m values for 8-ROA:G.

Melting temperatures also varied depending on the position of the 8-alkoxyA base in the guide strand. It is not clear why the 5'-end of the guide strand can tolerate modifications of variable size more effectively than the other end. The 5'-end of the guide strand is thermodynamically less stable and hence more prone to unzipping; introducing a destabilizing nucleoside in this region, as opposed to the other end, should further decrease the thermal stability of the duplex. Again at positions 10 and 15, the 8-alkoxyAs are between two purines, so a greater stacking interaction is expected. These siRNAs would be expected to have higher melting temperatures compared to siRNAs bearing these modifications at

positions 4 and 6. In practice however, the reverse trend is observed. In all cases, higher duplex stability was obtained when the modifications are closer to the 5'-end of the guide strand. Modifications in the middle position (10) or towards the 3'-end (15) of the guide strand always exhibited reduced thermal stability.

For single modifications, lowering of the T_m was moderate for modifications at positions 4 and 6 opposite U (e.g., Pg4:U, Pg6:U, Ce4:U); however single modifications at positions 10 and 15 exhibited significantly lower T_m irrespective of their placement opposite to G or U. As expected, reduction in duplex stability was more pronounced (7–14 °C) for doubly modified duplexes. Interestingly, except for position 15, the size of the alkoxy groups did not have much influence on the thermal stability of siRNA duplexes.

A similar trend was noticed in 8-ClA-containing RNA duplexes which also have lower melting temperatures compared to the unmodified duplexes.⁵³ Therefore, thermal destabilization cannot be primarily linked to the substituent size and shape effect, but also to substituent electronegativity values. Because both Cl and O have higher electronegativities compared to H, a strong electron withdrawing inductive effect may operate to reduce electron density in the purine rings significantly, and hence this can reduce the stability of hydrogen bonds formed by 8-ClA:U or 8-ROA:U in the Watson-Crick base pairing and affect base stacking as well.

Lowering of the T_m in the 8ROA:U base pair compared with the A:U base pair can also be explained in terms of a relatively pronounced *syn* preference of 8-substituted purines over the *anti* conformation, both in the context of DNA^{54,55} and RNA^{53,56–58} nucleosides and duplex oligonucleotides. X-ray crystallographic⁵⁶ and CD studies⁵⁷ of 8BrA and the corresponding 5'-monophosphate have shown that these molecules predominantly exist in the *syn* conformation around the glycosidic bond due to the presence of a sterically demanding Br atom at position 8 in the purine ring. Although the van der Waals radius of Cl (1.80 Å) is significantly smaller than Br (1.95 Å), theoretical calculations predicted preferentially the *syn* conformation even for 8-ClA nucleosides.⁵⁸ Importantly, a relative lowering of melting temperature of A:U vs. A:G combination was noticeably greater than the 8-ROA:U vs. 8ROA:G combinations. Interestingly, this difference shrank even more for the 8-PeOA and 8-CeOA containing duplexes, implying that with increasing the steric bulk of the alkoxy group, the *syn* preference of the 8-alkoxyadenosines increased. All these observations fit well with the preferential *syn* conformation of the 8-substituted-adenosines, such as 8-ClA and 8-ROA, in the duplex RNA context.

Caspase 2 mRNA Knockdown Studies with Modified Adenosine-Containing siRNAs

Singly substituted siRNAs, alkoxyated at several positions of the guide strand, exhibited almost equivalent or slightly lower silencing effects compared to unmodified caspase 2 siRNAs (Figure 6). Among singly modified siRNAs, 8-PgOA exhibited the highest mRNA knockdown efficiency. Surprisingly, 8-CeOA modifications, even though the largest among three modifications, showed almost equal knockdown efficiency to 8-PgOA, whereas, 8-PeOA containing siRNAs were found to have the lowest efficacy. Thus, the mRNA knockdown ability of the modified caspase 2 siRNAs does not directly correlate with the substituent size.

Interestingly, singly modified siRNAs with intermediate thermal stability (modifications at positions 6 and 10) exhibited higher RNAi efficacy than those having the highest (modification at position 4) or the lowest (modification at position 15) stability. This observation supports the basic rules of siRNA design and efficacy, which emphasizes that, for optimal efficacy, the thermal stability of the siRNA duplexes be intermediate, neither

very high, nor very low.^{59,60} Because the differences in T_m among the four sequence contexts is not very high, their influence on the RNAi efficacy engenders curiosity.

In all cases, modifications at position 10 of the guide strand exhibited the highest gene-silencing efficacy, followed by position 6. Modifications at position 15 seemed to reduce the potency of siRNAs, with the exception of the propargyloxy modification, which was found to be equally active when compared to other positions in the series. So, position 15 was found to tolerate smaller groups better as opposed to larger groups such as phenylethoxy and cyclohexylethoxy groups. This trend is somewhat unexpected, since it is already known that substitutions towards the 3'-end of the guide strand typically can tolerate chemical modifications and even mismatches rather well.^{32,34} Contrarily, the seed region and the cleavage site are more sensitive to chemical modifications and mismatches, in general. Sometimes, a single mismatch⁵⁹ at the center or chemical modifications distorting the A-form duplex structure² abolishes the efficacy of the corresponding siRNA. In this case, modifications in two important positions of the seed region with large substituents are quite well tolerated. More surprisingly, a large modification adjacent the cleavage site (position 10) actually rendered the siRNA more effective towards knocking down caspase 2 mRNA. This trend was observed for all modifications; however with larger modifications such as 8CeOA and 8PeOA, this trend was observed without any ambiguity. The presence of a hydrophobic pocket at the corresponding position in Argonaute2 in the RISC might be a plausible explanation for this observation.

Multiple 8-ROA substitutions at the guide strands showed significantly reduced silencing efficacy compared to their corresponding positive control siRNAs or single modifications (Figure 7). This trend holds irrespective of the substituent size, shape and position. These siRNAs only exhibit significant mRNA knock down at 50 nM concentrations. Modified purines at positions 6 and 10 in the guide strand opposite to G or U (in the passenger strand) did not substantially change the knock down efficiency (Figure 8). In most of the cases, delivering the modified purine opposite to G brought about almost equal or slightly higher mRNA knockdown than delivering it opposite to U. Only with Ce6, a reverse trend was observed, though it was only by a small margin. These data suggest that both 8-ROA:G and 8-ROA:U can be loaded into the RISC with equal efficiency.

Importance of Switching the Steric Blockade from the Minor to the Major Groove in the RISC

To explore the importance of 'base switching' in the context of RISC-mediated cleavage of caspase 2 mRNA, two mutant plasmids, P6 and P10 were synthesized using appropriate inserts by mutating specific positions of the wild-type caspase 2 insert. Two important sites in the guide strand of the caspase 2 siRNA, position 6 in the seed region and position 10 adjacent the cleavage site, were chosen for the construction of mutant plasmids, P6 and P10. In such cases, the 8-alkoxyAs (in the guide strand) face G as the complementary base both during their delivery opposite to the passenger strands and during interaction with the mRNA in the RISC. Therefore, alkoxy steric blockades will always be presented in the minor groove both in the guide-passenger and guide-mRNA constructs.

As expected, the mRNA knockdown efficiency was drastically reduced for all three (8-PgOA 8-PeOA and 8-CeOA) modifications; 8-PgOA being the smallest in size is relatively more efficacious (Figures 9 and 10). In the case of A:G mismatched guide siRNA:mRNA constructs, a similar decrease in mRNA knockdown efficiency was noticed with respect to both mutant plasmids, P6 and P10.

These results indicate that for the guide:mRNA duplex, both local widening³⁶ (due to the A:G pair) or minor groove crowding (due to the 8ROA:G pair) can lead to drastic reduction

of the mRNA knockdown efficacy. Probably these distortions in the guide:mRNA duplex do not allow the necessary protein side chain interactions in the guide:mRNA:RISC ternary complex.

SiRNA-PKR Binding Studies

To assess the effect of these modifications on siRNA binding to an off-pathway dsRNA-binding protein, we used a PKR binding assay involving immobilized biotinylated siRNAs as previously described.³² 5'-Biotinylated passenger strands with either U or G opposite to the modified purines were annealed to Pg6, Ce6, Pe6, Pg10, Ce10, Pe10 and unmodified guide strands. The biotinylated strands were then attached to magnetic streptavidin beads and used in affinity purification for PKR. The degree of PKR binding for each strand was then analyzed by western blot.

Our results show that different types of modifications, when placed in the guide strand opposite to G or U in the passenger strand, are significantly less prone to bind to PKR compared to the unmodified sequences (Figure 11). Again, unmodified siRNAs with A:G mismatches at position 6 and 10 are almost as susceptible to bind PKRs as the unmodified one. Hence prevention of protein binding is not a result of mere bulges due to A:G mismatches, but rather an outcome of placement of different types of steric blockades in the minor and major groove by means of placing 8-alkoxyA opposite to U or G.

With propargyloxy modifications being the smallest in size of those studied here, Pg6:G, Pg10:U and Pg10:G were the least effective in preventing PKR binding with the exception of Pg6:U; whereas all cyclohexylethoxy and phenylethoxy modifications reduced the PKR binding significantly. It seems that PKR is slightly more prone to bind to modified siRNAs where U in the passenger strand was placed opposite to the modification in the guide strand than to siRNAs where modifications are opposite to G. In some cases (for example, Pg10:U vs. Pg10:G, Pe10:U vs. Pe10:G) this discrimination is more pronounced, and in some cases (Pg6:U vs. Pg6:G, Pe6:U vs. Pe6:G) less. These data suggest that this type of steric occlusion of the minor groove is moderately or marginally advantageous over major groove occlusion. Surprisingly, when 8-CeO modifications were placed in position 6 and 10 opposite to U or G, no distinction was observed between the binding affinities towards PKR. Both Ce6:G and Ce6:U were equally effective in preventing PKR binding. A similar trend with lesser efficacy was observed with Ce10:G and Ce10:U.

There are other reports of prevention of PKR binding by manipulating minor groove modifications;^{28,32,33} however there is no report, so far, of any major groove modification that is equally efficient in attenuating siRNA-PKR interaction when placed in the same position of an siRNA. 8-PeOA and 8-CeOA modified siRNAs are the first examples to exhibit such a characteristic. From the thermal analysis data and RNAi studies with mutant plasmids, it is clear that 8-ROA:U and 8:ROA:G combinations are distinct, their differences likely related to different glycosidic bond conformations and placement of the steric blockades. Previously, Kannan et al. reported N^2 alkylated 8-oxoguanine-based switches are capable of preventing PKR binding only when the steric blockade is placed in the minor groove as opposed to the major groove.³³ In that study N^2 -propyl and N^2 -benzyl modified 8-oxoguanines were used. The phenylethoxy and cyclohexylethoxy groups used here are not only bulky, but also significantly longer than the propyl and benzyl groups used in the previous studies. Hence, irrespective of their position in either groove, these larger groups repel PKR to a much greater extent even when placed in the major groove.

Another probable rationale for the high activity of 8ROA:U might be the modification site (position 8) of adenosine itself. PKR is known to bind dsRNA by interacting with two segments of the minor groove and the intervening major groove.⁶¹ In the major groove,

lysine side chains of PKR interact with the negatively charged phosphate backbone of dsRNA. Hence, appropriate major groove modifications of the siRNA can potentially prevent this electrostatic interaction and block PKR binding. In the case of 8ROA_{anti}:U-containing siRNAs, the alkoxy group is suitably positioned to disrupt such major groove interactions with the siRNA.

Conclusions

A general solution to sequence-independent off-pathway protein binding of siRNAs, while maintaining mRNA knock down efficacy, was investigated here. This study also explored the scope of guide strand modifications by utilizing 8-alkoxyadenosines of variable alkyl group size. This is an unusual example in which a purine ribonucleoside is proposed to exist in both *syn* and *anti* conformations around the glycosidic bond depending on the base-pairing partner. 8-Alkoxyadenosine modifications were tolerated both in singly and doubly modified duplexes irrespective of G or U as the pairing partner, although lowering of the T_m is very pronounced in cases of multiple modifications. Singly modified siRNAs were almost the same or slightly less effective in RNAi, whereas multiply modified siRNAs exhibited much lower caspase 2 insert mRNA knock down activity. Singly modified siRNAs were capable of inhibiting PKR-siRNA interactions compared to the unmodified siRNA. The necessity of 'base switching' in the RISC was demonstrated by using two caspase 2 inserts, individually mutated at positions 6 and 10. To confirm the *syn* vs. *anti* conformational preference of 8-substituted adenosines in appropriate duplex RNA contexts, NMR spectroscopic studies are currently in progress.

Methods

Synthesis of 8-Alkoxyadenosine Phosphoramidites

Complete experimental details and spectra are given in the Supporting Information.

Synthesis and Purification of SiRNAs

All siRNA oligonucleotides, modified and unmodified, were synthesized by standard solid-phase RNA synthesis on DNA columns in the DNA/Peptide Core Facility of the University of Utah using an Applied Biosystems Model 394 DNA/RNA synthesizer. After synthesis, the 21-mer RNA oligonucleotides were cleaved from the column and deprotected using methanolic ammonia for 24 h at room temperature, and the 2'-OTBDMS group was deprotected by using TEA•3HF overnight at room temperature. The oligomers were then dialyzed at 4 °C for 6 h and lyophilized in a SpeedVac. SiRNA oligomers were purified by semi-preparative ion-exchange HPLC using ammonium acetate, and lyophilized to remove salt and water. The remaining salt was removed by dialysis. The samples were dialyzed again and stored at -20 °C under dry conditions.

SiRNA Duplex Formation

SiRNA oligomers were dissolved in annealing buffer (100 mM NaCl and 10 mM TRIS, pH 7.1), and appropriate guide and passenger strands were mixed in equal amounts in an Eppendorf tube. The nucleotide mixtures were placed in a 90 °C water bath for 5 min, and the bath was gradually cooled down to room temperature in about 2 h.

Thermal Analysis of SiRNA Duplexes

Melting temperature experiments of the hybridized duplexes were performed in a Beckman DU 650 spectrophotometer. In all experiments, the temperature was varied from 25 °C to 80 °C, and the rate of heating was 0.5 °C/min. Thermal denaturation was monitored at 260 nm. All experiments were performed with 1 nmol duplex RNA in siRNA annealing buffer

(containing 10mM Tris, 100 mM NaCl, pH 7.1) in triplicate and normalized to appropriate blank controls. The reported T_m value was the average of 3 independent experiments, and error bars represent standard deviation from the average value. Experimental data were analyzed by 'two point average' method. 4 cuvettes were filled with 325 μ L annealing buffer and were read as blanks. Then, 3 cuvettes were emptied, and 1 nmol siRNA duplex was added to the siRNA annealing buffer so that the total volume of the mixture became 975 μ L. This mixture was equally distributed in those 3 cuvettes, and the fourth cuvette was loaded with 325 μ L annealing buffer. Lids were closed tightly and the temperature was varied from 25 to 80 $^{\circ}$ C.

Synthesis of Plasmids

The appropriate caspase 2 insert was introduced into the psiCHECK2 vector using the multiple cloning region of the plasmid. One wild type and two mutant plasmids were synthesized. The plasmids were multiplied in *E. coli*, extracted by Qiagen mini plasmid extraction kit and sequenced at the Core Facility, University of Utah. The protocol for the recombinant plasmid synthesis is furnished in the supporting information.

Cell Culture

HeLa cells were cultured in Dubelco's DMEM cell culture medium with 10% FBS and maintained under 5% CO₂ in an incubator. Corning 75 mL and 225 mL cell culture flasks were used for growing cells. Water in the incubator pan was always autoclaved before use. The cells that were used in the experiments were between passage numbers 5 to 12. Cell confluence was kept between 50–70% for all RNAi assays.

RNAi Assay

During the day of the experiment, cells within the proper confluence level (~ 60%) were trypsinized and detached from the flask. Excess medium was added to inactivate trypsin. Cells were then centrifuged and resuspended in fresh medium. Cells were counted manually by using a standard hemocytometer and diluted so that 6000 cells are present per 80 μ L medium. Cells were kept in the incubator at 37 $^{\circ}$ C for about 45 minutes, while siRNA transfection complexes were being formed. siPORTTM NeoFXTM transfection agent was used as the siRNA delivery method in all cell culture studies. All siRNAs and plasmids were diluted in Opti-MEM[®] reduced serum medium, and siRNAs and plasmids were mixed together; similarly transfection agents were diluted in Opti-MEM[®] and transfection complexes were allowed to form. Then, siRNA-plasmid mixtures are added into diluted transfection agent solutions and mixed thoroughly by pipetting. In all the experiments, 96-well plates were used and 20 μ L siRNA-transfection complex was discharged into each well. The cell suspension was then taken out of the incubator and 80 μ L of the cell suspension (containing 6000 cells) was added to each well. The plate was shaken several times and tilted several times to ensure homogenization of the two solutions. Then the plate was kept inside the incubator and within 4 hours, the transfection of the cells was complete. Cells were allowed to grow for 36 hours and were assayed for caspase 2 mRNA insert silencing using the Dual-GloTM Luciferase Assay System from Promega. Initially firefly luciferase emission was recorded and then by adding the Dual Glo Stop and Glo reagent, the firefly luciferase was terminated and Renilla luminescence was recorded. Ratios of Renilla to firefly luminescence of samples were used to compute percent expression of caspase 2 insert mRNA. Each data point is the average of 6 independent experiments and error bars represent standard deviation from the average value.

SiRNA-PKR Binding Assay

U87 cells (approximately 8×10^6 in a 75 cm^2 flask) were treated 24 h before lysis with human interferon- α A (PBL Interferon Source) to a final concentration of $1 \times 10^6 \text{ U L}^{-1}$. The interferon-treated cells were washed twice with PBS and lysed by shaking with 3 mL solubilization buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 1% Triton X-100, supplemented with protease inhibitor cocktail (ProteoBlock, Fermentas) for 30 min on ice. The lysates were clarified by centrifugation at $14000 \times g$ at 4°C for 20 min and used directly in pull-down experiments. Magnetic streptavidin beads (0.5 mg, $50 \mu\text{L}$) (Dynabeads M-280, Invitrogen) were prepared for RNA manipulation according to the manufacturer's protocol. Briefly, beads were suspended in 350 μL binding buffer (5 mM Tris-HCl, pH 7.5, 0.5 mM EDTA, 1 M NaCl) and coated with 100 pmol of siRNA (100 μL) by gently rocking for 30 min at room temperature. The siRNA-coated beads were washed twice with binding buffer (500 μL) and once with solubilization buffer (500 μL) and incubated with 0.5 mL cell lysate for 20 min at room temperature. Beads without siRNA were also incubated with cell lysate and served as a control. After incubation, the beads were washed four times with 500 μL wash buffer (20 mM HEPES, pH 7.9, 2.5 mM MgCl_2 , 100 mM KCl, 20% glycerol, 0.5 mL DTT, 0.2 mg/mL yeast RNA, 0.2 mg/mL salmon sperm DNA). Bound protein was eluted from the beads by heating in loading buffer, separated on 6% SDS-PAGE and transferred to a PVDF membrane for Western blotting. Membranes were blocked with blotting-grade milk (Bio-Rad), incubated with PKR antibody (Santa Cruz Biotechnology, 1:1000 dilution), washed with TBS-Tween, and incubated with alkaline phosphatase-conjugated secondary antibody (Santa Cruz Biotechnology, 1:2000 dilution). The proteins were detected using ECF substrate (GE Healthcare) on a Typhoon Trio Variable Mode Imager (GE Healthcare) and band intensities were quantified using ImageQuant software (Molecular Dynamics). PKR binding affinity is reported as the average ratio of band intensities of modified siRNA to native siRNA for three independent experiments.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments


Support of this work by a grant from the National Institutes of Health (R01 GM080784) is gratefully acknowledged.

References

1. Braasch DA, Jensen S, Liu Y, Kaur K, Arar K, White MA, Corey DR. *Biochemistry*. 2003; 42:7967. [PubMed: 12834349]
2. Chiu Y-L, Rana TM. *RNA*. 2003; 9:1034. [PubMed: 12923253]
3. Allerson CR, Sioufi N, Jarres R, Prakash TP, Naik N, Berdeja A, Wanders L, Griffey RH, Swayze EE, Bhat B. *J Med Chem*. 2005; 48:901. [PubMed: 15715458]
4. Egli M, Minasov G, Tereshko V, Pallan PS, Teplova M, Inamati GB, Lesnik EA, Owens SR, Ross BS, Prakash TP, Manoharan M. *Biochemistry*. 2005; 44:9045. [PubMed: 15966728]
5. Prakash TP, Allerson CR, Dande P, Vickers TA, Sioufi N, Jarres R, Baker BF, Swayze EE, Griffey RH, Bhat B. *J Med Chem*. 2005; 48:4247. [PubMed: 15974578]
6. Hoerter JA, Walter NG. *RNA*. 2007; 13:1887. [PubMed: 17804643]
7. Parrish S, Fleenor J, Xu S, Mello C, Fire A. *Mol Cell*. 2000; 6:1077. [PubMed: 11106747]
8. Dowler T, Bergeron D, Tedeschi A-L, Paquet L, Ferrari N, Damha MJ. *Nucleic Acids Res*. 2006; 34:1669. [PubMed: 16554553]

9. Manoharan M, Akinc A, Pandey RK, Qin J, Hadwiger P, John M, Mills K, Charisse K, Maier MA, Nechev L, Greene EM, Pallan PS, Rozners E, Rajeev KG, Egli M. *Angew Chem Int Ed.* 2011; 50:2284.
10. Fauster K, Hartl M, Santner T, Aigner M, Kreutz C, Bister K, Ennifar E, Micura R. *ACS Chem Biol.* 2012; 7:581. [PubMed: 22273279]
11. Watts JK, Choubdar N, Sadalapure K, Robert F, Wahba AS, Pelletier J, Pinto BM, Damha MJ. *Nucleic Acids Res.* 2007; 35:1441. [PubMed: 17284457]
12. Gore KR, Nawale GN, Harikrishna S, Chittoor VG, Pandey SK, Höbartner C, Patankar S, Pradeepkumar PI. *J Org Chem.* 2012; 77:3233. [PubMed: 22372696]
13. Manoharan M. *Curr Opin Chem Biol.* 2004; 8:570. [PubMed: 15556399]
14. Rozners E. *Curr Org Chem.* 2006; 10:675.
15. Watts JK, Deleavey GF, Damha MJ. *Drug Discov Today.* 2008; 13:842. [PubMed: 18614389]
16. Manoharan, M.; Rajeev, KG. U.S. Patent 120878. 2009.
17. Deleavey GF, Watts JK, Alain T, Robert F, Kalota A, Aishwarya V, Pelletier J, Gewirtz AM, Sonenberg N, Damha MJ. *Nucleic Acids Res.* 2010; 38:4547. [PubMed: 20413581]
18. Corey DR. *The Journal of clinical investigation.* 2007; 117:3615. [PubMed: 18060019]
19. Bramsen JB, Pakula MM, Hansen TB, Bus C, Langkjaer N, Odadzic D, Smicius R, Wengel SL, Chattopadhyaya J, Engels JW, Herdewijn P, Wengel J, Kjems J. *Nucleic Acids Res.* 2010; 38:5761. [PubMed: 20453030]
20. Oishi M, Nagasaki Y, Itaka K, Nishiyama N, Kataoka K. *J Am Chem Soc.* 2005; 127:1624. [PubMed: 15700981]
21. Zhu L, Mahato RI. *Bioconjugate Chem.* 2010; 21:2119.
22. Meyer M, Dohmen C, Philipp A, Kiener D, Maiwald G, Scheu C, Ogris M, Wagner E. *Mol Pharm.* 2009; 6:752. [PubMed: 19348503]
23. Grijalvo S, Ocampo SM, Perales JC, Eritja R. *J Org Chem.* 2010; 75:6806. [PubMed: 20863084]
24. Yamada T, Peng CG, Matsuda S, Addepalli H, Jayaprakash KN, Alam MR, Mills K, Maier MA, Charisse K, Sekine M, Manoharan M, Rajeev KG. *J Org Chem.* 2011; 76:1198. [PubMed: 21299239]
25. Manoharan, M.; Elbashir, S.; Harborth, J. U.S. Patent 091515. 2004.
26. Soutschek J, Akinc A, Bramlage B, Charisse K, Constien R, Donoghue M, Elbashir S, Geick A, Hadwiger P, Harborth J, John M, Kesavan V, Lavine G, Pandey RK, Racie T, Rajeev KG, Roehl I, Toudjarska I, Wang G, Wuschko S, Bumcrot D, Koteliansky V, Limmer S, Manoharan M, Vornlocher H-P. *Nature.* 2004; 432:173. [PubMed: 15538359]
27. Cesarone G, Edupuganti OP, Chen C-P, Wickstrom E. *Bioconjugate Chem.* 2007; 18:1831.
28. Puthenveetil S, Whitby L, Ren J, Kelnar K, Krebs JF, Beal PA. *Nucleic Acids Res.* 2006; 34:4900. [PubMed: 16982647]
29. Somoza A, Chelliserrykattil J, Kool ET. *Angew Chem Int Ed.* 2006; 45:4994.
30. Xia J, Noronha A, Toudjarska I, Li F, Akinc A, Braich R, Frank-Kamenetsky M, Rajeev KG, Egli M, Manoharan M. *ACS Chem Biol.* 2006; 1:176. [PubMed: 17163665]
31. Terrazas M, Kool ET. *Nucleic Acids Res.* 2009; 37:346. [PubMed: 19042976]
32. Peacock H, Fostvedt E, Beal PA. *ACS Chem Biol.* 2010; 5:1115. [PubMed: 20863128]
33. Kannan A, Fostvedt E, Beal PA, Burrows CJ. *J Am Chem Soc.* 2011; 133:6343. [PubMed: 21452817]
34. Wahba AS, Azizi F, Deleavey GF, Brown C, Robert F, Carrier M, Kalota A, Gewirtz AM, Pelletier J, Hudson RHE, Damha MJ. *ACS Chem Biol.* 2011; 6:912. [PubMed: 21667942]
35. Peacock H, Kannan A, Beal PA, Burrows CJ. *J Org Chem.* 2011; 76:7295. [PubMed: 21834582]
36. Li F, Pallan PS, Maier MA, Rajeev KG, Mathieu SL, Kreutz C, Fan Y, Sanghvi J, Micura R, Rozners E, Manoharan M, Egli M. *Nucleic Acids Res.* 2007; 35:6424. [PubMed: 17881374]
37. Ibarra-Soza JM, Morris AA, Jayalath P, Peacock H, Conrad WE, Donald MB, Kurth MJ, Beal PA. *Org Biomol Chem.* 2012; 10:6491. [PubMed: 22766576]
38. Marques JT, Williams BR. *Nat Biotechnol.* 2005; 23:1399. [PubMed: 16273073]
39. Schlee M, Hornung V, Hartmann G. *Mol Ther.* 2006; 14:463. [PubMed: 16877044]

40. Judge AD, Sood V, Shaw JR, Fang D, McClintock K, MacLachlan I. *Nat Biotechnol.* 2005; 23:457. [PubMed: 15778705]
41. Agrawal S, Kandimalla ER. *Nat Biotechnol.* 2004; 22:1533. [PubMed: 15583662]
42. Jurk M, Chikh G, Schulte B, Kritzler A, Richardt-Pargmann D, Lampron C, Luu R, Krieg AM, Vicari AP, Vollmer J. *Nucleic Acid Ther.* 2011; 21:201. [PubMed: 21749297]
43. Sledz CA, Holko M, de Veer MJ, Silverman RH, Williams BR. *Nat Cell Biol.* 2003; 5:834. [PubMed: 12942087]
44. Yang W, Wang Q, Howell KL, Lee JT, Cho D-SC, Murray JM, Nishikura K. *J Biol Chem.* 2005; 280:3946. [PubMed: 15556947]
45. Li S, Peters GA, Ding K, Zhang X, Qin J, Sen GC. *Proc Natl Acad Sci USA.* 2006; 103:10005. [PubMed: 16785445]
46. Nishikura K. *Nat Rev Mol Cell Biol.* 2006; 7:919. [PubMed: 17139332]
47. Heale BSE, Keegan LP, McGurk L, Michlewski G, Brindle J, Stanton CM, Caceres JF, O'Connell MA. *EMBO J.* 2009; 28:3145. [PubMed: 19713932]
48. Nallagatla SR, Bevilacqua PC. *RNA.* 2008; 14:1201. [PubMed: 18426922]
49. Collins LV, Hajizadeh S, Holme E, Jonsson I-M, Tarkowski A. *J Leukocyte Biol.* 2004; 75:995. [PubMed: 14982943]
50. Pope BL, MacIntyre JP, Kimball E, Lee S, Zhou L, Taylor GR, Goodman MG. *Cel Immunol.* 1995; 162:333.
51. Ikehara M, Uesugi S. *Chem Pharm Bull.* 1969; 17:348. [PubMed: 5782965]
52. Trost BM, Caldwell CG. *Tetrahedron Lett.* 1981; 22:4999.
53. Chen LS, Sheppard TL. *Nucleosides Nucleotides Nucleic Acids.* 2002; 21:599. [PubMed: 12484453]
54. Dias E, Battiste JL, Williamson JR. *J Am Chem Soc.* 1994; 116:4479.
55. Eason RG, Burkhardt DM, Phillips SJ, Smith DP, David SS. *Nucleic Acids Res.* 1996; 24:890. [PubMed: 8600457]
56. Tavale SS, Sobell HM. *J Mol Biol.* 1970; 48:109. [PubMed: 5448585]
57. Ikehara M, Uesugi S, Yoshida K. *Biochemistry.* 1972; 11:830. [PubMed: 5059890]
58. Walker GA, Bhatia SC, Hall JH Jr. *J Am Chem Soc.* 1987; 109:7629.
59. Elbashir SM, Martinez J, Patkaniowska A, Lendeckel W, Tuschl T. *EMBO J.* 2001; 20:6877. [PubMed: 11726523]
60. Reynolds A, Leake D, Boese Q, Scaringe S, Marshall WS, Khvorova A. *Nat Biotechnol.* 2004; 22:326. [PubMed: 14758366]
61. Cole JL. *Trends Biochem Sci.* 2007; 32:57. [PubMed: 17196820]

**Figure 1.**

(A) Proposed ‘base switch’ showing flipping of a steric blockade from the minor to the major groove; steric occlusion in the minor groove prevents siRNA-PKR interaction in transit and major groove accommodation of the steric blockade allows siRNA-protein interactions in the RISC. (B) Proposed base pairs of the 8-alkoxyAs.




Figure 2.
Synthesis of the 8-alkoxyadenosine phosphoramidites.




Figure 3. SiRNA sequences used. A:U represents unmodified siRNA and modified (denoted by M:G) sequences have the standard A:U base pair replaced by 8-alkoxy(X):G in one or two positions in the siRNAs.




Figure 4. Thermal analysis of siRNA duplexes with single 8-alkoxyA modifications at four different positions, 4, 6, 10 and 15, in the guide strand. Modifications were placed opposite to U or G in the corresponding passenger strands. The sequences are shown in Figure 2; here X represents 8-PgOA, 8-PeOA or 8-CeOA, and N is either U or G.





Figure 5. Thermal analysis of siRNA duplexes with double 8-alkoxyA modifications at three different positions, 6, 10 and 15, in the guide strand. Modifications were placed opposite to U or G in the corresponding passenger strands. The sequences are shown in the Figure 2; here X is 8-PgOA, 8-PeOA or 8-CeOA, and N is either U or G.

**Figure 6.**

% Expression of Renilla luciferase relative to firefly luciferase when treated with siRNAs bearing single 8-alkoxyA modification at position 4, 6, 10 or 15 in the guide strand. 8-AlkoxyA modifications were placed opposite G (in the passenger strand) during delivery and targeted U (in the mRNA) in the RISC. The siRNA sequences are depicted in Figure 2; here N = U when X = A, and N = G when X = 8-PgOA, 8-PeOA or 8-CeOA. Experiments were conducted at three different concentrations: 50 nM, 0.5 nM and 0.01 nM.




Figure 7.

% Expression of the Renilla luciferase relative to the firefly luciferase when treated with siRNAs bearing double 8-alkoxyA modifications in three different combinations in the guide strand: (6, 10), (6, 15) and (10, 15). 8-AlkoxyA modifications were placed opposite to G in the passenger strand during delivery and targeted U (in mRNA) in the RISC. The siRNA sequences are depicted in Figure 2; here N = U when X = A, and N = G when X = 8-PgOA, 8-PeOA or 8-CeOA. Experiments were conducted at two different concentrations: 50 nM and 0.5 nM.




Figure 8.
% Expression of the Renilla luciferase relative to the firefly luciferase when treated with siRNAs bearing 8-alkoxyA modifications at positions 6 and 10 opposite to either G or U.




Figure 9. % Expression of the Renilla luciferase relative to the firefly luciferase, when treated with siRNAs bearing modifications at position 6 in the guide strand, and targeted U (from dA6:dT plasmid, depicted by solid bars) vs. G (from dC6:dG plasmid, depicted by hashed bars) in the caspase 2 insert and mutant caspase 2 insert mRNA respectively. Sequences are given in Figure 2. Experiments were conducted at two different concentrations: 50 nM and 0.5 nM. A:U, Pg6:U, Ce6:U and Pe6:U represents targeting of U in the caspase 2 insert mRNA (from dA6:dT plasmid) and C:G, A:G, Pg6:G, Ce6:G and Pe6:G represents targeting of G in the mutant mRNA where a U was mutated to G (from dC6:dG plasmid). Nomenclature of the plasmid comes from the corresponding modification position in the guide strand of caspase 2 insert mRNA.




Figure 10.

% Expression of the Renilla luciferase relative to the firefly luciferase, when treated with siRNAs bearing modifications at position 10 in the guide strand, and targeted U (from dA10:dT plasmid, depicted by solid bars) vs. G (from dC10:dG plasmid, depicted by hashed bars) in the caspase 2 insert and mutant caspase 2 insert mRNA, respectively. Sequences are given in Figure 2. Experiments were conducted at two different concentrations: 50 nM and 0.5 nM. A:U, Pg10:U, Ce10:U and Pe10:U represent targeting of U in the caspase 2 insert mRNA (from dA10:dT plasmid) and C:G, A10:G, Pg10:G, Ce10:G and Pe10:G represent targeting of G in the mutant mRNA where a U was mutated to G (from dC10:dG plasmid). Nomenclature of the plasmid comes from corresponding modification position in the guide strand of caspase 2 insert mRNA.




Figure 11.

PKR binding to modified siRNAs containing 8-alkoxyadenosine switches. (A) Biotinylated siRNAs were bound to magnetic streptavidin beads and treated with lysates from U87 cells treated with IFN- α . The amount of PKR retained was determined by western blotting. Cyclohexylethoxy (Ce), phenylethoxy (Pe), and propargyloxy (Pg) modifications were tested at positions 6 and 10 opposite either G (Blue) or U (Red). Strands with A:G mismatches were also tested at positions 6 and 10 as controls (Green). (B). A representative western blot consisting of 1. Unmodified siRNA 2. PgG6 3. PgU6, and 4,5. Pos6 in duplicate.