

Published in final edited form as:

Nat Immunol. 2011 May ; 12(5): 450–459. doi:10.1038/ni.2020.

## The transcription factor E4BP4 regulates the production of IL-10 and IL-13 in CD4<sup>+</sup> T cells

Yasutaka Motomura<sup>1,2</sup>, Hiroshi Kitamura<sup>3</sup>, Atsushi Hijikata<sup>3</sup>, Yuko Matsunaga<sup>4</sup>, Koichiro Matsumoto<sup>4</sup>, Hiromasa Inoue<sup>4</sup>, Koji Atarashi<sup>5</sup>, Shohei Hori<sup>6</sup>, Hiroshi Watarai<sup>7</sup>, Jinfang Zhu<sup>8</sup>, Masaru Taniguchi<sup>7</sup>, and Masato Kubo<sup>1,2</sup>

<sup>1</sup>Laboratory for Signal Network, Research Center for Allergy and Immunology, RIKEN Yokohama Institute, Yokohama, Japan

<sup>2</sup>Division of Molecular Pathology, Research Institute for Biological Science, Tokyo University of Science, Chiba, Japan

<sup>3</sup>Laboratory for Immunogenomics, Research Center for Allergy and Immunology, RIKEN Yokohama Institute, Yokohama, Japan

<sup>4</sup>Research Institute for Diseases of the Chest, Graduate School of Medical Sciences, Kyushu University, Fukuoka, Japan

<sup>5</sup>Department of Immunology, Graduate School of Medicine, The University of Tokyo, Tokyo, Japan

<sup>6</sup>Laboratory for Immune Homeostasis, Research Center for Allergy and Immunology, RIKEN Yokohama Institute, Yokohama, Japan

<sup>7</sup>Laboratory for Immune Regulation, Research Center for Allergy and Immunology, RIKEN Yokohama Institute, Yokohama, Japan

<sup>8</sup>National Institutes of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, Maryland, USA

### Abstract

The immunoregulatory cytokine interleukin 10 (IL-10) is expressed mainly by T helper type 2 (T<sub>H</sub>2) cells but also by T<sub>H</sub>1 cells during chronic infection. Here we observed plasticity in the expression of IL-10 and IL-13 after chronic T<sub>H</sub>1 stimulation; furthermore, the expression of *Il10* and *Il13* was regulated by the transcription factor E4BP4. Chronically stimulated E4BP4-deficient (*Nfil3*<sup>-/-</sup>; called '*E4bp4*<sup>-/-</sup>' here) T<sub>H</sub>1 cells, regulatory T cells (T<sub>reg</sub> cells) and natural killer T cells (NKT cells) had attenuated expression of IL-10 and IL-13. Enforced expression of *E4bp4* initiated the production of IL-10 and IL-13 by conventional T<sub>H</sub>1 cells. *E4bp4*<sup>-/-</sup> T<sub>H</sub>2 cells showed impairment of IL-10 production with no effect on IL-13. Our results indicate that E4BP4 has

© 2011 Nature America, Inc. All rights reserved.

Correspondence should be addressed to M.K. (raysolfc@rcai.riken.jp).

Note: Supplementary information is available on the Nature Immunology website.

#### AUTHOR CONTRIBUTIONS

Y. Motomura built the initial constructs, generated mouse lines and confirmed them *in vivo*, and did most of the experiments; H.K. and A.H. did microarray and bioinformatics analysis; Y. Matsunaga, K.M. and H.I. did airway hyper-responsiveness experiments; K.A. provided *Il10* reporter mice; S.H. analyzed T<sub>reg</sub> cells; H.W. and M.T. analyzed NKT cells; J.Z. provided materials; and M.K. designed and coordinated experiments, supervised the project and wrote the paper.

#### COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.

multiple functions in controlling the plasticity of IL-13 in T<sub>H</sub>1 cells and IL-10 in T<sub>H</sub>1 cells, T<sub>H</sub>2 cells, T<sub>reg</sub> cells and NKT cells.

After being activated by antigen-presenting cells (APC), naive CD4<sup>+</sup> T cells differentiate into helper T cells that acquire the ability to secrete effector cytokines. Helper T cells can be categorized into at least three subsets. T helper type 1 (T<sub>H</sub>1) cells, which secrete interleukin 2 (IL-2) and the proinflammatory cytokines interferon (IFN- $\gamma$ ) and tumor necrosis factor (TNF), promote cellular immune responses mainly to intracellular pathogens and viruses. T<sub>H</sub>2 cells secrete IL-4, IL-5, IL-6, IL-10 and IL-13 and promote humoral immune responses mainly to extracellular pathogens<sup>1</sup>. Cells of the T<sub>H</sub>17 helper T cell subset, which secrete IL-17A, IL-17F and IL-22, mediate immune responses mainly to extracellular pathogenic bacteria and fungi<sup>2</sup> or have been linked to autoimmune disease<sup>3</sup>.

IL-13 was originally identified as a T<sub>H</sub>2 cytokine<sup>4</sup>; however, it is now known that IL-13 is produced by a variety of cell types, such as natural killer T cells (NKT cells), natural killer cells (NK cells), eosinophils and basophils<sup>5</sup>. As all of these cells also produce IL-4, it has been speculated that both IL-13 expression and IL-4 expression are controlled by common transcriptional mechanisms<sup>6,7</sup>. However, T<sub>H</sub>1 cells produce IL-13 in response to a combination of antigenic stimulation and IL-18 (ref. 8). Human patients with atopic or nonatopic asthma who have high titers of allergen-specific immunoglobulin E (IgE) can have more production of the T<sub>H</sub>1 cytokine IFN- $\gamma$ , along with substantially more IL-13 production<sup>9,10</sup>. Therefore, T<sub>H</sub>1 cells could also be a source of IL-13 because of their plasticity, which is controlled by chronic antigen stimulation. However, the molecular mechanisms that regulate the plasticity of IL-13 production in T<sub>H</sub>1 cells is unclear.

IL-10 is also secreted by a wide variety of cells, such as T<sub>H</sub>2 cells, NKT cells, NK cells, natural regulatory T cells (T<sub>reg</sub> cells), regulatory B cells, macrophages, dendritic cells (DCs) and monocytes<sup>11</sup>. T<sub>H</sub>17 cells also express IL-10 in the presence of IL-27 (ref. 12). The mechanism by which IL-10 production is abrogated in T<sub>H</sub>1 cells has been thought to be stable and immutable. However, IL-10 production by T<sub>H</sub>1 cells has been reported in mice chronically infected with parasites such as *Toxoplasma gondii*<sup>13,14</sup> or *Leishmania major*<sup>15</sup>. T<sub>H</sub>1 cells also produce IL-10 in response to high-dose antigenic stimulation and hyperactivation of T cell antigen receptor (TCR)-mediated signaling leading to continuous phosphorylation of the mitogen-activated protein kinases Erk1 and Erk2. Mechanistically, activation of the transcription factor STAT4 seems to elicit this plasticity<sup>16</sup>. IL-10 is an immunoregulatory cytokine that can downregulate active immune responses and acts mainly on T cells or macrophages to attenuate inflammatory cytokine production and antigen presentation. IL-10-deficient mice die with spontaneous colitis<sup>17</sup>, and this phenotype is partly retained even in mice lacking IL-10 only in T<sub>reg</sub> cells<sup>18</sup>. IL-10 therefore has an important role regulating overactive responses that would otherwise result in autoinflammatory disease. The plasticity of IL-10 production by T<sub>H</sub>1 cells in response to excess antigen stimuli could be an effective failsafe mechanism by which immune homeostasis is maintained. The differentiation of T<sub>H</sub>1, T<sub>H</sub>2 and T<sub>H</sub>17 cells is regulated by distinct signaling pathways<sup>19</sup>. Despite differences in their developmental origin, these T cell subsets share the ability to make IL-10 (ref. 11), although the molecular mechanisms that control this remain unclear.

The expression of *Il10* and *Il13* is regulated by GATA-3, the master transcriptional regulator for T<sub>H</sub>2 differentiation. GATA-3 coordinately regulates a group of genes encoding T<sub>H</sub>2 cytokines. Indeed, GATA-3 acts directly on the *Il13* promoter<sup>20</sup> and initiates chromatin remodeling at the *Il10* locus<sup>21</sup>. Activated T cell-specific conditional deletion of *Gata3* markedly attenuates IL-13 expression in committed T<sub>H</sub>2 cells<sup>22</sup>. However, IL-10 is also expressed by T<sub>H</sub>1 cells and NK cells, which normally do not express GATA-3, and the

mechanism by which IL-10 is expressed in the absence of GATA-3 remains unclear<sup>16</sup>. Therefore, an unknown regulator of plasticity may substitute for the function of GATA-3, particularly in these cell types.

E4BP4 (E4 promoter-binding protein 4) was originally identified as a basic leucine zipper transcription factor. E4BP4 regulates circadian rhythm by competing for DNA binding with a member of the related PAR family of basic leucine zipper transcription factors that has a DNA-binding domain similar to that of E4BP4. PAR factors have been characterized as transcriptional activators, whereas E4BP4 is a repressor. E4BP4 and the PAR proteins may switch back and forth between the on state and off state on target genes to control the central circadian clock<sup>23</sup>. In the immune system, E4BP4 is also known as NFIL-3, as it regulates the *Il3* promoter and promotes IL-3-mediated cell survival<sup>24,25</sup>. E4BP4 is a critical regulator of NK cell development through its induction of the transcriptional inhibitor Id2 (refs. 26,27) and also regulates IgE class switching in B cells<sup>28</sup>. Expression of the gene encoding E4BP4 (*Nfil3*; called '*E4bp4*' here) is specific to T<sub>H2</sub> cells in both mice and humans<sup>29,30</sup>. However, the functional importance of T<sub>H2</sub> cell-specific expression of E4BP4 is unclear.

In this study, we demonstrate that E4BP4 is the unknown factor important for the plasticity in the production of IL-10 and IL-13 by T<sub>H1</sub> cells after chronic antigen stimulation and that E4BP4 induced IL-10 production in the absence of GATA-3. Furthermore, E4BP4 regulated the transcription of *Il10* and *Il13* in natural T<sub>reg</sub> cells and NKT cells. However, IL-13 expression by T<sub>H2</sub> cells was independent of E4BP4, because T<sub>H2</sub> cells lacking *E4bp4* did not produce IL-10 but IL-13 production was intact. Therefore, our data indicate that E4BP4 is a critical transcriptional regulator for IL-10, especially in CD4<sup>+</sup> T cells. In addition, our data provide evidence of the functional importance of the expression of E4BP4 by T<sub>H2</sub> cells, in which it is an essential regulator of IL-10 production.

## RESULTS

### Plasticity in IL-13 production by T<sub>H1</sub> cells

In an airway hyper-responsiveness model, *Il4*<sup>-/-</sup> and *Il4r*<sup>-/-</sup> mice lacking T<sub>H2</sub> cells still had substantial airway hyper-responsiveness, similar to that of control C57BL/6 mice. To confirm the involvement of IL-13 in this response, we isolated CD4<sup>+</sup> T cells from bronchoalveolar lavage fluid and mesenteric lymph nodes of *Il4*<sup>-/-</sup> mice. These T cells produced IL-13 (Supplementary Fig. 1) despite their lack of detectable expression of GATA-3 protein. These results demonstrate that noncanonical IL-13-producing CD4<sup>+</sup> T cells can be generated under T<sub>H1</sub>-priming conditions. We speculated that repetitive T<sub>H1</sub> priming would engender plasticity in T<sub>H1</sub> cells, enabling them to express IL-13. To test our hypothesis, we obtained naive CD4<sup>+</sup> T cells from mice transgenic for the ovalbumin (OVA)-specific TCR (DO11.10 mice), stimulated the cells weekly for 1–4 weeks (rounds 1–4 (R1–R4)) with OVA in the presence of APCs under T<sub>H1</sub>-skewing conditions and analyzed cytokine production every week up to 4 weeks. As expected, abundant IFN- $\gamma$  was induced after the initial antigen priming, and this gradually increased with serial stimulation. Although the cells never made IL-4 or IL-5, IL-13 was detectable after the third round of stimulation (R3), and this increased with further stimulation (Fig. 1a). We observed similar IL-13-producing T<sub>H1</sub> cells derived from *Il4r*<sup>-/-</sup> and *Stat6*<sup>-/-</sup> mice (Fig. 1b), both of which have very few T<sub>H2</sub> cells. Intracellular cytokine staining (ICS) analysis showed that 8–12% of the IFN- $\gamma$ -producing *Il4r*<sup>-/-</sup> T<sub>H1</sub> cells also expressed IL-13 after four rounds of stimulation (R4), although no cells produced IL-13 after the first round (R1; Fig. 1c). Repetitive stimulation therefore induces IL-13 production by T<sub>H1</sub>-skewed CD4<sup>+</sup> T cells.

## E4BP4 regulates IL-13 expression by T<sub>H</sub>1 cells

Although *Il13* expression by T<sub>H</sub>2 cells is regulated mainly by GATA-3 (refs. 20,22,31), our data suggested that *Il13* expression by T<sub>H</sub>1 cells is regulated by unknown mechanisms that are independent of this transcription factor. Activation of these mechanisms correlated with repetitive T cell activation; thus, we sought to identify the IL-13 regulators specifically expressed in R4 T<sub>H</sub>1 cells but not in conventional T<sub>H</sub>1 cells. Microarray analysis identified more than 50 genes with fivefold higher mRNA expression in R4 T<sub>H</sub>1 cells (Supplementary Fig. 2). Among these genes, 16 encoded molecules with features characteristic of transcription factors, and we confirmed their enhanced expression in R4 T<sub>H</sub>1 cells by quantitative RT-PCR (Fig. 2a). To determine which of the candidate genes encoded a molecule that could induce IL-13 expression, we ectopically expressed each in conventional T<sub>H</sub>1 cells by retroviral transduction. Only overexpression of *E4bp4* induced readily detectable IL-13 protein (Supplementary Fig. 3). In contrast, overexpression of *Maf*, *Junb* or *Nr2c1*, which all encode molecules thought to be regulators of T<sub>H</sub>2 differentiation, did not induce IL-13 production by T<sub>H</sub>1 cells (Fig. 2b). However, IL-4 was nearly below the level of detection in the *E4bp4*-transduced cells (Fig. 2b). The tempo of IL-13 production was accelerated in the transduced cells; single-cell analysis after intracellular cytokine staining demonstrated that more than 20% of the E4BP4-expressing cells produced IL-13 after only a single round of T<sub>H</sub>1 stimulation (Fig. 2c).

Published studies based on microarray data have shown *E4bp4* expression mainly in T<sub>H</sub>2 cells<sup>29,30</sup>. We confirmed that finding but also found that *E4bp4* mRNA was markedly enhanced in T<sub>H</sub>1 cells after chronic T<sub>H</sub>1 stimulation (Fig. 2d). This promiscuous expression in both subsets was unique to *E4bp4*, as expression of mRNA for the transcription factor T-bet (*Tbx21* mRNA) and *Gata3* mRNA was limited to T<sub>H</sub>1 and T<sub>H</sub>2 cells, respectively. Next, to determine whether *E4bp4* expression directly correlated with IL-13 expression at the single-cell level, we did immunofluorescence staining of R4 T<sub>H</sub>1 cells generated from DO11.10 mice that lacked T<sub>H</sub>2 cells because of *Il4r* deletion (*Il4r*<sup>-/-</sup> DO11.10 mice). E4BP4 protein was located mainly in the nucleus, and 90% of IL-13-producing cells had readily detectable E4BP4 expression (Fig. 2e). As expected, IL-13 was coexpressed with IFN- $\gamma$  in the stimulated cells (Fig. 2e).

To test the generality of our observations in mice, we examined naive (CD45RA<sup>hi</sup>) CD4<sup>+</sup> T cells from human peripheral blood (Supplementary Fig. 4a). Three rounds of stimulation with monoclonal antibody (mAb) to CD3 under T<sub>H</sub>1 conditions induced IL-13 production in ~9% of cells, of which ~97% were double producers of IL-13 and IFN- $\gamma$  (Supplementary Fig. 4b). These cells also had nuclear expression of E4BP4 protein, and about 90% of E4BP4-expressing cells CD4<sup>+</sup> T cells produced IL-13 (Supplementary Fig. 4c). Therefore, there was a correlation between E4BP4 protein and IL-13 expression in human T<sub>H</sub>1 cells after chronic TCR stimulation.

Expression of *E4bp4*, as well as of *Gata3*, was inducible in T<sub>H</sub>2 cells<sup>29,30</sup> (Fig. 2d). To begin to assess the mechanisms of regulation of *E4bp4* expression in T<sub>H</sub>2 cells, we first tested the importance of the IL-4–STAT6 axis and confirmed that both neutralization of IL-4 and deficiency in *Stat6* attenuated *E4bp4* expression in T<sub>H</sub>2-skewed cells (Supplementary Fig. 5). However, ectopic expression of *Gata3* failed to induce *E4bp4* expression (Supplementary Fig. 5), which suggested that *E4bp4* expression in T<sub>H</sub>2 cells is regulated by the IL-4–STAT6 pathway and not by GATA-3.

## *E4bp4* regulates the expression of IL-13 and IL-10 in CD4<sup>+</sup> T cells

To study the role of E4BP4 *in vivo*, we established mice expressing a transgene encoding Flag-tagged E4BP4. Three independent lines had expression of Flag-tagged E4BP4 in 1–6%

of CD4<sup>+</sup> T cells (Fig. 3a). Enforced expression of E4BP4 in CD4<sup>+</sup> T cells resulted in more IL-13 production at 48 h after initial activation without affecting the production of IL-4 or IL-5 (Fig. 3b). Furthermore, this *E4bp4* transgene also induced the expression of IL-3 as well as of IL-10 (Fig. 3b). The effect on IL-3 production was expected, as E4BP4 is a known inducer of this cytokine<sup>24</sup>; however, the effect on IL-10 expression was completely unexpected. To further examine the possible regulation of both *Il13* and *Il10* by E4BP4, we assessed cytokine production and transgene expression in single cells. Production of IL-10 and IL-13 was augmented in cells expressing the *E4bp4* transgene compared with that in transgene-negative CD4<sup>+</sup> T cells, whereas IL-4 production was the same in both cell types (Fig. 3c). Moreover, retroviral overexpression of *E4bp4* induced the production of IL-10 as well as of IL-13 in T<sub>H</sub>1 cells (Fig. 3d,e). These results suggested that E4BP4 ‘preferentially’ induces the expression of both *Il10* and *Il13*.

### E4BP4 is essential for the expression of *Il10* and *Il13*

Given their apparent coregulation by E4BP4, we next examined the production of both IL-10 and IL-13 in R4 T<sub>H</sub>1 cells. Repetitive T<sub>H</sub>1 priming induced both cytokines (Fig. 4a). To assess whether the expression of *Il10* and *Il13* required E4BP4 expression in T<sub>H</sub>1 cells, we used a retrovirus vector encoding short hairpin RNA (shRNA) specific for *E4bp4* to suppress its expression. Knockdown of *E4bp4* under T<sub>H</sub>1-skewing conditions markedly inhibited the production of IL-10 and IL-13 normally observed after the third or fourth round of stimulation but had no effect on IFN- $\gamma$  production during the first round of stimulation (Fig. 4b), which indicated that E4BP4 is the critical factor that regulates the production of both IL-10 and IL-13 in chronically activated T<sub>H</sub>1 cells.

To better understand the mechanistic effects of E4BP4 on T cell function, especially in terms of regulation of cytokine production, we generated mice lacking *E4bp4* (Supplementary Fig. 6a). The development of thymic T cells and splenic T cells and B cells in *E4bp4*-deficient mice was similar to that of their littermates (Supplementary Fig. 6b). Moreover, T cell proliferation after TCR engagement and the expression profiles of the T cell–activation markers CD25 and CD69 were similar for both wild-type and *E4bp4*-deficient splenic T cells (Supplementary Fig. 6c,d). In contrast, there was significantly less production of IL-10 and IL-13 by R4 T<sub>H</sub>1 cells derived from the *E4bp4*<sup>-/-</sup> mice than by those from their littermates (Fig. 4c). We next examined the effect of *E4bp4* deficiency on the differentiation of T<sub>H</sub>1, T<sub>H</sub>2, T<sub>H</sub>17 and induced T<sub>reg</sub> cells. *E4bp4*<sup>-/-</sup> T cells showed normal production of IFN- $\gamma$ , IL-4 and IL-17 as well as normal induction of T<sub>reg</sub> cells by transforming growth factor- $\beta$  (Supplementary Fig. 6e and data not shown).

Mouse memory T<sub>H</sub>1 cells and human T<sub>H</sub>1 cells produce IL-13 when they are generated in the presence of IL-18 (refs. 32,33). However, E4BP4 is dispensable for IL-18-induced IL-13 production, because *E4bp4*<sup>-/-</sup> T<sub>H</sub>1 cells expressed IL-13 after stimulation with IL-18 (Supplementary Fig. 7a). IL-10 production by T<sub>H</sub>1 cells after stimulation with IL-27 has also been reported<sup>34</sup>. Indeed, control T<sub>H</sub>1 cells expressed IL-10 in the presence of IL-27, whereas *E4bp4*<sup>-/-</sup> T<sub>H</sub>1 cells did not (Supplementary Fig. 7b). Moreover, although IL-10 production has been reported in T<sub>H</sub>17-polarizing conditions<sup>35</sup>, we were unable to confirm that result (Supplementary Fig. 7c). Therefore, all these data support the hypothesis that E4BP4 is required for IL-13 expression by T<sub>H</sub>1 cells in which plasticity has been induced by chronic stimulation and for IL-10 expression when plasticity is induced by IL-27 stimulation.

### E4BP4 regulates *Il10* expression by T<sub>H</sub>2 cells

Next we analyzed the role of E4BP4 in the production of IL-10 and IL-13 by T<sub>H</sub>2 cells. We obtained naive T cells from *E4bp4*<sup>-/-</sup> or wild-type mice and stimulated the cells under T<sub>H</sub>2-



polarizing conditions. About 50% of the *E4bp4*<sup>+/+</sup> T cells generated under T<sub>H</sub>2 conditions produced IL-13 or IL-10, ~20% of the IL-4<sup>+</sup> cells were also IL-10<sup>+</sup>, and ~25% of the IL-13<sup>+</sup> cells also produced IL-10. In contrast, the population of IL-10<sup>+</sup> cells among the *E4bp4*<sup>-/-</sup> T<sub>H</sub>2 cells, was much smaller than that of cells from the wild-type littermates (Fig. 5a), although the production of IL-4, IL-5 and IL-13 by these cells was similar to that of wild-type T cells (Fig. 5b,c).

To further confirm that the effect of E4BP4 on CD4<sup>+</sup> T cells was intrinsic, we crossed mice transgenic for *Cd4*-driven expression of Cre recombinase (*Cd4-Cre*) with our mouse line with conditional deficiency in *E4bp4* to generate mice with CD4<sup>+</sup> T cell-specific deletion of *E4bp4* (*E4bp4*<sup>fl/fl</sup>*Cd4-Cre* mice); although the deletion of *E4bp4* in CD4<sup>+</sup> T cells was partial, it was still greater than 80% (Supplementary Fig. 6a). Again, IFN- $\gamma$  production by *E4bp4*<sup>fl/fl</sup>*Cd4-Cre* T<sub>H</sub>1 cells and the expression of IL-4 and IL-13 by *E4bp4*<sup>fl/fl</sup>*Cd4-Cre* T<sub>H</sub>2 cells were similar to that of the *Cd4-Cre* control mice. However, IL-10 expression in T<sub>H</sub>2 cells was much lower than that of the *Cd4-Cre* control cells (Fig. 5d). As expected, neither wild-type nor *E4bp4*<sup>fl/fl</sup> *Cd4-Cre* T<sub>H</sub>1 cells had detectable production of IL-10.

We next determined whether the production of IL-10 or IL-13 by T cells was also dependent on GATA-3. To determine if E4BP4 requires GATA-3 for IL-10 production, we transduced *E4bp4*<sup>+/+</sup> and *E4bp4*<sup>-/-</sup> T cells with a *Gata3*-expressing retroviral vector. Exogenous *Gata3* induced the generation of IL-10-producing *E4bp4*<sup>+/+</sup> T cells slightly, but the response was completely abolished in *E4bp4*<sup>-/-</sup> T cells (Fig. 5e, top row). In addition, we transduced a retroviral vector expressing Cre into T cells with loxP-flanked *Gata3* (*Gata3*<sup>fl/fl</sup>) and into wild-type C57BL/6 T cells. We further transduced cells with a retroviral vector expressing *E4bp4* and found that ectopic expression of E4BP4 protein significantly induced the development of IL-10<sup>+</sup> T cells with or without *Gata3* expression (Fig. 5e, bottom). We further examined the production of IL-10 and IL-13 by T<sub>H</sub>1 cells using CD4<sup>+</sup> T cells derived from *Gata3*<sup>fl/fl</sup>*Cd4-Cre* mice. IL-13 production in R4 T<sub>H</sub>1 cells lacking *Gata3* was ~50% lower, but there was no effect on IL-10 production (Fig. 5f). These results indicate that E4BP4 is a critical factor for *Ii10* expression in CD4<sup>+</sup>T cells independently of *Gata3* expression and that *Ii13* expression in T<sub>H</sub>1 cells requires low *Gata3* expression.

### E4BP4 regulates the expression of *Ii13* and *Ii10* in NKT cells

Various T cell subsets, including NKT cells and T<sub>reg</sub> cells, produce IL-10 and IL-13, and we hypothesized that E4BP4 is a regulator of the expression of *Ii10* and *Ii13* in these cells as well. To test our hypothesis, we first determined if there was a correlation between the expression *E4bp4* and the expression of both *Ii10* and *Ii13* by NKT cells. We stimulated whole splenocytes with the NKT cell ligand  $\alpha$ -galactosyl ceramide ( $\alpha$ -GalCer) and analyzed cytokine production cells by ICS after gating on cells that recognized a dimer of  $\alpha$ -GalCer and the antigen-presenting molecule CD1d. About 55% of NKT cells produced IL-13, and all of the IL-10-producing cells also produced IL-13 (Fig. 6a). We also observed that *E4bp4* expression was induced by stimulation with  $\alpha$ -GalCer, which suggested a possible role for E4BP4 in the expression of *Ii10* and *Ii13* by NKT cells. It has been reported that the treatment of IL-17RB<sup>+</sup> NKT cells with IL-25 induces high *Ii13* expression<sup>36</sup>; therefore, we focused on a possible role for E4BP4 in inducing the expression of *Ii10* and *Ii13* in this NKT cell subset. We isolated IL-17RB<sup>+</sup> NKT cells from BALB/c mice and stimulated the cells with IL-25 in the presence of DCs. *E4bp4* expression was induced considerably in the IL-25-treated cells, and its expression correlated well with the expression of *Ii10* and *Ii13* (Fig. 6b). The frequency of splenic NKT cells was nearly identical in *E4bp4*<sup>-/-</sup> and wild-type mice (Fig. 6c); however, in response to stimulation with  $\alpha$ -GalCer, *E4bp4*<sup>-/-</sup> NKT cells had no detectable IL-10 expression and lower IL-13 expression, and in response to IL-25, *E4bp4*<sup>-/-</sup> NKT cells had lower IL-13 expression and both wild-type and *E4bp4*<sup>-/-</sup> NKT

cells had no detectable IL-10, whereas the expression of IFN- $\gamma$  and IL-4 was unaffected (Fig. 6d).

To study the involvement of E4BP4 in IL-10 production by T<sub>reg</sub> cells, we isolated naive CD4<sup>+</sup> T cells (Foxp3<sup>-</sup>CD44<sup>lo</sup>CD62L<sup>hi</sup>), memory CD4<sup>+</sup> T cells (Foxp3<sup>-</sup>CD44<sup>hi</sup>CD62L<sup>lo</sup>) and natural T<sub>reg</sub> cells (Foxp3<sup>+</sup>CD25<sup>-</sup> or Foxp3<sup>+</sup>CD25<sup>hi</sup>) from mice expressing the transcription factor Foxp3 from a human *CD2* reporter (Foxp3<sup>h</sup>CD2 mice)<sup>37</sup>, which allowed better separation of Foxp3<sup>+</sup> cells from other live cells on the basis of the expression of human CD2, and stimulated the cells through the TCR. Natural T<sub>reg</sub> cells expressed IL-10 but not IL-13 after stimulation of the TCR (Supplementary Fig. 8a). We found high expression of *E4bp4* in *Il10*-expressing natural T<sub>reg</sub> cells, particularly in the CD25<sup>-</sup> subset, and we also detected this in memory T cells, which also expressed *Il10*. In contrast, naive T cells had low expression of both *E4bp4* mRNA and *Il10* mRNA (Supplementary Fig. 8a). IL-10 production was impaired in natural T<sub>reg</sub> cells and memory T cells derived from *E4bp4*<sup>-/-</sup> mice, but IL-13 production was not (Fig. 6e,f and Supplementary Fig. 8b,c). These results indicate that *E4bp4* expression is correlated with the expression of *Il10* and *Il13* by NKT cells, memory T cells and IL-10-expressing natural T<sub>reg</sub> cells, which suggests that E4BP4 regulates *Il10* expression in all T cell subsets and *Il13* expression in NKT cells.

### E4BP4 binds specifically to the *Il13* and *Il10* loci

The *Il4* locus in T<sub>H1</sub> cells is thought to be maintained in a repressed state by epigenetic modifications<sup>38</sup>. However, as shown above, chronic T<sub>H1</sub> stimulation promoted IL-13 production, which suggests that the *Il13* and *Il4* loci may have distinct modes of regulation. Therefore, we studied methylation and histone modifications of the *Il13* locus in T<sub>H1</sub> cells. The *Il13* distal promoter contains two well characterized DNase I-hypersensitive sites, HS1 and HS2, which are highly conserved in mammals. We used a bisulfite-sequencing PCR assay to investigate the DNA-methylation status, especially of the HS1 region, in naive T cells and in developing T<sub>H1</sub> and T<sub>H2</sub> cells (Fig. 7a). Most CpG dinucleotides were equally unmethylated in naive CD4<sup>+</sup> cells; however, we observed a few methylated sites in both developing T<sub>H1</sub> cells and T<sub>H2</sub> cells (Fig. 7a). We also investigated histone modification of the *Il13* and *Il4* loci in various T<sub>H1</sub> cell developmental stages by chromatin immunoprecipitation (ChIP) and found considerable enrichment of histone H3 trimethylated at Lys4 in the HS1 region of the *Il13* promoter, even in T<sub>H1</sub> cells stimulated for 1 week (R1; Fig. 7b, top). We then used ChIP analysis with nuclear extracts prepared from conventional T<sub>H1</sub> and R4 T<sub>H1</sub> cells to determine whether E4BP4 binds to the *Il13* promoter. There was considerable enrichment of specific binding of E4BP4 to the HS1 site in R4 T<sub>H1</sub> cells but not in conventional T<sub>H1</sub> cells (Fig. 7b, bottom).

We then used electrophoretic mobility-shift assay to identify the putative E4BP4-binding site on the *Il13* promoter. We prepared nuclear extracts of R4 T<sub>H1</sub> cells and T<sub>H2</sub> cells and tested for binding of E4BP4 to oligonucleotides corresponding to the putative binding sites in the adenovirus *E4* promoter and the mouse *Il13* promoter. We found a distinctly shifted band in T<sub>H2</sub> cells and a slight shift even in T<sub>H1</sub> cells (Fig. 7c). This binding was blocked by treatment with either antibody to E4BP4 (anti-E4BP4) or an excess of unlabeled competitor oligonucleotide corresponding to the HS1 site of the *Il13* promoter (Fig. 7c). The HS1 site corresponds to a conserved GATA-3-response element (CGRE) identified as a distal promoter for *Il13*, and CGRE contains a consensus E4BP4-binding sequence as well as a GATA-3-binding site<sup>39</sup>. We found that 2-kilobase fragments including the CGRE had strong promoter activity in a TCR-stimulation assay, and deletion of the CGRE from the promoter region attenuated this activity even when *E4bp4* was overexpressed (Supplementary Fig. 9a). Thus, binding of E4BP4 to the CGRE regulates *Il13* promoter activity. Together these results demonstrate that binding of E4BP4 to the CGRE site of the *Il13* promoter promotes the expression of IL-13 in T cells. The HS1 site in the *Il13* promoter is transcriptionally

permissive even in T<sub>H</sub>1 cells, which suggests that E4BP4 expression may directly govern the plasticity of *Il13* in T<sub>H</sub>1 cells.

We next did a similar study of the *Il10* locus, in which we identified 15 possible binding sites on the basis of the predicted E4BP4-binding sequence (Fig. 7d). ChIP analysis showed enrichment for histone H3 acetylated at Lys9 or Lys14 in some of the binding sites. We observed such enrichment in these regions mainly in T<sub>H</sub>2 cells but not in conventional T<sub>H</sub>1 cells. Further ChIP analysis showed that E4BP4 bound to intron 4 and the 3' noncoding region of the *Il10* locus (Fig. 7d). Two 5' upstream fragments of *Il10* 400 and 1,000 in length that do not contain consensus binding sequences for E4BP4 had no promoter activity in the T cell-activation assay (Supplementary Fig. 9a), which indicated that E4BP4 does not regulate the activation-induced promoter activity. To unequivocally demonstrate a role for E4BP4 in the epigenetic modification of the *Il10* locus, we analyzed *E4bp4*<sup>-/-</sup> cells. The strong enrichment for acetylated histone H3 in the *Il10* locus noted above was completely absent from *E4bp4*<sup>-/-</sup> T<sub>H</sub>2 cells (Fig. 7e). From these results, we conclude that E4BP4 directly regulates the permissive status of the *Il10* locus at the chromatin level.

### E4BP4 in the control for autoimmune responses

Mice with complete ablation of *Il10* or T cell-specific ablation of *Il10* (*Il10*<sup>f/f</sup>*Cd4*-Cre mice) develop spontaneous intestinal inflammation<sup>40</sup>, and T<sub>reg</sub> cell-specific ablation of *Il10* also results in spontaneous colitis, albeit a milder version<sup>18</sup>. Such results demonstrate that IL-10 produced by T<sub>reg</sub> cells has an important role in suppressing immune inflammation in the intestine. Indeed, about half of the *E4bp4*<sup>-/-</sup> mice spontaneously developed mild diarrhea (data not shown), and induced T<sub>reg</sub> cells generated from *E4bp4*<sup>-/-</sup> mice had impaired IL-10 production after TCR stimulation even though control induced T<sub>reg</sub> cells had detectable, but low, expression of IL-10 (Fig. 8a). Given those findings, we examined the involvement of E4BP4 in T cells in a T cell-transfer colitis model in which we reconstituted lymphopenic mice deficient in recombination-activating gene 1 with CD4<sup>+</sup>CD44<sup>hi</sup>CD62L<sup>lo</sup>CD25<sup>-</sup> naive T cells from control littermate mice or *E4bp4*<sup>-/-</sup> mice and monitored the recipient mice for the development of inflammatory disease. Mice reconstituted with control T cells gained weight for the first 2 weeks but then began to gradually lose weight (Fig. 8b). In contrast, mice reconstituted with *E4bp4*<sup>-/-</sup> T cells gained less weight and had more severe weight loss and a worse clinical score, including death (two of ten mice; Fig. 8b). Five of the ten mice reconstituted with *E4bp4*<sup>-/-</sup> T cells began to have diarrhea after 4 weeks (data not shown). We confirmed the clinical symptoms by evaluating colonic inflammation morphologically and by histology (Fig. 8c) and also found that *E4bp4*<sup>-/-</sup> T cells derived from intestinal lymph nodes and colon did not express IL-10 (Fig. 8d). Thus, regulation of IL-10 in T cells by E4BP4 is important for tolerance induction to prevent the development of chronic inflammatory disease in the colon.

We further examined the importance of the regulation of IL-10 by E4BP4 in an experimental autoimmune encephalomyelitis (EAE) model. We immunized *E4bp4*<sup>-/-</sup> mice and their control littermates with a peptide of myelin oligodendrocyte glycoprotein (amino acids 35–55) emulsified in complete Freund's adjuvant. Control and *E4bp4*<sup>-/-</sup> mice showed notable EAE development that reached a maximum at 15 d (Fig. 8e). Control mice began to recover, showing improvement in their clinical scores after 25 d. In contrast, *E4bp4*<sup>-/-</sup> mice showed no evidence of recovery and had worse EAE scores after 30 d (Fig. 8e). *E4bp4*<sup>-/-</sup> T cells from the regional lymph node expressed no IL-10 and had a slightly more IL-17 production (Fig. 8f). We further monitored *Il10* transcription in mice with ongoing disease using a reporter system in which sequence encoding the yellow fluorescent protein Venus was knocked into the 3' untranslated region of the *Il10* locus<sup>41</sup>. Fewer *E4bp4*<sup>-/-</sup> T cells from the spleen, among intestinal intraepithelial lymphocytes or from the lamina propria or



bone marrow produced IL-10 than did those from control mice (Supplementary Fig. 10). Therefore, E4BP4 induction of IL-10 is also important for EAE remission.

## DISCUSSION

We have demonstrated here that chronic antigen stimulation resulted in plasticity of T<sub>H</sub>1 cells, which then secreted certain T<sub>H</sub>2 cytokines, including IL-10 and IL-13, but not IL-4 or IL-5. Acquisition by T<sub>H</sub>1 cells of the ability to express IL-10 and IL-13 was regulated by E4BP4, whose own expression was upregulated by chronic antigenic stimulation. E4BP4 also regulated the production of IL-10 and IL-13 in NKT cells, memory T cells and T<sub>reg</sub> cells. In contrast, *Il13* regulation in T<sub>H</sub>2 cells was totally independent of E4BP4, which indicated that E4BP4 regulates *Il13* differently in distinct CD4<sup>+</sup> T cell subsets but commonly regulates *Il10* in all CD4<sup>+</sup> T cells. Therefore, we propose that E4BP4 has multiple functions in the regulation of genes encoding cytokines and is a transcriptional factor essential to the regulation of IL-10 production. Our results provide the new insight that different T<sub>H</sub>2 cytokines are regulated independently by a distinct combination of transcription factors in different T cell subsets.

During the past two decades, it has become widely accepted that *Il10* and *Il13* are controlled together with *Il4* by common transcriptional mechanisms. This idea has been supported by many observations; for example, that upregulation of *Gata3* expression is essential for the expression of IL-4 and IL-13 in T<sub>H</sub>2 cells<sup>42</sup>, and that conditional deletion of *Gata3* in activated CD4<sup>+</sup> T cells attenuates the production of both cytokines<sup>22</sup>. However, we found no defect in IL-13-induced airway hyper-responsiveness in *Il4*<sup>-/-</sup> mice, which completely lack T<sub>H</sub>2 cells, which indicated that T<sub>H</sub>2 cells are not the source of IL-13 under certain conditions<sup>43,44</sup>. The source of IL-13 in *Il4*<sup>-/-</sup> mice has thus far remained unknown, but our study has shown that because of their plasticity, T<sub>H</sub>1 cells are an alternative source of this cytokine. In terms of IL-10, its production by T<sub>H</sub>1 cells has been reported after chronic infection with *T. gondii* or *L. major*<sup>13,15</sup>. IL-10 production is thus induced by relatively strong chronic stimulation, as in parasite infection, and this is consistent with our data showing that chronic antigen stimulation induced the expression of E4BP4, which controls the production of IL-10 and IL-13 by T<sub>H</sub>1 cells.

*E4bp4* is expressed mainly by T<sub>H</sub>2 cells<sup>29,30</sup>, and we have demonstrated that *E4bp4* expression was controlled by the IL-4–STAT6 pathway. However, the question of how chronic T<sub>H</sub>1 stimulation induces *E4bp4* expression remains. A published report has indicated that calcium signaling–mediated activation of the transcription factor NFAT induces *E4bp4* expression<sup>45</sup>, and *Nfat5* was one of the genes we found was induced by repetitive stimulation, although its overexpression did not induce *E4bp4* expression. None of the 15 other candidate genes that showed differences in upregulation in R4 T<sub>H</sub>1 cells affected *E4bp4* expression; therefore, further investigation is need to clarify how chronic T<sub>H</sub>1 stimulation induces *E4bp4* expression.

E4BP4 was originally identified as a negative regulator of the mammalian circadian oscillatory system<sup>23</sup>. *E4bp4*<sup>-/-</sup> mice were born at normal Mendelian ratios and grew normally, which indicated that deficiency in E4BP4 has no severe phenotypic consequences except the spontaneous development of mild diarrhea. In the absence of E4BP4, the phenotype of CD4<sup>+</sup> T cells, including NKT cells, was fairly normal, consistent with the published observation that deletion of *E4bp4* affects mainly NK cell development with no change in the proportion of other T cell lineages<sup>26,27</sup>. T cells lacking *E4bp4* had normal T<sub>H</sub>1, T<sub>H</sub>2, induced T<sub>reg</sub> and T<sub>H</sub>17 differentiation, although the proportion of T<sub>H</sub>1 cells and T<sub>H</sub>2 cells was slightly lower than that in wild-type mice. *E4bp4*-deficient T<sub>H</sub>2 cells were considerably impaired in IL-10 production but not in GATA-3 expression or the production

of other T<sub>H</sub>2 cytokines such as IL-4 and IL-5, and IL-13 production was even enhanced in *E4bp4*<sup>-/-</sup> T<sub>H</sub>2 cells.

GATA-3 directly regulates the *Il13* promoter<sup>20</sup> and, furthermore, conditional deletion of *Gata3* in T<sub>H</sub>2-committed T cells completely attenuates *Il13* expression<sup>22,31</sup>, which suggests that GATA-3 is indispensable for *Il13* transcription in T<sub>H</sub>2 cells but E4BP4 is not. In contrast, E4BP4 was absolutely required for IL-13 expression in T<sub>H</sub>1 cells. The distal *Il13* promoter, CGRE, 1.4–1.8 kilobases upstream of exon 1, contains 42 CpG dinucleotides<sup>46</sup>, and analysis of DNA dimethylation and methylation of histone H3 at Lys4 indicated that the CGRE is transcriptionally accessible in naive T cells, T<sub>H</sub>1 cells and T<sub>H</sub>2 cells. Reporter assays indicated the activity of this promoter is regulated by E4BP4. Therefore, IL-13 production by T<sub>H</sub>1 cells seems to be determined by the regulation of the promoter at the level of E4BP4 expression rather than by epigenetic regulation.

We have demonstrated that E4BP4 was also essential for IL-10 expression in most of the CD4<sup>+</sup> subsets we examined. The precise mechanism by which IL-10 expression is regulated in T cells has remained unclear, although a contribution by GATA-3 has been demonstrated by retroviral overexpression experiments<sup>21</sup>. Our data indicated that E4BP4 induced IL-10 production in the absence of GATA-3, which suggests that GATA-3 is dispensable for IL-10 production. Therefore, E4BP4 seems to be an essential regulator of IL-10 expression not only in T<sub>H</sub>2 cells but also in T<sub>H</sub>1 cells, which typically have very low expression of GATA-3. However, unlike its binding to *Il13*, E4BP4 did not bind to the *Il10* promoter region, binding instead to an intronic region and a 3' noncoding sequence, both of which are well conserved between humans and mice. T<sub>H</sub>2 cells lacking E4BP4 had no permissive chromatin structure in these regions, which suggests that *Il10* is epigenetically regulated by E4BP4 binding. However, T cells are not the only source of IL-10, and IL-10 is reported to be secreted by NK cells, regulatory B cells and regulatory DCs. Indeed, regulatory DCs from the lamina propria of *E4bp4*<sup>-/-</sup> mice had lower expression of IL-10 (data not shown), which suggests that E4BP4 may function as a common transcriptional regulator for IL-10 production in a variety of cell types.

## ONLINE METHODS

### Mice

DO11.10 mice on the BALB/c background were provided by K. Murphy; and *Stat6*<sup>-/-</sup> mice, *Il4*<sup>-/-</sup> mice and *Cd4*-Cre mice were gifts from S. Akira<sup>47</sup>, F. Brombacher<sup>48</sup> and C. Wilson, respectively. Foxp3<sup>hCD2</sup> reporter mice<sup>37</sup> and Venus reporter mice<sup>41</sup> were maintained on the C57BL/6 background. Mice were kept in specific pathogen-free conditions in accordance with the guidelines of the Institutional Animal Care Committee of the RIKEN Institute.

For the generation of *E4bp4*-transgenic mice, cDNA encoding a Flag-tagged E4BP4 fusion protein was inserted downstream of the E<sub>μ</sub> enhancer of the immunoglobulin heavy-chain locus and was injected into fertilized mouse eggs. Three independent transgenic mouse lines (1, 2 and 3) were selected for study on the basis of intracellular staining with anti-Flag (M-2; Sigma). The mice were backcrossed to C57BL/6 mice for more than ten generations.

*E4bp4*-deficient mice were generated by homologous recombination and replacement of *E4bp4*, in 129/C57BL/6 hybrid embryonic stem cells, with a loxP-flanked cassette (RIKEN Center for Developmental Biology) containing the coding sequence of mouse *E4bp4* (Supplementary Fig. 6). A neomycin-resistance expression cassette flanked by flippase recognition target sites was located 3' of the targeting site. After germline deletion of those elements by crossing of those mice with mice expressing Cre recombinase (CAGcre-

transgenic mice)<sup>49</sup>, the resulting mice were backcrossed with C57BL/6 mice for five generations.

### Cell preparation and reagents

CD4<sup>+</sup> T cells were isolated from spleen with the IMag magnetic bead system (BD Biosciences) and naive (CD44<sup>lo</sup> CD62L<sup>hi</sup>) cells were isolated by cell sorting with a FACSVantage (BD Biosciences). T cells ( $1 \times 10^6$  cells per ml) were stimulated with plate-bound mAb to TCR $\beta$  (H57-597) and mAb to CD28 (PV-1). For DO11.10 CD4<sup>+</sup> T cells, cells were stimulated with the antigenic peptide Loh15 (1  $\mu$ g/ml) in the presence of irradiated spleen cells ( $1 \times 10^7$  cells per ml). For the induction of T<sub>H</sub>2 cells, cells were stimulated in the presence of mouse IL-4 (100 U/ml) and mAb to IFN- $\gamma$  (R46A2). For T<sub>H</sub>1 polarization, cells were cultured in the presence of mouse IL-12 and mAb to IL-4 (11B11). For T<sub>H</sub>17 and T<sub>reg</sub> induction, cells were cultured with recombinant human transforming growth factor- $\beta$  (10 ng/ml) with or without recombinant mouse IL-6 (4 ng/ml) in the presence of mAb to IL-4 (11B11) and mAb to IFN- $\gamma$  (R46A2). For the preparation of NKT cells and T<sub>reg</sub> cells, cells were sorted with biotinylated mAb to IL-17RB (3H8)<sup>36</sup>, phycoerythrin-conjugated mAb to CD25 (PC61.5; 12-0251; eBioscience), biotinylated mAb to human CD2 (RPA2.10; 300203; BioLegend) and allophycocyanin-conjugated mAb to CD44 (1M7; 559250; BD Biosciences).

Human CD4<sup>+</sup> T cells were isolated from peripheral blood mononuclear cells with IMag beads, and cells were stimulated for 7 d with mAb to CD3 (OKT3; 14-0037-82; eBioscience) and mAb to CD28 (CD28.2; 555726; BD Biosciences) in the presence of human IL-12 (5 ng/ml) and mAb to IL-4 (8D48; 556917; BD Biosciences) for the induction of T<sub>H</sub>1 cells.

### Measurement of cytokine production

T cells were stimulated for 24 h with plate-bound mAb to TCR plus mAb to CD28. The concentrations of IL-4, IL-5, IL-13 and IFN- $\gamma$  in culture supernatants were measured by ELISA<sup>50</sup>; mAb to IFN- $\gamma$  (R4-6A2; 551216; BD Biosciences), mAb to IL-4 mAb(BVD4-1D11; 554387; BD Biosciences), mAb to IL-5 (JES6-1A12; 554393; BD Biosciences) and mAb to IL-13 (DY413; 38213; R&D Systems) were used for capture, and mAb to IFN- $\gamma$  (XMG1.2; 554410; BD Biosciences), mAb to IL-4 (BVD6-24G2; 13-7042; eBioscience), mAb to IL-5 (JES6-5H4; 554397; BD Biosciences) and mAb to IL-13 (DY413) were used for detection. Cytokines were measured by a suspension array system (Luminex 200; Bio-Rad Laboratories).

### Intracellular cytokine staining and confocal microscopy

CD4<sup>+</sup> T cells were restimulated for 6 h with mAb to TCR $\beta$  in the presence of 2  $\mu$ M monensin. Cells were fixed with 4% (vol/vol) paraformaldehyde and were made permeable with 0.5% (vol/vol) Triton X-100. After blocking, cells were stained with the following: fluorescein isothiocyanate-conjugated mAb to IFN- $\gamma$  (XMG1.2; 554410; BD Biosciences), allophycocyanin-conjugated mAb to IL-4 (11B11; 554436; BD Biosciences) and Alexa Fluor 647-conjugated mAb to IL-13 (eBio-13A; 51-7133; eBioscience). GATA-3 was detected with mAb to GATA-3 (L50-823; 560163; BD Biosciences). A FACSCalibur and CellQuest software (BD Biosciences) were used for flow cytometry. For confocal microscopy, cells were made permeable as described above and then stained with rabbit anti-E4BP4 (H-300; sc-28203; Santa Cruz) and fluorescein isothiocyanate-conjugated rat antibody to rabbit IgG (111-095-003; Jackson Laboratories). Fixed cells were imaged with an LSM 510 confocal microscopy system (Leica Microsystems).

### Microarray analysis and quantitative RT-PCR

Total RNA was isolated with TRIzol (Invitrogen). An Affymetrix microarray chip (Mouse Genome 430 2.0 array) was used for microarray analysis, and results were normalized with Genespring software (Silicon Genetics). A 7500 real-time PCR system (Applied Biosystems) was used for quantitative real-time PCR (primers, Supplementary Table 1).

### Retroviral transduction

The retroviral vector pMX-IRES-GFP was transfected into PLAT-E packaging cells (a gift from T. Kitamura) with FuGENE 6 (Roche), and culture supernatants were used as a source of viral particles. Naive CD4<sup>+</sup> T cells were stimulated with mAb to TCR and mAb to CD28. Viral particles were transduced into the purified CD4<sup>+</sup> T cells, and GFP<sup>+</sup> cells were sorted after 2 d.

### Generation of shRNA for knockdown for *E4bp4*

For generation of the *E4bp4*-specific shRNA plasmid construct, the targeting oligonucleotide 5'-GTTGCATCTCAGTCATCAAGC-3' was cloned into the pSIREN-RetroQ vector (Clontech); shRNA specific for luciferase was used as control. Each shRNA was introduced into activated T cells by retroviral transduction, and transduced cells were selected with puromycin (2 ng/ml).

### Bisulfite sequencing PCR assay

DNA methylation was analyzed with a bisulfite-sequencing PCR assay kit (Chemicon International). Genomic DNA was chemically modified by treatment with sodium bisulfite, then modified DNA was amplified with specific primers by PCR and DNA methylation was detected by sequencing.

### ChIP assay

Nuclear fractions were extracted as chromatin from sonicated cells. After preclearance with protein G agarose (Amersham), chromatin fractions were incubated with the appropriate antibody. Input samples were defined as DNA extracts prepared from untreated chromatin. DNA was extracted from immunoprecipitated chromatin, and concentrations were quantified with a PicoGreen fluorescence assay (Molecular Probes). Equivalent masses of immunoprecipitated and input DNA were compared by real-time PCR (primers, Supplementary Table 1), with results presented as the ratio of the change in threshold value after immunoprecipitation to value to the change in threshold value of input DNA.

### Electrophoretic mobility-shift assay

Nuclear proteins were incubated for 30 min with <sup>32</sup>P-labeled oligonucleotide probes. Samples were separated by 4% PAGE in a buffer of low ionic strength. Shifts in mobility were visualized with a BAS-2500 system (oligonucleotide probes, Supplementary Table 1).

### Colitis model

CD4<sup>+</sup> T cells were prepared from the spleen with IMag magnetic beads, and CD44<sup>lo</sup>CD25<sup>-</sup> naive cells were isolated by cell sorting. Cells ( $4 \times 10^5$ ) were transferred intravenously into mice deficient in recombination-activating gene 1, and body weight and symptoms of clinical disease were monitored. Mice were killed after 34 d. Colons were fixed with 4% (vol/vol) paraformaldehyde and sections were stained with hematoxylin and eosin.

## EAE induction

Mice were immunized subcutaneously on day 0 with a peptide of myelin oligodendrocyte glycoprotein amino acids 35–55 emulsified in complete Freund's adjuvant (100 µg per mouse), and were given intravenous injection of pertussis toxin (200 ng per mouse; List Biological Laboratories) on days 0 and 2. EAE was assessed as follows: 0, no disease; 1, limp tail; 2, weak hindlimbs; 3, partially paralyzed hindlimbs; 4, complete hindlimb paralysis; and 5, death.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

We thank K. Murphy (Washington University) for DO11.10 mice on the BALB/c background; S. Akira (Osaka University) for *Stat6*<sup>-/-</sup> mice; F. Brombacher (University of Cape Town) for *Il4*<sup>-/-</sup> mice; C. Wilson (Washington University) for *Cd4*-Cre mice; R. Abe (Tokyo University of Science) for the PV-1 mAb to CD28; T. Kitamura (Tokyo University) for Platinum-E packaging cells; H. Fujimoto, Y. Suzuki, K. Ikari and E. Hayashi for technical assistance; and P. Burrows for comments on the manuscript. Supported by RIKEN (Y. Motomura), the Ministry of Education, Culture, Sports, Science and Technology of Japan, and the Program for Promotion of Fundamental Studies in Health Sciences of the National Institute of Biomedical Innovation.


## References

1. Murphy KM, Reiner SL. The lineage decisions of helper T cells. *Nat. Rev. Immunol.* 2002; 2:933–944. [PubMed: 12461566]
2. Dong C. T<sub>H</sub>17 cells in development: an updated view of their molecular identity and genetic programming. *Nat. Rev. Immunol.* 2008; 8:337–348. [PubMed: 18408735]
3. Iwakura Y, Nakae S, Saijo S, Ishigame H. The roles of IL-17A in inflammatory immune responses and host defense against pathogens. *Immunol. Rev.* 2008; 226:57–79. [PubMed: 19161416]
4. McKenzie AN, et al. Interleukin 13, a T-cell-derived cytokine that regulates human monocyte and B-cell function. *Proc. Natl. Acad. Sci. USA.* 1993; 90:3735–3739. [PubMed: 8097324]
5. Wynn TA. IL-13 effector functions. *Annu. Rev. Immunol.* 2003; 21:425–456. [PubMed: 12615888]
6. Lee GR, Fields PE, Griffin TJ, Flavell RA. Regulation of the Th2 cytokine locus by a locus control region. *Immunity.* 2003; 19:145–153. [PubMed: 12871646]
7. Wilson CB, Rowell E, Sekimata M. Epigenetic control of T-helper-cell differentiation. *Nat. Rev. Immunol.* 2009; 9:91–105. [PubMed: 19151746]
8. Hayashi N, et al. T helper 1 cells stimulated with ovalbumin and IL-18 induce airway hyperresponsiveness and lung fibrosis by IFN-γ and IL-13 production. *Proc. Natl. Acad. Sci. USA.* 2007; 104:14765–14770. [PubMed: 17766435]
9. Li L, Xia Y, Nguyen A, Feng L, Lo D. Th2-induced eotaxin expression and eosinophilia coexist with Th1 responses at the effector stage of lung inflammation. *J. Immunol.* 1998; 161:3128–3135. [PubMed: 9743380]
10. Smart JM, Kemp AS. Increased Th1 and Th2 allergen-induced cytokine responses in children with atopic disease. *Clin. Exp. Allergy.* 2002; 32:796–802. [PubMed: 11994108]
11. Saraiva M, O'Garra A. The regulation of IL-10 production by immune cells. *Nat. Rev. Immunol.* 2010; 10:170–181. [PubMed: 20154735]
12. Batten M, et al. Cutting edge: IL-27 is a potent inducer of IL-10 but not FoxP3 in murine T cells. *J. Immunol.* 2008; 180:2752–2756. [PubMed: 18292493]
13. Jankovic D, et al. Conventional T-bet<sup>+</sup>Foxp3<sup>-</sup> Th1 cells are the major source of host-protective regulatory IL-10 during intracellular protozoan infection. *J. Exp. Med.* 2007; 204:273–283. [PubMed: 17283209]




14. Shaw MH, et al. Tyk2 negatively regulates adaptive Th1 immunity by mediating IL-10 signaling and promoting IFN- $\gamma$ -dependent IL-10 reactivation. *J. Immunol.* 2006; 176:7263–7271. [PubMed: 16751369]
15. Anderson CF, Oukka M, Kuchroo VJ, Sacks D. CD4<sup>+</sup>CD25<sup>-</sup>Foxp3<sup>-</sup> Th1 cells are the source of IL-10-mediated immune suppression in chronic cutaneous leishmaniasis. *J. Exp. Med.* 2007; 204:285–297. [PubMed: 17283207]
16. Saraiva M, et al. Interleukin-10 production by Th1 cells requires interleukin-12-induced STAT4 transcription factor and ERK MAP kinase activation by high antigen dose. *Immunity.* 2009; 31:209–219. [PubMed: 19646904]
17. Kuhn R, Lohler J, Rennick D, Rajewsky K, Muller W. Interleukin-10-deficient mice develop chronic enterocolitis. *Cell.* 1993; 75:263–274. [PubMed: 8402911]
18. Rubtsov YP, et al. Regulatory T cell-derived interleukin-10 limits inflammation at environmental interfaces. *Immunity.* 2008; 28:546–558. [PubMed: 18387831]
19. O’Shea JJ, Paul WE. Mechanisms underlying lineage commitment and plasticity of helper CD4<sup>+</sup> T cells. *Science.* 2010; 327:1098–1102. [PubMed: 20185720]
20. Kishikawa H, Sun J, Choi A, Miaw SC, Ho IC. The cell type-specific expression of the murine IL-13 gene is regulated by GATA-3. *J. Immunol.* 2001; 167:4414–4420. [PubMed: 11591766]
21. Shoemaker J, Saraiva M, O’Garra A. GATA-3 directly remodels the IL-10 locus independently of IL-4 in CD4<sup>+</sup> T cells. *J. Immunol.* 2006; 176:3470–3479. [PubMed: 16517715]
22. Zhu J, et al. Conditional deletion of GATA-3 shows its essential function in T<sub>H</sub>1–T<sub>H</sub>2 responses. *Nat. Immunol.* 2004; 5:1157–1165. [PubMed: 15475959]
23. Cowell IG. E4BP4/NFIL3, a PAR-related bZIP factor with many roles. *Bioessays.* 2002; 24:1023–1029. [PubMed: 12386933]
24. Zhang W, et al. Molecular cloning and characterization of NF-IL3A, a transcriptional activator of the human interleukin-3 promoter. *Mol. Cell. Biol.* 1995; 15:6055–6063. [PubMed: 7565758]
25. Ikushima S, et al. Pivotal role for the NFIL3/E4BP4 transcription factor in interleukin 3-mediated survival of pro-B lymphocytes. *Proc. Natl. Acad. Sci. USA.* 1997; 94:2609–2614. [PubMed: 9122243]
26. Gascoyne DM, et al. The basic leucine zipper transcription factor E4BP4 is essential for natural killer cell development. *Nat. Immunol.* 2009; 10:1118–1124. [PubMed: 19749763]
27. Kamizono S, et al. Nfil3/E4bp4 is required for the development and maturation of NK cells in vivo. *J. Exp. Med.* 2009; 206:2977–2986. [PubMed: 19995955]
28. Kashiwada M, et al. IL-4-induced transcription factor NFIL3/E4BP4 controls IgE class switching. *Proc. Natl. Acad. Sci. USA.* 2010; 107:821–826. [PubMed: 20080759]
29. Lund R, Aittokallio T, Nevalainen O, Lahesmaa R. Identification of novel genes regulated by IL-12, IL-4, or TGF- $\beta$  during the early polarization of CD4<sup>+</sup> lymphocytes. *J. Immunol.* 2003; 171:5328–5336. [PubMed: 14607935]
30. Lund R, et al. Identification of genes involved in the initiation of human Th1 or Th2 cell commitment. *Eur. J. Immunol.* 2005; 35:3307–3319. [PubMed: 16220538]
31. Pai SY, Truitt ML, Ho IC. GATA-3 deficiency abrogates the development and maintenance of T helper type 2 cells. *Proc. Natl. Acad. Sci. USA.* 2004; 101:1993–1998. [PubMed: 14769923]
32. Hata H, Yoshimoto T, Hayashi N, Hada T, Nakanishi K. IL-18 together with anti-CD3 antibody induces human Th1 cells to produce Th1- and Th2-cytokines and IL-8. *Int. Immunol.* 2004; 16:1733–1739. [PubMed: 15477227]
33. Sugimoto T, et al. Interleukin 18 acts on memory T helper cells type 1 to induce airway inflammation and hyperresponsiveness in a naive host mouse. *J. Exp. Med.* 2004; 199:535–545. [PubMed: 14970180]
34. Stumhofer JS, et al. Interleukins 27 and 6 induce STAT3-mediated T cell production of interleukin 10. *Nat. Immunol.* 2007; 8:1363–1371. [PubMed: 17994025]
35. McGeachy MJ, et al. TGF- $\beta$  and IL-6 drive the production of IL-17 and IL-10 by T cells and restrain T<sub>H</sub>-17 cell-mediated pathology. *Nat. Immunol.* 2007; 8:1390–1397. [PubMed: 17994024]

36. Terashima A, et al. A novel subset of mouse NKT cells bearing the IL-17 receptor B responds to IL-25 and contributes to airway hyperreactivity. *J. Exp. Med.* 2008; 205:2727–2733. [PubMed: 19015310]
37. Komatsu N, et al. Heterogeneity of natural Foxp3<sup>+</sup> T cells: a committed regulatory T-cell lineage and an uncommitted minor population retaining plasticity. *Proc. Natl. Acad. Sci. USA.* 2009; 106:1903–1908. [PubMed: 19174509]
38. Koyanagi M, et al. EZH2 and histone 3 trimethyl lysine 27 associated with Il4 and Il13 gene silencing in Th1 cells. *J. Biol. Chem.* 2005; 280:31470–31477. [PubMed: 16009709]
39. Tanaka S, et al. The enhancer HS2 critically regulates GATA-3-mediated Il4 transcription in T<sub>H</sub>2 cells. *Nat. Immunol.* 2011; 12:77–85. [PubMed: 21131966]
40. Roers A, et al. T cell-specific inactivation of the interleukin 10 gene in mice results in enhanced T cell responses but normal innate responses to lipopolysaccharide or skin irritation. *J. Exp. Med.* 2004; 200:1289–1297. [PubMed: 15534372]
41. Atarashi K, et al. Induction of colonic regulatory T cells by indigenous *Clostridium* species. *Science.* 2011; 331:337–341. [PubMed: 21205640]
42. Zheng W, Flavell RA. The transcription factor GATA-3 is necessary and sufficient for Th2 cytokine gene expression in CD4 T cells. *Cell.* 1997; 89:587–596. [PubMed: 9160750]
43. Grunig G, et al. Requirement for IL-13 independently of IL-4 in experimental asthma. *Science.* 1998; 282:2261–2263. [PubMed: 9856950]
44. Walter DM, et al. Critical role for IL-13 in the development of allergen-induced airway hyperreactivity. *J. Immunol.* 2001; 167:4668–4675. [PubMed: 11591797]
45. Nishimura Y, Tanaka T. Calcium-dependent activation of nuclear factor regulated by interleukin 3/adenovirus E4 promoter-binding protein gene expression by calcineurin/nuclear factor of activated T cells and calcium/calmodulin-dependent protein kinase signaling. *J. Biol. Chem.* 2001; 276:19921–19928. [PubMed: 11262393]
46. Smirnov DV, Smirnova MG, Korobko VG, Frolova EI. Tandem arrangement of human genes for interleukin-4 and interleukin-13: resemblance in their organization. *Gene.* 1995; 155:277–281. [PubMed: 7721105]
47. Takeda K, et al. Essential role of Stat6 in IL-4 signalling. *Nature.* 1996; 380:627–630. [PubMed: 8602263]
48. Mohrs M, et al. Differences between IL-4<sup>-</sup> and IL-4 receptor  $\alpha$ -deficient mice in chronic leishmaniasis reveal a protective role for IL-13 receptor signaling. *J. Immunol.* 1999; 162:7302–7308. [PubMed: 10358179]
49. Sakai K, Miyazaki J. A transgenic mouse line that retains Cre recombinase activity in mature oocytes irrespective of the cre transgene transmission. *Biochem. Biophys. Res. Commun.* 1997; 237:318–324. [PubMed: 9268708]
50. Yagi R, et al. The IL-4 production capability of different strains of naive CD4<sup>+</sup> T cells controls the direction of the T<sub>H</sub> cell response. *Int. Immunol.* 2002; 14:1–11. [PubMed: 11751746]

**Figure 1.**

IL-13 production induced by repetitive antigen stimulation in  $T_H1$  cells. **(a)** Enzyme-linked immunosorbent assay (ELISA) of IFN- $\gamma$ , IL-4, IL-5 and IL-13 in cells generated from naive  $CD4^+$  T cells (from DO11.10 mice on a recombination-activating gene 2-deficient background) repeatedly stimulated with OVA peptide in the presence of BALB/c APCs every week under  $T_H1$ -skewing conditions, then restimulated with mAb to TCR $\beta$  at 1–4 weeks after initial stimulation. **(b)** ELISA of cytokines in cells generated from naive  $CD4^+$  T cells obtained from *I4r*<sup>+/+</sup> (DO11.10) or *I4r*<sup>-/-</sup> (DO11.10 IL-4R-KO) DO11.10 mice (on the BALB/c background) and stimulated with OVA peptide in the context of APCs (far left and middle left), or generated from naive  $CD4^+$  T cells obtained from wild-type (WT) or *Stat6*<sup>-/-</sup> (STAT6-KO) mice and stimulated by mAb to TCR $\beta$  and mAb to CD28 under  $T_H1$ -skewing conditions (far right and middle right). **(c)** ICS detection of cells producing IFN- $\gamma$ , IL-13 and IL-4 among cells generated from  $CD4^+$  T cells (from DO11.10 mice on a recombination-activating gene 2-deficient or *I4r*<sup>-/-</sup> background) stimulated with OVA peptide in the context of APCs every week under  $T_H1$ -skewing conditions, followed by restimulation with mAb to TCR $\beta$ . Numbers in or below quadrants indicate percent cells in each throughout. Data are from three independent experiments (**a,b**; mean  $\pm$  s.e.m.) or are representative of three experiments with similar results (**c**).



**Figure 2.** E4BP4 regulates IL-13 expression in T<sub>H1</sub> cells. **(a)** Quantitative RT-PCR analysis of the expression of 16 candidate genes identified by comparative transcriptome analysis in T<sub>H1</sub> cells given one round (R1) or four rounds (R4) of simulation and then restimulated with mAb to TCR $\beta$ , and also in 23-1-8 cells, a T<sub>H1</sub> cell clone corresponding to R4 T<sub>H1</sub> cells (primer sequences, Supplementary Table 1). **(b)** ELISA of IL-4 and IL-13 in supernatants of cultures of BALB/c CD4<sup>+</sup> T cells given initial stimulation with mAb to TCR $\beta$  and mAb to CD28, then transduced for 7 d with retroviral plasmid encoding green fluorescent protein (GFP) alone (Vector) or plasmid encoding GFP plus E4BP4, c-Maf, JunB or NR2C1, followed by restimulation of GFP<sup>+</sup> cells for 24 h with mAb to TCR $\beta$ . **(c)** ICS detection (left) of IL-13-producing cells among BALB/c CD4<sup>+</sup> T cells given initial stimulation with mAb to TCR $\beta$  and mAb to CD28, then transduced for 7 d with retroviral plasmid alone or plasmid


encoding E4BP4, followed by restimulation of GFP<sup>+</sup> cells for 6 h with mAb to TCR $\beta$ ; right, frequency of IL-13-producing cells among all CD4<sup>+</sup> T cells. **(d)** Quantitative RT-PCR analysis of the expression of *E4bp4*, *Tbx21* and *Gata3* in R1–R4 T<sub>H</sub>1 cells and in T<sub>H</sub>2 cells (primers, Supplementary Table 1). **(e)** ICS detection (left) of IL-13- and IFN- $\gamma$ -producing cells among R4 T<sub>H</sub>1 cells from DO11.10 mice; immunocytochemistry analysis (middle) of E4BP4 and IL-13 in DO11.10 T cells; and frequency of IL-13<sup>+</sup> cells among E4BP4<sup>+</sup> or E4BP4<sup>-</sup> cells (right). Original magnification (middle),  $\times 640$ . DAPI, DNA-intercalating dye. \* $P < 0.01$  (Student's *t*-test). Data are representative of two experiments with 100 cells expressing both IFN- $\gamma$  and IL-13 **(a)** or are from three independent experiments **(b–e)**; mean and s.e.m.).

\$watermark-text


\$watermark-text

\$watermark-text




**Figure 3.**


*E4bp4* overexpression induces the expression of IL-10 and IL-13 by CD4<sup>+</sup> T cells. **(a)** Intracellular staining to detect transgene expression by CD4<sup>+</sup> T cells purified from the spleens of a wild-type mouse or C57BL/6 mice expressing a transgene (TG) encoding Flag-tagged E4BP4 (three independent lines, TG (1)–TG (3)), assessed with anti-Flag. **(b)** Concentration of cytokines and chemokines in wild-type and *E4bp4*-transgenic CD4<sup>+</sup> T cells stimulated for 7 d with mAb to TCR $\beta$  and mAb to CD28, then restimulated with mAb to TCR $\beta$ . \* $P$  < 0.05 (Student's  $t$ -test). **(c)** Intracellular staining to detect T cells producing IL-4, IL-10 and IL-13 among Flag-positive cells (TG<sup>+</sup>) or Flag-negative cells (TG<sup>-</sup>) generated from *E4bp4*-transgenic CD4<sup>+</sup> T cells stimulated for 7 d as in **b**, then restimulated with mAb to TCR $\beta$ . \* $P$  < 0.05 (Student's  $t$ -test). **(d)** Intracellular staining to detect T cells producing IL-10 and IFN- $\gamma$  (left) among GFP<sup>+</sup> T cells generated from C57BL/6 CD4<sup>+</sup> T cells transduced with plasmid encoding GFP alone or GFP plus E4BP4 and activated under T<sub>H</sub>1 conditions; right, frequency of IL-10<sup>+</sup> cells. **(e)** Production of IL-10 and IL-13 in the GFP<sup>+</sup> T cells isolated in **d**. \* $P$  < 0.01 (Student's  $t$ -test). Data are representative of two experiments **(a)** or are from three independent experiments **(b–d)**; mean and s.e.m.).

**Figure 4.**

E4BP4 regulates the expression of *Il10* and *Il13* in TH1 cells. **(a)** ICS analysis (top) and ELISA (bottom) of IL-13, IL-10 and IFN- $\gamma$  in cells generated from C57BL/6 CD4<sup>+</sup> T cells stimulated four times with mAb to TCR $\beta$  and mAb to CD28 under TH1 conditions. Data are representative of three independent experiments (mean  $\pm$  s.e.m.). **(b)** Immunoblot analysis (top left) of E4BP4 and STAT6 in CD4<sup>+</sup> T cells stimulated for 48 h under TH2 conditions and transduced with retrovirus encoding control shRNA (Ctrl) or *E4bp4*-specific shRNA (E4BP4), assessed 1 week later by probing with anti-E4BP4, followed by densitometry analysis (above; normalized to STAT6 values and presented relative to E4BP4 in cells transduced with the control shRNA, set as 100); and ELISA of IFN- $\gamma$  in R4 TH1 cells (top right) or of IL-10 and IL-13 in TH1 cells stimulated 1–4 weeks (R1–R4; bottom). \* $P < 0.01$  (Student's *t*-test). Data are from three independent experiments (mean  $\pm$  s.e.m.). **(c)** ICS analysis (top) and ELISA (bottom) of IL-13, IL-10 and IFN- $\gamma$  in CD4<sup>+</sup> T cells derived from *E4bp4*<sup>-/-</sup> mice (KO) and their wild-type littermates (WT), then stimulated for 4 weeks under TH1-skewing conditions. \* $P < 0.01$  (Student's *t*-test). Data are representative of three independent experiments (ICS) or are from three independent experiments (mean  $\pm$  s.e.m.).


**Figure 5.**

Impaired production of IL-10 and IL-13 in *E4bp4*-deficient T cells. (a–c) Intracellular staining analysis (a,b) and ELISA (c) of IL-4, IL-10 and IL-13 in CD4<sup>+</sup> T cells obtained from *E4bp4*<sup>-/-</sup> mice (KO) and their *E4bp4*<sup>+/+</sup> littermates (WT) and stimulated under T<sub>H2</sub> conditions. \*P < 0.01 (Student's *t*-test). Data are representative of (a) or from (b,c) three independent experiments (mean and s.e.m. in b,c). (d) Intracellular staining analysis of IL-4, IL-10 and IL-13 in cells generated from naive CD4<sup>+</sup> T cells obtained from *Cd4-Cre* mice or *E4bp4*<sup>fl/fl</sup>*Cd4-Cre* mice (E4BP4-CKO), then stimulated with mAb to TCRβ and mAb to CD28 under T<sub>H2</sub> conditions or T<sub>H1</sub> conditions. Data are representative of three independent experiments. (e) Intracellular staining analysis of IL-10 in cells generated from activated CD4<sup>+</sup> T cells derived from *E4bp4*<sup>-/-</sup> mice (E4BP4-KO) and their *E4bp4*<sup>+/+</sup> littermates (WT) and transduced with empty plasmid or plasmid encoding GATA-3 (top), or derived from *Gata3*<sup>fl/fl</sup> (GATA-3-KO) mice or C57BL/6 mice (WT) and transduced with plasmid encoding GFP and E4BP4 (E4BP4) or plasmid encoding human CD8 (Vector; bottom), followed by restimulation with mAb to TCRβ. GATA-3-deficient T cells (bottom) were isolated by gating on cells positive for human CD8. Data are representative of three independent experiments. (f) ICS analysis (top) of IL-10 and IL-13 in CD4<sup>+</sup> T cells prepared from *Cd4-Cre**Gata3*<sup>fl/fl</sup> mice (GATA-3-CKO) and *Cd4-Cre* mice, then stimulated for 4 weeks under T<sub>H1</sub>-skewing conditions; and frequency of IL-10<sup>+</sup> and IL-13<sup>+</sup> cells (bottom). \*P < 0.01 (Student's *t*-test). Data are representative of three independent experiments (ICS) or are from three independent experiments (mean and s.e.m.).




**Figure 6.**

Impaired production of IL-10 and IL-13 by CD4<sup>+</sup> T cell subsets lacking E4BP4. (a) ICS detection of IL-13- and IL-10-producing cells among whole BALB/c splenocytes stimulated for 48 h with the α-GalCer–CD1d dimer (CD1d dimer). (b) Quantitative RT-PCR analysis of gene expression among total RNA from IL-17RB<sup>+</sup> NKT cells generated from IL-17RB<sup>+</sup>α-GalCer–CD1d<sup>+</sup>TCRβ<sup>+</sup> cells sorted from spleen cells and cultured for 24 h in the presence (IL-25<sup>+</sup>) or absence (IL-25<sup>-</sup>) of IL-25 together with bone marrow–derived DCs induced by granulocyte-macrophage colony-stimulating factor. (c) Flow cytometry analysis of NKT cells from the spleens of *E4bp4*<sup>-/-</sup> mice and their *E4bp4*<sup>+/+</sup> littermates. Numbers adjacent to outlined areas indicate percent α-GalCer–CD1d<sup>+</sup>TCRβ<sup>+</sup> cells. (d) ELISA of cytokine production in IL-17RB<sup>+</sup> NKT cells derived from *E4bp4*<sup>-/-</sup> mice and their *E4bp4*<sup>+/+</sup> littermates (C57BL/6 background) and stimulated for 24 h with various concentrations of IL-25 (top) or α-GalCer (bottom) in the presence of bone marrow–derived DCs induced by granulocyte-macrophage colony-stimulating factor. ND, not detected. (e,f) ELISA of cytokines in culture supernatants of Foxp3<sup>+</sup>CD4<sup>+</sup> (CD25<sup>-</sup> or CD25<sup>hi</sup>) T cells (e) and naive (Foxp3<sup>-</sup>CD62L<sup>hi</sup>CD44<sup>lo</sup>) and memory (Foxp3<sup>-</sup>CD62L<sup>lo</sup>CD44<sup>hi</sup>) CD4<sup>+</sup> T cells (f) purified from reporter mice expressing Foxp3 from human *CD2* (hCD2), on an *E4bp4*<sup>+/+</sup> or *E4bp4*<sup>-/-</sup> background, then stimulated for 48 h with mAb to TCRβ and mAb to CD28. \**P* < 0.01 (Student's *t*-test). Data are representative of two experiments (a) or three experiments (b–f; mean and s.e.m.).



**Figure 7.** E4BP4 protein binds specifically to the *III3* promoter and in the *III0* locus. **(a)** Methylation of genomic DNA (bottom) isolated from naive CD4<sup>+</sup> T cells and from T<sub>H</sub>1 cells and T<sub>H</sub>2 cells given 1 week of stimulation (R1), analyzed by sequencing with primers specific for the *III3* promoter (top) after bisulfite treatment. **(b)** ChIP analysis of the enrichment of histone H3 trimethylated at Lys4 (H3K4me3) in R1 T<sub>H</sub>1 cells, R1 T<sub>H</sub>2 cells or R3 T<sub>H</sub>1 cells (top) and of E4BP4 in R1 T<sub>H</sub>1 and R4 T<sub>H</sub>1 cells (bottom), both at the *III3* locus (middle). CNS1, conserved noncoding sequence 1; HSS, hypersensitive site; *Rad50*, gene encoding a protein involved in the repair of DNA double-strand breaks. **(c)** Electrophoretic mobility-shift assay of nuclear extracts (NE) of T<sub>H</sub>1 and T<sub>H</sub>2 cells, with an oligonucleotide from a specific region of the *III3* promoter as the probe (far left); supershift analysis of nuclear extracts from T<sub>H</sub>2 cells with anti-E4BP4 (Ab; middle left); and competition analysis with 25-, 50- or 100-fold excess competitor (middle right) or mutant oligonucleotides (M1–M5) and the *III3* promoter (pIL-13; far right). **(d)** ChIP analysis of the enrichment of histone H3 acetylated at Lys9 and Lys14 (H3K9ac–H3K14ac; top) or of E4BP4 (bottom) at the *III0* locus (above) in chromatin fractions extracted from T<sub>H</sub>1 or T<sub>H</sub>2 cells; results are presented relative to those obtained with input DNA prepared from untreated chromatin. **(e)** ChIP analysis of the enrichment of histone H3 acetylated at Lys9 and Lys14 at the *III0* locus (above) in T<sub>H</sub>2 cells from *E4bp4*<sup>-/-</sup> and wild-type mice; results presented as in **d**. \**P* < 0.01 (Student’s *t*-test). Data are representative of two experiments (**a–c**) or are from three independent experiment (**d,e**; mean and s.e.m.).





**Figure 8.** Exacerbated colitis and EAE in *E4bp4*<sup>-/-</sup> mice. **(a)** Flow cytometry analysis of the expression of human CD2 (as a marker of Foxp3) and CD4 (left), and ELISA of IL-10 (right) in Foxp3<sup>+</sup> T<sub>reg</sub> cells induced from naive T cells from Foxp3<sup>hCD2</sup> reporter mice on the *E4bp4*<sup>-/-</sup> or wild-type background by culture with transforming growth factor-β. **(b)** Body weight of mice deficient in recombination-activating gene 1 (*n* = 10 per group) given intravenous transfer of naive CD4<sup>+</sup> T cells ( $4 \times 10^5$ ) from the spleens of *E4bp4*<sup>-/-</sup> mice (*n* = 10) or wild-type mice (*n* = 10), presented relative to initial body weight on day 0. † indicates death (two of ten mice died). **(c)** Hematoxylin and eosin staining of colon tissue sections at day 34 in the recipient mice in **b**. Scale bars, 200 μm (left) or 50 μm (right). **(d)** ELISA of IL-10 in CD4<sup>+</sup> T cells isolated at day 34 from the lymph nodes (LN) and colons of the recipient mice in **b**, then stimulated for 48 h with mAb to TCRβ and mAb to CD28. **(e)** EAE clinical scores of *E4bp4*<sup>-/-</sup> mice (*n* = 5) and wild-type mice (*n* = 5) immunized at day 0 with a peptide of myelin oligodendrocyte glycoprotein (amino acids 35–55) in complete Freund’s adjuvant. **(f)** ELISA of cytokines in CD4<sup>+</sup> T cells isolated at day 35 from the lymph nodes of mice immunized as in **e**, then stimulated for 48 h with mAb to TCRβ and mAb to CD28. \**P* < 0.01 (Student’s *t*-test). Data are from three (**a,c,d,f**) or ten (**b**) independent experiments (mean and s.e.m. in **a,b,d-f**).