

Genome Sequences of the Primary Endosymbiont “*Candidatus Portiera aleyrodidarum*” in the Whitefly *Bemisia tabaci* B and Q Biotypes

Zi-Feng Jiang,^a Fangfang Xia,^b Kipp W. Johnson,^a Elizabeth Bartom,^c Jigyasa H. Tuteja,^a Rick Stevens,^{a,b} Robert L. Grossman,^a Marina Brumin,^d Kevin P. White,^a and Murad Ghanim^d

Institute for Genomics and Systems Biology, The University of Chicago, Chicago, Illinois, USA^a; Argonne National Laboratory, Argonne, Illinois, USA^b; Center for Research Informatics, The University of Chicago, Chicago, Illinois, USA^c; and The Agricultural Research Organization (ARO), Department of Entomology, Volcani Center, Bet Dagan, Israel^d

“*Candidatus Portiera aleyrodidarum*” is the obligate primary endosymbiotic bacterium of whiteflies, including the sweet potato whitefly *Bemisia tabaci*, and provides essential nutrients to its host. Here we report two complete genome sequences of this bacterium from the B and Q biotypes of *B. tabaci*.

“*Candidatus Portiera aleyrodidarum*” is the obligate primary endosymbiotic bacterium hosted by whiteflies, including the sweet potato whitefly *Bemisia tabaci*, one of the most globally damaging insect pests in open fields and protected agricultural crops, causing annual losses estimated at 1 to 2 billion dollars. *B. tabaci* is one of the top 100 invasive species worldwide (5). Similar to other obligate bacteria living in sap-sucking insects, “*Ca. Portiera aleyrodidarum*” is thought to provide essential nutrients to whiteflies (2). *B. tabaci* is a species complex comprised of 11 high-level genetic groups that are well defined by DNA markers and at least 24 morphologically indistinguishable species (1,4). The most predominant and damaging biotypes are B and Q, which differ considerably with regard to various fitness parameters: while B is defined by high fecundity and a wide host range, Q is known to develop higher resistance to insecticides (7). Despite these significant differences in their biology, nothing is known about the comparative genome sequences of the B and Q primary endosymbionts. Here we report the assembly and comparison of “*Ca. Portiera aleyrodidarum*” genomes from these two biotypes.

B and Q biotype whiteflies were collected in Israel, and each strain was interbred over 200 generations under standard laboratory conditions. High-quality DNA (>40 kb) was extracted from whole bodies of multiple pooled insect adults of each species. Paired-end libraries with 180-bp insertions and 5-kb mate-pair libraries were constructed and sequenced. Adaptors and low-quality reads were removed before assembly. ALLPATHS-LG (3) and Velvet (8) were used for *de novo* assembly. Based on the depth of coverage, paired-end connections (180 ± 10 bp), and 5-kb mate-pair connections, “*Ca. Portiera aleyrodidarum*” contigs were selected and linked into a closed circular molecule by customized scripts. We observed a limited number of short repeats/homopolymers that appear polymorphic and have taken the major allele in these cases as the consensus sequence. Over 1 million reads were mapped to each of the final assemblies. Assembly accuracy and the coverage (at least 25× coverage for each nucleotide) were manually curated. Annotation was done on the RAST annotation server (6).

The “*Ca. Portiera aleyrodidarum*” genome assembly is a 351-kb circular molecule. The “*Ca. Portiera aleyrodidarum*” genome from whitefly B (here referred to as WB) and the “*Ca. Portiera aleyrodidarum*” genome from whitefly Q (here referred to as WQ) share high

similarity in sequence (99%). Both genomes are highly AT biased (73%) and have 36 RNA-coding genes. A total of 277 protein-coding genes were identified in the “*Ca. Portiera aleyrodidarum*” WB genome, and 281 were identified in the “*Ca. Portiera aleyrodidarum*” WQ genome. Similar to other genome-sequenced obligate endosymbionts from sap-feeding insects, enrichment for genes involved in essential amino acid biosynthesis was observed, while genes involved in processes such as membrane transport, cell wall/capsule, or motility were not found. These results support the hypotheses that primary endosymbionts primarily supply their hosts by amino acids. Comparing “*Ca. Portiera aleyrodidarum*” genome sequences from the B and Q biotypes showed a number of regions that have diverged between the two genomes. Such divergence might play a differential role in “*Ca. Portiera aleyrodidarum*” interactions with their whitefly host.

Nucleotide sequence accession numbers. The complete chromosome sequences have been deposited in the NCBI GenBank database under accession numbers CP003867 (for “*Ca. Portiera aleyrodidarum*” WQ) and CP003868 (for “*Ca. Portiera aleyrodidarum*” WB).

ACKNOWLEDGMENTS

This research was partially supported by grant IS-4062-07 from the United States-Israel Binational Agricultural Research and Development Fund (BARD) and by research grant 887/07 from the Israel Science Foundation to M.G. We also thank the Chicago Center for Systems Biology for a Research Experiences for Undergraduates (REU) fellowship (NIH P50 GM081892) to K.W.J.

REFERENCES

1. Alemandri V, et al. 2012. Species within the *Bemisia tabaci* (Hemiptera: Aleyrodidae) complex in soybean and bean crops in Argentina. *J. Econ. Entomol.* 105:48–53.
2. Baumann P. 2005. Biology bacteriocyte-associated endosymbionts of plant sap-sucking insects. *Annu. Rev. Microbiol.* 59:155–189.
3. Gnerre S, et al. 2011. High-quality draft assemblies of mammalian ge-

Received 25 September 2012 Accepted 28 September 2012

Address correspondence to Murad Ghanim, ghanim@volcani.agri.gov.il, or Kevin P. White, kpwhite@uchicago.edu.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JB.01841-12

- nomes from massively parallel sequence data. *Proc. Natl. Acad. Sci. U. S. A.* **108**:1513–1518.
4. **Hu J, et al.** 2011. An extensive field survey combined with a phylogenetic analysis reveals rapid and widespread invasion of two alien whiteflies in China. *PLoS One* **6**:e16061.
 5. **Kontsedalova S, et al.** 2012. *Bemisia tabaci* biotype dynamics and resistance to insecticides in Israel during the years 2008-2010. *J. Integr. Agric.* **11**:312–320.
 6. **Meyer F, et al.** 2008. The metagenomics RAST server—a public resource for the automatic phylogenetic and functional analysis of metagenomes. *BMC Bioinformatics* **9**:386.
 7. **Skaljic M, et al.** 2012. Diversity and localization of bacterial symbionts in three whitefly species (Hemiptera: Aleyrodidae) from the east coast of the Adriatic Sea. *Bull. Entomol. Res.* **15**:1–12.
 8. **Zerbino DR, Birney E.** 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res.* **18**:821–829.